Check for updates

RESEARCH ARTICLE

Socioeconomic gradients in general and oral health of primary school children in Shiraz, Iran [version 1; peer review: 3 approved]

Ali Golkari^{1,2}, Aira Sabokseir¹, Aubrey Sheiham², Richard G. Watt²

¹Department of Dental Public Health, School of Dentistry, Shiraz University of Medical Sciences, Shiraz, 71345-1836, Iran

²Research Department of Epidemiology and Public Health, University College London, London, WC1E 7HB, UK

V1 First published: 27 Apr 2016, 5:767
<https://doi.org/10.12688/f1000research.8641.1>

Latest published: 27 Apr 2016, 5:767
<https://doi.org/10.12688/f1000research.8641.1>

Abstract

Background: Health status is largely determined by socio-economic status. The general health of individuals at higher social hierarchy is better than people in lower levels. Likewise, people with higher socio-economic status have better oral health than lower socio-economic groups. There has not been much work regarding the influence of socio-economic status on the health conditions of children in developing countries, particularly in Iran. The aim of this study was to compare the oral and general health conditions of primary school children of three different socio-economic areas in the city of Shiraz, Iran.

Methods: This cross-sectional study was conducted on 335, 8- to 11-year-old primary schoolchildren in Shiraz. The children were selected by a three-stage cluster sampling method from three socio-economically different areas. Tools and methods used by the United Kingdom's Medical Research Council were used to obtain anthropometric variables as indicators of general health. The Decay, Missing, Filled Teeth (DMFT) Index for permanent teeth, dmft Index for primary teeth, the Modified Developmental Defects of Enamel (DDE) Index, the Gingival Index (GI) and the Debris Index-Simplified (DI-S) were used for oral health assessment.

Results: Height ($P<0.001$), weight ($P<0.001$), and BMI ($P=0.001$) significantly increased as the socio-economic status of area increased. GI score ($P<0.001$), DI-S score ($P<0.001$), number of permanent teeth with DDE ($P=0.008$), and number of DDE lesions in permanent teeth ($P=0.008$) significantly decreased as the socio-economic status of area increased.

Discussion: Findings of this study generally confirmed that social gradients exist in both general and oral health status of the primary schoolchildren of Shiraz. The influence of socio-economic status on health condition means children have different life chances based on their socio-economic conditions. These findings emphasize the significance of interventions for tackling socio-economic inequalities

Open Peer Review

Approval Status

	1	2	3
version 1			
27 Apr 2016	view	view	view

1. **Tayebeh Malek Mohammadi**, Kerman University of Medical Sciences, Kerman, Iran
2. **Myung Ki**, Eulji University, Daejeon, South Korea
3. **Ali Kazemian**, Mahshad University of Medical Sciences, Mashhad, Iran

Any reports and responses or comments on the article can be found at the end of the article.

in order to improve the health status of children in lower socio-economic areas.

Keywords

Health status disparities , Inequalities , Oral health , Socio-economic factors

This article is included in the **University College London** collection.

Corresponding author: Ali Golkari (golkaria@sums.ac.ir)

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2016 Golkari A *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#) , which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. Data associated with the article are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

How to cite this article: Golkari A, Sabokseir A, Sheiham A and Watt RG. **Socioeconomic gradients in general and oral health of primary school children in Shiraz, Iran [version 1; peer review: 3 approved]** F1000Research 2016, 5:767 <https://doi.org/10.12688/f1000research.8641.1>

First published: 27 Apr 2016, 5:767 <https://doi.org/10.12688/f1000research.8641.1>

Introduction

Health status is largely determined by social class and socio-economic status. Understanding the association between socio-economic status and health outcomes is of great importance for planning health promotion strategies. It is generally accepted that “there is a social gradient in health”¹⁻⁴. In the national trend, the gradient is the status of an individual in socio-economic hierarchy and shows that people at the top of social hierarchy have better health than those in lower levels⁵. In the international trend, the gradient indicates more affluent countries have better health outcomes compared to poorer countries⁶.

The effect of socio-economic status is shown on a wide range of health outcomes from drug misuse related diseases⁷ to age-specific mortality⁸. A study on 6 to 11 year-old children of the United States found that social class had positive association with minor and major physical disorders such as colds, infections, allergies, diabetes, and even epilepsy⁹. The relationship between household income and a range of health outcomes in children and adolescents has been shown in Britain¹⁰. A study in France has found a significant positive correlation between family income and some of the most important child health indicators such as anthropometric measurements¹¹.

In consistence with general health, social gradients are shown in oral health indicators. It seems that despite the remarkable improvements in the averages of oral health indices among communities in the last decades, inequities in oral health - mostly related to social inequalities - continue to exist^{12,13}.

A study on the relationship between work conditions and health inequalities in Switzerland has found clear social gradients for almost all adverse working status and the outcomes of oral health¹⁴. In a case-control study in Mexico, lower socioeconomic status was identified as a risk factor for non-syndromic orofacial clefts¹⁵. Another study on Chinese 5-year-old children found that there were notable gradients in carious primary teeth related to their household income¹⁶. A study on young children of Salem, Tamil Nadu, India, showed that those from lower socio-economic groups, especially whose parents' had lower education were more likely to suffer from Early Childhood Caries¹⁷. It was also shown that poor socio-economic background can be a strong predictor for poor oral hygiene¹⁸.

There has been limited evidence in Iran regarding the association between socio-economic status and general and oral health gradients, particularly among children. Therefore, this study was designed to assess the existence of social gradients in oral and general health of 8- to 11-year-old primary school children in three different socio-economic areas of Shiraz, a city in Southern Iran.

Material and methods

This cross-sectional study was conducted on 8- to 11-year-old children (third to fifth grade primary school children) of the city of Shiraz, Iran, in 2009. Approval and ethical permission were obtained from the National Ethical Committee in Medical Research of Iran (# 85/p/3/1095). Permission to enter the selected schools was then obtained from the Educational Head Office of the Province of Fars (# 17568/55). Based on the prevalence of

Developmental Defects of Enamel (DDE) and a confidence level of 95%, a sample size of 335 was estimated appropriate for this study.

A three-stage cluster sampling approach was used to select the children. The main Educational Affairs Office of Shiraz had a category which divided the city into three areas based on socio-economic status: upper, middle, and lower social class areas. This kind of division seemed more appropriate for this study than any other division, such as geographically separated zones. One boys' and one girls' school (primary schools are segregated in Iran) were selected from each of the three socio-economically different areas. This was done by simple randomization using the list of primary schools in each area. Then, two classes were randomly selected in each school from the third, fourth, and fifth grades. All children present in each selected class were included in the study, only if their parents provided written consent.

Although socio-economic differences between the three areas were well recognized by officials (Educational Head Office 2006), the number of parents having an occupation (any job), and the number of parents having a permanent occupation (as a job security indicator) were used as socio-economic related variables to compare the three areas and confirm the socio-economic difference between children chosen from these areas^{19,20}. Related information was obtained from schools' administration offices.

Height and weight are valuable indicators of present and past health status. Dividing the body height into trunk and leg length provides more precise information particularly on nutrition status of the early years of life^{21,22}. Standing height, leg length, weight, and BMI (Body Mass Index) were used in this study as indicators of general health.

To obtain children's height and weight, the tools and methods used by United Kingdom's Medical Research Council were adopted²². For weight, a digital scale (Beurer Electronic Weight Scale PS 07, Germany) with accuracy of 100 grams was used. It was calibrated in each school. A wall mounted height meter with accuracy of millimetres was used for measuring children's standing height. A similar metre adjusted to the seat of a chair was used to measure the length of the upper trunk, from head to the chair's seat (sitting height). The leg length was calculated by deduction of the sitting height from the standing height. BMI was calculated by dividing the weight by the square of the standing height. Identically calibrated tools were used for all cases in all schools.

Assessment of dental caries, DDE and gingival health determined the oral health status of children. Clinical intra-oral examinations were carried out using WHO screening criteria to record data on Decay, Missing, and Filled Teeth (dmft) for the primary dentition and DMFT for the permanent dentition²³. Permanent teeth were assessed from first molar to first molar in each arch. Therefore, second permanent molars, if present, were not assessed. DDEs of permanent teeth were recorded based on the Modified DDE Index²⁴. Gingival health and oral hygiene were assessed using the Gingival Index (GI)^{25,26} and the Debris Index – Simplified (DI-S)²⁷. All examinations were carried out in classrooms using natural light, disposable mirrors and tongue blades.

Normality of distribution of each outcome variable was assessed by a histogram. All continuous variables had a distribution close to normal. The difference between sexes was analysed by a t-test. Two socio-economic variables of number of parents with any occupation and with a permanent occupation were compared among the three areas using one-way ANOVA. Anthropometric measurements, gingival health, and oral hygiene status were also assessed among the three areas by one-way ANOVA. Anthropometric measurements (linear regression), gingival health status, and oral hygiene status (quantile regression) were also compared among the three areas after adjusting for sex and age. DMFT, dmft, number of teeth needed treatment, number of permanent teeth with DDE and total number of DDEs in children's permanent dentition were compared between areas using Poisson regression model after adjusting for sex and age, and number of permanent teeth present in the mouth. SPSS statistical package (version 14) and STATA statistical package (version 10 Intercooled) were used. The significance level was set at $\alpha=0.05$.

Results

Consent forms were sent for 376 parents, out of which 335 accepted to participate (response rate = 89.1%). Table 1 shows the number of cases examined in each area by sex.

The number of parents with an occupation ($P < 0.001$) and with a permanent occupation ($P < 0.001$) significantly increased from the lower socioeconomic to the higher socioeconomic area (Table 1). These results confirm that the three areas were socio-economically different.

Girls were slightly heavier and taller, but had shorter leg length than boys. All anthropometric variables had distributions close to normal. Children in lower socio-economic neighbourhoods were shorter and lighter, and had shorter legs. Even after adjustments were made for sex and age, still all anthropometric variables: height ($P < 0.001$), weight ($P < 0.001$), BMI ($P = 0.001$) and leg length

($P < 0.001$) of children significantly increased as the socio-economic status of area increased. These findings demonstrated clear social gradient in the general health status among Shiraz primary school children.

The average number ($\pm$ standard deviation) of present deciduous teeth was $6.2 (\pm 4.3)$. Number of permanent teeth present in the mouth ranged from four to 24 with an average of 16 fully erupted permanent teeth per child (Table 2). Number of present permanent teeth was significantly higher in girls (mean = 18.1) than in boys (mean = 13.6). The difference remained statistically significant after adjustment for age ($P < 0.001$).

Forty nine percent (164) of children had a DDE. There was no difference between the percentages of boys and girls having a DDE in permanent teeth. Table 2 shows the number of permanent teeth with DDE and the total number of DDEs in permanent teeth. Both variables increased as socio-economic status of area decreased. The trends were statistically significant for number of permanent teeth with DDE ($P = 0.018$) and the total number of DDEs in permanent teeth ($P = 0.025$) after adjusting for number of present permanent teeth in the mouth. The significance levels increased ($P = 0.008$ for both variables) after adjusting for age and sex in the regression model.

The average dmft (of the primary dentition) was 2.8 ± 2.5 . Three quarters of the cases (75.2%) had a dmft of 1 or more. The average DMFT (of the permanent dentition) was 1.22 ± 1.5 . Almost half of the sample (47.5%) had caries experience in their permanent teeth. DMFT increased from 0.7 in 8-year-olds to 1.4 in 11-year-olds. Although the mean DMFT was higher in girls (1.3) than in boys (1.0) ($P = 0.02$), the difference was not significant after adjustments for age and number of permanent teeth ($P = 0.42$).

There was no statistically significant trend in DMFT or dmft among areas. Children in the middle class area had the highest

Table 1. Sample size and parents' occupation status of children by area and sex (N=335).

Area (socio-economic level)	Sex	Number of examined children	Mean number of parents with any occupation	Odds ratio of having at least one jobless parent	Mean number of parents with a permanent job	Odds ratio of having no parent with a permanent job
Low	Girls	69	1.76	0.08	1.55	< 0.0001
	Boys	43				
Middle	Girls	47	1.37	0.45	1.01	0.12
	Boys	54				
High	Girls	61	1.17	1	0.70	1
	Boys	61				
Significance level of difference among areas			P < 0.001		P < 0.001	

caries prevalence and highest need for treatment among all three areas. Although a clear path of social gradients was not observed, total number of teeth with treatment need was significantly lower in children in high socio-economic area compared with other children ($P = 0.006$). There was also a significant difference in mean number of deciduous teeth with treatment need between the high socio-economic area and the two other areas ($P = 0.005$) (Table 3).

A considerable number of children had gingival inflammation. Average GI score was 0.18 with only 82% of children being

scored 0. The average DI-S was 0.52. The difference between the sexes was not statistically significant ($P = 0.43$ for GI and $P = 0.44$ for DI-S).

The gingival health significantly improved as the socio-economic status of area increased ($P < 0.001$). The significance did not change after adjustments for sex and age. There was also significant trend in oral hygiene status ($P < 0.001$). The number of children with dental plaque ($P < 0.001$), and also the average level of dental plaque in children ($P < 0.001$) decreased as the socio-economic status of area increased (Table 4).

Table 2. Permanent teeth, permanent teeth with DDE, and DDEs per child, by socio-economic areas.

Area	Average number of fully erupted permanent teeth per child	Average number of permanent teeth with DDE per child	Percent of permanent teeth with DDE (%)	Average number of DDEs per child
Low-socioeconomic	15.98	1.72	10.8	1.76
Middle-socioeconomic	15.35	1.19	7.8	1.26
High-socioeconomic	16.78	1.02	6.1	1.07
Average/total	16.01	1.30	8.1	1.36
Significance level (1)	--	$P = 0.018$		$P = 0.025$
Significance level (2)	--	$P = 0.008$		$P = 0.008$

(1) Adjustments made only for number of permanent teeth.

(2) Adjustments made for number of permanent teeth, sex, and age.

Table 3. DMFT, dmft, and number of teeth with treatment need, by socio-economic areas and sex.

Area	Sex	DMFT	Mean number of permanent teeth with treatment need	dmft	Mean number of deciduous teeth with treatment need	Total number of teeth with treatment need
Low-socioeconomic	Boys (N = 43)	0.56	0.33	3.0	2.9	3.2
	Girls (N = 69)	1.14	0.84	2.2	2.1	3.0
	Both (N = 112)	0.92	0.64	2.5	2.4	3.1
Middle-socioeconomic	Boys (N = 54)	1.54	1.13	4.7	4.2	5.3
	Girls (N = 47)	1.62	1.13	2.0	1.7	2.8
	Both (N = 101)	1.58	1.13	3.5	3.0	4.2
High-socioeconomic	Boys (N = 61)	0.97	0.54	3.4	2.5	3.0
	Girls (N = 61)	1.43	1.03	1.6	1.1	2.2
	Both (N = 122)	1.20	0.79	2.5	1.8	2.6
Total (N = 335)		1.22	0.84	2.8	2.4	3.2

Table 4. Gingival health and oral hygiene status among the three socio-economic areas, by sex.

Area	Sex	Number of children with GI=0 (%)	Number of children with GI> 0 (%)	Average DI-S score
Low-socioeconomic	Boys (N = 43)	29 (67.4)	14 (32.6)	0.82
	Girls (N = 69)	53 (76.8)	16 (23.2)	0.70
	Both (N = 112)	82 (73.2)	30 (26.8)	0.75
Middle-socioeconomic	Boys (N = 54)	44 (81.5)	10 (18.5)	0.52
	Girls (N = 47)	36 (76.6)	11 (23.4)	0.50
	Both (N = 101)	80 (79.2)	21 (20.8)	0.51
High-socioeconomic	Boys (N = 61)	53 (86.9)	8 (13.1)	0.44
	Girls (N = 61)	60 (98.4)	1 (1.6)	0.19
	Both (N = 122)	113 (92.6)	9 (7.4)	0.32
Total (N = 335)		275 (82.1)	60 (17.9)	0.52
Significance level of difference among areas		P< 0.001		P< 0.001

Dataset 1. Raw data

<http://dx.doi.org/10.5256/f1000research.8641.d120612>

Data on age, sex, socioeconomic, anthropometric and oral health variables are provided.

Discussion

Findings of this study generally confirmed that social gradients exist in both general and oral health status of the primary school children of Shiraz. Children in lower socio-economic areas were shorter and lighter, and had shorter legs. In terms of oral health, socio-economic status had more effects on gingival inflammation, levels of plaque, and specially occurrence of DDEs, but was less related to dental caries experience and caries treatment need.

Most findings of this study were consistent with those of similar studies in other parts of the world. As an example, the social gradients seen in the anthropometric measurements were similar to those presented among French children¹¹. One of the strong relationships found in this study was between the area of living and the state of oral hygiene determined by DI-S. This finding was in accordance with the findings of a recent study by Mathur *et al.* in India¹⁸. They had also divided a city, Delhi, into three socio-economically different areas and found that children in poorer areas were more susceptible for poorer oral hygiene.

In the current study, social gradient could not been shown in DMFT or dmft of studied children. Further research revealed that similar conditions had been reported by others. As an example, Sagheri *et al.* (2009) reported a social gradient in using preventive dental services among children of Ireland, but no such trend in DMFT of the same children²⁸. To justify such findings, one could blame the nature and shortcomings of the DMF Index or complexity of the risk factors for dental caries.

DDE is probably the most important oral health indicator assessed in this study. Presence of DDE on teeth can be an index of both general and oral health. Many early childhood adverse health conditions and diseases can increase the risk of developing a DDE on permanent dentition²⁹. Despite the importance of DDE as a health indicator, few studies have tried to assess the relationship of the prevalence of DDE, especially on permanent teeth, with socio-economic factors. One of such studies has been conducted by Basha *et al.* that reports a significant negative association between the presence of DDE and socio-economic status of children³⁰, a finding that is very similar to what is shown in the current study.

The influence of socio-economic status on health condition means children have different life chances based on their birthplace, or area of living, even inside one city. If they live in a higher socio-economic zone, they would have a higher chance for better oral and general health in comparison to their peers in neighboring lower

socio-economic zones. These findings could help policymakers to make intervention decisions that might help to improve the health status of children in lower socio-economic areas.

Bearing in mind that no clear social class determinant or assessment tool exists in Iran, the authors decided to use the school area as the best available indicator of subjects' social class. Assessing the job status of parents acted as a proof that the selected groups of children were really from different backgrounds. Therefore, it seems fair to say that this study has been able to illustrate that oral health follows the other aspects of health in having gradients according to socio-economic status.

As oral health and general health follow the social gradients, tackling health inequalities would need identifying and understanding the underlying causes of problems³¹. Upstream measures would be needed to build a society that could reinforce good oral and general health. Future assessment of the relationship between social gradients in oral health and general health is recommended.

Conclusion

There are social gradients in general and oral health among primary school children of Shiraz, Iran. The gradient in oral health seems to follow the same pattern in general health.

Consent

Written informed consent for participation in the study was obtained from the parents of the children.

Data availability

F1000Research: Dataset 1. Raw data, [10.5256/f1000research.8641.d120612](https://doi.org/10.5256/f1000research.8641.d120612)³².

Author contributions

The study was conceived and designed by AG, Ash, and RGW. AG and ASa carried out the research. AG analyzed the data. All authors were involved in preparation of the first draft. AG and ASa finalized the manuscript. The data for this study were extracted from a PhD thesis by AG under supervision of RGW and ASH. ASa was involved in data collection and preparation of the manuscript.

Competing interests

No competing interests were disclosed.

Grant information

The author(s) declared that no grants were involved in supporting this work.

References

- Macintyre S: **Understanding the social patterning of health: the role of the social sciences.** *J Public Health Med.* 1994; **16**(1): 53–9.
[PubMed Abstract](#)
- Marmot MG, Smith GD, Stansfeld S, *et al.*: **Health inequalities among British civil servants: the Whitehall II study.** *Lancet.* 1991; **337**(8754): 1387–93.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Wilkinson RG: **Unhealthy societies: the afflictions of inequality.** London: Routledge; 1996.
[Reference Source](#)
- Wilkinson RG, Marmot M: **Social determinants of health.** 2nd ed. Oxford: Oxford University Press, 2006.
[Reference Source](#)
- Matthews KA, Gallo LC: **Psychological perspectives on pathways linking socioeconomic status and physical health.** *Annu Rev Psychol.* 2011; **62**: 501–30.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Dasgupta P, Weale M: **On measuring the quality of life.** *World development.* 1992; **20**(1): 119–31.
[Publisher Full Text](#)
- Origer A, Le Bihan E, Baumann M: **A social gradient in fatal opioids and cocaine related overdoses?** *PLoS One.* 2015; **10**(5): e0125568.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Belon AP, Barros MB, Marín-León L: **Mortality among adults: gender and socioeconomic differences in a Brazilian city.** *BMC Public Health.* 2012; **12**: 39.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Starfield B, Robertson J, Riley AW: **Social class gradients and health in childhood.** *Ambul Pediatr.* 2002; **2**(4): 238–46.
[PubMed Abstract](#)
- Emerson E, Graham H, Hatton C: **Household income and health status in children and adolescents in Britain.** *Eur J Public Health.* 2006; **16**(4): 354–60.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Apouey BH, Geoffroy PY: **Child health and access to health care in France: Evidence on the role of family income.** *Rev Epidemiol Sante Publique.* 2014; **62**(3): 179–90.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Locker D: **Deprivation and oral health: a review.** *Community Dent Oral Epidemiol.* 2000; **28**(3): 161–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kumar S, Kroon J, Lalloo R: **A systematic review of the impact of parental socio-economic status and home environment characteristics on children's oral health related quality of life.** *Health Qual Life Outcomes.* 2014; **12**: 41.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Hämmig O, Bauer GF: **The social gradient in work and health: a cross-sectional study exploring the relationship between working conditions and health inequalities.** *BMC Public Health.* 2013; **13**: 1170.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Acuña-González G, Medina-Solís CE, Maupomé G, *et al.*: **Family history and socioeconomic risk factors for non-syndromic cleft lip and palate: a matched case-control study in a less developed country.** *Biomedica.* 2011; **31**(3): 381–91.
[PubMed Abstract](#)
- Guan Y, Zeng X, Tai B, *et al.*: **Socioeconomic inequalities in dental caries among 5-year-olds in four Chinese provinces.** *Community Dent Health.* 2015; **32**(3): 185–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Stephen A, Krishnan R, Ramesh M, *et al.*: **Prevalence of early childhood caries and its risk factors in 18–72 month old children in Salem, Tamil Nadu.** *J Int Soc Prev Community Dent.* 2015; **5**(2): 95–102.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Mathur MR, Tsakos G, Parmar P, *et al.*: **Socioeconomic inequalities and determinants of oral hygiene status among Urban Indian adolescents.** *Community Dent Oral Epidemiol.* 2016; **44**(3): 248–54.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Siegrist J, Theorell T: **Socioeconomic position and health: the role of work and employment.** In: Siegrist J, Marmot M, editors. *Social inequalities in health: new evidence and policy implication.* Oxford: Oxford University Press; 2006; 73–100.
- Wilkinson R, Marmot M: **Social determinants of health: the solid facts.** 2nd ed. Copenhagen: World Health Organization, Regional Office for Europe; 2003.
[Reference Source](#)
- Langenberg C, Hardy R, Kuh D, *et al.*: **Influence of height, leg and trunk length on pulse pressure, systolic and diastolic blood pressure.** *J Hypertens.* 2003; **21**(3): 537–43.
[PubMed Abstract](#)

22. Wadsworth ME, Hardy RJ, Paul AA, *et al.*: **Leg and trunk length at 43 years in relation to childhood health, diet and family circumstances; evidence from the 1946 national birth cohort.** *Int J Epidemiol.* 2002; **31**(2): 383–90.
[PubMed Abstract](#) | [Publisher Full Text](#)
23. World Health Organization: **Oral health surveys: basic methods.** 4th Edition, Geneva: World Health Organization Publication; 1997.
[Reference Source](#)
24. Clarkson J, O'Mullane D: **A modified DDE Index for use in epidemiological studies of enamel defects.** *J Dent Res.* 1989; **68**(3): 445–50.
[PubMed Abstract](#) | [Publisher Full Text](#)
25. Loe H: **The Gingival Index, the Plaque Index and the Retention Index Systems.** *J Periodontol.* 1967; **38**(6 Suppl): 610–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
26. Loe H, Silness J: **Periodontal disease in pregnancy. I. Prevalence and severity.** *Acta Odontol Scand.* 1963; **21**(6): 533–51.
[PubMed Abstract](#) | [Publisher Full Text](#)
27. Greene JG, Vermillion JR: **The simplified oral hygiene index.** *J Am Dent Assoc.* 1964; **68**(1): 7–13.
[PubMed Abstract](#) | [Publisher Full Text](#)
28. Sagheri D, McLoughlin J, Clarkson JJ: **The prevalence of dental caries and fissure sealants in 12 year old children by disadvantaged status in Dublin (Ireland).** *Community Dent Health.* 2009; **26**(1): 32–7.
[PubMed Abstract](#)
29. Memarpour M, Golkari A, Ahmadian R: **Association of characteristics of delivery and medical conditions during the first month of life with developmental defects of enamel.** *BMC Oral Health.* 2014; **14**: 122.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
30. Basha S, Mohamed RN, Swamy HS: **Prevalence and associated factors to developmental defects of enamel in primary and permanent dentition.** *Oral Health Dent Manag.* 2014; **13**(3): 588–94.
[PubMed Abstract](#)
31. Watt RG: **From victim blaming to upstream action: tackling the social determinants of oral health inequalities.** *Community Dent Oral Epidemiol.* 2007; **35**(1): 1–11.
[PubMed Abstract](#) | [Publisher Full Text](#)
32. Golkari A, Sabokseir A, Sheiham A, *et al.*: **Dataset 1 in: Socioeconomic gradients in general and oral health of primary school children in Shiraz, Iran.** *F1000Research.* 2016.
[Data Source](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 25 May 2016

<https://doi.org/10.5256/f1000research.9297.r13623>

© 2016 Kazemian A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Ali Kazemian

Department of Community Oral Health, Mashhad School of Dentistry, Mahshad University of Medical Sciences, Mashhad, Iran

I found this article worth indexing. Well organized, well written.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 24 May 2016

<https://doi.org/10.5256/f1000research.9297.r13930>

© 2016 Ki M. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Myung Ki

Department of Preventive Medicine, School of Medicine, Eulji University, Daejeon, South Korea

The theme of this paper is, to my mind, very interesting (socioeconomic inequalities in general and dental health) and various general and dental measures with trustable definition are plus. The design and data analysis seem acceptable to support the conclusion. Thus, I believe that this paper merits indexation, but there is a few points to be addressed / discussed more.

1. For anthropometric variables, authors describe: "Even after adjustments were made for sex and age, still all anthropometric variables: height ($P < 0.001$), weight ($P < 0.001$), BMI ($P = 0.001$) and leg length ($P < 0.001$) of children significantly increased as the socioeconomic

status of area increased.”

However, no actual results on this appear in the table and please check in which table the result is shown. If this finding is less important and authors decided not to present the data, please indicate that “data is not presented”.

2. It is not clear why two p values are presented in the table 1 and I think it would be four p-values per each column. Similarly, only two p-values were presented in the table 2, instead of four as to four measures. I am sorry, if misunderstanding is mine.
3. For presentation of table 1, I wonder whether the use of Odds Ratio is proper approach. I think instead of OR, presenting different proportions for three area groups and statistical test with Chi-square would be relevant and complementary to other results in table 1, for example, Chi-square test for “having at least one jobless parent” for three area levels.
4. Use of area level SEP could be more justifiable, if the parents are actual residents of the areas to which their siblings attend, if additional information of this kind is available.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 16 May 2016

<https://doi.org/10.5256/f1000research.9297.r13621>

© 2016 Malek Mohammadi T. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Tayebeh Malek Mohammadi

Oral and Dental Diseases Research Centre, Department of Dental Public Health, Kerman Dental School, Kerman University of Medical Sciences, Kerman, Iran

The manuscript reported very important subject in oral health' field, seems sound and very well written. It is acceptable for indexation. I suggest the authors should address some studies regarding oral health and general health and also oral health and socio-economic factors which have been reported in Iran.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research