REVIEW

Current techniques for visualizing RNA in cells [version 1; peer review: 2 approved]

Lilith V.J.C. Mannack^{1,2}, Sebastian Eising¹, Andrea Rentmeister^{1,2}

¹Department of Chemistry, Institute of Biochemistry, University of Münster, Münster, Germany

²Cells in Motion, Cluster of Excellence, Münster, Germany

V1 First published: 28 Apr 2016, 5(F1000 Faculty Rev):775
<https://doi.org/10.12688/f1000research.8151.1>

Latest published: 28 Apr 2016, 5(F1000 Faculty Rev):775
<https://doi.org/10.12688/f1000research.8151.1>

Abstract

Labeling RNA is of utmost interest, particularly in living cells, and thus RNA imaging is an emerging field. There are numerous methods relying on different concepts ranging from hybridization-based probes, over RNA-binding proteins to chemo-enzymatic modification of RNA. These methods have different benefits and limitations. This review aims to outline the current state-of-the-art techniques and point out their benefits and limitations.

Keywords

RNA, visualization, imaging, proteins

Open Peer Review

Approval Status

	1	2
version 1 28 Apr 2016

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Seergazhi G. Srivatsan**, Indian Institute of Science Education and Research, Pune, India
2. **Jiangyun Wang**, Institute of Biophysics, Chinese Academy of Sciences, Chaoyang District, Beijing, China

Any comments on the article can be found at the end of the article.

Corresponding author: Andrea Rentmeister (a.rentmeister@uni-muenster.de)

Competing interests: The authors declare that they have no competing interests.

Grant information: Our work is supported by the Deutsche Forschungsgemeinschaft (RE 2796/2-1 and EXC 1003 Cells in Motion – Cluster of Excellence, Münster, Germany). Andrea Rentmeister gratefully acknowledges the Fonds der Chemischen Industrie for a “Dozentenstipendium”.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2016 Mannack LVJC *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Mannack LVJC, Eising S and Rentmeister A. **Current techniques for visualizing RNA in cells [version 1; peer review: 2 approved]** F1000Research 2016, 5(F1000 Faculty Rev):775 <https://doi.org/10.12688/f1000research.8151.1>

First published: 28 Apr 2016, 5(F1000 Faculty Rev):775 <https://doi.org/10.12688/f1000research.8151.1>

Introduction

The localization of mRNA in the cell has been a topic of interest since the 1980s, when protein localization was linked to localized mRNA translation¹. At that time, the only method by which RNA could be visualized was *in situ* hybridization (ISH)². Since then, the options for RNA detection have expanded greatly. Labeling RNA—particularly mRNA—is of utmost interest, as mRNA localization has been shown to be important in a range of situations. For example, in a developing *Drosophila* oocyte, asymmetrically localized mRNA produces a *bicoid* protein gradient through localized translation, which specifies the anterior-posterior polarity of the developing larva³. In neurons, localization of mRNA is also particularly important, and localized mRNA leads to multiple rounds of translation at the synapse and activity-dependent changes⁴. Additionally, since the number of genes transcribed was found to surpass the amount of protein-coding genes, interest in non-coding RNAs has increased⁵. Thus, imaging microRNAs or small interfering RNAs (siRNAs) is also of interest⁶. Furthermore, defects in mRNA localization play an important role in some diseases such as fragile X syndrome⁷, and non-coding RNAs have been shown to be important in diseases such as cancers⁸.

Hybridization methods

ISH is used to visualize RNA containing a known sequence. A DNA or RNA strand of complementary sequence hybridizes to the RNA strand of interest via Watson-Crick base pairing. The probe bears features that enable its visualization (e.g., a fluorophore; Figure 1A). Fluorescence *in situ* hybridization (FISH) can distinguish between RNA molecules that differ in only a single base⁹. FISH is highly sequence-specific, and individual RNA strands may be detected when combined with various amplification procedures in fixed cells^{9,10}. A variety of derivatives of ISH that reduce

background signal have been developed, most notably molecular beacons (Figure 1B). Molecular beacons consist of a DNA probe that is linked to a fluorophore at one end and a quencher at the opposite end. When unbound, the probe folds into a hairpin structure, bringing the fluorophore and quencher together, thereby inhibiting fluorescence. Upon target recognition, the probe anneals and stretches out, separating the quencher from the fluorophore and enabling fluorescence¹¹. Molecular beacons have been advanced further since the 1990s. For example, the types of quencher used have been expanded to include nanoparticles¹². Microinjected molecular beacons can mislocalize to the nucleus in live cells; however, incorporating a tRNA sequence was shown to abrogate this problem¹³.

An alternative approach to increase the specific fluorescent signal upon binding is to use forced intercalation (FIT) probes (Figure 1C). FIT probes are peptide nucleic acid (PNA) or DNA single strands containing a base surrogate (typically, thiazole orange) that intercalates between the Watson-Crick base pairs and fluoresces only upon exact hybridization^{14,15}. Their strong turn-on effect (~30-fold) makes FIT probes an attractive improvement of ISH. FIT probes have been expanded to contain dyes that emit in the blue and green ranges^{16,17} and have been successfully used for mRNA visualization in cells and *Drosophila* embryos^{14,18}.

Hybridization-based RNA detection is an excellent tool for use in fixed samples and can be used in living cells and organisms when strong turn-on effects are achieved (e.g., molecular beacons and FIT probes). However, probes based on modified nucleic acids or derivatives thereof are neither cell-permeable nor can they be produced by the cell itself. Furthermore, hybridization has to occur in regions of the target RNA free of secondary structure, and

Figure 1. Hybridization-based methods for RNA imaging. (A) Standard fluorescence *in situ* hybridization (FISH): a fluorophore-linked RNA probe binds the target RNA sequence. (B) Molecular beacon: signal to noise is improved relative to a standard FISH probe because the fluorescence signal of the reporter probe is quenched when unbound. (C) Forced intercalation (FIT) probes: binding enforces intercalation of the dye molecule into the probe-target duplex, resulting in a strong turn-on effect of the fluorophore.

hybridization conditions are typically not optimized for the cellular milieu. Recently, probes and conditions have started to be developed for use in live cells; this approach is termed fluorescence *in vivo* hybridization (FIVH)^{19–21}. In particular, 2'-*O*-methylated oligonucleotides exhibit faster hybridization kinetics, increased melting temperatures, enhanced binding specificity, improved nuclease stability, and the ability to bind structured molecules—properties beneficial for FIVH probes²⁰. They were used to detect a variety of RNA types, such as snRNAs, rRNA, and poly(A) RNA²⁰. Nevertheless, these probes need to be introduced into the cells and thus FIVH requires transient permeabilization of cells.

Aptamers

RNA aptamers are another form of nucleotide-based probe but work on a different principle to the above mentioned hybridization probes. Aptamers are short, single-stranded oligonucleotides capable of binding specific target molecules based on their shape and can be obtained by *in vitro* selection²². Recently, an aptamer termed “Spinach” was selected that folds to allow binding of a small-molecule fluorophore that fluoresces only upon binding the RNA aptamer (Figure 2C)²³. Herein, the reporter and probe are contained within the same oligonucleotide. The aptamer sequence can be appended to the RNA of interest to enable visualization of that RNA upon binding of the fluorophore²³. RNA aptamers in conjunction with said small fluorophore are available in a range of colors from blue to red²³ and have been improved to further enhance binding efficiency and fluorescence strength²⁴. Additionally, the folding

properties have been optimized for the cellular milieu^{25,26}. A downside of RNA aptamers is the potential impediment in localization or function of some RNAs by the “Spinach” RNA tag.

Particle-associated hybridization-based imaging probes

The group of Mirkin describe a nanoparticle conjugated spherical nucleic acid that recognizes specific RNA targets and is capable of entering the cell without the need for transfection^{27,28}. However, there is considerable controversy surrounding this study, mainly concerning whether these sticky- or nano-flares mark specific RNAs or merely remain in endosomes after uptake by the cell²⁹. Gold nanoparticles bound to quantum dots via hybridizing DNA strands have been developed to detect specific microRNA³⁰. The microRNA triggers the dissociation of the quantum dots from the gold particle, resulting in the abrogation of quenching and thus a signal. These gold nanoparticle-quantum dot-probes bind target RNA quantitatively *in vitro*, and cell lines expressing a certain microRNA can be distinguished from cell lines that do not.

Covalent modification of RNA in cells

An alternative method to mark RNA is to incorporate visualizable moieties directly into the RNA. A convenient way to achieve marking RNA without introducing a large moiety that may interfere with RNA function is to incorporate a small chemical group that may be further reacted by using click chemistry to attach to a fluorophore. There are a number of different click reactions, the most prominent being the copper(I)-catalyzed azide alkyne cycloaddition (CuAAC). Here, an azide reacts with an alkyne in the presence of Cu(I) as a catalyst. CuAAC is rapid and extremely selective; however, Cu(I) at millimolar concentrations is toxic to cells and thus this approach is limited to fixed cell samples. Jao and Salic succeeded in incorporating ethynyl groups into total RNA by feeding cells with the uridine analog 5-ethynyluridine (EU), which is converted to the respective triphosphate inside the cell (Figure 3A)³¹. Using a similar approach—feeding cells with *N*⁶-propargyl adenosine—the poly(A) dynamics of mRNA could be monitored³².

There are also a number of copper-free click reactions, which are more suitable for live-cell imaging, termed bioorthogonal click reactions (reviewed in 33). Sawant *et al.* synthesized an azido-modified UTP analog that can be used in the bioorthogonal strain-promoted azide-alkyne cycloaddition (SPAAC)³⁴. This allowed the click reaction to proceed in live cells; however, this UTP analog had to be transfected into the cells as its uridine precursor was no longer cell-permeable or was not a good substrate for the ribonucleoside salvage pathway (Figure 3A).

A downside of incorporating modified nucleotides during transcription or poly(A) tail addition is that different subtypes of RNA cannot be distinguished. A possible method by which to obtain specific labeling of different subtypes is to attach chemical groups used for click reactions post-synthetically by using RNA-modifying enzymes. Subtypes of RNA may also be labeled by taking advantage of certain structures or modifications in an RNA type. For example, the 5' cap of mRNA may be specifically labeled by using an engineered methyltransferase that is only active on the mRNA cap (Figure 3B)^{35–37}. This approach should be suitable in live cells

Figure 2. Visualization based on reporter molecules binding to a specific RNA sequence. (A) A green fluorescent protein-fused-MS2 coat protein (GFP-MCP) binds a consensus sequence (MS2) appended to the RNA of interest. (B) Two pumilio variants fused to different halves of split-GFP recognize a target sequence within an RNA molecule of interest. (C) The aptamer “Spinach” folds to bind a turn-on fluorophore and can be appended to an RNA of interest.

Figure 3. Introducing covalent modifications into RNA and subsequent labeling by click chemistry allow visualization. (A) Incorporation of modified nucleotides into nascent RNA by endogenous RNA polymerases. Ethynyluridine is cell-permeable, and the respective triphosphate is made inside the cell; hence, feeding the cell with the nucleoside precursor is possible. In other examples (azido-U), the cells have to be transfected with the respective triphosphates. (B) Hallmarks of RNA subtypes, such as the 5' cap, can be selectively modified. A methyltransferase (MTase) variant can be used to modify the mRNA cap with the clickable group if the respective *S*-adenosylmethionine (AdoMet) analog is provided. (C) Transcript-specific installation of a click reactive moiety can be achieved by appending a tRNA-mimicking sequence to the RNA of interest. The enzyme tRNA^{lle2}-agmatidine synthetase (Tias) modifies the tag with a clickable group if appropriate agmatine analogs are provided.

because the *S*-adenosylmethionine (AdoMet) analog can be made from cell-permeable and stable methionine analogs by a variant of the methionine adenosyltransferase (MAT), which is responsible for AdoMet synthesis³⁷.

Sequence-specific RNA-modification with a propargyl group and subsequent labeling with a fluorophore have been achieved *in vitro* by using a box C/D methyltransferase-guide RNA complex and the respective propargyl-bearing analog of the cosubstrate AdoMet³⁸. Li *et al.* developed an RNA labeling system in which an RNA of interest was extended by a tRNA-derived sequence and an enzyme that specifically modifies this sequence (tRNA^{lle2}-agmatidine synthetase, or Tias) was introduced into a cell (Figure 3C)³⁹. This RNA-Tias combination can also accept agmatine analogs that are click-reactive and thus can be used to label RNA in cells³⁹. Similarly, a tRNA-derived recognition motif may be specifically marked by using an engineered transglycosylase that is able to transfer large visualizable groups⁴⁰.

RNA-binding proteins

A number of bacteriophage-derived RNA-binding proteins have been used to mark RNA in cells. The most notable of these is the MS2-MS2 coat protein (MS2-MCP) system (Figure 2A). This comprises a green fluorescent protein (GFP)-fused version of the

bacteriophage MCP (an RNA-binding protein that recognizes a specific RNA sequence-determined hairpin) and the RNA of interest extended by multiple MS2-binding sites (MBS)⁴¹. Recently, the MS2 system has been used to image single mRNA molecules in living mouse cells⁴². This study displays both the power and drawback of this method. On the one hand, the MS2 system allows tracking and resolution of single mRNA molecules; on the other, producing a transgenic organism is very time-consuming. Another drawback is that the size of the MS2-fusion tag and the appendages to the RNA might interfere with normal mRNA function or localization^{43,44}. Furthermore, the quantity of MS2 used must be sufficient to saturate the target RNA without raising background fluorescence, which may be difficult to achieve⁴⁵.

Another RNA-binding protein worth mentioning is pumilio. Pumilio is a member of the RNA-binding protein family PUF⁴⁶. Like many RNA-binding proteins, pumilio is modularly composed of domains that can be engineered to alter the specific RNA sequence bound^{47–49}. Pumilio is of particular interest as it can target RNA directly without the need to introduce an RNA tag into the target RNA.

An advantage of using RNA-binding proteins to visualize RNA is that two individual RNA sequences may be targeted by separate RNA-binding proteins, thus allowing the imaging of the

association of two RNA molecules of interest^{50,51}. The potential drawback of high background fluorescence due to unbound protein may be countered by using a split GFP, which fluoresces only upon dimerization (Figure 2B)^{52,53}.

Reporter protein expression by trans-splicing to visualize RNA

Two approaches have been developed by which a pre-mRNA may be spliced into a functional form, which allows the expression of a reporter protein. This enables tissue-specific localization of an mRNA of interest, although the resolution at a subcellular level is lost. Bhaumik *et al.* described a method based on trans-splicing that results in the expression of luciferase in cells of a living organism microinjected with an exogenous RNA that was processed to pre-mRNA⁵⁴. So *et al.*, employing a similar approach, developed an engineered ribozyme, which fuses a reporter gene to a specific gene of interest⁵⁵. The authors were able to detect p53 in a whole organism and on a cellular level. Despite theoretical expansion potential⁵⁶, the approach taken by the Gambhir lab has not been significantly developed since the publication of the original study, leaving it with the limitation that only exogenous RNA can be visualized. Similarly, the work of So *et al.* has not been further developed.

Conclusions: current applications and outlook

Imaging of RNA is of interest at the level of both single cells and the whole organism. Labeling RNA in a single cell can show the localization of a specific transcript, which may have important biological consequences⁵⁷. RNA imaging at the whole organism level is important to determine the tissue expression pattern of a specific transcript. RNA labeling has seen extensive use in imaging of infection by RNA viruses (e.g., 58). Another interesting

application of RNA imaging has been to monitor transcription and this has been used, for example, to determine the toxicity of certain substances that inhibit transcription⁵⁹.

In summary, RNA may be visualized by a variety of methods. RNA may be seen via hybridization of a reporter molecule, most commonly through FISH or variations thereof. Alternatively, RNA-binding proteins that bind specific sequences may mark an RNA molecule of interest, or an RNA aptamer that fluoresces upon binding of a fluorophore may be incorporated into the target molecule. RNA may be sequence- or subtype-specifically labeled by using click chemistry. Challenges facing the field of RNA imaging are the cell permeability of dyes used and the low abundance of target RNA. Furthermore, no method of RNA labeling is yet able to yield quantitative data on its target RNA. However, with continued development, RNA imaging will continue to provide important biological insights.

Competing interests

The authors declare that they have no competing interests.

Grant information

Our work is supported by the Deutsche Forschungsgemeinschaft (RE 2796/2-1 and EXC 1003 Cells in Motion – Cluster of Excellence, Münster, Germany). Andrea Rentmeister gratefully acknowledges the Fonds der Chemischen Industrie for a “Dozentenstipendium”.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- Lehmann R, Nüsslein-Volhard C: **Abdominal segmentation, pole cell formation, and embryonic polarity require the localized activity of oskar, a maternal gene in *Drosophila*.** *Cell*. 1986; 47(1): 141–52.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Weil TT, Parton RM, Davis I: **Making the message clear: visualizing mRNA localization.** *Trends Cell Biol*. 2010; 20(7): 380–90.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Driever W, Nüsslein-Volhard C: **The bicoid protein determines position in the *Drosophila* embryo in a concentration-dependent manner.** *Cell*. 1988; 54(1): 95–104.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kindler S, Wang H, Richter D, *et al.*: **RNA transport and local control of translation.** *Annu Rev Cell Dev Biol*. 2005; 21: 223–45.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kellis M, Wold B, Snyder MP, *et al.*: **Defining functional DNA elements in the human genome.** *Proc Natl Acad Sci U S A*. 2014; 111(17): 6131–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Moore A, Medarova Z: **Imaging of siRNA Delivery and Silencing.** In: Sioud M, editor. *siRNA and miRNA Gene Silencing*. Totowa, NJ: Humana Press, 2009; 487: 1–18.
[Publisher Full Text](#)
- Bassell GJ, Kelic S: **Binding proteins for mRNA localization and local translation, and their dysfunction in genetic neurological disease.** *Curr Opin Neurobiol*. 2004; 14(5): 574–81.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Jansson MD, Lund AH: **MicroRNA and cancer.** *Mol Oncol*. 2012; 6(6): 590–610.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Larsson C, Grundberg I, Söderberg O, *et al.*: **In situ detection and genotyping of individual mRNA molecules.** *Nat Methods*. 2010; 7(5): 395–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Paré A, Lemons D, Kosman D, *et al.*: **Visualization of individual Scr mRNAs during *Drosophila* embryogenesis yields evidence for transcriptional bursting.** *Curr Biol*. 2009; 19(23): 2037–42.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Tyagi S, Kramer FR: **Molecular beacons: probes that fluoresce upon hybridization.** *Nat Biotechnol*. 1996; 14(3): 303–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Zheng J, Yang R, Shi M, *et al.*: **Rationally designed molecular beacons for bioanalytical and biomedical applications.** *Chem Soc Rev*. 2015; 44(10): 3036–55.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Mhlanga MM, Vargas DY, Fung CW, *et al.*: **tRNA-linked molecular beacons for imaging mRNAs in the cytoplasm of living cells.** *Nucleic Acids Res*. 2005; 33(6): 1902–12.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Hövelmann F, Gaspar I, Ephrussi A, *et al.*: **Brightness enhanced DNA FIT-probes for wash-free RNA imaging in tissue.** *J Am Chem Soc*. 2013; 135(50): 19025–32.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Köhler O, Jarikote DV, Seitz O: **Forced intercalation probes (FIT Probes): thiazole orange as a fluorescent base in peptide nucleic acids for homogeneous single-nucleotide-polymorphism detection.** *Chembiochem*. 2005; 6(1): 69–77.
[PubMed Abstract](#) | [Publisher Full Text](#)

16. Kovaliov M, Segal M, Kafri P, *et al.*: **Detection of cyclin D1 mRNA by hybridization sensitive NIC-oligonucleotide probe.** *Bioorg Med Chem.* 2014; 22(9): 2613–21.
[PubMed Abstract](#) | [Publisher Full Text](#)
17. Hövelmann F, Gaspar I, Chamiolo J, *et al.*: **LNA-enhanced DNA FIT-probes for multicolour RNA imaging.** *Chem Sci.* 2016; 7: 128–35.
[Publisher Full Text](#) | [F1000 Recommendation](#)
18. Kummer S, Knoll A, Socher E, *et al.*: **Fluorescence imaging of influenza H1N1 mRNA in living infected cells using single-chromophore FIT-PNA.** *Angew Chem Int Ed Engl.* 2011; 50(8): 1931–4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
19. Fontenete S, Guimarães N, Leite M, *et al.*: **Hybridization-based detection of *Helicobacter pylori* at human body temperature using advanced locked nucleic acid (LNA) probes.** *PLoS One.* 2013; 8(11): e81230.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
20. Dirks RW, Molenaar C, Tanke HJ: **Visualizing RNA molecules inside the nucleus of living cells.** *Methods.* 2003; 29(1): 51–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
21. Paillisson S, Van De Corput M, Dirks RW, *et al.*: **In situ hybridization in living cells: detection of RNA molecules.** *Exp Cell Res.* 1997; 231(1): 226–33.
[PubMed Abstract](#) | [Publisher Full Text](#)
22. Famulok M, Hartig JS, Mayer G: **Functional aptamers and aptazymes in biotechnology, diagnostics, and therapy.** *Chem Rev.* 2007; 107(9): 3715–43.
[PubMed Abstract](#) | [Publisher Full Text](#)
23. Paige JS, Wu KY, Jaffrey SR: **RNA mimics of green fluorescent protein.** *Science.* 2011; 333(6042): 642–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
24. Dolgoshina EV, Jeng SC, Panchapakesan SS, *et al.*: **RNA mango aptamer-fluorophore: a bright, high-affinity complex for RNA labeling and tracking.** *ACS Chem Biol.* 2014; 9(10): 2412–20.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
25. Filonov GS, Moon JD, Svensen N, *et al.*: **Broccoli: rapid selection of an RNA mimic of green fluorescent protein by fluorescence-based selection and directed evolution.** *J Am Chem Soc.* 2014; 136(46): 16299–308.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
26. Strack RL, Disney MD, Jaffrey SR: **A superfolding Spinach2 reveals the dynamic nature of trinucleotide repeat-containing RNA.** *Nat Methods.* 2013; 10(12): 1219–24.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
27. Briley WE, Bondy MH, Randeria PS, *et al.*: **Quantification and real-time tracking of RNA in live cells using Sticky-flares.** *Proc Natl Acad Sci U S A.* 2015; 112(31): 9591–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
28. Seferos DS, Giljohann DA, Hill HD, *et al.*: **Nano-flares: probes for transfection and mRNA detection in living cells.** *J Am Chem Soc.* 2007; 129(50): 15477–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
29. Mason D, Levy R: **Sticky-flares: real-time tracking of mRNAs... or of endosomes?** *BioRxiv.* 2015.
[Publisher Full Text](#)
30. He X, Zeng T, Li Z, *et al.*: **Catalytic Molecular Imaging of MicroRNA in Living Cells by DNA-Programmed Nanoparticle Disassembly.** *Angew Chem Int Ed Engl.* 2016; 55(9): 3073–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
31. Jao CY, Salic A: **Exploring RNA transcription and turnover in vivo by using click chemistry.** *Proc Natl Acad Sci U S A.* 2008; 105(41): 15779–84.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
32. Grammel M, Hang H, Conrad NK: **Chemical reporters for monitoring RNA synthesis and poly(A) tail dynamics.** *ChemBiochem.* 2012; 13(8): 1112–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
33. McKay CS, Finn MG: **Click chemistry in complex mixtures: bioorthogonal bioconjugation.** *Chem Biol.* 2014; 21(9): 1075–101.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
34. Sawant AA, Tanpure AA, Mukherjee PP, *et al.*: **A versatile toolbox for posttranscriptional chemical labeling and imaging of RNA.** *Nucleic Acids Res.* 2016; 44(2): e16.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
35. Holstein JM, Stummer D, Rentmeister A: **Engineering *Giardia lamblia* trimethylguanosine synthase (GlaTgs2) to transfer non-natural modifications to the RNA 5'-cap.** *Protein Eng Des Sel.* 2015; 28(6): 179–86.
[PubMed Abstract](#) | [Publisher Full Text](#)
36. Schulz D, Holstein JM, Rentmeister A: **A chemo-enzymatic approach for site-specific modification of the RNA cap.** *Angew Chem Int Ed Engl.* 2013; 52(30): 7874–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
37. Muttach F, Rentmeister A: **A Biocatalytic Cascade for Versatile One-Pot Modification of mRNA Starting from Methionine Analogues.** *Angew Chem Int Ed Engl.* 2016; 55(5): 1917–20.
[PubMed Abstract](#) | [Publisher Full Text](#)
38. Tomkuvienė M, Clouet-d'Orval B, Cerniauskas I, *et al.*: **Programmable sequence-specific click-labeling of RNA using archaeal box C/D RNP methyltransferases.** *Nucleic Acids Res.* 2012; 40(14): 6765–73.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
39. Li F, Dong J, Hu X, *et al.*: **A covalent approach for site-specific RNA labeling in Mammalian cells.** *Angew Chem Int Ed Engl.* 2015; 54(15): 4597–602.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
40. Alexander SC, Busby KN, Cole CM, *et al.*: **Site-Specific Covalent Labeling of RNA by Enzymatic Transglycosylation.** *J Am Chem Soc.* 2015; 137(40): 12756–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
41. Bertrand E, Chartrand P, Schaefer M, *et al.*: **Localization of *ASH1* mRNA particles in living yeast.** *Mol Cell.* 1998; 2(4): 437–45.
[PubMed Abstract](#) | [Publisher Full Text](#)
42. Park HY, Lim H, Yoon YJ, *et al.*: **Visualization of dynamics of single endogenous mRNA labeled in live mouse.** *Science.* 2014; 343(6169): 422–4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
43. Buxbaum AR, Haimovich G, Singer RH: **In the right place at the right time: visualizing and understanding mRNA localization.** *Nat Rev Mol Cell Biol.* 2015; 16(2): 95–109.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
44. Garcia JF, Parker R: **MS2 coat proteins bound to yeast mRNAs block 5' to 3' degradation and trap mRNA decay products: implications for the localization of mRNAs by MS2-MCP system.** *RNA.* 2015; 21(8): 1393–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
45. Wu B, Chao JA, Singer RH: **Fluorescence fluctuation spectroscopy enables quantitative imaging of single mRNAs in living cells.** *Biophys J.* 2012; 102(12): 2936–44.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
46. Spassov DS, Jurecic R: **The PUF family of RNA-binding proteins: does evolutionarily conserved structure equal conserved function?** *IUBMB Life.* 2003; 55(7): 359–66.
[PubMed Abstract](#) | [Publisher Full Text](#)
47. Lunde BM, Moore C, Varani G: **RNA-binding proteins: modular design for efficient function.** *Nat Rev Mol Cell Biol.* 2007; 8(6): 479–90.
[PubMed Abstract](#) | [Publisher Full Text](#)
48. Cheong CG, Hall TM: **Engineering RNA sequence specificity of Pumilio repeats.** *Proc Natl Acad Sci U S A.* 2006; 103(37): 13635–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
49. Kellermann SJ, Rath AK, Rentmeister A: **Tetramolecular fluorescence complementation for detection of specific RNAs in vitro.** *Chembiochem.* 2013; 14(2): 200–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
50. Chen J, Nikolaitchik O, Singh J, *et al.*: **High efficiency of HIV-1 genomic RNA packaging and heterozygote formation revealed by single virion analysis.** *Proc Natl Acad Sci U S A.* 2009; 106(32): 13535–40.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
51. Dilley KA, Ni N, Nikolaitchik OA, *et al.*: **Determining the frequency and mechanisms of HIV-1 and HIV-2 RNA copackaging by single-virion analysis.** *J Virol.* 2011; 85(20): 10499–508.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
52. Wu B, Chen J, Singer RH: **Background free imaging of single mRNAs in live cells using split fluorescent proteins.** *Sci Rep.* 2014; 4: 3615.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
53. Magliery TJ, Wilson CG, Pan W, *et al.*: **Detecting protein-protein interactions with a green fluorescent protein fragment reassembly trap: scope and mechanism.** *J Am Chem Soc.* 2005; 127(1): 146–57.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
54. Bhaumik S, Walls Z, Puttaraju M, *et al.*: **Molecular imaging of gene expression in living subjects by spliceosome-mediated RNA trans-splicing.** *Proc Natl Acad Sci U S A.* 2004; 101(23): 8693–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
55. So MK, Gowrishankar G, Hasegawa S, *et al.*: **Imaging target mRNA and siRNA-mediated gene silencing in vivo with ribozyme-based reporters.** *Chembiochem.* 2008; 9(16): 2682–91.
[PubMed Abstract](#) | [Publisher Full Text](#)
56. Walls ZF, Puttaraju M, Temple GF, *et al.*: **A generalizable strategy for imaging pre-mRNA levels in living subjects using spliceosome-mediated RNA trans-splicing.** *J Nucl Med.* 2008; 49(7): 1146–54.
[PubMed Abstract](#) | [Publisher Full Text](#)
57. Chen KH, Boettiger AN, Moffitt JR, *et al.*: **RNA imaging. Spatially resolved, highly multiplexed RNA profiling in single cells.** *Science.* 2015; 348(6233): aaa6090.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
58. Neef AB, Pernot L, Schreier VN, *et al.*: **A Bioorthogonal Chemical Reporter of Viral Infection.** *Angew Chem Int Ed Engl.* 2015; 54(27): 7911–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
59. Kametani Y, Iwai S, Kuraoka I: **An RNA synthesis inhibition assay for detecting toxic substances using click chemistry.** *J Toxicol Sci.* 2014; 39(2): 293–9.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. **Jiangyun Wang**

Institute of Biophysics, Chinese Academy of Sciences, Chaoyang District, Beijing, China

Competing Interests: No competing interests were disclosed.

2. **Seergazhi G. Srivatsan**

Indian Institute of Science Education and Research, Pune, India

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research