REVIEW

Central control of body temperature [version 1; peer review: 3 approved]

Shaun F. Morrison

Department of Neurological Surgery, Oregon Health & Science University, Portland, OR, USA

V1 First published: 12 May 2016, 5(F1000 Faculty Rev):880
<https://doi.org/10.12688/f1000research.7958.1>

Latest published: 12 May 2016, 5(F1000 Faculty Rev):880
<https://doi.org/10.12688/f1000research.7958.1>

Abstract

Central neural circuits orchestrate the behavioral and autonomic repertoire that maintains body temperature during environmental temperature challenges and alters body temperature during the inflammatory response and behavioral states and in response to declining energy homeostasis. This review summarizes the central nervous system circuit mechanisms controlling the principal thermoeffectors for body temperature regulation: cutaneous vasoconstriction regulating heat loss and shivering and brown adipose tissue for thermogenesis. The activation of these thermoeffectors is regulated by parallel but distinct efferent pathways within the central nervous system that share a common peripheral thermal sensory input. The model for the neural circuit mechanism underlying central thermoregulatory control provides a useful platform for further understanding of the functional organization of central thermoregulation, for elucidating the hypothalamic circuitry and neurotransmitters involved in body temperature regulation, and for the discovery of novel therapeutic approaches to modulating body temperature and energy homeostasis.

Keywords

Brown adipose tissue, shiver, cutaneous vasoconstriction, thermogenesis, fever, sympathetic nerve activity, preoptic hypothalamus, rostral raphe pallidus, dorsomedial hypothalamus, therapeutic hypothermia, obesity

Open Peer Review

Approval Status

	1	2	3
version 1 12 May 2016			

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Robin M McAllen**, University of Melbourne, Parkville, Australia
University of Melbourne, Parkville, Australia
Michael J McKinley, University of Melbourne, Parkville, Australia
University of Melbourne, Parkville, Australia
2. **Matteo Cerri**, University of Bologna, Bologna, Italy
3. **Andrej A Romanovsky**, St. Joseph's Hospital and Medical Center, Phoenix, USA

Any comments on the article can be found at the end of the article.

Corresponding author: Shaun F. Morrison (morrisos@ohsu.edu)

Competing interests: The author declares that he has no competing interests.

Grant information: This work was supported by National Institutes of Health grants R01NS040987 and R01NS091066 to Shaun F. Morrison.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2016 Morrison SF. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Morrison SF. **Central control of body temperature [version 1; peer review: 3 approved]** F1000Research 2016, 5(F1000 Faculty Rev):880 <https://doi.org/10.12688/f1000research.7958.1>

First published: 12 May 2016, 5(F1000 Faculty Rev):880 <https://doi.org/10.12688/f1000research.7958.1>

Body temperature (T_{core}) is a critical homeostatic parameter influencing cellular function and organismal survival. Life-threatening protein denaturation looms as T_{core} increases, and reductions in membrane fluidity, ion fluxes, and enzyme performance accompany significant reductions in T_{core} . The fundamental central neural circuits for thermoregulation orchestrate behavioral and autonomic repertoires that maintain T_{core} during thermal challenges, sensed by thermal receptors, that arise from either the ambient or the internal (e.g., during exercise) environment. A variety of other neural circuits and neurochemical modulators impinge on the fundamental thermoregulatory pathways to produce the alterations in T_{core} that occur with circadian and ultradian periodicities^{1,2} and those that accompany many behavioral states, such as fever during sickness, increased T_{core} during psychological stress and at ovulation, and reductions in T_{core} during sleep, sepsis, or exposure to metabolic distress (e.g., hypoxia and starvation). Recent research in the central nervous system (CNS) control of T_{core} focuses principally on: (a) elaborating the CNS pathways and neurotransmitter systems involved in the fundamental central thermoregulatory network; (b) the modulation of activity within the fundamental central thermoregulatory network by non-thermal factors and behavioral states; and (c) pharmacological manipulation of the CNS thermoregulatory network for a variety of therapeutic goals (e.g., reducing T_{core} to produce therapeutic hypothermia in stroke victims).

In its simplest form, the fundamental thermoregulatory network can be modeled as a reflex^{3,4} in which a central integrative circuit alters the activity of thermoeffector mechanisms in response to an input from the combination of peripheral (i.e., skin) and central (i.e., visceral and brain) thermoreceptors that provides a consolidated assessment of T_{core} and, importantly, of imminent threats to T_{core} . The primary thermoeffector mechanisms recruited for both cold defense and centrally driven hyperthermias (e.g., fever) include: (a) thermoregulatory behaviors to reduce the loss of heat produced during basal metabolism; (b) cutaneous vasoconstriction (CVC) to conserve heat in the body core and limit heat loss to the environment; and (c) heat production (thermogenesis). The principal sources of metabolic heat production, beyond those contributing to basal metabolic rate (e.g., pumping ions across membranes), are brown adipose tissue (BAT), whose sympathetic neural input fuels mitochondria that shunt proton fluxes into heat production, and shivering behavior in skeletal muscle, dependent on the inefficiency of ATP utilization to generate heat. Effector mechanisms for heat defense include: (a) thermoregulatory behavior to increase heat loss; (b) cutaneous vasodilation, which in humans includes a sympathetic vasodilator outflow^{5,6}, combined with increased cardiac output⁷ and visceral vasoconstriction^{8–10} to facilitate core heat loss from the body surface; and (c) evaporative cooling (e.g., sweating).

Apart from important “first responders”, the CNS pathways controlling and mediating thermoregulatory behaviors remain incompletely defined^{11–15}. Along these same lines, most research subjects are furry mammals that are strongly dependent on non-sweating mechanisms for evaporative cooling. Although this has limited the available detail on the CNS pathways regulating human sweating^{6,16,17}, considerable insight has been gained on the CNS regulation of other mechanisms of evaporative cooling, such as

panting^{18–20}. Additionally, since most of the basic neuroscience of CNS thermoregulatory pathways has been derived from experiments in rodents, including the exclusively *in vitro* studies of neurons with intrinsic thermosensitivity, the translation of the conclusions, including the circuit model in Figure 1, to humans must be cautiously undertaken (e.g., 21). Finally, although considerable progress has been achieved in the relatively young field of the neuroscience of thermoregulation, the synthesis (Figure 1) of our understanding of this multifaceted neural network controlling multiple thermoeffectors represents a working model, with the expectation of revisions and added detail.

The CNS thermoregulatory control of the sympathetic outflows mediating CVC and BAT thermogenesis and of the somatic motoneurons producing shivering is effected through parallel but distinct, effector-specific, integrative/efferent circuits (Figure 1, and reviewed in 22–25) that share common peripheral thermal sensory inputs. The hypothalamus contains the primary integrative and rostral efferent components of these circuits. Although many details of the preoptic area (POA) microcircuitry for thermoregulation remain to be elucidated, neurons in the POA are postulated to integrate ascending peripheral thermosensory signals with local thermosensitivity to regulate the output of BAT and shivering thermogenesis-promoting neurons in the dorsomedial hypothalamus (DMH)^{26,27} and of CVC-promoting neurons in the median preoptic nucleus (MnPO)^{28,29}.

The POA regulation of DMH thermogenesis-promoting neurons represents the balance between a GABAergic inhibition^{30,31} and a glutamatergic excitation³²; the latter inputs, potentially arising from neurons in the MnPO that project to the DMH, are synaptically connected to BAT³³ and express the leptin receptor³⁴. These glutamatergic inputs to DMH³² could provide the excitation required to drive the BAT sympathoexcitatory neurons and the shivering-promoting neurons in DMH when their POA inhibitory input is reduced during skin cooling or fever³⁵. Although intrinsically warm-sensitive (W-S) (Figure 1), POA neurons^{36–38}, generally located in the medial preoptic area (MPA)³⁹, are postulated to play a key role in central thermoregulation by providing a prominent core temperature-modulated, GABAergic³⁸ regulation of thermogenesis-promoting neurons in DMH (Figure 1), the considerable direct functional evidence required to establish this attractive hypothesis has yet to be obtained. Different thermal sensitivities or neurochemical modulation among populations of temperature-sensitive POA neurons may underlie the differential responsiveness of different effectors to changes in cutaneous versus brain temperatures⁴⁰ as well as the significant alterations in thermoeffector activation during different sleep phases⁴¹. Through their responses to immune signaling molecules, neurons in the POA are also the primary site for the organization and maintenance of the febrile response to inflammation and infection, which includes the stimulation of CVC, and shivering (“chills”) and BAT thermogenesis mediated by the action of prostaglandin E_2 (PGE_2) on its EP3 receptors^{42–45}. Similarly, the fundamental thermoregulatory network mediates stress-induced hyperthermia^{46–48}. Unraveling the complexity of the thermoregulatory circuitry in the hypothalamus^{20,28,29,49–52}, including the phenotypic characterization of the projection neurons³⁴ and their synaptic interactions that mediate the circadian¹³ and many behavioral^{53,54}

Figure 1. Functional neuroanatomical model for the fundamental pathways providing the thermoregulatory control and pyrogenic activation of cutaneous vasoconstriction (CVC) and brown adipose tissue (BAT) and shivering thermogenesis. Cool and warm cutaneous thermoreceptors transmit signals to respective primary sensory neurons in the dorsal root ganglia (DRG) which relay this information to second-order thermal sensory neurons in the dorsal horn (DH). Cool sensory DH neurons glutamatergically activate third-order sensory neurons in the external lateral subnucleus of the lateral parabrachial nucleus (LPB), while warm sensory DH neurons project to third-order sensory neurons in the dorsal subnucleus of the LPB. Thermosensory signals driving thermoregulatory responses are transmitted from the LPB to the preoptic area (POA), which contains the microcircuitry through which cutaneous and core thermal signals are integrated to regulate the balance of POA outputs that are excitatory (dashed green) and inhibitory (dashed red) to thermogenesis-promoting neurons in the dorsomedial hypothalamus (DMH) and to CVC sympathetic premotor neurons in the rostral raphe pallidus (rRPa). Within the POA, GABAergic interneurons (red) in the median preoptic (MnPO) subnucleus are postulated to receive a glutamatergic input from skin cooling-activated neurons in LPB and inhibit each of the distinct populations of warm-sensitive (W-S) neurons in the medial preoptic area (MPA) that control CVC, BAT, and shivering. In contrast, glutamatergic interneurons (dark green) in the MnPO are postulated to be excited by glutamatergic inputs from skin warming-activated neurons in LPB and, in turn, excite the populations of W-S neurons in MPA. Prostaglandin E₂ (PGE₂) binds to EP3 receptors, which are postulated to inhibit the activity of each of the classes of W-S neurons in the POA. Preoptic W-S neurons may provide inhibitory control of CVC by inhibiting CVC sympathetic premotor neurons in the rostral ventromedial medulla, including the rRPa, that project to CVC sympathetic preganglionic neurons (SPNs) in the intermediolateral nucleus (IML). Preoptic W-S neurons may provide inhibitory thermoregulatory control of BAT and shivering thermogenesis by inhibiting BAT sympathoexcitatory neurons and shivering-promoting neurons, respectively, in the DMH, which, when disinhibited during skin and core cooling, provide respective excitatory drives to BAT sympathetic premotor neurons and to skeletal muscle shivering premotor neurons in the rRPa. These, in turn, project, respectively, to BAT SPNs in the IML and to alpha (α) and gamma (γ) motoneurons in the ventral horn (VH) of the spinal cord.

modulations in T_{core} , continues to pose significant research challenges for understanding the “heart” of the CNS thermoregulatory network. The downstream targets of the hypothalamic projection neurons for thermoregulation are the sympathetic and somatic premotor neurons in the rostral ventromedial medulla, centered on the rostral raphe pallidus (rRPa). Midbrain (e.g., periaqueductal gray^{55–57} and retrorubral field⁵⁸) and medullary (e.g., ventrolateral medulla and nucleus of the solitary tract (NTS)^{59,60}) pathways to these premotor neurons exert important modulatory influences on the thermoregulatory activation of thermoeffector organs (reviewed in 22,24). These premotor neurons, in turn, excite CVC and BAT sympathetic preganglionic neurons and α -motoneurons and γ -motoneurons^{61,62} in the spinal cord.

Cold and warm thermoreceptors in the skin and viscera provide the extracranial thermal signals relating to skin temperature and T_{core} . These are integrated with the brain temperature information potentially derived from the discharge of W-S, GABAergic preoptic neurons of the central thermoregulatory network to regulate thermoeffector activities. The membranes of thermoreceptor afferent neurons contain transient receptor potential (TRP) cation channels whose temperature-dependent conductances transduce skin temperature into primary thermoreceptor afferent neuronal activity. The TRPM8 channel, activated by menthol and cooling, is the strongest candidate for the cutaneous cold receptor (reviewed in 63). The identity of the peripheral warm receptor remains to be established, but the warm-sensing mechanism of preoptic neurons, though still debated^{64–66}, is unlikely to involve a transient receptor potential (TRP) channel⁶⁷. Primary thermoreceptor dorsal root ganglion neurons synapse on thermoreceptive-specific, lamina I spinal (or trigeminal) dorsal horn cells⁶⁸, and these, in turn, collateralize to innervate the thalamus, providing the neural substrate for cutaneous thermosensory perception and localization^{68,69}, and the pontine lateral parabrachial nucleus (LPB)^{70,71}, which is responsible for triggering involuntary (e.g., autonomic and shivering) thermoregulatory responses. Spinal lamina I skin cooling-responsive neurons provide a glutamatergic excitation to neurons in the external lateral subdivision of the lateral parabrachial nucleus (LPBel), which, in turn, project principally to the MnPO of the POA^{72,73}, while a parallel, spinoparabrachial glutamatergic pathway excites POA-projecting neurons in the dorsal subnucleus of the LPB (LPBd)^{72,74} in response to skin warming⁷⁴. Thus, activations of POA-projecting LPBd and LPBel neurons, driven respectively by cutaneous, and likely visceral, warm and cold thermoreceptor stimuli, initiate heat defense and cold defense inhibitions and excitations, respectively, in CVC sympathetic outflow and cutaneous blood flow, in BAT sympathetic outflow and BAT thermogenesis, and in shivering EMGs and shivering thermogenesis to maintain a homeostatic T_{core} .

Synaptic integration sites throughout the core thermoregulatory network provide the substrate for a wide variety of non-thermal physiological parameters, disease processes, neuromodulators, and drugs to influence the central regulation of T_{core} (reviewed in 25). For example, the high metabolic rate of BAT and shivering skeletal muscles during thermogenesis cannot be sustained without a dependable supply of metabolic fuels, particularly oxygen⁷⁵, lipolytic by-products³⁴, and glucose⁷⁶. Thus, the CNS networks driving

cold-defensive and behavioral BAT activation or shivering are strongly inhibited by signals reflecting a reduction in the short- and long-term availability of these fuel molecules essential for BAT and skeletal muscle metabolism. Similarly, as hypovolemia progresses during dehydration, the ensuing hyperosmolarity reduces the thermoregulatory drive for sweating^{20,77,78}, which serves to prevent cardiovascular collapse⁷⁹. Some viscerosensory afferents with axons in the vagus nerve and synapsing on second-order neurons in the NTS can also influence BAT activity^{25,80} and shivering responses and thus are expected to influence the regulation of T_{core} . For instance, vagal afferents convey the “metabolic” signals that produce the inhibition of BAT activity induced by upregulation of hepatic glucokinase⁸¹ and the BAT activation following either intragastric delivery of the TRP agonist, capsiate⁸², or the presence of lipids in the duodenum⁸³. Another influence on T_{core} includes a prominent hypothermia during motion sickness and nausea⁸⁴, for which the pathways relating vestibular stimulation to inhibition of CVC and thermogenesis remain to be elaborated.

Providing an additional layer of complexity in the CNS regulation of T_{core} is the presence of the thermally responsive TRPV1 channel in the membranes of a variety of classically non-thermal, unmyelinated afferents^{85–88} that may have access to central thermoregulatory circuits, including via NTS neurons that can inhibit thermogenesis⁵⁹. The finding that the TRPV1 responsiveness to local brain temperature alters spontaneous glutamate release from the terminals of unmyelinated vagal afferents^{88,89} provides a potential substrate for T_{core} to modulate the activity of second-order sensory neurons in NTS that modulate thermoeffector activation. Importantly, TRPV1 channels are also activated by non-thermal factors, including low (or high) pH, inorganic cations, or endovanilloids, thereby providing a basis for such factors to influence thermoeffector activation and thus T_{core} . For instance, TRPV1 channels, potentially on the terminals of afferents in the peritoneum, are stimulated by an endogenous ligand to tonically inhibit BAT thermogenesis, which results in a hyperthermic response to TRPV1 antagonist administration^{85,90}. Although these afferents were not directly tested for their thermal responsiveness, the hyperthermic response to TRPV1 antagonism was not altered by changes in T_{core} . The relative influences and the interactions of thermal and non-thermal stimuli on the conductance of the relevant TRPV1 channels could play a role in determining their effect on the level of thermoeffector activation to thermoreceptor stimuli.

Interest in pharmacological modulation of the central thermoregulatory network has focused: (a) on reducing CVC and thermogenesis to lower T_{core} ; and (b) on augmenting thermogenesis to elevate energy expenditure with the goal of weight loss through consumption of the high-energy lipid stores in white adipose tissue. Novel approaches to reducing T_{core} would have immediate benefits in treating intractable fevers that are unresponsive to cyclooxygenase inhibitors. Therapeutic hypothermia can have beneficial effects on survival and on reducing brain and tissue damage in ischemic insults such as cardiac arrest, stroke, and neonatal encephalopathy^{91–94}. Extended space travel (e.g., Mars One) also may require pharmacological induction of a hypothermic, hibernation-like state to reduce energy consumption and psychological stress.

Under basal metabolic and movement conditions, changes in T_{core} must arise from changes in the level of activation of thermoeffector tissues. Although modulating neuronal discharge at any site within the central thermoregulatory network would be expected to alter T_{core} , centrally generated hyperthermias, such as fever, that arise from altered activity in hypothalamic thermoregulatory neurons could be most effectively reduced by manipulating thermoeffector efferent pathways^{95,96}. Indeed, directly inhibiting the discharge of neurons in the rRPa area, including the functionally significant premotor neurons that control the principal thermoeffectors, produced a fall in brain temperature of approximately 14°C in an ambient environment of 15°C⁹⁷, and stimulation of $\alpha 2$ adrenergic receptors in the rRPa could completely block or prevent a lipopolysaccharide-evoked fever⁹⁵. Similarly, central administration of an adenosine A1 receptor (A1AR) agonist reduced rat T_{core} by 10°C in an ambient temperature of 15°C⁶⁰ and mouse T_{core} by 5°C in an ambient temperature of 4°C⁹⁸. Of particular interest, in rats, this hypothermia, which was produced by a blockade of the cold-evoked activation of BAT and shivering thermogenesis, was long-lasting and paralleled by marked reductions in heart and respiratory rate, in EEG, and in behavior⁶⁰—all reminiscent of those occurring in hibernation and torpor, which require central stimulation of the A1AR^{99–101}. The discovery of a role for TRP channels in thermal sensation and in thermoeffector activation has stimulated research into the pharmacological modulation of TRP channels in the central thermoregulatory network to abrogate the normal cold defense mechanisms and allow T_{core} to fall in a cool ambient environment. By reducing the activation of cooling-responsive skin thermoreceptors, a TRPM8 antagonist reduced rat T_{core} by approximately 1°C when the rats were in an ambient temperature of 19°C⁶³. Stimulating TRPV1 reduced mouse T_{core} by approximately 12°C when mice were exposed to a 10°C ambient temperature¹⁰², and this effect was potentiated by the addition of a TRPM8 antagonist¹⁰³. The location of the relevant, hypothermic TRPV1 channels remains unknown.

Not only is BAT a thermogenic thermoeffector, including in adult humans^{104–107}, but through its consumption of lipid and glucose energy stores and oxygen, thermogenic metabolism in BAT is a neurally regulated contributor to energy homeostasis. Thus, particularly in the face of an elevated consumption of energy-rich food (e.g., a high-fat diet), a chronic reduction in cooling-evoked BAT thermogenesis would contribute to the augmented adipose energy stores that characterize obesity. Indeed, mice without BAT exhibit a propensity for obesity and diabetes^{108,109}; conversely, overexpression

of uncoupling protein-1 (UCP-1), principally responsible for thermogenesis in BAT, mitigates obesity induced by a high-fat diet¹¹⁰. Several anti-obesity therapies currently being explored are based on increased activation of BAT thermogenesis, through either activation of the central thermoregulatory network to increase the sympathetic outflow to BAT¹¹¹ or an alteration in the cellular biochemical pathways in brown adipocytes or a hyperplasia of BAT to augment thermogenesis. The consistent findings that obese humans have significantly reduced cooling-activated BAT^{104–106,112}, and that the basal (i.e., principally cooling-evoked) sympathetic activation of BAT is reduced in rats fed a high-fat diet^{113,114}, and that a vagal afferent input to the NTS mediates the reduced cooling-evoked BAT activity in rats fed a high-fat diet¹¹⁵ not only support a role for reduced BAT activity in the excess adipose accumulation of obesity but also highlight the significance of non-thermal inputs to the central thermoregulatory network^{25,81,83} in influencing even the most basic thermoregulatory responses.

Considerable progress has been achieved in revealing the functional organization of the dedicated thermoregulatory network within the CNS that provides the fundamental neural control of the thermoregulatory effectors: thermoregulatory behavior, CVC, and BAT and shivering thermogenesis, although many of the details of the neurophysiology and neuroanatomy of the central thermoregulatory network remain active areas of investigation. The changes in T_{core} that accompany a wide range of behaviors and in response to many hormones and drugs arise through altered non-thermal inputs to, or neurochemical modulation of, the neural activity within the fundamental thermoregulatory network. The latter, as well as thermoreceptor-based strategies, are being researched as therapeutic approaches in which the central thermoregulatory networks are recruited to alter T_{core} and metabolism.

Competing interests

The author declares that he has no competing interests.

Grant information

This work was supported by National Institutes of Health grants R01NS040987 and R01NS091066 to Shaun F. Morrison.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- Ootsuka Y, de Menezes RC, Zaretsky DV, *et al.*: **Brown adipose tissue thermogenesis heats brain and body as part of the brain-coordinated ultradian basic rest-activity cycle.** *Neuroscience*. 2009; **164**(2): 849–61.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Vujovic N, Gooley JJ, Zhou TC, *et al.*: **Projections from the subparaventricular zone define four channels of output from the circadian timing system.** *J Comp Neurol*. 2015; **523**(18): 2714–37.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Romanovsky AA: **Do fever and anaprexia exist? Analysis of set point-based definitions.** *Am J Physiol Regul Integr Comp Physiol*. 2004; **287**(4): R992–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Werner J: **System properties, feedback control and effector coordination of human temperature regulation.** *Eur J Appl Physiol*. 2010; **109**(1): 13–25.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Wallin BG, Charkoudian N: **Sympathetic neural control of integrated cardiovascular function: insights from measurement of human sympathetic nerve activity.** *Muscle Nerve*. 2007; **36**(5): 595–614.
[PubMed Abstract](#) | [Publisher Full Text](#)

6. Smith CJ, Johnson JM: **Responses to hyperthermia. Optimizing heat dissipation by convection and evaporation: Neural control of skin blood flow and sweating in humans.** *Auton Neurosci.* 2016; **196**: 25–36.
[PubMed Abstract](#) | [Publisher Full Text](#)
7. Crandall CG: **Heat stress and baroreflex regulation of blood pressure.** *Med Sci Sports Exerc.* 2008; **40**(12): 2063–70.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
8. Escourrou P, Freund PR, Rowell LB, *et al.*: **Splanchnic vasoconstriction in heat-stressed men: role of renin-angiotensin system.** *J Appl Physiol Respir Environ Exerc Physiol.* 1982; **52**(6): 1438–43.
[PubMed Abstract](#)
9. Kregel KC, Wall PT, Gisolfi CV: **Peripheral vascular responses to hyperthermia in the rat.** *J Appl Physiol (1985).* 1988; **64**(6): 2582–8.
[PubMed Abstract](#)
10. Minson CT, Wladkowski SL, Pawelczyk JA, *et al.*: **Age, splanchnic vasoconstriction, and heat stress during tilting.** *Am J Physiol.* 1999; **276**(1 Pt 2): R203–12.
[PubMed Abstract](#)
11. Satinoff E, Rutstein J: **Behavioral thermoregulation in rats with anterior hypothalamic lesions.** *J Comp Physiol Psychol.* 1970; **71**(1): 77–82.
[PubMed Abstract](#) | [Publisher Full Text](#)
12. Nagashima K, Nakai S, Tanaka M, *et al.*: **Neuronal circuitries involved in thermoregulation.** *Auton Neurosci.* 2000; **85**(1–3): 18–25.
[PubMed Abstract](#) | [Publisher Full Text](#)
13. Almeida MC, Steiner AA, Branco LG, *et al.*: **Cold-seeking behavior as a thermoregulatory strategy in systemic inflammation.** *Eur J Neurosci.* 2006; **23**(12): 3359–67.
[PubMed Abstract](#) | [Publisher Full Text](#)
14. **F** Almeida MC, Steiner AA, Branco LG, *et al.*: **Neural substrate of cold-seeking behavior in endotoxin shock.** *PLoS One.* 2006; **1**(1): e1.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
15. Terrien J, Perret M, Aujard F: **Behavioral thermoregulation in mammals: a review.** *Front Biosci (Landmark Ed).* 2011; **16**: 1428–44.
[PubMed Abstract](#) | [Publisher Full Text](#)
16. Shibasaki M, Crandall CG: **Mechanisms and controllers of eccrine sweating in humans.** *Front Biosci (Schol Ed).* 2010; **2**: 685–96.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
17. Farrell MJ, Treva D, Taylor NA, *et al.*: **Regional brain responses associated with thermogenic and psychogenic sweating events in humans.** *J Neurophysiol.* 2015; **114**(5): 2578–87.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
18. McKinley MJ, McAllen RM, Whyte D, *et al.*: **Central osmoregulatory influences on thermoregulation.** *Clin Exp Pharmacol Physiol.* 2008; **35**(5–6): 701–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
19. Poon CS: **Optimal interaction of respiratory and thermal regulation at rest and during exercise: role of a serotonin-gated spinoparabrachial thermoafferent pathway.** *Respir Physiol Neurobiol.* 2009; **169**(3): 234–42.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
20. McKinley MJ, Yao ST, Uschakov A, *et al.*: **The median preoptic nucleus: front and centre for the regulation of body fluid, sodium, temperature, sleep and cardiovascular homeostasis.** *Acta Physiol (Oxf).* 2015; **214**(1): 8–32.
[PubMed Abstract](#) | [Publisher Full Text](#)
21. McAllen RM, Farrell M, Johnson JM, *et al.*: **Human medullary responses to cooling and rewarming the skin: a functional MRI study.** *Proc Natl Acad Sci U S A.* 2006; **103**(3): 809–13.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
22. Morrison SF, Nakamura K: **Central neural pathways for thermoregulation.** *Front Biosci (Landmark Ed).* 2011; **16**: 74–104.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
23. Nakamura K: **Central circuitries for body temperature regulation and fever.** *Am J Physiol Regul Integr Comp Physiol.* 2011; **301**(5): R1207–28.
[PubMed Abstract](#) | [Publisher Full Text](#)
24. Morrison SF, Madden CJ: **Central nervous system regulation of brown adipose tissue.** *Compr Physiol.* 2014; **4**(4): 1677–713.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
25. Morrison SF, Madden CJ, Tupone D: **Central neural regulation of brown adipose tissue thermogenesis and energy expenditure.** *Cell Metab.* 2014; **19**(5): 741–56.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
26. Nakamura K, Morrison SF: **Central efferent pathways mediating skin cooling-evoked sympathetic thermogenesis in brown adipose tissue.** *Am J Physiol Regul Integr Comp Physiol.* 2007; **292**(1): R127–36.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
27. Nakamura K, Morrison SF: **Central efferent pathways for cold-defensive and febrile shivering.** *J Physiol.* 2011; **589**(Pt 14): 3641–58.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
28. Tanaka M, McKinley MJ, McAllen RM: **Preoptic-raphé connections for thermoregulatory vasomotor control.** *J Neurosci.* 2011; **31**(13): 5078–88.
[PubMed Abstract](#) | [Publisher Full Text](#)
29. Tanaka M, McKinley MJ, McAllen RM: **Role of an excitatory preoptic-raphé pathway in febrile vasoconstriction of the rat's tail.** *Am J Physiol Regul Integr Comp Physiol.* 2013; **305**(12): R1479–89.
[PubMed Abstract](#) | [Publisher Full Text](#)
30. Cao WH, Fan W, Morrison SF: **Medullary pathways mediating specific sympathetic responses to activation of dorsomedial hypothalamus.** *Neuroscience.* 2004; **126**(1): 229–40.
[PubMed Abstract](#) | [Publisher Full Text](#)
31. **F** Nakamura Y, Nakamura K, Matsumura K, *et al.*: **Direct pyrogenic input from prostaglandin EP3 receptor-expressing preoptic neurons to the dorsomedial hypothalamus.** *Eur J Neurosci.* 2005; **22**(12): 3137–46.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
32. Madden CJ, Morrison SF: **Excitatory amino acid receptors in the dorsomedial hypothalamus mediate prostaglandin-evoked thermogenesis in brown adipose tissue.** *Am J Physiol Regul Integr Comp Physiol.* 2004; **286**(2): R320–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
33. **F** Dimitrov EL, Kim YY, Usdin TB: **Regulation of hypothalamic signaling by tuberoinfundibular peptide of 39 residues is critical for the response to cold: a novel peptidergic mechanism of thermoregulation.** *J Neurosci.* 2011; **31**(49): 18166–79.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
34. **F** Zhang Y, Kerman IA, Laque A, *et al.*: **Leptin-receptor-expressing neurons in the dorsomedial hypothalamus and median preoptic area regulate sympathetic brown adipose tissue circuits.** *J Neurosci.* 2011; **31**(5): 1873–84.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
35. Zhang YH, Yanase-Fujiwara M, Hosono T, *et al.*: **Warm and cold signals from the preoptic area: which contribute more to the control of shivering in rats?** *J Physiol.* 1995; **485**(Pt 1): 195–202.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
36. Boulant JA, Dean JB: **Temperature receptors in the central nervous system.** *Annu Rev Physiol.* 1986; **48**: 639–54.
[PubMed Abstract](#) | [Publisher Full Text](#)
37. Griffin JD, Kaple ML, Chow AR, *et al.*: **Cellular mechanisms for neuronal thermosensitivity in the rat hypothalamus.** *J Physiol.* 1996; **492**(Pt 1): 231–42.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
38. Lundius EG, Sanchez-Alavez M, Ghochani Y, *et al.*: **Histamine influences body temperature by acting at H1 and H3 receptors on distinct populations of preoptic neurons.** *J Neurosci.* 2010; **30**(12): 4369–81.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
39. Griffin JD, Saper CB, Boulant JA: **Synaptic and morphological characteristics of temperature-sensitive and -insensitive rat hypothalamic neurons.** *J Physiol.* 2001; **537**(Pt 2): 521–35.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
40. Ootsuka Y, McAllen RM: **Comparison between two rat sympathetic pathways activated in cold defense.** *Am J Physiol Regul Integr Comp Physiol.* 2006; **291**(3): R589–95.
[PubMed Abstract](#) | [Publisher Full Text](#)
41. Glotzbach SF, Heller HC: **Central nervous regulation of body temperature during sleep.** *Science.* 1976; **194**(4264): 537–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
42. Scammell TE, Elmquist JK, Griffin JD, *et al.*: **Ventromedial preoptic prostaglandin E2 activates fever-producing autonomic pathways.** *J Neurosci.* 1996; **16**(19): 6246–54.
[PubMed Abstract](#)
43. Nakamura K, Kaneko T, Yamashita Y, *et al.*: **Immunohistochemical localization of prostaglandin EP3 receptor in the rat nervous system.** *J Comp Neurol.* 2000; **421**(4): 543–69.
[PubMed Abstract](#) | [Publisher Full Text](#)
44. Nakamura K, Matsumura K, Kaneko T, *et al.*: **The rostral raphe pallidus nucleus mediates pyrogenic transmission from the preoptic area.** *J Neurosci.* 2002; **22**(11): 4600–10.
[PubMed Abstract](#)
45. **F** Lazarus M, Yoshida K, Coppari R, *et al.*: **EP3 prostaglandin receptors in the median preoptic nucleus are critical for fever responses.** *Nat Neurosci.* 2007; **10**(9): 1131–3.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
46. **F** Lkhagvasuren B, Nakamura Y, Oka T, *et al.*: **Social defeat stress induces hyperthermia through activation of thermoregulatory sympathetic premotor neurons in the medullary raphe region.** *Eur J Neurosci.* 2011; **34**(9): 1442–52.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
47. **F** Kataoka N, Hioki H, Kaneko T, *et al.*: **Psychological stress activates a dorsomedial hypothalamus-medullary raphe circuit driving brown adipose tissue thermogenesis and hyperthermia.** *Cell Metab.* 2014; **20**(2): 346–58.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
48. **F** Mohammed M, Ootsuka Y, Blessing W: **Brown adipose tissue thermogenesis contributes to emotional hyperthermia in a resident rat suddenly confronted with an intruder rat.** *Am J Physiol Regul Integr Comp Physiol.* 2014; **306**(6): R394–400.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
49. Nakamura K, Morrison SF: **Preoptic mechanism for cold-defensive responses to skin cooling.** *J Physiol.* 2008; **586**(10): 2611–20.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
50. Tanaka M, McKinley MJ, McAllen RM: **Roles of two preoptic cell groups in tonic and febrile control of rat tail sympathetic fibers.** *Am J Physiol Regul Integr Comp Physiol.* 2009; **296**(4): R1248–57.
[PubMed Abstract](#) | [Publisher Full Text](#)
51. Yoshida K, Li X, Cano G, *et al.*: **Parallel preoptic pathways for thermoregulation.**

- J Neurosci.* 2009; **29**(38): 11954–64.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
52. Osaka T: **Thermoregulatory responses elicited by microinjection of L-glutamate and its interaction with thermogenic effects of GABA and prostaglandin E₂ in the preoptic area.** *Neuroscience.* 2012; **226**: 156–64.
[PubMed Abstract](#) | [Publisher Full Text](#)
 53. Mohammed M, Kulasekara K, De Menezes RC, *et al.*: **Inactivation of neuronal function in the amygdaloid region reduces tail artery blood flow alerting responses in conscious rats.** *Neuroscience.* 2013; **228**: 13–22.
[PubMed Abstract](#) | [Publisher Full Text](#)
 54. Ootsuka Y, Mohammed M: **Activation of the habenula complex evokes autonomic physiological responses similar to those associated with emotional stress.** *Physiol Rep.* 2015; **3**(2): pii: e12297.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 55. **F** Zhang YH, Hosono T, Yanase-Fujiwara M, *et al.*: **Effect of midbrain stimulations on thermoregulatory vasomotor responses in rats.** *J Physiol.* 1997; **503**(Pt 1): 177–86.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 56. **F** Yoshida K, Maruyama M, Hosono T, *et al.*: **Fos expression induced by warming the preoptic area in rats.** *Brain Res.* 2002; **933**(2): 109–17.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 57. **F** Yoshida K, Konishi M, Nagashima K, *et al.*: **Fos activation in hypothalamic neurons during cold or warm exposure: projections to periaqueductal gray matter.** *Neuroscience.* 2005; **133**(4): 1039–46.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 58. Shibata M, Uno T, Hashimoto M: **Disinhibition of lower midbrain neurons enhances non-shivering thermogenesis in anesthetized rats.** *Brain Res.* 1999; **833**(2): 242–50.
[PubMed Abstract](#) | [Publisher Full Text](#)
 59. Cao WH, Madden CJ, Morrison SF: **Inhibition of brown adipose tissue thermogenesis by neurons in the ventrolateral medulla and in the nucleus tractus solitarius.** *Am J Physiol Regul Integr Comp Physiol.* 2010; **299**(1): R277–90.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 60. Tupone D, Madden CJ, Morrison SF: **Central activation of the A1 adenosine receptor (A1AR) induces a hypothalamic, torpor-like state in the rat.** *J Neurosci.* 2013; **33**(36): 14512–25.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 61. Sato H, Hashitani T, Isobe Y, *et al.*: **Descending influences from nucleus raphe magnus on fusimotor neurone activity in rats.** *J Therm Biol.* 1990; **15**(3–4): 259–65.
[Publisher Full Text](#)
 62. Tanaka M, Owens NC, Nagashima K, *et al.*: **Reflex activation of rat fusimotor neurons by body surface cooling, and its dependence on the medullary raphe.** *J Physiol.* 2006; **572**(Pt 2): 569–83.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 63. Almeida MC, Hew-Butler T, Soriano RN, *et al.*: **Pharmacological blockade of the cold receptor TRPM8 attenuates autonomic and behavioral cold defenses and decreases deep body temperature.** *J Neurosci.* 2012; **32**(6): 2086–99.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 64. Tabarean IV, Behrens MM, Bartfai T, *et al.*: **Prostaglandin E₂-increased thermosensitivity of anterior hypothalamic neurons is associated with depressed inhibition.** *Proc Natl Acad Sci U S A.* 2004; **101**(8): 2590–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 65. Boulant JA: **Counterpoint: Heat-induced membrane depolarization of hypothalamic neurons: an unlikely mechanism of central thermosensitivity.** *Am J Physiol Regul Integr Comp Physiol.* 2006; **290**(5): R1481–4; discussion R1484.
[PubMed Abstract](#)
 66. Kobayashi S, Hori A, Matsumura K, *et al.*: **Point: Heat-induced membrane depolarization of hypothalamic neurons: a putative mechanism of central thermosensitivity.** *Am J Physiol Regul Integr Comp Physiol.* 2006; **290**(5): R1479–80; discussion R1484.
[PubMed Abstract](#) | [Publisher Full Text](#)
 67. Wechselberger M, Wright CL, Bishop GA, *et al.*: **Ionic channels and conductance-based models for hypothalamic neuronal thermosensitivity.** *Am J Physiol Regul Integr Comp Physiol.* 2006; **291**(3): R518–29.
[PubMed Abstract](#) | [Publisher Full Text](#)
 68. Craig AD: **How do you feel? Interoception: the sense of the physiological condition of the body.** *Nat Rev Neurosci.* 2002; **3**(8): 655–66.
[PubMed Abstract](#) | [Publisher Full Text](#)
 69. Craig AD, Bushnell MC, Zhang ET, *et al.*: **A thalamic nucleus specific for pain and temperature sensation.** *Nature.* 1994; **372**(6508): 770–3.
[PubMed Abstract](#) | [Publisher Full Text](#)
 70. Hylden JL, Anton F, Nahin RL: **Spinal lamina I projection neurons in the rat: collateral innervation of parabrachial area and thalamus.** *Neuroscience.* 1989; **28**(1): 27–37.
[PubMed Abstract](#) | [Publisher Full Text](#)
 71. Li J, Xiong K, Pang Y, *et al.*: **Medullary dorsal horn neurons providing axons to both the parabrachial nucleus and thalamus.** *J Comp Neurol.* 2006; **498**(4): 539–51.
[PubMed Abstract](#) | [Publisher Full Text](#)
 72. **F** Bratincsák A, Palkovits M: **Activation of brain areas in rat following warm and cold ambient exposure.** *Neuroscience.* 2004; **127**(2): 385–97.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 73. **F** Nakamura K, Morrison SF: **A thermosensory pathway that controls body temperature.** *Nat Neurosci.* 2008; **11**(1): 62–71.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 74. **F** Nakamura K, Morrison SF: **A thermosensory pathway mediating heat-defense responses.** *Proc Natl Acad Sci U S A.* 2010; **107**(19): 8848–53.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 75. Madden CJ, Morrison SF: **Hypoxic activation of arterial chemoreceptors inhibits sympathetic outflow to brown adipose tissue in rats.** *J Physiol.* 2005; **566**(Pt 2): 559–73.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 76. **F** Madden CJ: **Glucoprivation in the ventrolateral medulla decreases brown adipose tissue sympathetic nerve activity by decreasing the activity of neurons in raphe pallidus.** *Am J Physiol Regul Integr Comp Physiol.* 2012; **302**(2): R224–32.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 77. Baker MA, Doris PA: **Control of evaporative heat loss during changes in plasma osmolality in the cat.** *J Physiol.* 1982; **328**(1): 535–45.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 78. Takamata A, Mack GW, Gillen CM, *et al.*: **Osmoregulatory modulation of thermal sweating in humans: reflex effects of drinking.** *Am J Physiol.* 1995; **268**(2 Pt 2): R414–22.
[PubMed Abstract](#)
 79. Kenney WL, Staniewicz AE, Bruning RS, *et al.*: **Blood pressure regulation III: what happens when one system must serve two masters: temperature and pressure regulation?** *Eur J Appl Physiol.* 2014; **114**(3): 467–79.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 80. Székely M: **The vagus nerve in thermoregulation and energy metabolism.** *Auton Neurosci.* 2000; **85**(1–3): 26–38.
[PubMed Abstract](#) | [Publisher Full Text](#)
 81. **F** Tsukita S, Yamada T, Uno K, *et al.*: **Hepatic glucokinase modulates obesity predisposition by regulating BAT thermogenesis via neural signals.** *Cell Metab.* 2012; **16**(6): 825–32.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 82. **F** Ono K, Tsukamoto-Yasui M, Hara-Kimura Y, *et al.*: **Intragastric administration of capsiate, a transient receptor potential channel agonist, triggers thermogenic sympathetic responses.** *J Appl Physiol (1985).* 2011; **110**(3): 789–98.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 83. **F** Blouet C, Schwartz GJ: **Duodenal lipid sensing activates vagal afferents to regulate non-shivering brown fat thermogenesis in rats.** *PLoS One.* 2012; **7**(12): e51898.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 84. Ngampramuan S, Cerri M, Del Vecchio F, *et al.*: **Thermoregulatory correlates of nausea in rats and musk shrews.** *Oncotarget.* 2014; **5**(6): 1565–75.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 85. Romanovsky AA, Almeida MC, Garami A, *et al.*: **The transient receptor potential vanilloid-1 channel in thermoregulation: a thermosensor it is not.** *Pharmacol Rev.* 2009; **61**(3): 228–61.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 86. **F** Peters JH, McDougall SJ, Fawley JA, *et al.*: **Primary afferent activation of thermosensitive TRPV1 triggers asynchronous glutamate release at central neurons.** *Neuron.* 2010; **65**(5): 657–69.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 87. **F** Fawley JA, Hofmann ME, Andresen MC: **Cannabinoid 1 and transient receptor potential vanilloid 1 receptors discretely modulate evoked glutamate separately from spontaneous glutamate transmission.** *J Neurosci.* 2014; **34**(24): 8324–32.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 88. **F** Fawley JA, Hofmann ME, Largent-Milnes TM, *et al.*: **Temperature differentially facilitates spontaneous but not evoked glutamate release from cranial visceral primary afferents.** *PLoS One.* 2015; **10**(5): e0127764.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 89. **F** Shoudai K, Peters JH, McDougall SJ, *et al.*: **Thermally active TRPV1 tonically drives central spontaneous glutamate release.** *J Neurosci.* 2010; **30**(43): 14470–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 90. **F** Steiner AA, Turek VF, Almeida MC, *et al.*: **Nonthermal activation of transient receptor potential vanilloid-1 channels in abdominal viscera tonically inhibits autonomic cold-defense effectors.** *J Neurosci.* 2007; **27**(28): 7459–68.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 91. **F** Hypothermia after Cardiac Arrest Study Group: **Mild therapeutic hypothermia to improve the neurologic outcome after cardiac arrest.** *N Engl J Med.* 2002; **346**(8): 549–56.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 92. **F** Shankaran S, Laptook AR, Ehrenkranz RA, *et al.*: **Whole-body hypothermia for neonates with hypoxic-ischemic encephalopathy.** *N Engl J Med.* 2005; **353**(15): 1574–84.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 93. van der Worp HB, van Gijn J: **Clinical practice. Acute ischemic stroke.** *N Engl J Med.*

- 2007; **357**(6): 572–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
94. Hemmen TM, Lyden PD: **Hypothermia after acute ischemic stroke**. *J Neurotrauma*. 2009; **26**(3): 387–91.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 95. Madden CJ, Tupone D, Cano G, *et al.*: **$\alpha 2$ Adrenergic receptor-mediated inhibition of thermogenesis**. *J Neurosci*. 2013; **33**(5): 2017–28.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 96. Tupone D, Madden CJ, Morrison SF: **Autonomic regulation of brown adipose tissue thermogenesis in health and disease: potential clinical applications for altering BAT thermogenesis**. *Front Neurosci*. 2014; **8**: 14.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 97. **F** Cerri M, Mastrotto M, Tupone D, *et al.*: **The inhibition of neurons in the central nervous pathways for thermoregulatory cold defense induces a suspended animation state in the rat**. *J Neurosci*. 2013; **33**(7): 2984–93.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 98. **F** Muzzi M, Blasi F, Masi A, *et al.*: **Neurological basis of AMP-dependent thermoregulation and its relevance to central and peripheral hyperthermia**. *J Cereb Blood Flow Metab*. 2013; **33**(2): 183–90.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 99. **F** Miyazawa S, Shimizu Y, Shiina T, *et al.*: **Central A1-receptor activation associated with onset of torpor protects the heart against low temperature in the Syrian hamster**. *Am J Physiol Regul Integr Comp Physiol*. 2008; **295**(3): R991–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 100. **F** Jinka TR, Tøien Ø, Drew KL: **Season primes the brain in an arctic hibernator to facilitate entrance into torpor mediated by adenosine A₁ receptors**. *J Neurosci*. 2011; **31**(30): 10752–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 101. **F** Iliff BW, Swoap SJ: **Central adenosine receptor signaling is necessary for daily torpor in mice**. *Am J Physiol Regul Integr Comp Physiol*. 2012; **303**(5): R477–84.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 102. **F** Feketa VV, Balasubramanian A, Flores CM, *et al.*: **Shivering and tachycardic responses to external cooling in mice are substantially suppressed by TRPV1 activation but not by TRPM8 inhibition**. *Am J Physiol Regul Integr Comp Physiol*. 2013; **305**(9): R1040–50.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 103. **F** Feketa VV, Zhang Y, Cao Z, *et al.*: **Transient receptor potential melastatin 8 channel inhibition potentiates the hypothermic response to transient receptor potential vanilloid 1 activation in the conscious mouse**. *Crit Care Med*. 2014; **42**(5): e355–63.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 104. **F** Cypess AM, Lehman S, Williams G, *et al.*: **Identification and importance of brown adipose tissue in adult humans**. *N Engl J Med*. 2009; **360**(15): 1509–17.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 105. **F** Saito M, Okamatsu-Ogura Y, Matsushita M, *et al.*: **High incidence of metabolically active brown adipose tissue in healthy adult humans: effects of cold exposure and adiposity**. *Diabetes*. 2009; **58**(7): 1526–31.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 106. **F** van Marken Lichtenbelt WD, Vanhommerig JW, Smulders NM, *et al.*: **Cold-activated brown adipose tissue in healthy men**. *N Engl J Med*. 2009; **360**(15): 1500–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 107. Orava J, Nuutila P, Lidell ME, *et al.*: **Different metabolic responses of human brown adipose tissue to activation by cold and insulin**. *Cell Metab*. 2011; **14**(2): 272–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
 108. Lowell BB, S-Susulic V, Hamann A, *et al.*: **Development of obesity in transgenic mice after genetic ablation of brown adipose tissue**. *Nature*. 1993; **366**(6457): 740–2.
[PubMed Abstract](#) | [Publisher Full Text](#)
 109. Hamann A, Flier JS, Lowell BB: **Decreased brown fat markedly enhances susceptibility to diet-induced obesity, diabetes, and hyperlipidemia**. *Endocrinology*. 1996; **137**(1): 21–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
 110. Kopecky J, Clarke G, Enerbäck S, *et al.*: **Expression of the mitochondrial uncoupling protein gene from the aP2 gene promoter prevents genetic obesity**. *J Clin Invest*. 1995; **96**(6): 2914–23.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 111. **F** Hanssen MJ, van der Lans AA, Brans B, *et al.*: **Short-term cold acclimation recruits brown adipose tissue in obese humans**. *Diabetes*. 2015; pii: db151372.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 112. Orava J, Nuutila P, Nojonen T, *et al.*: **Blunted metabolic responses to cold and insulin stimulation in brown adipose tissue of obese humans**. *Obesity (Silver Spring)*. 2013; **21**(11): 2279–87.
[PubMed Abstract](#) | [Publisher Full Text](#)
 113. Sakaguchi T, Arase K, Fislser JS, *et al.*: **Effect of a high-fat diet on firing rate of sympathetic nerves innervating brown adipose tissue in anesthetized rats**. *Physiol Behav*. 1989; **45**(6): 1177–82.
[PubMed Abstract](#) | [Publisher Full Text](#)
 114. Levin BE: **Reduced norepinephrine turnover in organs and brains of obesity-prone rats**. *Am J Physiol*. 1995; **268**(2 Pt 2): R389–94.
[PubMed Abstract](#)
 115. Madden CJ: **Consumption of a high fat diet inhibits sympathetic outflow to brown adipose tissue (BAT) via vagal afferent activation of neurons in the Nucleus Tractus Solitarius (NTS)**. *Auton Neurosci*. 2015; **192**: 13.
[Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. **Andrej A Romanovsky**
Systemic Inflammation Laboratory (FeverLab), Trauma Research, St. Joseph's Hospital and Medical Center, Phoenix, AZ, USA
Competing Interests: No competing interests were disclosed.
2. **Matteo Cerri**
Department of Biomedical and NeuroMotor Sciences, University of Bologna, Bologna, Italy
Competing Interests: No competing interests were disclosed.
3. **Robin M McAllen**
¹ The Florey Institute of Neuroscience and Mental Health, University of Melbourne, Parkville, Victoria, Australia
² Department of Anatomy and Neuroscience, University of Melbourne, Parkville, Victoria, Australia
Michael J McKinley
¹ The Florey Institute of Neuroscience and Mental Health, University of Melbourne, Parkville, Victoria, Australia
² Department of Physiology, University of Melbourne, Parkville, Victoria, Australia
Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research