REVIEW

Recent advances in microscopic techniques for visualizing leukocytes *in vivo* [version 1; peer review: 3 approved]

Rohit Jain^{1,2}, Shweta Tikoo ^{1,2}, Wolfgang Weninger¹⁻³

¹Immune Imaging Program, The Centenary Institute, University of Sydney, Newtown, NSW 2042, Australia

²Discipline of Dermatology, Sydney Medical School, University of Sydney, NSW 2006, Australia

³Department of Dermatology, Royal Prince Alfred Hospital, Camperdown, NSW 2050, Australia

V1 First published: 19 May 2016, 5(F1000 Faculty Rev):915
<https://doi.org/10.12688/f1000research.8127.1>

Latest published: 19 May 2016, 5(F1000 Faculty Rev):915
<https://doi.org/10.12688/f1000research.8127.1>

Abstract

Leukocytes are inherently motile and interactive cells. Recent advances in intravital microscopy approaches have enabled a new vista of their behavior within intact tissues in real time. This brief review summarizes the developments enabling the tracking of immune responses *in vivo*.

Keywords

leukocytes, visualizing, *in vivo*

Open Peer Review

Approval Status

	1	2	3
version 1 19 May 2016

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Tim Lammermann**, Max Planck Institute of Immunobiology and Epigenetics, Freiburg, Germany
2. **Peter Friedl**, Radboud University Medical Centre, 6525GA Nijmegen, The Netherlands
University of Texas MD Anderson Cancer Center, Houston, USA
3. **Tri Phan**, Garvan Institute of Medical research, Darlinghurst, Australia

Any comments on the article can be found at the end of the article.

Corresponding authors: Rohit Jain (r.jain@centenary.org.au), Wolfgang Weninger (w.weninger@centenary.org.au)

Competing interests: The authors declare that they have no competing interests.

Grant information: Wolfgang Weninger was supported by project grants 1066477, 1102398, 1104876 and 1106439, from the National Health and Medical Research Council Australia. Rohit Jain was supported by an Early Career Researcher grant from the Sydney Medical School, Sydney, Australia. Shewta Tikoo was supported by an Early Career Researcher grant from the Sydney Medical School and PdCCRS grant from Cure Cancer Australia.

Copyright: © 2016 Jain R *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Jain R, Tikoo S and Weninger W. **Recent advances in microscopic techniques for visualizing leukocytes *in vivo* [version 1; peer review: 3 approved]** F1000Research 2016, 5(F1000 Faculty Rev):915 <https://doi.org/10.12688/f1000research.8127.1>

First published: 19 May 2016, 5(F1000 Faculty Rev):915 <https://doi.org/10.12688/f1000research.8127.1>

Introduction

The elicitation of an immune response against invading pathogens or tumor cells relies on a complex and highly orchestrated interplay among leukocytes, blood vessels, and stromal cell populations. During this process, immune cells need to identify and interpret structural and molecular guidance cues such as extracellular matrix fibers and chemokines to navigate through diverse tissue microenvironments towards foci of tissue injury, where they finally mediate their effector functions. Microscopic investigations into this highly orchestrated behavior of immune cells have provided insight into the workings of innate and adaptive immunity in live mice at a level of resolution previously unattained.

To visualize the behavior of immune cells *in vivo*, initial studies relied on bright field and epifluorescence video microscopy. Although limited in resolution and restricted by tissue clarity, these studies were instrumental for dissecting the molecular mechanisms regulating leukocyte homing during the process of extravasation^{1–4}. However, the advent of advanced confocal⁵ and two-photon^{6,7} imaging modalities revolutionized our understanding of lymphocyte behavior within the interstitial space of intact organs, initially within re-aggregated thymic organ cultures⁶ and explanted intact lymph nodes^{5,7}. These studies illuminated the complex *in situ* behavior of T cells within lymph nodes in the absence or presence of antigen^{5,7} and during their selection within the thymus⁶. These advancements in imaging technologies further catalyzed the emerging field of intravital imaging of the immune system. Indeed, shortly after the explant models, imaging of intact organs *in situ* in living mice was reported^{8,9}. Since then, a large number of tissues/organ systems have been adapted for *in vivo* imaging, revealing insights into the functioning of the immune system in the steady state, during initial pathogen encounter, during the priming phase of adaptive immune responses, and within effector organs during infection, inflammation, and within tumors. The impact of intravital imaging on the field of leukocyte biology and immunology has recently been reviewed in detail elsewhere^{10–13}.

In summary, the development of new imaging tools and molecular probes, coupled with a wealth of transgenic fluorescent reporter mice, has ushered in a new era for understanding the cellular and molecular regulators of leukocyte function *in situ*. In this review, we highlight some of the technical advances in this field of research, with a focus on multi-photon imaging.

Advances in two-photon imaging methodologies

Without doubt, the introduction of two-photon microscopy into biological imaging has enormously furthered our understanding of physiological and pathological processes at the level of intact organs. Unlike single-photon excitation used in confocal microscopy, two-photon excitation is spatially restricted to the focal plane^{10,12,14,15}. This results in reduced levels of phototoxicity and photobleaching, enabling deep tissue imaging for prolonged periods of time^{10,12}. Nevertheless, several technical challenges still persist. For example, physiological tissue movement due to pulsation of blood vessels, contraction of muscles, respiration, or heartbeat can be detrimental to the overall quality of the data. In addition, surgical and anesthetic requirements may alter the behavior of cells due to the induction of hypoxia, changes in tissue pH,

or the generation of pro-inflammatory mediators. It is therefore imperative to develop standardized, reproducible procedures for the exposure of organs during the imaging session.

For many peripheral tissues like skin¹⁶ and draining lymph nodes^{8,9,17}, and internal organs such as the liver^{18–21} and spleen^{22–24}, tissue stabilization during imaging has been carried out using relatively simple methodologies and devices. Similarly, intravital imaging of bone marrow cavity within the skull calvaria has been adapted for two-photon microscopy²⁵ and has played a significant role in highlighting immune cell behavior within this environment^{25–28}. Akin to this, various methodologies have been developed for longitudinal imaging of tumors¹⁵. Cumulatively, these new imaging methodologies have provided unprecedented understanding of leukocyte behavior *in vivo* in a variety of tissues^{11–13,29–31}.

Nonetheless, stable imaging of certain tissues, such as the brain, heart, and lung, still presents a significant challenge. In the case of the brain, the two most frequently used methodologies involve either bone thinning^{32,33} or removal of the cranial bone^{34–36} for imaging of the underlying meninges and brain parenchyma. Similarly, longitudinal imaging of the brain has been made possible by implantation of an optical window^{34,37,38}. These methodologies allow studying the behavior of various immune cells like microglia^{36,39,40}, monocytes^{41,42}, and T cells^{13,43–46} within the intact brain and spine under homeostasis, infection, or inflammation. Newer techniques, such as implantation of glass prisms⁴⁷ or micro-optical probes⁴⁸, have been developed for imaging of deeper structures within the brain^{47,48}, but they are yet to be harnessed for understanding leukocyte behavior.

In vivo imaging of lungs presents a significant technical challenge due to tissue movement and its localization within the pleural cavity. Nonetheless, several strategies have been developed that involve the closing of ventilation to the imaged lung⁴⁹ or synchronizing image acquisition with respiration⁵⁰. Recently, several new methodologies^{51–53} have described the creation of a small thoracic window, wherein the tissue is stabilized via application of a mild negative pressure^{51,52} or surgical glue⁵³. These approaches maintain ventilation and perfusion within the tissue and permit imaging throughout the respiratory cycle^{51,52}. These recent advances have highlighted the dynamic nature of T cells^{51,54}, neutrophils^{28,51,53}, monocytes⁵³, macrophages⁵⁵, and dendritic cells^{54,55} within the pulmonary tissue in the steady state and during inflammatory conditions. For instance, *in vivo* imaging of the lungs during influenza virus infection has identified a unique cooperative behavior between neutrophils and CD8⁺ effector T cells. Thus, within influenza-infected lungs, migrating neutrophils leave behind a trail of the chemokine CXCL12, which is used by effector CD8⁺ T cells as a guidance cue within the interstitium⁵⁶. This study enforces the role of *in vivo* imaging in understanding the dynamic spatiotemporal cooperation between immune cells required for generating an efficient immune response.

By using endoscopic time-lapse imaging⁵⁷ or optical windows^{58–60}, researchers have recently addressed the challenges associated with *in vivo* imaging of leukocytes within the beating heart⁶¹. Endoscopic imaging of the heart under physiological conditions revealed

the patrolling behavior of monocytes within coronary vessels⁵⁷. Post-infarction, rapid recruitment of both monocytes and neutrophils was observed^{57,58}. Surprisingly, unlike most tissues, large coronary veins but not post-capillary venules¹³ were the predominant sites for neutrophil extravasation post-ischemia/reperfusion injury⁵⁸. Most neutrophils migrated via “hotspots” within the inflamed coronary veins⁵⁸. Although this technique is still in its early days, *in vivo* imaging of the heart nonetheless holds immense potential in understanding leukocyte/lymphocyte behavior post-cardiac transplant and during organ rejection.

In addition to the intravital imaging of the beating heart, several methodologies have been developed to image blood vessels *in vivo*^{62–65}. Studies have highlighted, for example, the dynamic migratory behavior of neutrophils^{63,64} and T lymphocytes⁶³ within atherosclerotic lesions. Recently, the use of cardiac triggered acquisition coupled with image registration and post-processing image correction has provided further advancement of stable imaging of blood vessels⁶⁵. This methodology enables improved temporal resolution, which has been used to define the dynamic scanning behavior of dendritic processes of leukocytes within the atherosclerotic plaques of pulsatile blood vessels⁶⁵.

The use of two-photon excitation can also be exploited for inducing localized injury without physical manipulation of the surrounding tissue. This provides opportunities for studying the behavior of tissue-resident and recruited leukocytes during sterile inflammation *in situ*. This methodology has been used, for example, to highlight the dynamic behavior of microglia within the brain post injury^{36,39} and to delineate the cellular and molecular players required for interstitial migration of neutrophils^{66,67}, monocytes⁶⁷, and dendritic cells⁶⁸ within the skin^{13,69,70}. Thus, laser-induced injury has been crucial for understanding the cellular and molecular interplay required for interstitial migration of neutrophils and the damage response within the inflamed dermis^{66,67}. Following injury, scarce initial neutrophils rapidly migrate towards the site of injury following chemokine gradients. This has been dubbed the “neutrophil scouting phase”, which is followed by an “amplification phase” wherein a large number of neutrophils display directed migration towards the injury site. This second phase is dependent on cyclic-adenosine diphosphate ribose (c-ADPR) and leukotriene B4 (LTB4) signaling^{66,67}. The large influx of neutrophils at the injury site leads to the formation of a stable neutrophil cluster also referred to as the “stabilization phase”. The stability of the cluster is governed by signaling via G-protein-coupled receptors such as C-X-C chemokine receptor 2 (CXCR-2), *N*-formyl peptide receptor 2 (FPR-2), and LTB4 receptor-1 (BLT-1). Surprisingly, although integrins were dispensable for neutrophil migration within the inflamed dermis, they were required for migrating within the injury foci devoid of extracellular matrix. Together, these studies uncover the crucial role of intravital imaging in understanding the spatiotemporal dynamics of the innate immune response *in vivo*.

The use of *in vivo* imaging has also been instrumental for understanding immune responses against invading pathogens^{13,71,72} and has highlighted the dynamic behavior of innate and adaptive immune cells during various infections including *Leishmania major*^{73–77}, *Leishmania donovani*^{78–80}, *Toxoplasma gondii*^{81–83}, *Borrelia burgdorferi*^{84,85}, *Staphylococcus aureus*^{86–88}, *Plasmodium*

spp.^{42,46,89–93}, and *Mycobacterium*-induced granulomas^{19,21}. These studies have identified various strategies used by pathogens to evade immune responses; for instance, post-intradermal inoculation, *Plasmodium berghei* sporozoites rapidly migrate into the blood vessels and lymphatics⁸⁹. In contrast, experiments performed with another vector-borne parasite, *Leishmania major*, revealed that these parasites do not actively migrate but rather induce neutrophil recruitment to aid parasite survival^{73,74,94–97}. Similarly, *in vivo* imaging of mycobacterial granulomas within the liver have highlighted the presence of numerous antigen-presenting cells (APCs) and CD4⁺ T cells within the granulomatous tissue^{19,21}. A paucity of mycobacterial antigens limited the number of stable interactions between APCs and antigen-specific CD4⁺ T cells and thereby suppressed the release of immunomodulatory/protective cytokines²¹. Cumulatively, *in vivo* imaging of host-pathogen interactions have identified novel cellular and molecular mechanisms involved in regulating pathogen uptake and transport, generation of adaptive immune responses, identification of new migratory strategies used for scanning infected tissues, and elicitation of effector immune responses.

Similarly, two-photon imaging of tumors has provided fascinating insights into leukocyte behavior within the tumor milieu: for example, effector CD8⁺ T cell-mediated scanning of tumor tissue^{98–100}, induction of tumor cell apoptosis¹⁰¹, generation of T cell tolerance¹⁰², co-migration of macrophages and tumor cells¹⁰³, and the role of perivascular macrophages in tumor cell dissemination¹⁰⁴. Using experimental models that recapitulate the pathophysiology of multiple sclerosis, investigators have revealed fibrinogen-mediated perivascular clustering of microglia⁴⁰, formation of immunological synapse between APCs and effector CD4⁺ T cells⁴⁴, and neuronal cell death¹⁰⁵ with the use of *in vivo* imaging.

The latest development in the use of optical phase-locked ultrasound lens provides a new methodology for high-speed volumetric imaging *in vivo*¹⁰⁶. This allows for the imaging of Ca²⁺ signaling as well as alterations in cell morphology at high speeds¹⁰⁶. This methodology has been used to highlight intravascular and extravascular neutrophil dynamics at a temporal resolution previously unattainable¹⁰⁶.

Altogether, the advancements in imaging instrumentation and development of new and improved physiological models have been crucial in developing a deeper and dynamic understanding of innate and adaptive immunity.

Transgenic fluorescent reporter mouse models for *in vivo* imaging

Visualization of cells and structures within tissues via laser excitation relies, on the one hand, on their inherent (auto)-fluorescent features; for example, second and third harmonic generation facilitate highlighting large extracellular molecules such as collagen and elastin fibers without the need for counterstaining^{107–110}. In addition, certain auto-fluorescent metabolites such as NADPH enable the interrogation of metabolic states of cells^{111,112}. On the other hand, most leukocyte subsets require external or genetic tagging for their detection. Initial imaging studies made use of *ex vivo* labeled, adoptively transferred immune cells. Although manipulation of cells may alter their behavior, these studies were crucial in

understanding migratory and interactive cellular dynamics *in vivo*. Nevertheless, advances in genetic engineering technology coupled with the availability of a wide array of genetically encoded fluorophores have significantly expanded the scope of *in vivo* imaging. Introduction of transgenic fluorescent proteins at specific genetic loci or under the control of tissue- or cell-specific promoters has been crucial for following cells of interest under physiologic conditions. Numerous fluorescent reporter mouse strains are now available for *in vivo* imaging, and the choice of a particular reporter relies on the experimental design and the scientific question being pursued^{113,114}. Although the detailed description of these strains is beyond the scope of this review, a few of the novel and versatile reporter mouse strains are detailed below.

The dynamic nature of immune cells presents a significant challenge while tracking the individual cell longitudinally. This challenge can be overcome by using photoactivatable fluorescent reporters, for example photoactivatable GFP¹¹⁵. Using such an approach, Victora and colleagues deciphered B cell¹¹⁶ and T follicular helper cell¹¹⁷ dynamics within the germinal center of murine lymph nodes, providing valuable information regarding the regulation of humoral immunity. This strategy is useful for tracking the migration and distribution of immune cells *in vivo*. However, the degradation of photoactivated GFP within the marked cells limits the timespan available for tracking, usually in the order of days ($t_{1/2}$ - 30 hours for naive B cells¹¹⁶). More recently, photo-switchable transgenic reporters like KikGR^{118,119} and Kaede¹²⁰ mice have become available. Briefly, in Kaede mice, ubiquitously expressed photoconvertible GFP¹²¹ undergoes peptide cleavage and alteration within the chromophore upon exposure to ultraviolet light, resulting in a shift from green to red fluorescence¹²⁰. This conversion is irreversible and thereby marks the treated cell until the photoconverted Kaede is degraded and replaced by new Kaede protein¹²⁰. Kaede mice have been used to decipher the migratory kinetics and behavior of T follicular helper cells under homeostasis or during memory responses¹²². In brief, the above-mentioned strategies coupled with *in vivo* imaging have revealed the dynamic behavior of T follicular helper cells during primary and secondary immune responses. During the primary response, T follicular helper cells are restricted within the germinal center, where they provide cognate help to developing B cells. However, during secondary immune responses (akin to a recurrent infection), the memory T follicular helper cells show unrestricted migration in and out of the germinal centers, a strategy that might be useful in eliciting rapid humoral responses¹²².

Although both KikGR and Kaede mice have been used for tracking the dynamics of various immune cell subsets like neutrophils¹²³, T cells^{124,125}, innate lymphoid cells¹²⁶, and dendritic cells¹²⁷ at a population level, great opportunities still persist in understanding the behavior of these immune cells at single-cell resolution.

Another strategy to irreversibly mark individual cells *in vivo* relies on the Cre/lox recombination system. Using this strategy, Livet and colleagues generated the Brainbow mouse strain where Cre-mediated recombination stochastically permutes multiple copies of the construct containing several different fluorescent proteins^{128,129}. This recombination strategy results in a mosaic of fluorescent colors leading to the detection of nearly 90 separate

colors that can be used to mark individual cells. This strategy was initially used to identify individual neurons within the CNS and for tracing neuronal circuitry¹²⁸. The generation of the Brainbow mouse paved the way for the development of several new approaches like LeGO¹³⁰, Confetti¹³¹, and Ubow¹³² to mark cells for tracing the fate of individual cells. Ubow mice have been used for fate mapping of Langerhans cells¹³² and follicular dendritic cells¹³³ within the skin and lymph node, respectively. Recently, a combination of fluorescent reporters and the Cre/lox system has been used exquisitely to highlight *in vivo* transfer of metastatic information between tumor cells¹³⁴. Use of the Cre/lox system provides a significant advantage over other strategies such as photoactivation or photoconversion by genetically/permanently labeling the cells, thereby enabling tracking throughout the lifetime of the cell. However, in most scenarios, the genetic labeling is targeted to a subset of cells and not to spatiotemporally selected individual cells.

Taken together, the recent advances in recombinant DNA technology like TALEN and the CRISPR/Cas9 genome editing system along with the availability of a multitude of reporter and knockout mouse strains promise a more colorful and dynamic future for the *in vivo* imaging of leukocytes.

Other methodologies

Although the use of transgenic fluorescent reporter strains for *in vivo* imaging has been gaining momentum, the transduction of immune cells using various fluorescent reporters still provides a valuable tool for understanding leukocyte function. These strategies have been used to delineate activation^{45,102,135}, calcium signaling¹³⁶, and apoptosis¹³⁷ within T cells *in vivo*. Similarly, several fluorescent probes are available to highlight cellular and molecular events crucial for leukocyte function *in vivo*. These fluorescent probes have been utilized to decipher various cellular processes: for example, NETosis^{88,138}, the generation of reactive oxygen species¹³⁹, and cell death^{67,140}. The use of fluorescent probes to mark DNA coupled with *in vivo* imaging has been crucial in deciphering the physiological role of DNA-NETs (DNA-neutrophil extracellular traps). Intravital imaging in various infectious model systems has uncovered the important role of DNA-NETs in ensnaring and killing pathogens^{88,141–143}. Although DNA-NETs are an essential arsenal of the innate immune response, release of DNA-NETs *in vivo* can trap circulating tumor cells and enhance metastasis¹⁴⁴.

Likewise, novel approaches have been developed to couple immune cell behavior with receptor-based signaling *in vivo* via administration of labeled antibodies. In this methodology, labeled Fab fragments are administered *in vivo* and the distribution of cell surface receptors and the behavior of labeled cells are recorded, segmented, and analyzed as flow cytometry plots¹⁴⁵. This methodology is of immense importance where the migratory dynamics of immune cells need to be analyzed in context with the receptor signals delivered to individual immune cells *in situ*.

Concluding remarks

In the past decade, *in vivo* two-photon imaging has provided researchers with unparalleled views of immune cell behavior under homeostasis or pathology. This has resulted in the identification of new molecular and cellular regulators of the immune system.

In our laboratory, *in vivo* imaging has been instrumental in highlighting the dynamic behavior of neutrophils⁶⁶, monocytes⁴², T cells^{146,147}, dendritic cells⁷³, type-2 innate lymphoid cells¹⁴⁸, and perivascular macrophages⁸⁶ under homeostasis or during various infections or inflammatory conditions.

Nonetheless, the use of *in vivo* imaging to understand cell-to-cell communication and the generation of complex interaction networks still requires extensive investigation. Intravital multiphoton imaging in most tissues is restricted to approximately 500 µm from the surface, which limits the visualization of processes within deeper tissues *in vivo*, therefore leaving certain aspects of the immune response in the dark. Various other strategies such as optical frequency domain imaging (OFDI)¹⁴⁹ and speckle-variance optical coherence tomography (svOCT)¹⁵⁰ have been developed for overcoming this limitation; however, these methodologies currently lack the necessary resolution for imaging immune cell behavior and are mostly useful for imaging tissue architecture and the vasculature^{151,152}.

We believe that the increased availability of fluorescent reporters for *in vivo* marking of cellular and molecular signaling, technical improvements in instrumentation (for example, adaptive optics), and better analytical capability will be crucial for understanding the complex behavior of leukocytes *in vivo*.

Competing interests

The authors declare that they have no competing interests.

Grant information

Wolfgang Weninger was supported by project grants 1066477, 1102398, 1104876 and 1106439, from the National Health and Medical Research Council Australia. Rohit Jain was supported by an Early Career Researcher grant from the Sydney Medical School, Sydney, Australia. Shweta Tikoo was supported by an Early Career Researcher grant from the Sydney Medical School and PdCCRS grant from Cure Cancer Australia.

References

- Butcher EC, Picker LJ: **Lymphocyte homing and homeostasis.** *Science.* 1996; 272(5258): 60–66.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Schmidt EE, MacDonald IC, Groom AC: **Interactions of leukocytes with vessel walls and with other blood cells, studied by high-resolution intravital videomicroscopy of spleen.** *Microvasc Res.* 1990; 40(1): 99–117.
[PubMed Abstract](#) | [Publisher Full Text](#)
- von Andrian UH: **Intravital microscopy of the peripheral lymph node microcirculation in mice.** *Microcirculation.* 1996; 3(3): 287–300.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Springer TA: **Traffic signals for lymphocyte recirculation and leukocyte emigration: the multistep paradigm.** *Cell.* 1994; 76(2): 301–14.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Stoll S, Delon J, Brotz TM, *et al.*: **Dynamic imaging of T cell-dendritic cell interactions in lymph nodes.** *Science.* 2002; 296(5574): 1873–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Bousso P, Bhakta NR, Lewis RS, *et al.*: **Dynamics of thymocyte-stromal cell interactions visualized by two-photon microscopy.** *Science.* 2002; 296(5574): 1876–80.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Miller MJ, Wei SH, Parker I, *et al.*: **Two-photon imaging of lymphocyte motility and antigen response in intact lymph node.** *Science.* 2002; 296(5574): 1869–73.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Miller MJ, Wei SH, Cahalan MD, *et al.*: **Autonomous T cell trafficking examined *in vivo* with intravital two-photon microscopy.** *Proc Natl Acad Sci U S A.* 2003; 100(5): 2604–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Mempel TR, Henrickson SE, Von Andrian UH: **T-cell priming by dendritic cells in lymph nodes occurs in three distinct phases.** *Nature.* 2004; 427(6970): 154–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Jain R, Weninger W: **Shedding light on cutaneous innate immune responses: the intravital microscopy approach.** *Immunol Cell Biol.* 2013; 91(4): 263–70.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Bousso P, Moreau HD: **Functional immunoimaging: the revolution continues.** *Nat Rev Immunol.* 2012; 12(12): 858–64.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Germain RN, Robey EA, Cahalan MD: **A decade of imaging cellular motility and interaction dynamics in the immune system.** *Science.* 2012; 336(6089): 1676–81.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Weninger W, Biro M, Jain R: **Leukocyte migration in the interstitial space of non-lymphoid organs.** *Nat Rev Immunol.* 2014; 14(4): 232–46.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Cahalan MD, Parker I, Wei SH, *et al.*: **Two-photon tissue imaging: seeing the immune system in a fresh light.** *Nat Rev Immunol.* 2002; 2(11): 872–80.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Conway JR, Carragher NO, Timpson P: **Developments in preclinical cancer imaging: innovating the discovery of therapeutics.** *Nat Rev Cancer.* 2014; 14(5): 314–28.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Li JL, Goh CC, Keeble JL, *et al.*: **Intravital multiphoton imaging of immune responses in the mouse ear skin.** *Nat Protoc.* 2012; 7(2): 221–34.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Liou HL, Myers JT, Barkauskas DS, *et al.*: **Intravital imaging of the mouse popliteal lymph node.** *J Vis Exp.* 2012; (60): pii: 3720.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Wong CH, Jenne CN, Lee WY, *et al.*: **Functional innervation of hepatic INKT cells is immunosuppressive following stroke.** *Science.* 2011; 334(6052): 101–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Egen JG, Rothfuchs AG, Feng CG, *et al.*: **Macrophage and T cell dynamics during the development and disintegration of mycobacterial granulomas.** *Immunity.* 2008; 28(2): 271–84.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Marques PE, Antunes MM, David BA, *et al.*: **Imaging liver biology *in vivo* using conventional confocal microscopy.** *Nat Protoc.* 2015; 10(2): 258–68.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Egen JG, Rothfuchs AG, Feng CG, *et al.*: **Intravital imaging reveals limited antigen presentation and T cell effector function in mycobacterial granulomas.** *Immunity.* 2011; 34(5): 807–19.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Ferrer M, Martin-Jaular L, Calvo M, *et al.*: **Intravital microscopy of the spleen: quantitative analysis of parasite motility and blood flow.** *J Vis Exp.* 2012; (59): pii: 3609.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Amon TI, Horton RM, Grigorova IL, *et al.*: **Visualization of splenic marginal zone B-cell shuttling and follicular B-cell egress.** *Nature.* 2013; 493(7434): 684–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Waite JC, Leiner I, Lauer P, *et al.*: **Dynamic imaging of the effector immune response to *listeria* infection *in vivo*.** *PLoS Pathog.* 2011; 7(3): e1001326.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Cavanagh LL, Bonasio R, Mazo IB, *et al.*: **Activation of bone marrow-resident memory T cells by circulating, antigen-bearing dendritic cells.** *Nat Immunol.*

- 2005; **6**(10): 1029–37.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
26. Jacquelin S, Licata F, Dorgham K, *et al.*: CX3CR1 reduces Ly6C^{high}-monocyte motility within and release from the bone marrow after chemotherapy in mice. *Blood*. 2013; **122**(5): 674–83.
[PubMed Abstract](#) | [Publisher Full Text](#)
 27. Evrard M, Chong SZ, Devi S, *et al.*: Visualization of bone marrow monocyte mobilization using Cx3cr1^{tdp}+Flt3L^{-/-} reporter mouse by multiphoton intravital microscopy. *J Leukoc Biol*. 2015; **97**(3): 611–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
 28. Devi S, Wang Y, Chew WK, *et al.*: Neutrophil mobilization via plerixafor-mediated CXCR4 inhibition arises from lung demargination and blockade of neutrophil homing to the bone marrow. *J Exp Med*. 2013; **210**(11): 2321–36.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 29. **F** Nourshargh S, Alon R: Leukocyte migration into inflamed tissues. *Immunity*. 2014; **41**(5): 694–707.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 30. Qi H, Kastenmüller W, Germain RN: Spatiotemporal basis of innate and adaptive immunity in secondary lymphoid tissue. *Annu Rev Cell Dev Biol*. 2014; **30**: 141–67.
[PubMed Abstract](#) | [Publisher Full Text](#)
 31. Mandl JN, Torabi-Parizi P, Germain RN: Visualization and dynamic analysis of host-pathogen interactions. *Curr Opin Immunol*. 2014; **29**: 8–15.
[PubMed Abstract](#) | [Publisher Full Text](#)
 32. **F** Grutzendler J, Kasthuri N, Gan WB: Long-term dendritic spine stability in the adult cortex. *Nature*. 2002; **420**(6917): 812–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 33. Yang G, Pan F, Parkhurst CN, *et al.*: Thinned-skull cranial window technique for long-term imaging of the cortex in live mice. *Nat Protoc*. 2010; **5**(2): 201–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 34. Mostany R, Portera-Cailliau C: A craniotomy surgery procedure for chronic brain imaging. *J Vis Exp*. 2008; **12**(2): pii: 680.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 35. Pai S, Danne KJ, Qin J, *et al.*: Visualizing leukocyte trafficking in the living brain with 2-photon intravital microscopy. *Front Cell Neurosci*. 2013; **6**: 67.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 36. **F** Davalos D, Grutzendler J, Yang G, *et al.*: ATP mediates rapid microglial response to local brain injury in vivo. *Nat Neurosci*. 2005; **8**(6): 752–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 37. Holtmaat A, Bonhoeffer T, Chow DK, *et al.*: Long-term, high-resolution imaging in the mouse neocortex through a chronic cranial window. *Nat Protoc*. 2009; **4**(8): 1128–44.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 38. Marker DF, Tremblay ME, Lu SM, *et al.*: A thin-skull window technique for chronic two-photon in vivo imaging of murine microglia in models of neuroinflammation. *J Vis Exp*. 2010; **43**(2): pii: 2059.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 39. **F** Nimmerjahn A, Kirchhoff F, Helmchen F: Resting microglial cells are highly dynamic surveillants of brain parenchyma in vivo. *Science*. 2005; **308**(5726): 1314–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 40. Davalos D, Ryu JK, Merlini M, *et al.*: Fibrinogen-induced perivascular microglial clustering is required for the development of axonal damage in neuroinflammation. *Nat Commun*. 2012; **3**: 1227.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 41. Michaud JP, Bellavance MA, Préfontaine P, *et al.*: Real-time in vivo imaging reveals the ability of monocytes to clear vascular amyloid beta. *Cell Rep*. 2013; **5**(3): 646–53.
[PubMed Abstract](#) | [Publisher Full Text](#)
 42. **F** Pai S, Qin J, Cavanagh L, *et al.*: Real-time imaging reveals the dynamics of leukocyte behaviour during experimental cerebral malaria pathogenesis. *PLoS Pathog*. 2014; **10**(7): e1004236.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 43. **F** Harris TH, Banigan EJ, Christian DA, *et al.*: Generalized Lévy walks and the role of chemokines in migration of effector CD8⁺ T cells. *Nature*. 2012; **486**(7404): 545–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 44. Kawakami N, Nägerl UV, Odoardi F, *et al.*: Live imaging of effector cell trafficking and autoantigen recognition within the unfolding autoimmune encephalomyelitis lesion. *J Exp Med*. 2005; **201**(11): 1805–14.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 45. **F** Pesic M, Bartholomäus I, Kyrtasous NI, *et al.*: 2-photon imaging of phagocyte-mediated T cell activation in the CNS. *J Clin Invest*. 2013; **123**(3): 1192–201.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 46. **F** Shaw TN, Stewart-Hutchinson PJ, Strangward P, *et al.*: Perivascular Arrest of CD8⁺ T Cells Is a Signature of Experimental Cerebral Malaria. *PLoS Pathog*. 2015; **11**(11): e1005210.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 47. Andermann ML, Gilfoy NB, Goldey GJ, *et al.*: Chronic cellular imaging of entire cortical columns in awake mice using microprisms. *Neuron*. 2013; **80**(4): 900–13.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 48. Barretto RP, Ko TH, Jung JC, *et al.*: Time-lapse imaging of disease progression in deep brain areas using fluorescence microendoscopy. *Nat Med*. 2011; **17**(2): 223–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 49. Hasegawa A, Hayashi K, Kishimoto H, *et al.*: Color-coded real-time cellular imaging of lung T-lymphocyte accumulation and focus formation in a mouse asthma model. *J Allergy Clin Immunol*. 2010; **125**(2): 461–468.e6.
[PubMed Abstract](#) | [Publisher Full Text](#)
 50. Tabuchi A, Mertens M, Kuppe H, *et al.*: Intravital microscopy of the murine pulmonary microcirculation. *J Appl Physiol (1985)*. 2008; **104**(2): 338–46.
[PubMed Abstract](#) | [Publisher Full Text](#)
 51. **F** Looney MR, Thornton EE, Sen D, *et al.*: Stabilized imaging of immune surveillance in the mouse lung. *Nat Methods*. 2011; **8**(1): 91–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 52. Presson RG Jr, Brown MB, Fisher AJ, *et al.*: Two-photon imaging within the murine thorax without respiratory and cardiac motion artifact. *Am J Pathol*. 2011; **179**(1): 75–82.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 53. Kreisel D, Nava RG, Li W, *et al.*: In vivo two-photon imaging reveals monocyte-dependent neutrophil extravasation during pulmonary inflammation. *Proc Natl Acad Sci U S A*. 2010; **107**(42): 18073–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 54. **F** Thornton EE, Looney MR, Bose O, *et al.*: Spatiotemporally separated antigen uptake by alveolar dendritic cells and airway presentation to T cells in the lung. *J Exp Med*. 2012; **209**(6): 1183–99.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 55. Fiole D, Deman P, Trescos Y, *et al.*: Two-photon intravital imaging of lungs during anthrax infection reveals long-lasting macrophage-dendritic cell contacts. *Infect Immun*. 2014; **82**(2): 864–72.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 56. **F** Lim K, Hyun YM, Lambert-Emo K, *et al.*: Neutrophil trails guide influenza-specific CD8⁺ T cells in the airways. *Science*. 2015; **349**(6252): aaa4352.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 57. Jung K, Kim P, Leuschner F, *et al.*: Endoscopic time-lapse imaging of immune cells in infarcted mouse hearts. *Circ Res*. 2013; **112**(6): 891–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 58. Li W, Nava RG, Bribiesco AC, *et al.*: Intravital 2-photon imaging of leukocyte trafficking in beating heart. *J Clin Invest*. 2012; **122**(7): 2499–508.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 59. Lee S, Vinegoni C, Feruglio PF, *et al.*: Real-time in vivo imaging of the beating mouse heart at microscopic resolution. *Nat Commun*. 2012; **3**: 1054.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 60. Vinegoni C, Aguirre AD, Lee S, *et al.*: Imaging the beating heart in the mouse using intravital microscopy techniques. *Nat Protoc*. 2015; **10**(11): 1802–19.
[PubMed Abstract](#) | [Publisher Full Text](#)
 61. Li W, Goldstein DR, Kreisel D: Intravital 2-photon imaging, leukocyte trafficking, and the beating heart. *Trends Cardiovasc Med*. 2013; **23**(8): 287–93.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 62. Megens RT, Reitsma S, Prinzen L, *et al.*: In vivo high-resolution structural imaging of large arteries in small rodents using two-photon laser scanning microscopy. *J Biomed Opt*. 2010; **15**(1): 11108.
[PubMed Abstract](#) | [Publisher Full Text](#)
 63. Chèvre R, González-Granado JM, Megens RT, *et al.*: High-resolution imaging of intravascular atherogenic inflammation in live mice. *Circ Res*. 2014; **114**(5): 770–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
 64. Drechsler M, Megens RT, van Zandvoort M, *et al.*: Hyperlipidemia-triggered neutrophilia promotes early atherosclerosis. *Circulation*. 2010; **122**(18): 1837–45.
[PubMed Abstract](#) | [Publisher Full Text](#)
 65. McArdle S, Chodaczek G, Ray N, *et al.*: Intravital live cell triggered imaging system reveals monocyte patrolling and macrophage migration in atherosclerotic arteries. *J Biomed Opt*. 2015; **20**(2): 26005.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 66. Ng LG, Qin JS, Roediger B, *et al.*: Visualizing the neutrophil response to sterile tissue injury in mouse dermis reveals a three-phase cascade of events. *J Invest Dermatol*. 2011; **131**(10): 2058–68.
[PubMed Abstract](#) | [Publisher Full Text](#)
 67. **F** Lämmermann T, Afonso PV, Angermann BR, *et al.*: Neutrophil swarms require LTB4 and integrins at sites of cell death in vivo. *Nature*. 2013; **498**(7454): 371–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 68. Goh CC, Li JL, Devi S, *et al.*: Real-time imaging of dendritic cell responses to sterile tissue injury. *J Invest Dermatol*. 2015; **135**(4): 1181–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
 69. Lämmermann T: In the eye of the neutrophil swarm-navigation signals that bring neutrophils together in inflamed and infected tissues. *J Leukoc Biol*. 2015; pii: jlb.1MR0915-403.
[PubMed Abstract](#) | [Publisher Full Text](#)

70. Li JL, Ng LG: **Peeking into the secret life of neutrophils.** *Immunol Res.* 2012; 53(1–3): 168–81.
[PubMed Abstract](#) | [Publisher Full Text](#)
71. Coombes JL, Robey EA: **Dynamic imaging of host-pathogen interactions *in vivo*.** *Nat Rev Immunol.* 2010; 10(5): 353–64.
[PubMed Abstract](#) | [Publisher Full Text](#)
72. Garside P, Brewer J: ***In vivo* imaging of infection immunology—41's!** *Semin Immunopathol.* 2010; 32(3): 289–96.
[PubMed Abstract](#) | [Publisher Full Text](#)
73. **F** Ng LG, Hsu A, Mandell MA, *et al.*: **Migratory dermal dendritic cells act as rapid sensors of protozoan parasites.** *PLoS Pathog.* 2008; 4(11): e1000222.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
74. **F** Peters NC, Egen JG, Secundino N, *et al.*: ***In vivo* imaging reveals an essential role for neutrophils in leishmaniasis transmitted by sand flies.** *Science.* 2008; 321(5891): 970–4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
75. Filipe-Santos O, Pescher P, Beart B, *et al.*: **A dynamic map of antigen recognition by CD4 T cells at the site of *Leishmania major* infection.** *Cell Host Microbe.* 2009; 6(1): 23–33.
[PubMed Abstract](#) | [Publisher Full Text](#)
76. **F** Müller AJ, Filipe-Santos O, Eberl G, *et al.*: **CD4⁺ T cells rely on a cytokine gradient to control intracellular pathogens beyond sites of antigen presentation.** *Immunity.* 2012; 37(1): 147–57.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
77. Olekhovitch R, Ryffel B, Müller AJ, *et al.*: **Collective nitric oxide production provides tissue-wide immunity during *Leishmania* infection.** *J Clin Invest.* 2014; 124(4): 1711–22.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
78. Beattie L, Peltan A, Maroof A, *et al.*: **Dynamic imaging of experimental *Leishmania donovani*-induced hepatic granulomas detects Kupffer cell-restricted antigen presentation to antigen-specific CD8 T cells.** *PLoS Pathog.* 2010; 6(3): e1000805.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
79. Moore JW, Beattie L, Dalton JE, *et al.*: **B cell: T cell interactions occur within hepatic granulomas during experimental visceral leishmaniasis.** *PLoS One.* 2012; 7(3): e34143.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
80. Beattie L, d'El-Rei Hermida M, Moore JW, *et al.*: **A transcriptomic network identified in uninfected macrophages responding to inflammation controls intracellular pathogen survival.** *Cell Host Microbe.* 2013; 14(3): 357–68.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
81. **F** Chtanova T, Schaeffer M, Han SJ, *et al.*: **Dynamics of neutrophil migration in lymph nodes during infection.** *Immunity.* 2008; 29(3): 487–96.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
82. Schaeffer M, Han SJ, Chtanova T, *et al.*: **Dynamic imaging of T cell-parasite interactions in the brains of mice chronically infected with *Toxoplasma gondii*.** *J Immunol.* 2009; 182(10): 6379–93.
[PubMed Abstract](#) | [Publisher Full Text](#)
83. **F** Wilson EH, Harris TH, Mrass P, *et al.*: **Behavior of parasite-specific effector CD8⁺ T cells in the brain and visualization of a kinesis-associated system of reticular fibers.** *Immunity.* 2009; 30(2): 300–11.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
84. Lee WY, Moriarty TJ, Wong CH, *et al.*: **An intravascular immune response to *Borrelia burgdorferi* involves Kupffer cells and iNKT cells.** *Nat Immunol.* 2010; 11(4): 295–302.
[PubMed Abstract](#) | [Publisher Full Text](#)
85. Lee WY, Sanz MJ, Wong CH, *et al.*: **Invariant natural killer T cells act as an extravascular cytotoxic barrier for joint-invasive Lyme *Borrelia*.** *Proc Natl Acad Sci U S A.* 2014; 111(38): 13936–41.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
86. **F** Abtin A, Jain R, Mitchell AJ, *et al.*: **Perivascular macrophages mediate neutrophil recruitment during bacterial skin infection.** *Nat Immunol.* 2014; 15(1): 45–53.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
87. Liese J, Rooijackers SH, van Strijp JA, *et al.*: **Intravital two-photon microscopy of host-pathogen interactions in a mouse model of *Staphylococcus aureus* skin abscess formation.** *Cell Microbiol.* 2013; 15(6): 891–909.
[PubMed Abstract](#) | [Publisher Full Text](#)
88. **F** Yipp BG, Petri B, Salina D, *et al.*: **Infection-induced NETosis is a dynamic process involving neutrophil multitasking *in vivo*.** *Nat Med.* 2012; 18(9): 1386–93.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
89. **F** Amino R, Thiberge S, Martin B, *et al.*: **Quantitative imaging of *Plasmodium* transmission from mosquito to mammal.** *Nat Med.* 2006; 12(2): 220–4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
90. **F** Frevet U, Engelmann S, Zougbedé S, *et al.*: **Intravital observation of *Plasmodium berghei* sporozoite infection of the liver.** *PLoS Biol.* 2005; 3(6): e192.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
91. Nacer A, Movila A, Sohet F, *et al.*: **Experimental cerebral malaria pathogenesis—hemodynamics at the blood brain barrier.** *PLoS Pathog.* 2014; 10(12): e1004528.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
92. Cockburn IA, Amino R, Kelemen RK, *et al.*: ***In vivo* imaging of CD8⁺ T cell-mediated elimination of malaria liver stages.** *Proc Natl Acad Sci U S A.* 2013; 110(22): 9090–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
93. **F** Radtke AJ, Kastenmüller W, Espinosa DA, *et al.*: **Lymph-node resident CD8^α⁺ dendritic cells capture antigens from migratory malaria sporozoites and induce CD8⁺ T cell responses.** *PLoS Pathog.* 2015; 11(2): e1004637.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
94. Peters NC, Kimblin N, Secundino N, *et al.*: **Vector transmission of *leishmania* abrogates vaccine-induced protective immunity.** *PLoS Pathog.* 2009; 5(6): e1000484.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
95. Ritter U, Frischknecht F, van Zandbergen G: **Are neutrophils important host cells for *Leishmania* parasites?** *Trends Parasitol.* 2009; 25(11): 505–10.
[PubMed Abstract](#) | [Publisher Full Text](#)
96. **F** van Zandbergen G, Klinger M, Mueller A, *et al.*: **Cutting edge: neutrophil granulocyte serves as a vector for *Leishmania* entry into macrophages.** *J Immunol.* 2004; 173(11): 6521–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
97. Hurrell BP, Regli IB, Tacchini-Cottier F: **Different *Leishmania* Species Drive Distinct Neutrophil Functions.** *Trends Parasitol.* 2016; 32(5): 392–401.
[PubMed Abstract](#) | [Publisher Full Text](#)
98. Boissonnas A, Fetter L, Zeelenberg IS, *et al.*: ***In vivo* imaging of cytotoxic T cell infiltration and elimination of a solid tumor.** *J Exp Med.* 2007; 204(2): 345–56.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
99. Mrass P, Kinjyo I, Ng LG, *et al.*: **CD44 mediates successful interstitial navigation by killer T cells and enables efficient antitumor immunity.** *Immunity.* 2008; 29(6): 971–85.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
100. Mrass P, Takano H, Ng LG, *et al.*: **Random migration precedes stable target cell interactions of tumor-infiltrating T cells.** *J Exp Med.* 2006; 203(12): 2749–61.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
101. Beart B, Lemaître F, Celli S, *et al.*: **Two-photon imaging of intratumoral CD8⁺ T cell cytotoxic activity during adoptive T cell therapy in mice.** *J Clin Invest.* 2008; 118(4): 1390–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
102. **F** Marangoni F, Murooka TT, Manzo T, *et al.*: **The transcription factor NFAT exhibits signal memory during serial T cell interactions with antigen-presenting cells.** *Immunity.* 2013; 38(2): 237–49.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
103. Wyckoff J, Wang W, Lin EY, *et al.*: **A paracrine loop between tumor cells and macrophages is required for tumor cell migration in mammary tumors.** *Cancer Res.* 2004; 64(19): 7022–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
104. Wyckoff JB, Wang Y, Lin EY, *et al.*: **Direct visualization of macrophage-assisted tumor cell intravasation in mammary tumors.** *Cancer Res.* 2007; 67(6): 2649–56.
[PubMed Abstract](#) | [Publisher Full Text](#)
105. **F** Siffrin V, Radbruch H, Glumm R, *et al.*: ***In vivo* imaging of partially reversible th17 cell-induced neuronal dysfunction in the course of encephalomyelitis.** *Immunity.* 2010; 33(3): 424–36.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
106. **F** Kong L, Tang J, Little JP, *et al.*: **Continuous volumetric imaging via an optical phase-locked ultrasound lens.** *Nat Methods.* 2015; 12(8): 759–62.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
107. Chu SW, Tai SP, Ho CL, *et al.*: **High-resolution simultaneous three-photon fluorescence and third-harmonic-generation microscopy.** *Microsc Res Tech.* 2005; 66(4): 193–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
108. Cox G, Kable E, Jones A, *et al.*: **3-dimensional imaging of collagen using second harmonic generation.** *J Struct Biol.* 2003; 141(1): 53–62.
[PubMed Abstract](#) | [Publisher Full Text](#)
109. Franken PA, Hill AE, Peters CW, *et al.*: **Generation of Optical Harmonics.** *Phys Rev Lett.* 1961; 7(4): 118–9.
[Publisher Full Text](#)
110. Friedl P, Wolf K, von Andrian UH, *et al.*: **Biological second and third harmonic generation microscopy.** *Curr Protoc Cell Biol.* In: Bonifacio JS, Dasso M, Harford JB, Lippincott-Schwartz J, Yamada KM, editors. Hoboken, NJ USA: John Wiley & Sons, Inc; 2007; Chapter 4: Unit 4.15.
[PubMed Abstract](#) | [Publisher Full Text](#)
111. Rochelleau JV, Head WS, Piston DW: **Quantitative NAD(P)H/flavoprotein autofluorescence imaging reveals metabolic mechanisms of pancreatic islet pyruvate response.** *J Biol Chem.* 2004; 279(30): 31780–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
112. Blacker TS, Mann ZF, Gale JE, *et al.*: **Separating NADH and NADPH fluorescence in live cells and tissues using FLIM.** *Nat Commun.* 2014; 5: 3936.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
113. Abe T, Fujimori T: **Reporter mouse lines for fluorescence imaging.** *Dev Growth Differ.* 2013; 55(4): 390–405.
[PubMed Abstract](#) | [Publisher Full Text](#)
114. Pittet MJ, Weissleder R: **Intravital imaging.** *Cell.* 2011; 147(5): 983–91.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

115. **F** Patterson GH, Lippincott-Schwartz J: **A photoactivatable GFP for selective photolabeling of proteins and cells.** *Science*. 2002; 297(5588): 1873–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
116. **F** Vitorica GD, Schwickert TA, Fooksman DR, *et al.*: **Germinal center dynamics revealed by multiphoton microscopy with a photoactivatable fluorescent reporter.** *Cell*. 2010; 143(4): 592–605.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
117. **F** Shulman Z, Gitlin AD, Targ S, *et al.*: **T follicular helper cell dynamics in germinal centers.** *Science*. 2013; 341(6146): 673–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
118. Kurotaki Y, Hatta K, Nakao K, *et al.*: **Blastocyst axis is specified independently of early cell lineage but aligns with the ZP shape.** *Science*. 2007; 316(5825): 719–23.
[PubMed Abstract](#) | [Publisher Full Text](#)
119. Nowotshin S, Hadjantonakis AK: **Use of KikGR a photoconvertible green-to-red fluorescent protein for cell labeling and lineage analysis in ES cells and mouse embryos.** *BMC Dev Biol*. 2009; 9: 49.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
120. Tomura M, Yoshida N, Tanaka J, *et al.*: **Monitoring cellular movement in vivo with photoconvertible fluorescence protein “Kaede” transgenic mice.** *Proc Natl Acad Sci U S A*. 2008; 105(31): 10871–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
121. **F** Ando R, Hama H, Yamamoto-Hino M, *et al.*: **An optical marker based on the UV-induced green-to-red photoconversion of a fluorescent protein.** *Proc Natl Acad Sci U S A*. 2002; 99(20): 12651–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
122. Suan D, Nguyen A, Moran I, *et al.*: **T follicular helper cells have distinct modes of migration and molecular signatures in naive and memory immune responses.** *Immunity*. 2015; 42(4): 704–18.
[PubMed Abstract](#) | [Publisher Full Text](#)
123. **F** Hampton HR, Bailey J, Tomura M, *et al.*: **Microbe-dependent lymphatic migration of neutrophils modulates lymphocyte proliferation in lymph nodes.** *Nat Commun*. 2015; 6: 7139.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
124. **F** Tomura M, Honda T, Tanizaki H, *et al.*: **Activated regulatory T cells are the major T cell type emigrating from the skin during a cutaneous immune response in mice.** *J Clin Invest*. 2010; 120(3): 883–93.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
125. Bromley SK, Yan S, Tomura M, *et al.*: **Recirculating memory T cells are a unique subset of CD4⁺ T cells with a distinct phenotype and migratory pattern.** *J Immunol*. 2013; 190(3): 970–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
126. Mackley EC, Houston S, Marriott CL, *et al.*: **CCR7-dependent trafficking of ROR γ ILCs creates a unique microenvironment within mucosal draining lymph nodes.** *Nat Commun*. 2015; 6: 5862.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
127. Tomura M, Hata A, Matsuoka S, *et al.*: **Tracking and quantification of dendritic cell migration and antigen trafficking between the skin and lymph nodes.** *Sci Rep*. 2014; 4: 6030.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
128. **F** Livet J, Weissman TA, Kang H, *et al.*: **Transgenic strategies for combinatorial expression of fluorescent proteins in the nervous system.** *Nature*. 2007; 450(7166): 56–62.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
129. Weissman TA, Sanes JR, Lichtman JW, *et al.*: **Generating and imaging multicolor Brainbow mice.** *Cold Spring Harb Protoc*. 2011; 2011(7): 763–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
130. Weber K, Thomaschewski M, Warlich M, *et al.*: **RGB marking facilitates multicolor clonal cell tracking.** *Nat Med*. 2011; 17(4): 504–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
131. **F** Snippert HJ, van der Flier LG, Sato T, *et al.*: **Intestinal crypt homeostasis results from neutral competition between symmetrically dividing Lgr5 stem cells.** *Cell*. 2010; 143(1): 134–44.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
132. Ghigo C, Mondor I, Jorquera A, *et al.*: **Multicolor fate mapping of Langerhans cell homeostasis.** *J Exp Med*. 2013; 210(9): 1657–64.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
133. Jarjour M, Jorquera A, Mondor I, *et al.*: **Fate mapping reveals origin and dynamics of lymph node follicular dendritic cells.** *J Exp Med*. 2014; 211(6): 1109–22.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
134. **F** Zomer A, Maynard C, Verweij FJ, *et al.*: **In Vivo imaging reveals extracellular vesicle-mediated phenocopying of metastatic behavior.** *Cell*. 2015; 161(5): 1046–57.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
135. **F** Lodygin D, Odoardi F, Schläger C, *et al.*: **A combination of fluorescent NFAT and H2B sensors uncovers dynamics of T cell activation in real time during CNS autoimmunity.** *Nat Med*. 2013; 19(6): 784–90.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
136. Mues M, Bartholomäus I, Thestrup T, *et al.*: **Real-time in vivo analysis of T cell activation in the central nervous system using a genetically encoded calcium indicator.** *Nat Med*. 2013; 19(6): 778–83.
[PubMed Abstract](#) | [Publisher Full Text](#)
137. Garrod KR, Moreau HD, Garcia Z, *et al.*: **Dissecting T cell contraction in vivo using a genetically encoded reporter of apoptosis.** *Cell Rep*. 2012; 2(5): 1438–47.
[PubMed Abstract](#) | [Publisher Full Text](#)
138. Kolaczowska E, Jenne CN, Surewaard BG, *et al.*: **Molecular mechanisms of NET formation and degradation revealed by intravital imaging in the liver vasculature.** *Nat Commun*. 2015; 6: 6673.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
139. Devi S, Li A, Westhorpe CL, *et al.*: **Multiphoton imaging reveals a new leukocyte recruitment paradigm in the glomerulus.** *Nat Med*. 2013; 19(1): 107–12.
[PubMed Abstract](#) | [Publisher Full Text](#)
140. **F** McDonald B, Pittman K, Menezes GB, *et al.*: **Intravascular danger signals guide neutrophils to sites of sterile inflammation.** *Science*. 2010; 330(6002): 362–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
141. Yipp BG, Kubes P: **NETosis: how vital is it?** *Blood*. 2013; 122(16): 2784–94.
[PubMed Abstract](#) | [Publisher Full Text](#)
142. Kolaczowska E, Kubes P: **Neutrophil recruitment and function in health and inflammation.** *Nat Rev Immunol*. 2013; 13(3): 159–75.
[PubMed Abstract](#) | [Publisher Full Text](#)
143. Jain R, Mitchell AJ, Tay SS, *et al.*: **Neutrophils.** *Immunology of the Skin*. Edited by: Kabashima K. Tokyo: Springer Japan; 2016; 147–167.
[Publisher Full Text](#)
144. **F** Cools-Lartigue J, Spicer J, McDonald B, *et al.*: **Neutrophil extracellular traps sequester circulating tumor cells and promote metastasis.** *J Clin Invest*. 2013; 123(8): 3446–3458.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
145. Moreau HD, Lemaître F, Terriac E, *et al.*: **Dynamic in situ cytometry uncovers T cell receptor signaling during immunological synapses and kinapses in vivo.** *Immunity*. 2012; 37(2): 351–63.
[PubMed Abstract](#) | [Publisher Full Text](#)
146. Prakash MD, Munoz MA, Jain R, *et al.*: **Granzyme B promotes cytotoxic lymphocyte transmigration via basement membrane remodeling.** *Immunity*. 2014; 41(6): 960–72.
[PubMed Abstract](#) | [Publisher Full Text](#)
147. **F** Shulman Z, Cohen SJ, Roediger B, *et al.*: **Transendothelial migration of lymphocytes mediated by intraendothelial vesicle stores rather than by extracellular chemokine depots.** *Nat Immunol*. 2011; 13(1): 67–76.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
148. **F** Roediger B, Kyle R, Yip KH, *et al.*: **Cutaneous immunosurveillance and regulation of inflammation by group 2 innate lymphoid cells.** *Nat Immunol*. 2013; 14(6): 564–73.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
149. Vakoc BJ, Lanning RM, Tyrrell JA, *et al.*: **Three-dimensional microscopy of the tumor microenvironment in vivo using optical frequency domain imaging.** *Nat Med*. 2009; 15(10): 1219–23.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
150. Conroy L, DaCosta RS, Vitkin IA: **Quantifying tissue microvasculature with speckle variance optical coherence tomography.** *Opt Lett*. 2012; 37(15): 3180–2.
[PubMed Abstract](#) | [Publisher Full Text](#)
151. Maeda A, Kulbatski I, DaCosta RS: **Emerging Applications for Optically Enabled Intravital Microscopic Imaging in Radiobiology.** *Mol Imaging*. 2015; 14: 452–74.
[PubMed Abstract](#)
152. Vakoc BJ, Fukumura D, Jain RK, *et al.*: **Cancer imaging by optical coherence tomography: preclinical progress and clinical potential.** *Nat Rev Cancer*. 2012; 12(5): 363–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. **Tri Phan**
Garvan Institute of Medical research, Darlinghurst, Australia
Competing Interests: No competing interests were disclosed.
2. **Peter Friedl**
¹ Department of Cell Biology, Radboud University Medical Centre, 6525GA Nijmegen, The Netherlands
² David H. Koch Center for Applied Research of Genitourinary Cancers, University of Texas MD Anderson Cancer Center, Houston, Texas, 77030, USA
Competing Interests: No competing interests were disclosed.
3. **Tim Lammermann**
Max Planck Institute of Immunobiology and Epigenetics, Freiburg, Germany
Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research