REVIEW

Chronic urticaria: a focus on pathogenesis [version 1; peer review: 3 approved]

Riccardo Asero ¹, Alberto Tedeschi², Angelo Valerio Marzano³, Massimo Cugno⁴

¹Ambulatorio di Allergologia, Clinica San Carlo, Paderno Dugnano (Milano), Italy

²Ambulatorio di Allergologia e Immunologia Clinica, Ospedale Fatebenefratelli e Oftalmico, Milano, Italy

³Unità di Dermatologia, Dipartimento di Fisiopatologia Medico Chirurgica e dei Trapianti, Università degli Studi di Milano, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milano, Italy

⁴Medicina Interna, Dipartimento di Fisiopatologia Medico Chirurgica e dei Trapianti, Università degli Studi di Milano, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milano, Italy

V1 First published: 11 Jul 2017, 6(F1000 Faculty Rev):1095
<https://doi.org/10.12688/f1000research.11546.1>

Latest published: 11 Jul 2017, 6(F1000 Faculty Rev):1095
<https://doi.org/10.12688/f1000research.11546.1>

Abstract

Chronic urticaria is a spontaneous or inducible group of diseases characterized by the occurrence of wheals (and, in about half of cases, angioedema) for more than 6 weeks. These are rather frequent conditions that may severely affect patients' quality of life and sometimes represent a challenge for doctors as well. The causes of chronic urticaria are still poorly defined, although there is growing evidence that different biologic systems including immunity, inflammation, and coagulation may take part in the pathomechanism eventually leading to mast cell and basophil degranulation and hence to wheal formation. This review will discuss the main findings that are (slowly) shedding light on the pathogenesis of this disorder.

Keywords

Chronic Urticaria, wheals, autologous serum skin test, Omalizumab

Open Peer Review

Approval Status

	1	2	3
version 1 11 Jul 2017

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Simon Francis Thomsen**, Bispebjerg Hospital, Copenhagen, Denmark
University of Copenhagen, Copenhagen, Denmark
2. **Young-Min Ye**, Ajou University School of Medicine, Suwon, South Korea
3. **Malcolm W Greaves**, St John's Institute of Dermatology, St Thomas' Hospital, London, UK

Any comments on the article can be found at the

.....
end of the article.

Corresponding author: Riccardo Asero (r.asero@libero.it)

Competing interests: The authors declare that they have no competing interests.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2017 Asero R *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Asero R, Tedeschi A, Marzano AV and Cugno M. **Chronic urticaria: a focus on pathogenesis [version 1; peer review: 3 approved]** F1000Research 2017, 6(F1000 Faculty Rev):1095 <https://doi.org/10.12688/f1000research.11546.1>

First published: 11 Jul 2017, 6(F1000 Faculty Rev):1095 <https://doi.org/10.12688/f1000research.11546.1>

Introduction

Chronic urticaria, defined as the recurrent occurrence of wheals with or without angioedema for longer than 6 weeks, can be classified into spontaneous or inducible based on the absence or presence of identifiable physical stimuli able to elicit the skin lesions¹. While there is little doubt that the release of histamine by mast cells and basophils represents the final stage in the pathomechanisms involved in this group of diseases, there is still some uncertainty about the factors activating these cells and eventually inducing their degranulation. Several lines of evidence indicate that different biologic systems like immunity, inflammation, and coagulation may contribute to a common mechanism leading to wheal generation. This short commentary will summarize the current knowledge about the mechanisms theoretically leading to histamine release in chronic urticaria.

Established facts

The bulk of studies carried out during the last 30 years have shown that different biologic systems (i.e. autoimmunity, auto-allergy, inflammation, and coagulation) may concur to produce the wheal and flare lesions as well as the angioedema that characterize chronic urticaria. For clarity, these biologic systems, which are largely linked with each other, along with the main cell lines involved in either their activation or their behavior as effector cells will be considered singularly in the following subsections.

Autoreactivity/autoimmunity

More than 30 years ago, Grattan and co-workers demonstrated that the intradermal injection of autologous serum induces a wheal-and-flare reaction in a large proportion of patients with chronic urticaria²; this led them to conclude that chronic urticaria is characterized by circulating histamine-releasing factors. Since then, a positive autologous serum skin test (ASST) has been considered as a marker of “autoreactivity”. This *in vivo* test is quite easy to perform and seems sufficiently specific for chronic urticaria, as normal subjects do not show any skin reactivity³, although it may score positive in inducible urticarias such as cold urticaria⁴ as well as in some other “allied” conditions, such as the multiple drug allergy syndrome and multiple hypersensitivity to non-steroidal anti-inflammatory drugs^{5,6}. However, the ASST scores positive in only about one half of chronic urticaria patients^{7,8}, leaving a large proportion of patients who do not show any autoreactivity despite their maybe severe and ongoing disease.

Following the first observation on skin reactivity to autologous serum that remains a cornerstone in urticaria research, the second important step in the definition of the pathogenic mechanism was the demonstration of histamine-releasing autoantibodies about 20 years ago, suggesting an autoimmune origin. This was based on the detection of circulating and functionally active IgG autoantibodies directed against either the high-affinity IgE receptor (FcεRI) present on both mast cells and basophils (in most cases) or membrane-bound IgE (in a minority of patients)^{9–12}.

Although this was a fascinating explanation for the ongoing histamine release, several subsequent studies found that this mechanism is in fact viable in only a minority of patients^{8,13}. Furthermore, both types of autoantibodies have been detected with similar frequency in chronic urticaria patients and controls without chronic

urticaria¹³. Nonetheless, in more recent times, the autoimmune pathogenesis hypothesis has received new, indirect but strong support from the detection of circulating autoreactive CD4⁺ T cells that proliferate in response to FcεRI in >50% of patients with chronic urticaria examined¹⁴.

Basophils: their pathogenic role and use in chronic urticaria diagnosis

Although there is little doubt that the mast cell is the main final effector cell in chronic urticaria, basophils and their role in the disease have received much attention during the last decade. It is well known that active chronic urticaria is characterized by basopenia due to the elective recruitment of circulating basophils into the skin lesions¹⁵ and that basopenia is directly related to disease severity. Recent studies found that chronic urticaria blood basophils show a reduced surface expression of CCR2 (the chemoattractant receptor-homologous molecule expressed on TH2 cells), the receptor for prostaglandin D2 (PGD2). It is therefore possible that decreased CCR2 levels on basophils in patients with chronic urticaria are due to *in vivo* PGD2-mediated activation, suggesting the engagement of CCR2 in patients with chronic urticaria¹⁶. Furthermore, basophils from chronic urticaria patients show different phenotypes (either responder or non-responder, based on the degranulation response to polyclonal goat anti-human IgE) that seem independent of the presence of IgG autoantibodies to IgE or FcεRI and seem stable during active disease^{13,17}, and their IgE receptor-mediated degranulation is enhanced during disease remission^{13,17}.

The finding that at least a proportion of chronic urticaria patients show autoimmune phenomena targeting mast cells and basophils has led to the investigation of the diagnostic accuracy of basophil-based *in vitro* tests aiming to diagnose autoimmunity. The first test used was the time-consuming and cumbersome basophil histamine release assay (BHRA), first introduced in the 1960s and subsequently marketed as a more convenient testing kit¹⁸; on the other hand, the observation that chronic urticaria sera were able to induce the surface expression of basophil activation markers such as CD63¹⁹ led to the investigation of the diagnostic usefulness of the basophil activation test (BAT) measuring CD63 and CD203c surface expression by flow cytometry^{20,21}. However, in view of the fact that no more than 50% of chronic urticaria patients have an autoimmune disease and of the intrinsic variability of basophil releasability, the clinical usefulness of these tests has been questioned²² and they have not reached the status of routine investigation for this disease.

Auto-allergy

Another important step was the discovery of specific IgE for thyroid peroxidase (TPO) for dsDNA and ssDNA, which has recently provided a different mechanism potentially able to promote the survival, proliferation, and activation of mast cells²³. This pathogenic mechanism could be classified as “auto-allergic” and autoimmune at the same time. However, the proportion of patients in whom such IgE autoantibodies can be detected is limited, and TPO-specific IgE autoantibodies are generally associated with the more common TPO IgG autoantibodies that characterize patients with Hashimoto’s thyroiditis which are found in a minority of patients. However, research in this field is very active, and it cannot be

excluded that other “auto-allergens” will be detected in larger proportions of patients with chronic urticaria in the near future²⁴.

Inflammation

It is increasingly clear that chronic urticaria is characterized by a systemic pro-inflammatory state. The studies showing increased levels of different single markers of inflammation (e.g. matrix metalloproteinase and others) that have appeared in the medical literature over the years will not be reviewed in detail here. Recent studies showed the co-existence of chronic urticaria and metabolic syndrome (MS) in a Korean population²⁵; patients with both chronic urticaria and MS were older, had a more severe disease, had higher levels of inflammation markers (i.e. eosinophil cationic protein [ECP], tumor necrosis factor [TNF]-alpha, and complement), and, interestingly, more frequently scored negative on the ASST than those without MS. In another Asian-based population study, chronic urticaria was more frequent among subjects with a prior diagnosis of hyperlipidemia²⁶. Recently, chronic urticaria patients were found to show increased levels of a series of adipokines (lipocalin-2, TNF-alpha, IL-6, and IL-10) but lower levels of adiponectin²⁷. The findings with IL-6 are in keeping with a previous study²⁸ showing an association between IL-6 and disease severity.

Coagulation

The observation that the autologous plasma skin test (APST) may score positive in some ASST-negative patients²⁹ has prompted investigation of the coagulation system in patients with chronic urticaria. Specific studies have shown that the coagulation cascade is activated in chronic urticaria and involves the extrinsic pathway first and the intrinsic pathway second^{29–32}. The process seems to be triggered by the hyper-expression of tissue factor by activated eosinophils³³ and parallels disease activity. IgG autoantibodies to the low-affinity IgE receptor FcεRII, which is present on the membrane of eosinophils, have been detected in quite a large proportion of patients (about 65%) with chronic urticaria³⁴ and might be involved in the activation of such cells with subsequent release of major basic protein and other mediators causing mast cell degranulation. The activation of the coagulation cascade might potentially play a relevant pathogenic role if one considers that thrombin can markedly increase the vascular permeability and is a potent inducer of mast cell degranulation in experimental models. The activation of the coagulation cascade occurs in other skin disorders characterized by an increase of vascular permeability, such as angioedema due to C1-inhibitor deficiency and bullous pemphigoid^{35,36}, but also in a wide spectrum of systemic diseases, such as disseminated intravascular coagulation³⁷, deep venous thrombosis³⁸, and endotoxemia³⁹, all prothrombotic conditions that are not characterized by urticaria or edema. However, the activation of coagulation should not necessarily be considered a non-specific phenomenon but rather a common intermediate step occurring in the pathophysiology of different diseases. Thus, the involvement of coagulation is another piece of truth unable to explain the whole story.

Other mechanisms

Quite recently, we were able to show that sera from patients with chronic urticaria induce significant activation of mast cells lacking the high-affinity IgE receptor, irrespective of the autoreactivity status or of the presence/absence of circulating autoantibodies⁴⁰. This fact suggests a pathomechanism that bypasses the IgE receptor mechanism. In subsequent studies, we showed that

low-molecular-weight circulating factors (molecular weight about 30 kDa) may be involved in the activation of mast cells in chronic urticaria patients⁴¹, thus confirming very old data by Grattan and co-workers⁴².

Using the response to omalizumab as a means to distinguish different chronic urticaria phenotypes

Omalizumab was recently licensed for the treatment of chronic urticaria. All three phase III studies as well as the real-life studies carried out in large numbers of patients have demonstrated that this humanized monoclonal IgG antibody to IgE is highly effective in a large proportion of patients with chronic urticaria who don't show any clinical response to second-generation antihistamines given at doses even higher than licensed ones. The pattern of response to this drug has indirectly led to the categorization of patients with severe disease into “fast responders” (subjects who show a complete response as short as 3–7 days after the first administration of the drug; these represent about 60–70% of patients), “slow responders” (subjects who need three to five monthly administrations before showing a significant clinical response; about 10–20% of cases), and “non-responders” (about 10–20% of cases)⁴³. It is easy to speculate that different pathogenic mechanisms might underlie the different patterns of response. Several potential markers of response to omalizumab treatment have been investigated so far. In one study looking at the potential use of a CD203c assay (the so-called BAT) as a biomarker of responsiveness to omalizumab, it was found that the lack of basophil CD203 upregulating activity was associated with a higher likelihood of response to the drug⁴⁴. This observation seems to agree with the results of another study finding that a positive response to omalizumab was associated with a negative histamine release assay⁴⁵. On the other hand, Gericke and co-workers found that a positive BHRA, which was associated with a positive ASST, was frequently observed in “slow responders” to omalizumab⁴³. Another interesting observation is that omalizumab has been found to be effective not only in the spontaneous form but also in several inducible forms of chronic urticaria^{46,47}. This suggests that the drug interferes with a common pathogenic mechanism underlying a proportion of both spontaneous and inducible urticarias. Notably, the fact that the symptomatic cold urticaria, a frequent chronic inducible urticaria, can be passively transmitted to normal subjects via the serum of affected individuals has been known for many years. Since the mechanism of action of omalizumab in urticaria is all but elucidated, the next few years will probably shed some light on these questions.

Conclusions

The pathogenesis of chronic urticaria is probably characterized by a multiplicity of mechanisms, including autoimmunity, autoallergy, and coagulation, each of which may carry different weight in each patient. Much research is in progress, and the overall perception is that the solution to the puzzle is not too far away.

Competing interests

The authors declare that they have no competing interests.

Grant information

The author(s) declared that no grants were involved in supporting this work.

References

1. Zuberbier T, Aberer W, Asero R, *et al.*: **The EAACI/GA(2) LEN/EDF/WAO Guideline for the definition, classification, diagnosis, and management of urticaria: the 2013 revision and update.** *Allergy*. 2014; **69**(7): 868–87.
[PubMed Abstract](#) | [Publisher Full Text](#)
2. Grattan CE, Wallington TB, Warin RP, *et al.*: **A serological mediator in chronic idiopathic urticaria—a clinical, immunological and histological evaluation.** *Br J Dermatol*. 1986; **114**(5): 583–90.
[PubMed Abstract](#) | [Publisher Full Text](#)
3. Konstantinou GN, Asero R, Maurer M, *et al.*: **EAACI/GA(2)LEN task force consensus report: the autologous serum skin test in urticaria.** *Allergy*. 2009; **64**(9): 1256–68.
[PubMed Abstract](#) | [Publisher Full Text](#)
4. Asero R, Tedeschi A, Lorini M: **Histamine release in idiopathic cold urticaria.** *Allergy*. 2002; **57**(12): 1211–2.
[PubMed Abstract](#) | [Publisher Full Text](#)
5. Asero R, Tedeschi A, Lorini M: **Autoreactivity is highly prevalent in patients with multiple intolerances to NSAIDs.** *Ann Allergy Asthma Immunol*. 2002; **88**(5): 468–72.
[PubMed Abstract](#) | [Publisher Full Text](#)
6. Asero R, Tedeschi A, Lorini M, *et al.*: **Sera from patients with multiple drug allergy syndrome contain circulating histamine-releasing factors.** *Int Arch Allergy Immunol*. 2003; **131**(3): 195–200.
[PubMed Abstract](#) | [Publisher Full Text](#)
7. Greaves M: **Chronic urticaria.** *J Allergy Clin Immunol*. 2000; **105**(4): 664–72.
[PubMed Abstract](#) | [Publisher Full Text](#)
8. Asero R, Tedeschi A, Lorini M, *et al.*: **Chronic urticaria: novel clinical and serological aspects.** *Clin Exp Allergy*. 2001; **31**(7): 1105–10.
[PubMed Abstract](#) | [Publisher Full Text](#)
9. Grattan CE, Francis DM, Hide M, *et al.*: **Detection of circulating histamine releasing autoantibodies with functional properties of anti-IgE in chronic urticaria.** *Clin Exp Allergy*. 1991; **21**(6): 695–704.
[PubMed Abstract](#) | [Publisher Full Text](#)
10. Hide M, Francis DM, Grattan CE, *et al.*: **Autoantibodies against the high-affinity IgE receptor as a cause of histamine release in chronic urticaria.** *N Engl J Med*. 1993; **328**(22): 1599–604.
[PubMed Abstract](#) | [Publisher Full Text](#)
11. Fiebiger E, Maurer D, Holub H, *et al.*: **Serum IgG autoantibodies directed against the alpha chain of Fc epsilon RI: a selective marker and pathogenetic factor for a distinct subset of chronic urticaria patients?** *J Clin Invest*. 1995; **96**(6): 2606–12.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
12. Ferrer M, Kiné JP, Kaplan AP: **Comparative studies of functional and binding assays for IgG anti-Fc(epsilon)RIalpha (alpha-subunit) in chronic urticaria.** *J Allergy Clin Immunol*. 1998; **101**(5): 672–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
13. **E** Eckman JA, Hamilton RG, Gober LM, *et al.*: **Basophil phenotypes in chronic idiopathic urticaria in relation to disease activity and autoantibodies.** *J Invest Dermatol*. 2008; **128**(8): 1956–63.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
14. **F** Auyeung P, Mittag D, Hodgkin PD, *et al.*: **Autoreactive T cells in chronic spontaneous urticaria target the IgE Fc receptor 1α subunit.** *J Allergy Clin Immunol*. 2016; **138**(3): 761–768.e4.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
15. **F** Saini SS: **Chronic spontaneous urticaria: etiology and pathogenesis.** *Immunol Allergy Clin North Am*. 2014; **34**(1): 33–52.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
16. **F** Oliver ET, Sterba PM, Devine K, *et al.*: **Altered expression of chemoattractant receptor-homologous molecule expressed on T_H2 cells on blood basophils and eosinophils in patients with chronic spontaneous urticaria.** *J Allergy Clin Immunol*. 2016; **137**(1): 304–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
17. Vasagar K, Vonakis BM, Gober LM, *et al.*: **Evidence of *in vivo* basophil activation in chronic idiopathic urticaria.** *Clin Exp Allergy*. 2006; **36**(6): 770–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
18. Platzer MH, Grattan CE, Poulsen LK, *et al.*: **Validation of basophil histamine release against the autologous serum skin test and outcome of serum-induced basophil histamine release studies in a large population of chronic urticaria patients.** *Allergy*. 2005; **60**(9): 1152–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
19. Wedi B, Novacovic V, Koerner M, *et al.*: **Chronic urticaria serum induces histamine release, leukotriene production, and basophil CD63 surface expression—inhibitory effects of anti-inflammatory drugs.** *J Allergy Clin Immunol*. 2000; **105**(3): 552–60.
[PubMed Abstract](#) | [Publisher Full Text](#)
20. **F** Szegedi A, Irinyi B, Gál M, *et al.*: **Significant correlation between the CD63 assay and the histamine release assay in chronic urticaria.** *Br J Dermatol*. 2006; **155**(1): 67–75.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
21. Lourenço FD, Azor MH, Santos JC, *et al.*: **Activated status of basophils in chronic urticaria leads to interleukin-3 hyper-responsiveness and enhancement of histamine release induced by anti-IgE stimulus.** *Br J Dermatol*. 2008; **158**(5): 979–86.
[PubMed Abstract](#) | [Publisher Full Text](#)
22. Eckman JA, Hamilton RG, Saini SS: **Independent evaluation of a commercial test for “autoimmune” urticaria in normal and chronic urticaria subjects.** *J Invest Dermatol*. 2009; **129**(6): 1584–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
23. Altrichter S, Peter HJ, Pisarevskaja D, *et al.*: **IgE mediated autoallergy against thyroid peroxidase—a novel pathomechanism of chronic spontaneous urticaria?** *PLoS One*. 2011; **6**(4): e14794.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
24. Chang TW, Chen C, Lin CJ, *et al.*: **The potential pharmacologic mechanisms of omalizumab in patients with chronic spontaneous urticaria.** *J Allergy Clin Immunol*. 2015; **135**(2): 337–42.
[PubMed Abstract](#) | [Publisher Full Text](#)
25. **F** Ye YM, Jin HJ, Hwang EK, *et al.*: **Co-existence of chronic urticaria and metabolic syndrome: clinical implications.** *Acta Derm Venereol*. 2013; **93**(2): 156–60.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
26. **F** Chung SD, Wang KH, Tsai MC, *et al.*: **Hyperlipidemia Is Associated with Chronic Urticaria: A Population-Based Study.** *PLoS One*. 2016; **11**(3): e0150304.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
27. **F** Trinh HK, Pham DL, Ban GY, *et al.*: **Altered Systemic Adipokines in Patients with Chronic Urticaria.** *Int Arch Allergy Immunol*. 2016; **171**(2): 102–10.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
28. Kasperska-Zajac A, Sztylec J, Machura E, *et al.*: **Plasma IL-6 concentration correlates with clinical disease activity and serum C-reactive protein concentration in chronic urticaria patients.** *Clin Exp Allergy*. 2011; **41**(10): 1386–91.
[PubMed Abstract](#) | [Publisher Full Text](#)
29. Asero R, Tedeschi A, Riboldi P, *et al.*: **Plasma of patients with chronic urticaria shows signs of thrombin generation, and its intradermal injection causes wheal-and-flare reactions much more frequently than autologous serum.** *J Allergy Clin Immunol*. 2006; **117**(5): 1113–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
30. Asero R, Tedeschi A, Coppola R, *et al.*: **Activation of the tissue factor pathway of blood coagulation in patients with chronic urticaria.** *J Allergy Clin Immunol*. 2007; **119**(3): 705–10.
[PubMed Abstract](#) | [Publisher Full Text](#)
31. Asero R, Tedeschi A, Riboldi P, *et al.*: **Severe chronic urticaria is associated with elevated plasma levels of D-dimer.** *Allergy*. 2008; **63**(2): 176–80.
[PubMed Abstract](#) | [Publisher Full Text](#)
32. **F** Takahagi S, Mihara S, Iwamoto K, *et al.*: **Coagulation/fibrinolysis and inflammation markers are associated with disease activity in patients with chronic urticaria.** *Allergy*. 2010; **65**(5): 649–56.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
33. Cugno M, Marzano AV, Tedeschi A, *et al.*: **Expression of tissue factor by eosinophils in patients with chronic urticaria.** *Int Arch Allergy Immunol*. 2009; **148**(2): 170–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
34. Puccetti A, Bason C, Simeoni S, *et al.*: **In chronic idiopathic urticaria autoantibodies against Fc epsilonRII/CD23 induce histamine release via eosinophil activation.** *Clin Exp Allergy*. 2005; **35**(12): 1599–607.
[PubMed Abstract](#) | [Publisher Full Text](#)
35. Marzano AV, Tedeschi A, Fanoni D, *et al.*: **Activation of blood coagulation in bullous pemphigoid: role of eosinophils, and local and systemic implications.** *Br J Dermatol*. 2009; **160**(2): 266–72.
[PubMed Abstract](#) | [Publisher Full Text](#)
36. Cugno M, Cicardi M, Bottasso B, *et al.*: **Activation of the coagulation cascade in C1-inhibitor deficiencies.** *Blood*. 1997; **89**(9): 3213–8.
[PubMed Abstract](#)
37. Teitel JM, Bauer KA, Lau HK, *et al.*: **Studies of the prothrombin activation pathway utilizing radioimmunoassays for the F2/F1 + 2 fragment and thrombin–antithrombin complex.** *Blood*. 1982; **59**(5): 1086–97.
[PubMed Abstract](#)
38. Boneu B, Bes G, Pelzer H, *et al.*: **D-Dimers, thrombin antithrombin III complexes and prothrombin fragments 1+2: diagnostic value in clinically suspected deep vein thrombosis.** *Thromb Haemost*. 1991; **65**(1): 28–31.
[PubMed Abstract](#)
39. van Deventer SJ, Büller HR, ten Cate JW, *et al.*: **Experimental endotoxemia in humans: analysis of cytokine release and coagulation, fibrinolytic, and complement pathways.** *Blood*. 1990; **76**(12): 2520–6.
[PubMed Abstract](#)
40. Bossi F, Frossi B, Radillo O, *et al.*: **Mast cells are critically involved in serum-mediated vascular leakage in chronic urticaria beyond high-affinity IgE receptor stimulation.** *Allergy*. 2011; **66**(12): 1538–45.
[PubMed Abstract](#) | [Publisher Full Text](#)
41. Cugno M, Tedeschi A, Frossi B, *et al.*: **Detection of Low-Molecular-Weight Mast**

Cell-Activating Factors in Serum From Patients With Chronic Spontaneous Urticaria. *J Investig Allergol Clin Immunol.* 2016; 26(5): 310–3.

[PubMed Abstract](#) | [Publisher Full Text](#)

42. Grattan CE, Hamon CG, Cowan MA, *et al.*: **Preliminary identification of a low molecular weight serological mediator in chronic idiopathic urticaria.** *Br J Dermatol.* 1988; 119(2): 179–83.

[PubMed Abstract](#) | [Publisher Full Text](#)

43. Gericke J, Metz M, Ohanyan T, *et al.*: **Serum autoreactivity predicts time to response to omalizumab therapy in chronic spontaneous urticaria.** *J Allergy Clin Immunol.* 2017; 139(3): 1059–1061.e1.

[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)

44. Palacios T, Stillman L, Borish L, *et al.*: **Lack of basophil CD203c-upregulating activity as an immunological marker to predict response to treatment with omalizumab in patients with symptomatic chronic urticaria.** *J Allergy Clin*

Immunol Pract. 2016; 4(3): 529–30.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)

45. Ghazanfar MN, Sand C, Thomsen SF: **Effectiveness and safety of omalizumab in chronic spontaneous or inducible urticaria: evaluation of 154 patients.** *Br J Dermatol.* 2016; 175(2): 404–6.

[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)

46. Maurer M, Schütz A, Weller K, *et al.*: **Omalizumab is effective in symptomatic dermographism-results of a randomized placebo-controlled trial.** *J Allergy Clin Immunol.* 2017; pii: S0091-6749(17)30514-6.

[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)

47. Metz M, Schütz A, Weller K, *et al.*: **Omalizumab is effective in cold urticaria-results of a randomized placebo-controlled trial.** *J Allergy Clin Immunol.* 2017; pii: S0091-6749(17)30516-X.

[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. **Malcolm W Greaves**

Cutaneous Allergy Clinic, St John's Institute of Dermatology, St Thomas' Hospital, London, UK

Competing Interests: No competing interests were disclosed.

2. **Young-Min Ye**

Department of Allergy and Clinical Immunology, Ajou University School of Medicine, Suwon, South Korea

Competing Interests: No competing interests were disclosed.

3. **Simon Francis Thomsen**

¹ Department of Dermatology, Bispebjerg Hospital, Copenhagen, Denmark

² Department of Biomedical Sciences, University of Copenhagen, Copenhagen, Denmark

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research