REVIEW

Progress toward an integrated understanding of Parkinson's disease [version 1; peer review: 2 approved]

Maxime W.C. Rousseaux ^{1,2}, Joshua M. Shulman¹⁻⁴, Joseph Jankovic³

¹Jan and Dan Duncan Neurological Research Institute, Texas Children's Hospital, 1250 Moursund St, Houston, TX, 77030, USA

²Department of Molecular and Human Genetics, Baylor College of Medicine, One Baylor Plaza, Houston, TX, 77030, USA

³Parkinson's Disease Center and Movement Disorders Clinic, Department of Neurology, Baylor College of Medicine, 7200 Cambridge, Houston, TX, 77030-4202, USA

⁴Department of Neuroscience, Baylor College of Medicine, One Baylor Plaza, Houston, TX, 77030, USA

V1 First published: 12 Jul 2017, 6(F1000 Faculty Rev):1121
<https://doi.org/10.12688/f1000research.11820.1>

Latest published: 12 Jul 2017, 6(F1000 Faculty Rev):1121
<https://doi.org/10.12688/f1000research.11820.1>

Abstract

Parkinson's disease (PD) is the second most common neurodegenerative disorder after Alzheimer's disease, affecting over 10 million individuals worldwide. While numerous effective symptomatic treatments are currently available, no curative or disease-modifying therapies exist. An integrated, comprehensive understanding of PD pathogenic mechanisms will likely address this unmet clinical need. Here, we highlight recent progress in PD research with an emphasis on promising translational findings, including (i) advances in our understanding of disease susceptibility, (ii) improved knowledge of cellular dysfunction, and (iii) insights into mechanisms of spread and propagation of PD pathology. We emphasize connections between these previously disparate strands of PD research and the development of an emerging systems-level understanding that will enable the next generation of PD therapeutics.

Keywords

Parkinson's disease, PD, neurodegenerative disorders, α -Synuclein, Parkinson's disease genetics

Open Peer Review

Approval Status

	1	2
version 1
12 Jul 2017		

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Roger A Barker**, University of Cambridge, Cambridge, UK

2. **David G Standaert**, , Department of Neurology, The University of Alabama at Birmingham, Birmingham, USA

Any comments on the article can be found at the end of the article.

Corresponding author: Joseph Jankovic (josephj@bcm.edu)

Author roles: Rousseaux MWC: Writing – Original Draft Preparation, Writing – Review & Editing; Shulman JM: Writing – Original Draft Preparation, Writing – Review & Editing; Jankovic J: Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: Maxime W.C. Rousseaux and Joshua M. Shulman declare that they have no disclosures. Joseph Jankovic has received research or training grants and/or has served as a consultant for the Michael J. Fox Foundation for Parkinson's Research, Prothena Biosciences Inc, and Teva Pharmaceutical Industries Ltd.

Grant information: Maxime W.C. Rousseaux is supported in part by Grant No. PF-JFA-1762 from the Parkinson's Disease Foundation. Joshua M. Shulman is supported by the Huffington Foundation, the Jan and Dan Duncan Neurological Research Institute at Texas Children's Hospital, and a Career Award for Medical Scientists from the Burroughs Wellcome Fund. Joseph Jankovic has received research and/or educational support from the Michael J. Fox Foundation for Parkinson's Research, Parkinson's Foundation, the Parkinson Study Group, Prothena Biosciences Inc, and Teva Pharmaceutical Industries Ltd.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2017 Rousseaux MWC *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Rousseaux MWC, Shulman JM and Jankovic J. **Progress toward an integrated understanding of Parkinson's disease [version 1; peer review: 2 approved]** F1000Research 2017, 6(F1000 Faculty Rev):1121 <https://doi.org/10.12688/f1000research.11820.1>

First published: 12 Jul 2017, 6(F1000 Faculty Rev):1121 <https://doi.org/10.12688/f1000research.11820.1>

Introduction

Parkinson's disease (PD) is the most common movement disorder, affecting 2–3% of individuals over the age of 65¹. It is clinically characterized by a core set of motor manifestations, including tremor, slow movement (bradykinesia), increased muscle tone (rigidity), and gait and postural impairment as well as a variety of other motor and non-motor features, including cognitive impairment, depression, pain and other sensory symptoms, autonomic dysfunction, and others². PD is characterized pathologically by the loss of predominantly dopaminergic neurons, associated with intracellular, insoluble α -synuclein (α -Syn) aggregates, largely localized to cytoplasmic inclusions termed Lewy bodies and within neuronal processes termed Lewy neurites. Whereas current treatments can ameliorate the cardinal motor symptoms, no disease-modifying therapies exist³. A large body of research has therefore focused on understanding the biological mechanisms that underlie disease onset and progression, with the goal of developing effective pathogenesis-targeted, disease-modifying therapies.

As this year marks the 200-year anniversary of the recognition of PD by James Parkinson⁴, we also celebrate the remarkable progress in understanding PD etiology and pathogenesis⁵. Here, we review recent advances in the study of the genetics, cell biology, and pathology of PD, highlighting emerging areas of overlap. Where these areas were previously studied in isolation, results from these disparate strands of research are beginning to converge, providing a more unified understanding of PD pathogenesis. We argue that this integrative approach to PD, in which seemingly disconnected results are re-examined as components of a cohesive whole, is also creating exciting opportunities for clinical applications.

Functional genomics and the promise of personalized medicine

Although only a small minority of patients with PD have thus far been found to have responsible pathogenic gene mutations, genetic discoveries have nevertheless been a driving force in the elucidation of PD mechanisms. Following on the early success from studies of rare families with Mendelian forms of PD, more than 20 PD genes and variants have now been implicated, including from large-scale genome-wide association studies (GWAS) and, more recently, whole exome sequencing (WES) studies in population-based cohorts^{6–8}. A common challenge with either approach entails definitive confirmation of the responsible genes and elucidation of implicated disease mechanisms. In GWAS, implicated genomic regions often contain several equally plausible gene candidates. Whereas WES studies usually single out specific gene candidates, the implicated alleles may be too rare to definitively confirm a causal link to PD based on currently available sample sizes. Medium- to high-throughput screening assays in cellular or animal models can connect promising candidate genes to PD-relevant biologic mechanisms, prioritizing a subset for further study. For example, Jansen *et al.*⁹ evaluated 27 promising candidate genes with homozygous or compound heterozygous loss-of-function alleles based on WES in 1,148 unrelated young-onset PD cases. Since nearly all of the gene candidates were observed only once in the cohort, the investigators probed each gene for roles in mitochondrial dynamics or α -Syn-mediated toxicity using cellular and fruit fly experimental models. Ultimately, five

genes were supported by both functional data and additional human genetic analyses consistent with replication. Beyond accelerating the discovery of novel PD genes, related approaches are also revealing the function of many other established genes/variants, grouping discrete susceptibility loci into common pathways and thereby consolidating our understanding of PD pathogenesis. For example, the recently identified PD gene *CHCHD2* may mediate its activity through a mitochondrial pathway like other recessive PD genes (see below)^{10–12}. Moreover, *VPS35* and *EIF4G1*, both implicated in autosomal dominant forms of PD^{13–17}, were recently found to genetically interact with one another and converge on α -Syn toxicity in yeast, worm, and mouse models of synucleinopathy¹⁸.

By contrast, with the sequence-based discovery of rare variant PD risk factors, the susceptibility alleles identified by GWAS⁷ usually do not alter protein-coding regions, making functional follow-up more challenging. For example, one of the earliest discovered PD risk polymorphisms at the human *MAPT* locus may primarily impact alternative mRNA splicing^{19,20}. In another recent study, Soldner *et al.* used human pluripotent cell-derived neurons containing an intronic PD-related variant in the gene encoding α -Syn (*SNCA*)²¹. The authors found that this common polymorphism, which is present in about half of the population, coincided with a distal enhancer element resulting in an approximately 10% increase in *SNCA* transcript levels. It was therefore suggested that a mild increase of α -Syn over the course of decades renders individuals susceptible to PD. This is consistent with the findings from rare families with *SNCA* locus multiplication. In these cases, individuals with *SNCA* gene multiplication present with clinical features typical of PD, including a similar age at onset to sporadic PD^{22,23}, whereas individuals with *SNCA* gene triplication present with a more early onset, aggressive form of PD²⁴. Thus, *SNCA* gene dosage may be an integral feature of PD pathogenesis. In the case of the more common *SNCA* polymorphism, the modest increase in α -Syn protein levels may interact with other genetic risk variants or environmental exposures to cause PD. For example, Goldman *et al.* found that head injury was significantly associated with increased PD risk, but only in the context of a disease-associated *SNCA* promoter polymorphism²⁵.

One of the great hopes for advances in PD genetics is to realize goals for personalized medicine, including improved risk prediction and even targeted therapies. It has long been speculated that much of PD's clinical heterogeneity may be genetically encoded^{26,27}. For example, besides their potent impact on risk of PD²⁸, *GBA* mutations have been reported by several groups to cause an earlier age-at-onset, more rapid progression, and an increased risk of cognitive impairment and dementia in carriers^{29–34}. Moreover, additional studies have looked at the effect of allelic heterogeneity on modifying PD clinical presentations^{35–37}. In the future, it may be important to couple such studies examining the clinical impact of selected allelic variants with experimental investigations to define functional consequences in well-defined cellular or animal models. Moreover, once characterized, these models can serve as a platform for testing putative “personalized” treatments^{38–42}. For example, Sanofi Genzyme is currently supporting a study (MOVES-PD) of GZ/SAR402671, a glucosylceramide synthase inhibitor, in PD patients

carrying a *GBA* gene mutation (ClinicalTrials.gov, NCT02906020). Another study (AiM-PD) is examining the effects of oral Ambroxol, a glucocerebrosidase-modulating chaperone, in patients with PD (ClinicalTrials.gov, NCT02941822). Lastly, the use of biomarkers in stratifying clinical populations and understanding the biological underpinning of PD subtypes will be critically important when developing personalized medicine approaches. Specifically, profiling blood and CSF biomarkers may enhance disease subtyping based on clinical manifestations alone. This hypothesis is currently being tested in the Parkinson's Progression Markers Initiative (PPMI) led by the Michael J. Fox Foundation for Parkinson's Research^{43,44}.

From genes to organelles and cellular homeostasis

The maintenance and function of cellular organelles, including mitochondria and lysosomes, are critical for functional neuronal integrity^{45–47}. Interestingly, several previously identified PD genes such as *PARK2*, *PARK6*, and *PARK7* (encoding Parkin, PTEN-induced putative kinase 1 [PINK1], and DJ-1, respectively) have been linked to mitochondrial function^{48–54}. Since their initial discovery, a large body of work has elucidated a cellular pathway through which dysfunctional mitochondria can be recycled by way of autophagy or, more specifically, “mitophagy”^{55–59}. Damaged mitochondria promote the phosphorylation of ubiquitin and Parkin by PINK1 and are subsequently degraded by the autophagosomal system^{59–62}. Additionally, studies of these genes have provided insight into their mitochondrial functions in healthy and diseased contexts. For example, a recent study suggests that Parkin acts as an endogenous buffer for mitochondrial stress and its loss sensitizes dopaminergic neurons to mitochondrial mutations over time^{63,64}. In addition, using a fly model, Vos and colleagues⁶⁵ discovered a new pathway that could suppress the motor and biochemical abnormalities caused by *PINK1* loss of function. Specifically, PINK1 genetically interacts with the enzyme responsible for the conversion of vitamin K1 into vitamin K2⁶⁵. Remarkably, supplementation of vitamin K2 could reverse *PINK1* mutant phenotypes, suggesting a potential therapeutic approach. However, patient selection will be critical for potential clinical trials, as *PARK6* mutations result in rare, autosomal recessive juvenile parkinsonism⁶⁶, and it is uncertain whether a similar functional deficiency in vitamin K2 may apply in the general PD patient population. Identifying the subcellular impact of other PD genes may similarly lead to other targeted therapies.

Beyond discovering the primary targets of genetic abnormalities, it is essential to understand the subsequent cascade of cellular injury, such as how damaged mitochondria impact other cellular constituents, leading to neuronal dysfunction and death. In other words, discrete targets must be understood in the context of a cohesive, dynamic system. One example comes from recent studies that strongly implicate defects in the vesicular trafficking system⁶⁷, which mediates cellular secretion and endocytosis as well as vesicle docking and fusion and is critically important for synaptic transmission, lysosomal degradation, and autophagy. Several PD genes, including *VPS35*, *LRRK2*, *RAB7L1*, *GBA*, and *SNCA*, have functions that converge on the vesicular trafficking system^{67–70}. Leucine-rich repeat kinase 2 (LRRK2) and Rab-7-like protein 1 (RAB7L1),

for example, act coordinately to regulate endolysosomal protein sorting via Rab GTPases, and several studies also support a key role for the cargo-shuttling retromer protein vacuolar protein sorting 35 (VPS35)^{71–73}. In fact, the retromer is implicated as a critical downstream effector whose dysfunction may lead to neuronal toxicity and death⁷⁴. Importantly, VPS35 may also play a role in mitophagy, perhaps via trafficking of mitochondria-derived vesicles to lysosomes⁷⁵. The dense interconnections between PD genes and other regulators of vesicular trafficking were highlighted by recent work that combined an α -Syn protein–protein interaction network with suppressor-enhancer screening in yeast^{76,77}. Based on these and other findings, chemical modulators of autophagy and/or the retromer, such as rapamycin (or similar “rapalogues”)^{78,79} and R55^{80,81}, respectively, may be promising therapeutic avenues for targeting vesicular sorting defects in PD. However, since these pathways mediate essential functions in most tissues, successful dose titration to achieve selective action in the nervous system while minimizing potentially deleterious off-target effects is one anticipated challenge⁸². Nevertheless, these studies illustrate how the emerging, systems-based understanding of PD can highlight vulnerable “nodes” within complex cellular networks, creating promising therapeutic opportunities.

α -Syn toxicity and propagation: from cells to systems

The centrality of α -Syn in PD pathogenesis was established nearly two decades ago with the dual finding that (i) this protein is the principal constituent of the hallmark Lewy body pathology⁸³ and (ii) *SNCA* gene mutations cause familial forms of PD⁸⁴. Since then, α -Syn genomic locus multiplication^{22–24} or promoter polymorphisms that increase protein expression²¹ have also been confirmed as causal factors. Thus, intensive investigation has probed the relationship between α -Syn with PD pathogenesis. α -Syn was first described as a member of the synuclein family, which is associated with the synapse and the nucleus⁸⁵. While studies on the physiological function of α -Syn suggest that it may play a role in synaptic transmission⁸⁶ and aid in curving cellular membranes⁸⁷, less is known regarding the mechanism(s) through which its gain-of-function causes neurodegeneration. Thus, one aspect of α -Syn research has focused on understanding how this protein causes toxicity within the cell. Studies in both animal models and human tissue have highlighted a role for α -Syn at the outer mitochondrial membrane^{88,89}, the nucleus^{90–92}, and the synapse^{93,94} as putative toxic mechanisms. Moreover, mechanisms involving proteostasis^{95,96} and lysosomal dysfunction⁴⁵ that collectively lead to increased α -Syn levels are also tantalizing.

Given the direct relationship between α -Syn abundance and its role in PD pathogenesis and propagation, immunotherapy against α -Syn has emerged as one of the most promising therapeutic approaches for PD^{97,98}. Results from ongoing and future immunotherapeutic trials by Prothena/Roche, Biogen, AFFiRiS, and other biotech companies will provide information on whether aggregated α -Syn is an important therapeutic target⁹⁹. Nevertheless, caution is warranted given previous negative outcomes from immunotherapeutic trials of other neurodegenerative diseases, such as Alzheimer's disease, characterized by protein aggregation (proteinopathies)¹⁰⁰.

Another avenue for therapeutic intervention that is actively being investigated is based on the observation that hyperactivity of the non-receptor tyrosine kinase c-Abl contributes to α -Syn phosphorylation, accumulation, and neurodegeneration¹⁰¹. This has led to preliminary investigation of Nilotinib, a c-Abl inhibitor previously approved for the treatment of chronic myeloid leukemia, as a potential therapeutic agent for PD¹⁰². Further studies, however, are needed before this potent drug can be recommended as a symptomatic or disease-modifying therapy for PD¹⁰³. A single-center trial at Georgetown University (ClinicalTrials.gov, NCT02954978) and a multicenter trial (NILO-PD) (ClinicalTrials.gov, NCT02281474), sponsored by the Michael J. Fox Foundation for Parkinson's Research and the Parkinson Study Group, are currently under way.

Since the seminal observation by Braak and colleagues that most idiopathic PD cases fit a relatively predictable, caudal to rostral pathologic staging progression¹⁰⁴, there has been great interest in understanding the mechanism by which α -Syn pathology may spread from the peripheral organs (e.g. the enteric or pericardial tissue) via the vagus nerve to the lower brainstem and eventually involve the neocortex. The intriguing possibility that α -Syn pathology can spread from cell to cell^{105–107} was suggested by observations of Lewy-like pathology in engrafted neurons from PD patients receiving fetal dopaminergic cell transplants^{108,109}. First studies indicated that aggregates of α -Syn could enter cells that express transgenic α -Syn and seed the formation of new aggregates^{110,111}. More recently, research findings have indicated that even synthetic, wild-type forms of α -Syn, if improperly folded and injected in the mouse brain, can induce the misfolding of otherwise normal endogenous α -Syn, thereby amplifying and propagating pathological forms^{112–114}, properties consistent with those of prions (infectious proteinaceous agents)¹¹⁵. It appears that α -Syn can adopt a variety of different misfolded/aggregated oligomeric conformations that correspond to distinct profiles of toxicity in experimental assays^{116,117}. These findings raise the intriguing, though yet unproven, hypothesis that certain α -Syn “strains” may contribute to clinical and pathologic heterogeneity among PD and other synucleinopathies. Interestingly, Mao *et al.*¹¹⁸ recently identified lymphocyte-activation gene 3 (LAG3) as a candidate receptor for α -Syn oligomeric “seeds”, and genetic manipulation of LAG3 in cells and mouse models altered pathologic progression. These findings raise the possibility of diagnostic and therapeutic advance based on the detection of α -Syn strains in patient populations as well as potential pharmacologic blockade of propagation¹¹⁸. It is also possible that neuroanatomic variation among key co-factors or receptors for α -Syn seeding or spread might contribute to the selective vulnerability of specific neuronal subpopulations in PD. Despite remarkable recent progress, the clinical relevance of α -Syn seeding and propagation (if any) remains to be fully understood^{105,119,120}. For example, autopsies from some PD patients receiving fetal grafts were devoid of pathology two decades following transplantation^{121,122}. Another recent study found that individuals receiving human growth hormone derived from human cadavers were at no greater risk of developing PD¹²³. The extent of pathological spread of α -Syn does not always follow a defined trans-synaptic pattern nor does the brain areas it affects fully correlate with clinical measures¹⁰⁵. Thus, a continued investigation

into the mechanism through which α -Syn pathological assemblies form and how these are tied to toxicity is warranted.

Gastrointestinal dysmotility is a common early complaint in PD patients^{124,125}, and the enteric nervous system has been implicated as an early target of α -Syn pathology^{104,126}. Along with the growing interest in immunologic and inflammatory disease mechanisms, the gut microbiome has recently come under investigation in studies of both PD patients^{127–130} and animal models¹³¹. Sampson and colleagues¹³¹ found that α -Syn transgenic mice living in a germ-free environment were less vulnerable to neurodegeneration, but when the mice were inoculated with fecal bacteria taken from patients with PD their motor function deteriorated. While this study suggests the possibility that the gut microbiome may influence PD manifestations, it will be important to define the specific microbial contributors to the disease as well as recapitulate these findings in other disease models before moving forward into humans¹³². Beyond its potential role in disease modulation, the gastrointestinal tract might even be a trigger point for disease based on the findings of early enteric nervous system α -Syn pathology in some pathologic series¹⁰⁴. Long before the current excitement concerning α -Syn propagation, Braak *et al.* speculated about a possible gastrointestinal pathogen (or other enteric exposure) as an initiating event, followed by pathologic spread via the vagus nerve to its medullary dorsal motor nucleus where PD changes first appear in the brainstem^{104,126,133}. Following up on this provocative hypothesis, investigators recently found that subjects undergoing vagotomy (transection of the vagal nerve for the treatment of peptic ulcer disease) are at a modest but significantly reduced risk of PD^{134,135}. While intriguing, the finding requires further confirmation, and animal model studies will be essential to definitively prove that the mechanism of protection is indeed based on the disruption of spread from the enteric to the central nervous system.

Conclusion

Recent advances have clearly enhanced our knowledge of the fundamental processes underlying PD pathogenesis. As connections are recognized among the disparate domains of PD inquiry, the broader patterns begin to emerge. As discussed above, we now recognize the cellular targets of many PD genes, such as how *PARKIN* and *GBA* regulate mitochondrial or lysosomal function, respectively. Additionally, the field has made progress toward understanding how dynamic interactions between such organelles impact overall cellular health, as in mitophagy. Lastly, studies of α -Syn illustrate the convergence of classical histopathologic analysis of PD with genetic investigations, and more recent investigations demonstrate how synucleinopathy not only impacts single cells or tissues but also may propagate throughout the nervous system. In sum, we are rapidly making progress toward a more cohesive model of PD pathogenesis, and this systems-level understanding is likely to accelerate therapeutic inroads. With the broad outlines of the “PD puzzle” now apparent (Figure 1), we predict that new insights can be more rapidly integrated within this framework. For example, forthcoming discoveries of new genetic risk loci can be understood within the context of known functional pathways within both neurons and other cell types, such as astrocytes or microglia. An integrated understanding of PD will

Figure 1. Putting the Parkinson's disease (PD) puzzle together. Three pieces of the "PD puzzle" covered in this short review are highlighted within the context of the brain. Additional elements remain to be fully elucidated. The integrated, systems-level understanding of PD that emerges from understanding how these components fit together will likely accelerate therapeutic advances. α -Syn, α -synuclein.

also enable more effective multi- and inter-disciplinary collaboration among scientists and clinicians, driving next-generation therapeutic trials targeting disease mechanisms and fulfilling the promise of personalized medicine. Advances in understanding the cellular mechanism underlying PD-related neurodegeneration will undoubtedly lead to better symptomatic and novel pathogenesis-targeted, disease-modifying therapies¹³⁶.

Abbreviations

α -Syn, α -synuclein; GWAS, genome-wide association study; LAG3, lymphocyte-activation gene 3; PD, Parkinson's disease; PINK1, PTEN-induced putative kinase 1; VPS35, vacuolar protein sorting 35; WES, whole exome sequencing.

Competing interests

Maxime W.C. Rousseaux and Joshua M. Shulman declare that they have no disclosures. Joseph Jankovic has received research or

training grants and/or has served as a consultant for the Michael J. Fox Foundation for Parkinson's Research, Prothena Biosciences Inc, and Teva Pharmaceutical Industries Ltd.

Grant information

Maxime W.C. Rousseaux is supported in part by Grant No. PF-JFA-1762 from the Parkinson's Disease Foundation. Joshua M. Shulman is supported by the Huffington Foundation, the Jan and Dan Duncan Neurological Research Institute at Texas Children's Hospital, and a Career Award for Medical Scientists from the Burroughs Wellcome Fund. Joseph Jankovic has received research and/or educational support from the Michael J. Fox Foundation for Parkinson's Research, Parkinson's Foundation, the Parkinson Study Group, Prothena Biosciences Inc, and Teva Pharmaceutical Industries Ltd.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

1. Poewe W, Seppi K, Tanner CM, *et al.*: **Parkinson disease.** *Nat Rev Dis Primers.* 2017; **3**: 17013.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
2. Postuma RB, Berg D, Stern M, *et al.*: **MDS clinical diagnostic criteria for Parkinson's disease.** *Mov Disord.* 2015; **30**(12): 1591–601.
[PubMed Abstract](#) | [Publisher Full Text](#)
3. Lotia M, Jankovic J: **New and emerging medical therapies in Parkinson's disease.** *Expert Opin Pharmacother.* 2016; **17**(7): 895–909.
[PubMed Abstract](#) | [Publisher Full Text](#)
4. Parkinson J: **An essay on the shaking palsy. 1817.** *J Neuropsychiatry Clin Neurosci.* 2002; **14**(2): 223–36; discussion 222.
[PubMed Abstract](#) | [Publisher Full Text](#)
5. Jankovic J: **Movement disorders in 2016: progress in Parkinson disease and other movement disorders.** *Nat Rev Neurol.* 2017; **13**(2): 76–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
6. Farlow JL, Robak LA, Hetrick K, *et al.*: **Whole-exome sequencing in familial Parkinson disease.** *JAMA Neurol.* 2016; **73**(1): 68–75.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
7. Nalls MA, Pankratz N, Lill CM, *et al.*: **Large-scale meta-analysis of genome-wide association data identifies six new risk loci for Parkinson's disease.** *Nat Genet.* 2014; **46**(9): 989–93.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
8. Lill CM: **Genetics of Parkinson's disease.** *Mol Cell Probes.* 2016; **30**(6): 386–96.
[PubMed Abstract](#) | [Publisher Full Text](#)

9. Jansen IE, Ye H, Heetveld S, *et al.*: **Discovery and functional prioritization of Parkinson's disease candidate genes from large-scale whole exome sequencing.** *Genome Biol.* 2017; **18**(1): 22.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
10. Funayama M, Ohe K, Amo T, *et al.*: **CHCHD2 mutations in autosomal dominant late-onset Parkinson's disease: a genome-wide linkage and sequencing study.** *Lancet Neurol.* 2015; **14**(3): 274–82.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
11. Singleton A: **A new gene for Parkinson's disease: should we care?** *Lancet Neurol.* 2015; **14**(3): 238–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
12. Ogaki K, Koga S, Heckman MG, *et al.*: **Mitochondrial targeting sequence variants of the CHCHD2 gene are a risk for Lewy body disorders.** *Neurology.* 2015; **85**(23): 2016–25.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
13. Deng H, Gao K, Jankovic J: **The VPS35 gene and Parkinson's disease.** *Mov Disord.* 2013; **28**(5): 569–75.
[PubMed Abstract](#) | [Publisher Full Text](#)
14. Vilarinho-Güell C, Wider C, Ross OA, *et al.*: **VPS35 mutations in Parkinson disease.** *Am J Hum Genet.* 2011; **89**(1): 162–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
15. Zimprich A, Benet-Pagès A, Struhal W, *et al.*: **A mutation in VPS35, encoding a subunit of the retromer complex, causes late-onset Parkinson disease.** *Am J Hum Genet.* 2011; **89**(1): 168–75.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
16. Deng H, Wu Y, Jankovic J: **The EIF4G1 gene and Parkinson's disease.** *Acta Neurol Scand.* 2015; **132**(2): 73–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
17. Chartier-Harlin MC, Dachsel JC, Vilarinho-Güell C, *et al.*: **Translation initiator EIF4G1 mutations in familial Parkinson disease.** *Am J Hum Genet.* 2011; **89**(3): 398–406.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
18. Dhungel N, Eleuteri S, Li LB, *et al.*: **Parkinson's disease genes VPS35 and EIF4G1 interact genetically and converge on α -synuclein.** *Neuron.* 2015; **85**(1): 76–87.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
19. Trabzuni D, Wray S, Vandrovcova J, *et al.*: **MAPT expression and splicing is differentially regulated by brain region: relation to genotype and implication for tauopathies.** *Hum Mol Genet.* 2012; **21**(18): 4094–103.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
20. Valencia GT, Srivastava GP, Oliveira-Filho J, *et al.*: **The role of MAPT haplotype H2 and isoform 1N/4R in parkinsonism of older adults.** *PLoS One.* 2016; **11**(7): e0157452.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
21. Soldner F, Stelzer Y, Shivalila CS, *et al.*: **Parkinson-associated risk variant in distal enhancer of α -synuclein modulates target gene expression.** *Nature.* 2016; **533**(7601): 95–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
22. Chartier-Harlin MC, Kachergus J, Roumier C, *et al.*: **Alpha-synuclein locus duplication as a cause of familial Parkinson's disease.** *Lancet.* 2004; **364**(9440): 1167–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
23. Ibáñez P, Bonnet AM, Débarges B, *et al.*: **Causal relation between alpha-synuclein gene duplication and familial Parkinson's disease.** *Lancet.* 2004; **364**(9440): 1169–71.
[PubMed Abstract](#) | [Publisher Full Text](#)
24. Singleton AB, Farrer M, Johnson J, *et al.*: **Alpha-synuclein locus triplication causes Parkinson's disease.** *Science.* 2003; **302**(5646): 841.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
25. Goldman SM, Kamel F, Ross GW, *et al.*: **Head injury, α -synuclein Rep1, and Parkinson's disease.** *Ann Neurol.* 2012; **71**(1): 40–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
26. Thenganatt MA, Jankovic J: **Parkinson disease subtypes.** *JAMA Neurol.* 2014; **71**(4): 499–504.
[PubMed Abstract](#) | [Publisher Full Text](#)
27. Nussbaum RL, Polymeropoulos MH: **Genetics of Parkinson's disease.** *Hum Mol Genet.* 1997; **6**(10): 1687–91.
[PubMed Abstract](#) | [Publisher Full Text](#)
28. Sidransky E, Nalls MA, Aasly JO, *et al.*: **Multicenter analysis of glucocerebrosidase mutations in Parkinson's disease.** *N Engl J Med.* 2009; **361**(17): 1651–61.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
29. Migdalska-Richards A, Schapira AH: **The relationship between glucocerebrosidase mutations and Parkinson disease.** *J Neurochem.* 2016; **139**(Suppl 1): 77–90.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
30. Blanz J, Saftig P: **Parkinson's disease: acid-glucocerebrosidase activity and alpha-synuclein clearance.** *J Neurochem.* 2016; **139**(Suppl 1): 198–215.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
31. Brockmann K, Srulijes K, Pflederer S, *et al.*: **GBA-associated Parkinson's disease: reduced survival and more rapid progression in a prospective longitudinal study.** *Mov Disord.* 2015; **30**(3): 407–11.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
32. Clark LN, Ross BM, Wang Y, *et al.*: **Mutations in the glucocerebrosidase gene are associated with early-onset Parkinson disease.** *Neurology.* 2007; **69**(12): 1270–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
33. Davis MY, Johnson CO, Leverenz JB, *et al.*: **Association of GBA mutations and the E326K polymorphism with motor and cognitive progression in Parkinson disease.** *JAMA Neurol.* 2016; **73**(10): 1217–24.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
34. Winder-Rhodes SE, Evans JR, Ban M, *et al.*: **Glucocerebrosidase mutations influence the natural history of Parkinson's disease in a community-based incident cohort.** *Brain.* 2013; **136**(Pt 2): 392–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
35. Nalls MA, McLean CY, Rick J, *et al.*: **Diagnosis of Parkinson's disease on the basis of clinical and genetic classification: a population-based modelling study.** *Lancet Neurol.* 2015; **14**(10): 1002–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
36. Pihlström L, Mosset KR, Grimstad E, *et al.*: **A cumulative genetic risk score predicts progression in Parkinson's disease.** *Mov Disord.* 2016; **31**(4): 487–90.
[PubMed Abstract](#) | [Publisher Full Text](#)
37. Mata IF, Leverenz JB, Weintraub D, *et al.*: **GBA variants are associated with a distinct pattern of cognitive deficits in Parkinson's disease.** *Mov Disord.* 2016; **31**(1): 95–102.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
38. Migdalska-Richards A, Daly L, Bezaud E, *et al.*: **Ambroxol effects in glucocerebrosidase and α -synuclein transgenic mice.** *Ann Neurol.* 2016; **80**(5): 766–75.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
39. Ishay Y, Zimran A, Szer J, *et al.*: **Combined beta-glucosylceramide and amroxol hydrochloride in patients with Gaucher related Parkinson disease: from clinical observations to drug development.** *Blood Cells Mol Dis.* 2016; pii: S1079-9796(16)30229-7.
[PubMed Abstract](#) | [Publisher Full Text](#)
40. Migdalska-Richards A, Ko WKD, Li Q, *et al.*: **Oral amroxol increases brain glucocerebrosidase activity in a nonhuman primate.** *Synapse.* 2017; **71**(7).
[PubMed Abstract](#) | [Publisher Full Text](#)
41. Wang G, McCain ML, Yang L, *et al.*: **Modeling the mitochondrial cardiomyopathy of Barth syndrome with induced pluripotent stem cell and heart-on-chip technologies.** *Nat Med.* 2014; **20**(6): 616–23.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
42. Zhang YS, Aleman J, Shin SR, *et al.*: **Multisensor-integrated organs-on-chips platform for automated and continual in situ monitoring of organoid behaviors.** *Proc Natl Acad Sci U S A.* 2017; **114**(12): E2293–E2302.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
43. Espay AJ, Schwarzschild MA, Tanner CM, *et al.*: **Biomarker-driven phenotyping in Parkinson's disease: a translational missing link in disease-modifying clinical trials.** *Mov Disord.* 2017; **32**(3): 319–24.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
44. Parkinson Progression Marker Initiative: **The Parkinson Progression Marker Initiative (PPMI).** *Prog Neurobiol.* 2011; **95**(4): 629–35.
[PubMed Abstract](#) | [Publisher Full Text](#)
45. Manzoni C, Lewis PA: **Dysfunction of the autophagy/lysosomal degradation pathway is a shared feature of the genetic synucleinopathies.** *FASEB J.* 2013; **27**(9): 3424–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
46. Schon EA, Manfredi G: **Neuronal degeneration and mitochondrial dysfunction.** *J Clin Invest.* 2003; **111**(3): 303–12.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
47. Van Laar VS, Berman SB: **The interplay of neuronal mitochondrial dynamics and bioenergetics: implications for Parkinson's disease.** *Neurobiol Dis.* 2013; **51**: 43–55.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
48. Haelterman NA, Yoon WH, Sandoval H, *et al.*: **A mitocentric view of Parkinson's disease.** *Annu Rev Neurosci.* 2014; **37**: 137–59.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
49. Irrcher I, Aleyasin H, Seifert EL, *et al.*: **Loss of the Parkinson's disease-linked gene DJ-1 perturbs mitochondrial dynamics.** *Hum Mol Genet.* 2010; **19**(19): 3734–46.
[PubMed Abstract](#) | [Publisher Full Text](#)
50. Park J, Lee SB, Lee S, *et al.*: **Mitochondrial dysfunction in *Drosophila* PINK1 mutants is complemented by parkin.** *Nature.* 2006; **441**(7097): 1157–61.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
51. Clark IE, Dodson MW, Jiang C, *et al.*: ***Drosophila* pink1 is required for mitochondrial function and interacts genetically with parkin.** *Nature.* 2006; **441**(7097): 1162–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
52. Yang Y, Gehrke S, Imai Y, *et al.*: **Mitochondrial pathology and muscle and dopaminergic neuron degeneration caused by inactivation of *Drosophila* pink1 is rescued by parkin.** *Proc Natl Acad Sci U S A.* 2006; **103**(28): 10793–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
53. Canet-Avilés RM, Wilson MA, Miller DW, *et al.*: **The Parkinson's disease protein**

- DJ-1 is neuroprotective due to cysteine-sulfinic acid-driven mitochondrial localization.** *Proc Natl Acad Sci U S A.* 2004; 101(24): 9103–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
54. Andres-Mateos E, Perier C, Zhang L, *et al.*: **DJ-1 gene deletion reveals that DJ-1 is an atypical peroxiredoxin-like peroxidase.** *Proc Natl Acad Sci U S A.* 2007; 104(37): 14807–12.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 55. **F** Kinghorn KJ, Asghari AM, Castillo-Quan JI: **The emerging role of autophagolysosomal dysfunction in Gaucher disease and Parkinson's disease.** *Neural Regen Res.* 2017; 12(3): 380–4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 56. Rahman MA, Rhim H: **Therapeutic implication of autophagy in neurodegenerative diseases.** *BMB Rep.* 2017; pii: 3831.
[PubMed Abstract](#)
 57. Youle RJ, Narendra DP: **Mechanisms of mitophagy.** *Nat Rev Mol Cell Biol.* 2011; 12(1): 9–14.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 58. **F** Narendra DP, Jin SM, Tanaka A, *et al.*: **PINK1 is selectively stabilized on impaired mitochondria to activate parkin.** *PLoS Biol.* 2010; 8(1): e1000298.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 59. **F** Lazarou M, Sliter DA, Kane LA, *et al.*: **The ubiquitin kinase PINK1 recruits autophagy receptors to induce mitophagy.** *Nature.* 2015; 524(7565): 309–14.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 60. **F** Kane LA, Lazarou M, Fogel AI, *et al.*: **PINK1 phosphorylates ubiquitin to activate parkin E3 ubiquitin ligase activity.** *J Cell Biol.* 2014; 205(2): 143–53.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 61. **F** Kazlauskaitė A, Kondapalli C, Gourlay R, *et al.*: **Parkin is activated by PINK1-dependent phosphorylation of ubiquitin at Ser⁶⁵.** *Biochem J.* 2014; 460(1): 127–39.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 62. **F** Koyano F, Okatsu K, Kosako H, *et al.*: **Ubiquitin is phosphorylated by PINK1 to activate parkin.** *Nature.* 2014; 510(7503): 162–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 63. **F** Pickrell AM, Youle RJ: **The roles of PINK1, parkin, and mitochondrial fidelity in Parkinson's disease.** *Neuron.* 2015; 85(2): 257–73.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 64. **F** Pickrell AM, Huang CH, Kennedy SR, *et al.*: **Endogenous parkin preserves dopaminergic substantia nigral neurons following mitochondrial DNA mutagenic stress.** *Neuron.* 2015; 87(2): 371–81.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 65. **F** Vos M, Esposito G, Edirisinghe JN, *et al.*: **Vitamin K2 is a mitochondrial electron carrier that rescues pink1 deficiency.** *Science.* 2012; 336(6086): 1306–10.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 66. **F** Valente EM, Abou-Sleiman PM, Caputo V, *et al.*: **Hereditary early-onset Parkinson's disease caused by mutations in PINK1.** *Science.* 2004; 304(5674): 1158–60.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 67. **F** Abeliovich A, Gitler AD: **Defects in trafficking bridge Parkinson's disease pathology and genetics.** *Nature.* 2016; 539(7628): 207–16.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 68. Wang S, Bellen HJ: **The retromer complex in development and disease.** *Development.* 2015; 142(14): 2392–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 69. **F** McLelland G, Soubannier V, Chen CX, *et al.*: **Parkin and PINK1 function in a vesicular trafficking pathway regulating mitochondrial quality control.** *EMBO J.* 2014; 33(4): 282–95.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 70. Lashuel HA, Hirling H: **Rescuing defective vesicular trafficking protects against alpha-synuclein toxicity in cellular and animal models of Parkinson's disease.** *ACS Chem Biol.* 2006; 1(7): 420–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
 71. **F** MacLeod DA, Rhinn H, Kuwahara T, *et al.*: **RAB7L1 interacts with LRRK2 to modify intraneuronal protein sorting and Parkinson's disease risk.** *Neuron.* 2013; 77(3): 425–39.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 72. **F** Steger M, Tonelli F, Ito G, *et al.*: **Phosphoproteomics reveals that Parkinson's disease kinase LRRK2 regulates a subset of Rab GTPases.** *eLife.* 2016; 5: pii: e12813.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 73. Beilina A, Rudenko IN, Kaganovich A, *et al.*: **Unbiased screen for interactors of leucine-rich repeat kinase 2 supports a common pathway for sporadic and familial Parkinson disease.** *Proc Natl Acad Sci U S A.* 2014; 111(7): 2626–31.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 74. **F** Williams ET, Chen X, Moore DJ: **VPS35, the retromer complex and Parkinson's disease.** *J Parkinsons Dis.* 2017; 7(2): 219–33.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 75. **F** Wang W, Wang X, Fujioka H, *et al.*: **Parkinson's disease-associated mutant VPS35 causes mitochondrial dysfunction by recycling DLP1 complexes.** *Nat Med.* 2016; 22(1): 54–63.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 76. **F** Khurana V, Peng J, Chung CY, *et al.*: **Genome-scale networks link neurodegenerative disease genes to α -synuclein through specific molecular pathways.** *Cell Syst.* 2017; 4(2): 157–170.e14.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 77. **F** Chung CY, Khurana V, Yi S, *et al.*: **In situ peroxidase labeling and mass-spectrometry connects alpha-synuclein directly to endocytic trafficking and mRNA metabolism in neurons.** *Cell Syst.* 2017; 4(2): 242–250.e4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 78. Malagelada C, Jin ZH, Jackson-Lewis V, *et al.*: **Rapamycin protects against neuron death in *in vitro* and *in vivo* models of Parkinson's disease.** *J Neurosci.* 2010; 30(3): 1166–75.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 79. Moors TE, Hoozemans JJ, Ingrassia A, *et al.*: **Therapeutic potential of autophagy-enhancing agents in Parkinson's disease.** *Mol Neurodegener.* 2017; 12(1): 11.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 80. Follett J, Bugarcic A, Yang Z, *et al.*: **Parkinson disease-linked Vps35 R524W mutation impairs the endosomal association of retromer and induces α -synuclein aggregation.** *J Biol Chem.* 2016; 291(35): 18283–98.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 81. **F** Mecozzi VJ, Berman DE, Simoes S, *et al.*: **Pharmacological chaperones stabilize retromer to limit APP processing.** *Nat Chem Biol.* 2014; 10(6): 443–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 82. Salmon AB: **About-face on the metabolic side effects of rapamycin.** *Oncotarget.* 2015; 6(5): 2585–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 83. Spillantini MG, Schmidt ML, Lee VM, *et al.*: **Alpha-synuclein in Lewy bodies.** *Nature.* 1997; 388(6645): 839–40.
[PubMed Abstract](#) | [Publisher Full Text](#)
 84. Polymeropoulos MH, Lavedan C, Leroy E, *et al.*: **Mutation in the alpha-synuclein gene identified in families with Parkinson's disease.** *Science.* 1997; 276(5321): 2045–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
 85. Maroteaux L, Campanelli JT, Scheller RH: **Synuclein: a neuron-specific protein localized to the nucleus and presynaptic nerve terminal.** *J Neurosci.* 1988; 8(8): 2804–15.
[PubMed Abstract](#)
 86. Wales P, Pinho R, Lázaro DF, *et al.*: **Limelight on alpha-synuclein: pathological and mechanistic implications in neurodegeneration.** *J Parkinsons Dis.* 2013; 3(4): 415–59.
[PubMed Abstract](#) | [Publisher Full Text](#)
 87. **F** West A, Brummel BE, Braun AR, *et al.*: **Membrane remodeling and mechanics: experiments and simulations of α -synuclein.** *Biochim Biophys Acta.* 2016; 1858(7 Pt B): 1594–609.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 88. **F** Guardia-Laguarta C, Area-Gomez E, Schon EA, *et al.*: **Novel subcellular localization for α -synuclein: possible functional consequences.** *Front Neuroanat.* 2015; 9: 17.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 89. **F** Di Maio R, Barrett PJ, Hoffman EK, *et al.*: **α -Synuclein binds to TOM20 and inhibits mitochondrial protein import in Parkinson's disease.** *Sci Transl Med.* 2016; 8(342): 342ra78.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 90. **F** Kontopoulos E, Parvin JD, Feany MB: **Alpha-synuclein acts in the nucleus to inhibit histone acetylation and promote neurotoxicity.** *Hum Mol Genet.* 2006; 15(20): 3012–23.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 91. Rousseaux MW, de Haro M, Lasagna-Reeves CA, *et al.*: **TRIM28 regulates the nuclear accumulation and toxicity of both alpha-synuclein and tau.** *eLife.* 2016; 5: pii: e19809.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 92. **F** Surguchov A: **Intracellular dynamics of synucleins: "here, there and everywhere".** *Int Rev Cell Mol Biol.* 2015; 320: 103–69.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 93. **F** Rockenstein E, Nuber S, Overk CR, *et al.*: **Accumulation of oligomer-prone α -synuclein exacerbates synaptic and neuronal degeneration *in vivo*.** *Brain.* 2014; 137(Pt 5): 1496–513.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 94. Garcia-Reitböck P, Anichtchik O, Bellucci A, *et al.*: **SNARE protein redistribution and synaptic failure in a transgenic mouse model of Parkinson's disease.** *Brain.* 2010; 133(Pt 7): 2032–44.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 95. Schildknecht S, Gerding HR, Karremans C, *et al.*: **Oxidative and nitrative alpha-synuclein modifications and proteostatic stress: implications for disease mechanisms and interventions in synucleinopathies.** *J Neurochem.* 2013; 125(4): 491–511.
[PubMed Abstract](#) | [Publisher Full Text](#)
 96. Cook C, Stettler C, Petrucci L: **Disruption of protein quality control in Parkinson's disease.** *Cold Spring Harb Perspect Med.* 2012; 2(5): a009423.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 97. Valera E, Spencer B, Masliah E: **Immunotherapeutic approaches targeting amyloid- β , α -synuclein, and tau for the treatment of neurodegenerative**

- disorders. *Neurotherapeutics*. 2016; **13**(1): 179–89.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
98. George S, Brundin P: **Immunotherapy in Parkinson's disease: micromanaging alpha-synuclein aggregation.** *J Parkinsons Dis*. 2015; **5**(3): 413–24.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 99. Jankovic J, Godman I, Safirstein B, *et al.*: **Results from a phase 1b multiple ascending-dose study of PRX002, an anti-alpha-synuclein monoclonal antibody, in patients with Parkinson's disease.** *Neurodegener Dis*. 2017; **17**(suppl 1)(567). [Abstract].
 100. Wang Y: **Alzheimer disease: lessons from immunotherapy for Alzheimer disease.** *Nat Rev Neurol*. 2014; **10**(4): 188–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
 101. **F** Brahmachari S, Ge P, Lee SH, *et al.*: **Activation of tyrosine kinase c-Abl contributes to α -synuclein-induced neurodegeneration.** *J Clin Invest*. 2016; **126**(8): 2970–88.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 102. **F** Pagan F, Hebron M, Valadez EH, *et al.*: **Nilotinib effects in Parkinson's disease and dementia with Lewy bodies.** *J Parkinsons Dis*. 2016; **6**(3): 503–17.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 103. Robledo I, Jankovic J: **Media hype: patient and scientific perspectives on misleading medical news.** *Mov Disord*. 2017.
[PubMed Abstract](#) | [Publisher Full Text](#)
 104. Braak H, Del Tredici K, Rüb U, *et al.*: **Staging of brain pathology related to sporadic Parkinson's disease.** *Neurobiol Aging*. 2003; **24**(2): 197–211.
[PubMed Abstract](#) | [Publisher Full Text](#)
 105. **F** Surmeier DJ, Obeso JA, Halliday GM: **Selective neuronal vulnerability in Parkinson disease.** *Nat Rev Neurosci*. 2017; **18**(2): 101–13.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 106. Hasegawa M, Nonaka T, Masuda-Suzukake M: **α -Synuclein: experimental pathology.** *Cold Spring Harb Perspect Med*. 2016; **6**(9): pii: a024273.
[PubMed Abstract](#) | [Publisher Full Text](#)
 107. Luna E, Luk KC: **Bent out of shape: α -synuclein misfolding and the convergence of pathogenic pathways in Parkinson's disease.** *FEBS Lett*. 2015; **589**(24 Pt A): 3749–59.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 108. Kordower JH, Chu Y, Hauser RA, *et al.*: **Lewy body-like pathology in long-term embryonic nigral transplants in Parkinson's disease.** *Nat Med*. 2008; **14**(5): 504–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
 109. **F** Li JY, Englund E, Holton JL, *et al.*: **Lewy bodies in grafted neurons in subjects with Parkinson's disease suggest host-to-graft disease propagation.** *Nat Med*. 2008; **14**(5): 501–3.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 110. **F** Desplats P, Lee HJ, Bae EJ, *et al.*: **Inclusion formation and neuronal cell death through neuron-to-neuron transmission of alpha-synuclein.** *Proc Natl Acad Sci U S A*. 2009; **106**(31): 13010–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 111. Luk KC, Song C, O'Brien P, *et al.*: **Exogenous alpha-synuclein fibrils seed the formation of Lewy body-like intracellular inclusions in cultured cells.** *Proc Natl Acad Sci U S A*. 2009; **106**(47): 20051–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 112. **F** Volpicelli-Daley LA, Luk KC, Patel TP, *et al.*: **Exogenous α -synuclein fibrils induce Lewy body pathology leading to synaptic dysfunction and neuron death.** *Neuron*. 2011; **72**(1): 57–71.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 113. **F** Luk KC, Kehm V, Carroll J, *et al.*: **Pathological α -synuclein transmission initiates Parkinson-like neurodegeneration in nontransgenic mice.** *Science*. 2012; **338**(6109): 949–53.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 114. **F** Rey NL, Steiner JA, Maroof N, *et al.*: **Widespread transneuronal propagation of α -synucleinopathy triggered in olfactory bulb mimics prodromal Parkinson's disease.** *J Exp Med*. 2016; **213**(9): 1759–78.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 115. Brundin P, Ma J, Kordower JH: **How strong is the evidence that Parkinson's disease is a prion disorder?** *Curr Opin Neurol*. 2016; **29**(4): 459–66.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 116. **F** Peelaerts W, Bousset L, van der Perren A, *et al.*: **α -Synuclein strains cause distinct synucleinopathies after local and systemic administration.** *Nature*. 2015; **522**(7556): 340–4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 117. **F** Guo JL, Covell DJ, Daniels JP, *et al.*: **Distinct α -synuclein strains differentially promote tau inclusions in neurons.** *Cell*. 2013; **154**(1): 103–17.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 118. **F** Mao X, Ou MT, Karuppagounder SS, *et al.*: **Pathological α -synuclein transmission initiated by binding lymphocyte-activation gene 3.** *Science*. 2016; **353**(6307): pii: aah3374.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 119. McCann H, Cartwright H, Halliday GM: **Neuropathology of α -synuclein propagation and braak hypothesis.** *Mov Disord*. 2016; **31**(2): 152–60.
[PubMed Abstract](#) | [Publisher Full Text](#)
 120. Hansen C, Li JY: **Beyond α -synuclein transfer: pathology propagation in Parkinson's disease.** *Trends Mol Med*. 2012; **18**(5): 248–55.
[PubMed Abstract](#) | [Publisher Full Text](#)
 121. Hallett PJ, Cooper O, Sadi D, *et al.*: **Long-term health of dopaminergic neuron transplants in Parkinson's disease patients.** *Cell Rep*. 2014; **7**(6): 1755–61.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 122. Mendez I, Viñuela A, Astradsson A, *et al.*: **Dopamine neurons implanted into people with Parkinson's disease survive without pathology for 14 years.** *Nat Med*. 2008; **14**(5): 507–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 123. Irwin DJ, Abrams JY, Schonberger LB, *et al.*: **Evaluation of potential infectivity of Alzheimer and Parkinson disease proteins in recipients of cadaver-derived human growth hormone.** *JAMA Neurol*. 2013; **70**(4): 462–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 124. Su A, Gandhi R, Barlow C, *et al.*: **A practical review of gastrointestinal manifestations in Parkinson's disease.** *Parkinsonism Relat Disord*. 2017; **39**: 17–26.
[PubMed Abstract](#) | [Publisher Full Text](#)
 125. Postuma RB, Berg D: **Advances in markers of prodromal Parkinson disease.** *Nat Rev Neurol*. 2016; **12**(11): 622–34.
[PubMed Abstract](#) | [Publisher Full Text](#)
 126. Braak H, Rüb U, Gai WP, *et al.*: **Idiopathic Parkinson's disease: possible routes by which vulnerable neuronal types may be subject to neuroinvasion by an unknown pathogen.** *J Neural Transm (Vienna)*. 2003; **110**(5): 517–36.
[PubMed Abstract](#) | [Publisher Full Text](#)
 127. **F** Hill-Burns EM, Debelius JW, Morton JT, *et al.*: **Parkinson's disease and Parkinson's disease medications have distinct signatures of the gut microbiome.** *Mov Disord*. 2017; **32**(5): 739–49.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 128. **F** Malkki H: **Parkinson disease: could gut microbiota influence severity of Parkinson disease?** *Nat Rev Neurol*. 2017; **13**(2): 66–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 129. **F** Keshavarzian A, Green SJ, Engen PA, *et al.*: **Colonic bacterial composition in Parkinson's disease.** *Mov Disord*. 2015; **30**(10): 1351–60.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 130. Wood H: **Parkinson disease. Gut reactions—can changes in the intestinal microbiome provide new insights into Parkinson disease?** *Nat Rev Neurol*. 2015; **11**(2): 66.
[PubMed Abstract](#) | [Publisher Full Text](#)
 131. **F** Sampson TR, Debelius JW, Thron T, *et al.*: **Gut microbiota regulate motor deficits and neuroinflammation in a model of Parkinson's disease.** *Cell*. 2016; **167**(6): 1469–1480.e12.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 132. Parashar A, Udayabanu M: **Gut microbiota: implications in Parkinson's disease.** *Parkinsonism Relat Disord*. 2017; **38**: 1–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
 133. Braak H, Del Tredici K: **Neuropathological staging of brain pathology in sporadic Parkinson's disease: separating the wheat from the chaff.** *J Parkinsons Dis*. 2017; **7**(s1): S73–S87.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 134. **F** Liu B, Fang F, Pedersen NL, *et al.*: **Vagotomy and Parkinson disease: a Swedish register-based matched-cohort study.** *Neurology*. 2017; **88**(21): 1996–2002.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 135. **F** Svensson E, Horváth-Puhó E, Thomsen RW, *et al.*: **Vagotomy and subsequent risk of Parkinson's disease.** *Ann Neurol*. 2015; **78**(4): 522–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 136. Tarakad A, Jankovic J: **Diagnosis and management of Parkinson's disease.** *Semin Neurol*. 2017; **37**(2): 118–26.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. **David G Standaert**
, Department of Neurology, The University of Alabama at Birmingham, Birmingham, Alabama, USA
Competing Interests: No competing interests were disclosed.
2. **Roger A Barker**
University of Cambridge, Cambridge, UK
Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research