REVIEW

Coronary CT angiography in acute chest pain [version 1; peer review: 3 approved]

Nikhil Goyal , Arthur Stillman

Department of Radiology and Imaging Sciences, Emory University, Atlanta, GA, USA

V1 First published: 13 Jul 2017, 6(F1000 Faculty Rev):1125
<https://doi.org/10.12688/f1000research.11250.1>

Latest published: 13 Jul 2017, 6(F1000 Faculty Rev):1125
<https://doi.org/10.12688/f1000research.11250.1>

Abstract

Coronary computed tomographic angiography has become a reliable diagnostic tool in the evaluation of patients with chest pain. Studies have shown this modality to be accurate and safe when compared with conventional methods of assessing patients with chest pain. We review the recent developments with coronary computed tomographic angiography and devote particular attention toward its application to triage patients in the emergency department.

Keywords

coronary computed tomographic angiography, acute chest pain, ischemic disease, acute coronary syndrome

Open Peer Review

Approval Status

	1	2	3
version 1 13 Jul 2017

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Marc Dewey**, Charité – Universitätsmedizin Berlin, Berlin, Germany
2. **Pamela Douglas**, Duke University, Durham, USA
3. **Ron Blankstein**, Brigham and Women's Hospital, Boston, USA
Harvard Medical School, Boston, USA

Any comments on the article can be found at the end of the article.

Corresponding author: Nikhil Goyal (nikhil.goyal@emory.edu)

Competing interests: AS is the principal investigator for the RESCUE (Randomized Evaluation of patients with Stable angina Comparing Utilization of noninvasive Examinations) trial.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2017 Goyal N and Stillman A. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Goyal N and Stillman A. **Coronary CT angiography in acute chest pain [version 1; peer review: 3 approved]** F1000Research 2017, 6(F1000 Faculty Rev):1125 <https://doi.org/10.12688/f1000research.11250.1>

First published: 13 Jul 2017, 6(F1000 Faculty Rev):1125 <https://doi.org/10.12688/f1000research.11250.1>

Introduction

Coronary computed tomographic angiography (CCTA) has emerged in the last 15 years as a non-invasive method to evaluate the coronary arteries, cardiac chambers, and valves. The use of electrocardiography gating permits synchronization of data acquisition to the cardiac cycle to minimize motion of the coronary arteries and cardiac structures.

Recent emphasis has been on the application of CCTA in the triage of low-risk chest pain patients presenting acutely to the emergency room. Randomized trials have shown that coronary angiography in the emergency department (ED) setting reduces length of stay, initial cost in the ED, and time to discharge when compared with the current standard of care with equivalent safety and accuracy. However, there are data showing that the initial cost savings may be offset by additional downstream testing and sometimes overall higher radiation dose when CCTA is used. Despite this, the potential use of CCTA in the ED setting is promising, especially given some of the newest developments that may curb the need for additional testing.

With advancements such as increasing number of detectors and x-ray tubes as well as advances in software (perfusion imaging and computed tomographic–fractional flow reserve, or CT-FFR), CCTA may offer the opportunity of having a “one-stop shop” for evaluating select patient populations.

Background and evolution of coronary computed tomographic angiography

Current CT scanners used for coronary angiography have superior spatial and temporal resolution when compared with older-model scanners with acceptable radiation exposure. With detector ranges of 64 to 320 detectors and a detector width of 0.5 to 0.7 mm, current scanners can provide wide coverage and thin slice acquisition. With the advent of dual-source scanners and the use of multi-segment reconstruction algorithms, temporal resolution has also improved, thus allowing scanning of faster and (if necessary) irregular heart rates. However, it should be stated that the use of beta-blockers remains common practice and should be used when necessary to obtain optimal images at the lowest possible radiation dose. The accuracy of CCTA when compared with conventional angiography has been well established in multi-center trials as well as in meta-analysis studies. The high negative predictive value (NPV) of 83–99%^{1–3} and the corresponding negative likelihood ratio of 0.02⁴ most importantly can rule out obstructive coronary artery disease or myocardial ischemia. The positive predictive value (PPV) of CCTA to detect myocardial ischemia is less⁵, and about 50% of cases with plaque causing more than 50% narrowing have an abnormal result on myocardial perfusion imaging or demonstrate ischemia by FFR. Thus, the strength of this modality currently lies in its ability to rule out ischemic disease rather than rule it in.

In addition to investigating its diagnostic accuracy, several studies have investigated the prognostic efficacy of CCTA. Computed tomographic angiography (CTA) is a good predictor of risk, and its most notable feature is predicting excellent outcomes for people with normal results. Long-term data now exist for

excellent prognosis beyond 5 years for patients with normal CTA results with a negative likelihood ratio of 0.008 for major adverse cardiac event (MACE) for a normal CTA result^{6–9}. Not surprisingly, risk for MACE increases incrementally as plaque burden increases^{6–9} and the more proximal a plaque is within the coronary tree, the worse the prognosis.^{5,10,11} This has been documented up to 7 years post-scan¹². Overall, these findings are not unexpected and reflect what has already been known from conventional angiography.

The chest pain problem

While the role of CTA in the outpatient setting has been established, much of the recent work has centered on the use of CTA in patients presenting to the ED with acute chest pain. The problem of acute chest pain is a major source of resource utilization in the ED. It is estimated that over 8 million visits to the emergency room are for chest pain¹³. The current model of admitting patients to the hospital to “rule out acute coronary syndrome (ACS)” with serial biomarkers and functional assessment is both time- and cost-inefficient and leads to overcrowding in hospitals. The majority of patients who are admitted under these circumstances do not have ACS. Additionally, a small fraction of patients are incorrectly discharged¹³ with a corresponding increase in mortality¹⁴. Given that the strength of CCTA is its high NPV, several investigators have applied the use of CCTA to low- to intermediate-risk chest pain patients presenting to the emergency room.

To date, three large clinical trials have demonstrated that CCTA is a safe and effective means to discharge low-risk patients from the ED^{15–17}. The CT-STAT (Coronary Computed Tomographic Angiography for Systematic Triage of Acute Chest Pain Patients to Treatment) trial demonstrated a 54% reduction in time to diagnosis and a 38% reduction in ED costs between CCTA and myocardial perfusion imaging with no difference in a safety endpoint of MACE at 60 days¹⁵. Litt *et al.*, in the ACRIN-PA trial¹⁶, demonstrated a 50% increase in discharge home and no increase in MACE at 30 days in patients with a normal CCTA when comparing CCTA with standard of care. Finally, the ROMICAT II (Rule Out Myocardial Infarction/Ischemia Using Computer Assisted Tomography II) trial demonstrated decreased length of stay (8 versus 26 hours), fourfold increase in rates of direct discharge from the ED, and no difference in MACE at 28 days¹⁷. However, it should be noted that cumulative costs at 28 days were higher in the CCTA group, as more patients in the CTA group underwent additional tests after initial discharge. A meta-analysis by Hulten *et al.* demonstrated that the use of CCTA in the ED led to an overall increase in downstream use of conventional angiography and revascularization by 2% when compared with usual care¹⁸. This is likely because of the high sensitivity of CTA in finding obstructive disease that required further testing to clarify whether these lesions were flow-limiting. In patients with subclinical atherosclerosis identified on CCTA in the ED, there may be potential benefit in the form of increased use of preventive measures, although initiating these measures remains challenging in the ED setting.

Longer-term data now exist for the “warranty” period of a “normal” CCTA reading for patients presenting to the ED.

Hollander *et al.* showed that, at 1 year following a CTA in the ED, none of the 481 patients with less than 50% stenosis and an ejection fraction of more than 30% had a non-fatal myocardial infarction or needed revascularization¹⁹. Schlett *et al.* published 2-year data for patients from the original ROMICAT trial and showed that patients with a normal ED CTA had a 0% 2-year risk for MACE²⁰. It is important to recognize that patients with non-obstructive coronary disease in this study had a 4% 2-year risk for MACE risk (which is non-trivial and requires appropriate therapy).

Additional data also show that there are reduced return visits to the ED in patients who undergo CCTA²¹. In the 2010 appropriate use criteria, CCTA is considered an appropriate indication for acute chest pain in low- to intermediate-risk patients.

It is fairly evident that patients who have normal (or near normal) CTA results can be safely discharged from the ED as not having ACS. CCTA has an NPV of nearly 100% in low-risk patients in this patient group^{22,23}. Meta-analysis demonstrates an associated negative likelihood ratio of 0.06²⁴. Most physicians also agree that patients with obstructive disease should go on for further testing. However, it remains unclear what to do with patients who exhibit intermediate lesions that *may* be flow-limiting. A lot of recent focus has centered on obtaining functional data as part of the CT exam in order to further assess these “intermediate” lesions. After all, combined anatomical assessment of the coronaries and their corresponding effect on perfusion remain the most comprehensive and desirable evaluation of the heart. The two forms of investigation recently developed to address this issue are CT-FFR and CT perfusion scanning.

In the catheterization lab, intermediate lesions are assessed by passing a pressure transducer wire across the lesion and recording pressures distal and proximal to the lesion in question. Adenosine is administered to induce hyperemia. This process is called measuring FFR. This is important because FFR measurement has been shown to better guide percutaneous coronary intervention therapy with more favorable outcomes than angiographic appearance alone²⁵. For reference, an FFR value below 0.80 is considered flow-limiting.

Given the invasive nature of conventional FFR, investigators sought out a non-invasive equivalent^{26–28}. The physics of simulating invasive pressure measurements is complex and involves the application of computational fluid dynamics equations to blood flowing within the coronary vessels. CT output data can be transmitted to supercomputers which can perform the necessary mathematics and generate a CT-FFR measurement. Two large trials have investigated the effect of computing FFR measurements from CTA data on sensitivity, specificity, PPV, and NPV profiles of CTA while using invasive FFR as the gold-standard comparison. The DISCOVER-FLOW (Diagnosis of Ischemia-Causing Stenoses Obtained Via Noninvasive Fractional Flow Reserve) trial²⁹ demonstrated improvements in specificity (25→82%) and PPV (58→85%). The NXT trial³⁰ showed improvements in specificity and PPV (32→84% and 33→67%, respectively).

Since then, several vendor- or institutional-specific FFR computational programs have been devised and some have been tested in single-center trials with similar results. Overall, the data from the multi-center and single-center local software trials demonstrate improvements in the area under the receiver-operator characteristic curves when evaluating CT-FFR versus CTA alone compared with invasive FFR. However, it should be noted that the computational power currently required to generate CT-FFR results necessitates a long (for example, up to 24 hours by some vendors) calculation time and thus is not suitable for use in EDs. Further work may reveal faster computation algorithms, making this modality more useful in the ED setting.

Hemodynamic evaluation via assessment of myocardial perfusion is also a rapidly developing area of study. CT perfusion (CTP), like a provocative nuclear test, involves both a rest and a stress phase, and several vasodilator agents, including dipyridamole, adenosine, and regadenoson, can be used. Myocardial perfusion is determined by visual analysis assessing for areas of myocardial hypoattenuation, although several semi-quantitative techniques are now available. Numerous studies have shown the efficacy of perfusion CT alone with sensitivity ranges from 50% to 96%, specificity ranges from 68% to 98%, and PPV and NPV ranges from 55% to 94% and from 79% to 98%, respectively, using various modalities as the reference gold standard²⁶. When combined with CCTA for corresponding anatomic evaluation, CTA+CTP demonstrates a higher sensitivity and specificity profile than CTA alone^{31–34}. It is important to consider overall radiation dose when considering CT hemodynamic assessment. CT-FFR requires no additional acquisitions but CTP does. Additional requirements of contrast, along with scanning twice (rest and stress), and drug administration should also be considered. In particular, the effective workflow requires immediate review of the images so that stress images are obtained only when necessary. Weininger *et al.*³⁵ have demonstrated that this is feasible in the ED setting. Nevertheless, for the above-mentioned reasons, implementing stress CTP in the ED setting may be logistically difficult.

There is a significant amount of evidence demonstrating that myocardial infarctions occur not only from severely stenotic vessels but from rupture of “unstable” plaque regardless of the degree of stenosis. It has been shown that plaque features in addition to degree of stenosis are predictors of MACE. But what features make a plaque unstable? Recent work has also centered on identifying plaque characteristics that may deem it “high risk” for rupture. Features such as positive remodeling, low attenuation, and spotty calcification are associated with higher probability of having an acute coronary syndrome³⁶.

Along with recent advances in imaging, there have been advances in the clinical evaluation of patients with chest pain. Improved biomarkers, in the form of high-sensitivity troponins, are under investigation in many centers and may be in widespread use in the coming years. These biomarkers demonstrate a higher sensitivity than conventional troponin assays and have a higher NPV for

excluding myocardial infarction within 3 hours^{37,38}. At least one study³⁹ has shown that length of stay and rates of direct discharge from the ED are equivalent between CCTA and standard of care if high-sensitivity troponins are used in triage.

Conclusions

CCTA is emerging as a complementary modality in the workup of patients with chest pain. Advances in CT hardware along with software have led to our understanding beyond just anatomical evaluation of stenosis. The ultimate goal of both anatomic and hemodynamic assessment of the coronaries by CT may be widely realized in the not-so-distant future. This will be particularly

important in the setting of evaluating patients presenting with acute chest pain in the ED.

Competing interests

AS is the principal investigator for the RESCUE (Randomized Evaluation of patients with Stable angina Comparing Utilization of noninvasive Examinations) trial.

Grant information

The author(s) declared that no grants were involved in supporting this work.

References

- Budoff MJ, Dowe D, Jollis JG, *et al.*: **Diagnostic performance of 64-multidetector row coronary computed tomographic angiography for evaluation of coronary artery stenosis in individuals without known coronary artery disease: results from the prospective multicenter ACCURACY (Assessment by Coronary Computed Tomographic Angiography of Individuals Undergoing Invasive Coronary Angiography) trial.** *J Am Coll Cardiol.* 2008; 52(21): 1724–32. [PubMed Abstract](#) | [Publisher Full Text](#)
- Miller JM, Rochitte CE, Dewey M, *et al.*: **Diagnostic performance of coronary angiography by 64-row CT.** *N Engl J Med.* 2008; 359(22): 2324–36. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Meijboom WB, Meijis MF, Schuijf JD, *et al.*: **Diagnostic accuracy of 64-slice computed tomography coronary angiography: a prospective, multicenter, multivendor study.** *J Am Coll Cardiol.* 2008; 52(25): 2135–44. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Li M, Du XM, Jin ZT, *et al.*: **The diagnostic performance of coronary artery angiography with 64-MSCT and post 64-MSCT: systematic review and meta-analysis.** *PLoS One.* 2014; 9(1): e84937. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Min JK, Shaw LJ, Devereux RB, *et al.*: **Prognostic value of multidetector coronary computed tomographic angiography for prediction of all-cause mortality.** *J Am Coll Cardiol.* 2007; 50(12): 1161–70. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Andreini D, Pontone G, Mushtaq S, *et al.*: **A long-term prognostic value of coronary CT angiography in suspected coronary artery disease.** *JACC Cardiovasc Imaging.* 2012; 5(7): 690–701. [PubMed Abstract](#) | [Publisher Full Text](#)
- Ostrom MP, Gopal A, Ahmadi N, *et al.*: **Mortality incidence and the severity of coronary atherosclerosis assessed by computed tomography angiography.** *J Am Coll Cardiol.* 2008; 52(16): 1335–43. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Fazel P, Peterman MA, Schussler JM: **Three-year outcomes and cost analysis in patients receiving 64-slice computed tomographic coronary angiography for chest pain.** *Am J Cardiol.* 2009; 104(4): 498–500. [PubMed Abstract](#) | [Publisher Full Text](#)
- Hulten EA, Carbonaro S, Petrillo SP, *et al.*: **Prognostic value of cardiac computed tomography angiography for prediction of cardiac events in patients with suspected coronary artery disease: a systematic review and meta-analysis.** *J Am Coll Cardiol.* 2011; 57(10): 1237–47. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Min JK, Shaw LJ, Berman DS: **The present state of coronary computed tomography angiography a process in evolution.** *J Am Coll Cardiol.* 2010; 55(10): 957–65. [PubMed Abstract](#) | [Publisher Full Text](#)
- Hadamitzky M, Freissmuth B, Meyer T, *et al.*: **Prognostic value of coronary computed tomographic angiography for prediction of cardiac events in patients with suspected coronary artery disease.** *JACC Cardiovasc Imaging.* 2009; 2(4): 404–11. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Dougoud S, Fuchs TA, Stehli J, *et al.*: **Prognostic value of coronary CT angiography on long-term follow-up of 6.9 years.** *Int J Cardiovasc Imaging.* 2014; 30(5): 969–76. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Seifarth H, Schlett CL, Truong QA, *et al.*: **Cardiac CT in the emergency department.** *Cardiol Clin.* 2012; 30(1): 117–33. [PubMed Abstract](#) | [Publisher Full Text](#)
- Pope JH, Aufderheide TP, Ruthazer R, *et al.*: **Missed diagnoses of acute cardiac ischemia in the emergency department.** *N Engl J Med.* 2000; 342(16): 1163–70. [PubMed Abstract](#) | [Publisher Full Text](#)
- Goldstein JA, Chinnaiyan KM, Abidov A, *et al.*: **The CT-STAT (Coronary Computed Tomographic Angiography for Systematic Triage of Acute Chest Pain Patients to Treatment) trial.** *J Am Coll Cardiol.* 2011; 58(14): 1414–22. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Litt HI, Gatsonis C, Snyder B, *et al.*: **CT angiography for safe discharge of patients with possible acute coronary syndromes.** *N Engl J Med.* 2012; 366(15): 1393–403. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Hoffmann U, Truong QA, Schoenfeld DA, *et al.*: **Coronary CT angiography versus standard evaluation in acute chest pain.** *N Engl J Med.* 2012; 367(4): 299–308. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Hulten E, Pickett C, Bittencourt MS, *et al.*: **Outcomes after coronary computed tomography angiography in the emergency department: a systematic review and meta-analysis of randomized, controlled trials.** *J Am Coll Cardiol.* 2013; 61(8): 880–92. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Hollander JE, Chang AM, Shofer FS, *et al.*: **One-year outcomes following coronary computerized tomographic angiography for evaluation of emergency department patients with potential acute coronary syndrome.** *Acad Emerg Med.* 2009; 16(8): 693–8. [PubMed Abstract](#) | [Publisher Full Text](#)
- Schlett CL, Banerji D, Siegel E, *et al.*: **Prognostic value of CT angiography for major adverse cardiac events in patients with acute chest pain from the emergency department: 2-year outcomes of the ROMICAT trial.** *JACC Cardiovasc Imaging.* 2011; 4(5): 481–91. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- El-Hayek G, Benjo AM, Cavalcante JL, *et al.*: **The Use of Coronary Computed Tomography Angiography in Chest Pain Patients in the Emergency Departments Reduces Future Hospital Admissions and Does Not Impact the Need for Cardiac Catheterizations: A busting meta-analysis.** *Circulation: Cardiovascular Quality and Outcomes.* 6(3 Supplement): A268. [Reference Source](#)
- Miller AH, Pepe PE, Peshock R, *et al.*: **Is coronary computed tomography angiography a resource sparing strategy in the risk stratification and evaluation of acute chest pain? Results of a randomized controlled trial.** *Acad Emerg Med.* 2011; 18(5): 458–67. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Cury RC, Feuchtner GM, Battle JC, *et al.*: **Triage of patients presenting with chest pain to the emergency department: implementation of coronary CT angiography in a large urban health care system.** *AJR Am J Roentgenol.* 2013; 200(1): 57–65. [PubMed Abstract](#) | [Publisher Full Text](#)
- Samad Z, Hakeem A, Mahmood SS, *et al.*: **A meta-analysis and systematic review of computed tomography angiography as a diagnostic triage tool for patients with chest pain presenting to the emergency department.** *J Nucl Cardiol.* 2012; 19(2): 364–76. [PubMed Abstract](#) | [Publisher Full Text](#)
- Tonino PA, De Bruyne B, Pijls NH, *et al.*: **Fractional flow reserve versus angiography for guiding percutaneous coronary intervention.** *N Engl J Med.* 2009; 360(3): 213–24. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)

26. Min JK, Leipsic J, Pencina MJ, *et al.*: **Diagnostic accuracy of fractional flow reserve from anatomic CT angiography.** *JAMA*. 2012; **308**(12): 1237–45.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
27. Nakanishi R, Budoff MJ: **Noninvasive FFR derived from coronary CT angiography in the management of coronary artery disease: technology and clinical update.** *Vasc Health Risk Manag*. 2016; **12**: 269–78.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
28. Hecht HS, Narula J, Fearon WF: **Fractional Flow Reserve and Coronary Computed Tomographic Angiography: A Review and Critical Analysis.** *Circ Res*. 2016; **119**(2): 300–16.
[PubMed Abstract](#) | [Publisher Full Text](#)
29. **F** Koo BK, Erglis A, Doh JH, *et al.*: **Diagnosis of ischemia-causing coronary stenoses by noninvasive fractional flow reserve computed from coronary computed tomographic angiograms. Results from the prospective multicenter DISCOVER-FLOW (Diagnosis of Ischemia-Causing Stenoses Obtained Via Noninvasive Fractional Flow Reserve) study.** *J Am Coll Cardiol*. 2011; **58**(19): 1989–97.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
30. **F** Norgaard BL, Leipsic J, Gaur S, *et al.*: **Diagnostic performance of noninvasive fractional flow reserve derived from coronary computed tomography angiography in suspected coronary artery disease: the NXT trial (Analysis of Coronary Blood Flow Using CT Angiography: Next Steps).** *J Am Coll Cardiol*. 2014; **63**(12): 1145–55.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
31. **F** Varga-Szemes A, Meinel FG, De Cecco CN, *et al.*: **CT myocardial perfusion imaging.** *AJR Am J Roentgenol*. 2015; **204**(3): 487–97.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
32. **F** Osawa K, Miyoshi T, Koyama Y, *et al.*: **Additional diagnostic value of first-pass myocardial perfusion imaging without stress when combined with 64-row detector coronary CT angiography in patients with coronary artery disease.** *Heart*. 2014; **100**(13): 1008–15.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
33. Rocha-Filho JA, Blankstein R, Shturman LD, *et al.*: **Incremental value of adenosine-induced stress myocardial perfusion imaging with dual-source CT at cardiac CT angiography.** *Radiology*. 2010; **254**(2): 410–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
34. Nasis A, Ko BS, Leung MC, *et al.*: **Diagnostic accuracy of combined coronary angiography and adenosine stress myocardial perfusion imaging using 320-detector computed tomography: pilot study.** *Eur Radiol*. 2013; **23**(7): 1812–21.
[PubMed Abstract](#) | [Publisher Full Text](#)
35. Weininger M, Schoepf UJ, Ramachandra A, *et al.*: **Adenosine-stress dynamic real-time myocardial perfusion CT and adenosine-stress first-pass dual-energy myocardial perfusion CT for the assessment of acute chest pain: initial results.** *Eur J Radiol*. 2012; **81**(12): 3703–10.
[PubMed Abstract](#) | [Publisher Full Text](#)
36. **F** Puchner SB, Liu T, Mayrhofer T, *et al.*: **High-risk plaque detected on coronary CT angiography predicts acute coronary syndromes independent of significant stenosis in acute chest pain: results from the ROMICAT-II trial.** *J Am Coll Cardiol*. 2014; **64**(7): 684–92.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
37. **F** Reichlin T, Hochholzer W, Bassetti S, *et al.*: **Early diagnosis of myocardial infarction with sensitive cardiac troponin assays.** *N Engl J Med*. 2009; **361**(9): 858–67.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
38. Keller T, Zeller T, Peetz D, *et al.*: **Sensitive troponin I assay in early diagnosis of acute myocardial infarction.** *N Engl J Med*. 2009; **361**(9): 868–77.
[PubMed Abstract](#) | [Publisher Full Text](#)
39. **F** Dedic A, Lubbers MM, Schaap J, *et al.*: **Coronary CT Angiography for Suspected ACS in the Era of High-Sensitivity Troponins: Randomized Multicenter Study.** *J Am Coll Cardiol*. 2016; **67**(1): 16–26.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. **Ron Blankstein**

¹ Brigham and Women's Hospital, Boston, MA, USA

² Harvard Medical School, Boston, MA, USA

Competing Interests: No competing interests were disclosed.

2. **Pamela Douglas**

Department of Medicine, Duke University, Durham, NC, USA

Competing Interests: No competing interests were disclosed.

3. **Marc Dewey**

Department of Radiology, Charité – Universitätsmedizin Berlin, Berlin, Germany

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research