RESEARCH ARTICLE

REVISED Optimal cutoff for the evaluation of insulin resistance through triglyceride-glucose index: A cross-sectional study in a Venezuelan population [version 2; peer review: 2 approved]

Juan Salazar ¹, Valmore Bermúdez ^{1,2}, María Calvo¹, Luis Carlos Olivar¹, Eliana Luzardo¹, Carla Navarro¹, Heysa Mencia¹, María Martínez¹, José Rivas-Ríos¹, Sandra Wilches-Durán², Marcos Cerda², Modesto Graterol², Rosemily Graterol², Carlos Garicano², Juan Hernández², Joselyn Rojas^{1,3}

¹Endocrine-Metabolic Research Center, , University of Zulia, Maracaibo, Venezuela

²Grupo de Investigación Altos Estudios de Frontera (ALEF), Universidad Simón Bolívar, Cúcuta, Colombia

³Division of Pulmonary and Critical Care Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA, 02115, USA

V2 First published: 07 Aug 2017, 6:1337
<https://doi.org/10.12688/f1000research.12170.1>
 Second version: 20 Oct 2017, 6:1337
<https://doi.org/10.12688/f1000research.12170.2>
 Latest published: 30 Jan 2018, 6:1337
<https://doi.org/10.12688/f1000research.12170.3>

Abstract

Background: Insulin resistance (IR) evaluation is a fundamental goal in clinical and epidemiological research. However, the most widely used methods are difficult to apply to populations with low incomes. The triglyceride-glucose index (TGI) emerges as an alternative to use in daily clinical practice. Therefore the objective of this study was to determine an optimal cutoff point for the TGI in an adult population from Maracaibo, Venezuela.

Methods: This is a sub-study of Maracaibo City Metabolic Syndrome Prevalence Study, a descriptive, cross-sectional study with random and multi-stage sampling. For this analysis, 2004 individuals of both genders ≥ 18 years old with basal insulin determination and triglycerides < 500 mg/dl were evaluated.. A reference population was selected according to clinical and metabolic criteria to plot ROC Curves specific for gender and age groups to determine the optimal cutoff point according to sensitivity and specificity. The TGI was calculated according to the equation: $\ln [\text{Fasting triglyceride (mg / dl)} \times \text{Fasting glucose (mg / dl)}] / 2$.

Results: The TGI in the general population was 4.6 ± 0.3 (male: 4.66 ± 0.34 vs. female: 4.56 ± 0.33 , $p = 8.93 \times 10^{-10}$). The optimal cutoff point was 4.49, with a sensitivity of 82.6% and specificity of 82.1% (AUC=0.889, 95% CI: 0.854-0.924). There were no significant differences in the predictive capacity of the index when evaluated according to gender and age groups. Those individuals with $\text{TGI} \geq 4.5$

Open Peer Review

Approval Status

	1	2
version 3 (revision) 30 Jan 2018		
version 2 (revision) 20 Oct 2017		 view
version 1 07 Aug 2017	 view	

1. **Silvia Fabiana Benozzi**, Universidad Nacional del Sur, Bahía Blanca, Argentina

Gisela Unger, Universidad Nacional del Sur, Bahía Blanca, Argentina

2. **Muthuswamy Balasubramanyam** ,
 Madras Diabetes Research Foundation
 (MDRF), Chennai, India

Any reports and responses or comments on the article can be found at the end of the article.

had higher HOMA2-IR averages than those with TGI <4.5 (2.48 vs 1.74, respectively, $p < 0.001$).

Conclusions: The TGI is a measure of interest to identify IR in the general population. We propose a single cutoff point of 4.5 to classify individuals with IR. Future studies should evaluate the predictive capacity of this index to determine atypical metabolic phenotypes, type 2 diabetes mellitus and even cardiovascular risk in our population.

Keywords

insulin resistance, triglycerides, glycemia, Metabolic syndrome, cutoff

Corresponding authors: Juan Salazar (juanjsv18@hotmail.com), Valmore Bermúdez (valmore@gmail.com)

Author roles: **Salazar J:** Formal Analysis, Writing – Original Draft Preparation; **Bermúdez V:** Conceptualization, Writing – Review & Editing; **Calvo M:** Data Curation, Investigation; **Olivar LC:** Data Curation, Investigation; **Luzardo E:** Data Curation, Investigation; **Navarro C:** Data Curation, Investigation; **Mencia H:** Investigation; **Martínez M:** Data Curation, Investigation; **Rivas-Ríos J:** Investigation, Writing – Original Draft Preparation; **Wilches-Durán S:** Funding Acquisition, Writing – Review & Editing; **Cerda M:** Funding Acquisition, Writing – Original Draft Preparation; **Graterol M:** Formal Analysis, Funding Acquisition; **Graterol R:** Funding Acquisition, Writing – Original Draft Preparation; **Garicano C:** Investigation, Writing – Original Draft Preparation; **Hernández J:** Formal Analysis, Investigation; **Rojas J:** Conceptualization, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by the Technological, Humanistic, and Scientific Development Council (Consejo de Desarrollo Científico, Humanístico y Tecnológico; CONDES), University of Zulia (grant n° CC-0437-10-21-09-10).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2017 Salazar J *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. Data associated with the article are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

How to cite this article: Salazar J, Bermúdez V, Calvo M *et al.* **Optimal cutoff for the evaluation of insulin resistance through triglyceride-glucose index: A cross-sectional study in a Venezuelan population [version 2; peer review: 2 approved]** F1000Research 2017, 6:1337 <https://doi.org/10.12688/f1000research.12170.2>

First published: 07 Aug 2017, 6:1337 <https://doi.org/10.12688/f1000research.12170.1>

REVISED Amendments from Version 1

We want to thank the reviewers for the valuable time spent in the analysis and review of our manuscript. In regards of the given suggestions:

1. The typos in the abstract were corrected
2. Keywords were added
3. The abbreviations section were deleted
4. As interpreted, all criteria were necessary to classify subjects as "healthy" or "unhealthy" This was specified in the text and [Figure 1](#)
5. Figure 1 legend was adjusted as suggested
6. The fasting time and the coefficients of variation of glycemia and insulin were described
7. ROC curve formula was edited
8. In all tables and figures the adjustment was made relative to the reference population
9. In the results by age, reference is made to the similarity between cut-off points. It is unnecessary to obtain the same result for each age group
10. The "n" value was added in the corresponding tables
11. The effect size analysis did not show significant results by age and sex, which shows that it is not necessary to determine specific cutoff points according to these characteristics in our population. This was already described in discussion section. However, factors related to the index should be evaluated in the next analysis
12. The "p" values were adjusted to 3 decimals. The word "indices" were unified and grammatical recommendations were added

Finally, we would like to thank our referees once again for their valuable suggestions.

See referee reports

Introduction

Insulin resistance (IR) is a metabolic condition in which insulin-dependent tissues become less sensitive to insulin action, leading to an imbalance in the metabolism of carbohydrates, lipids and proteins¹. This condition is caused by the influence of different risk factors in the population, such as aging, alcohol consumption, smoking, hypercaloric diets, sedentary lifestyle and obesity². Its role in the development of different pathologies, as cardiovascular disease (CVD)³ and cerebrovascular disease⁴, is now recognized, as well as playing an important role in the pathogenesis and clinical outcomes of the metabolic syndrome (MS)⁵ and type 2 diabetes mellitus (DM2)⁶.

In 1963, Randle and colleagues were among the first to investigate the pathophysiology of IR⁷, suggesting the elevation of free fatty acids (FFA) in the splenic circulation as the cornerstone of this disorder. They proposed the glucose fatty acid cycle, called the Randle cycle. Years later, a theory proposed by Shulman⁸ surfaced, which continued to support the role of FFA in the pathophysiology of IR. However, these authors suggested that FFA and its products, such as diacylglycerol, acyl-coA and ceramides, activate serin-threonine kinases, which phosphorylate important proteins, inhibit the

insulin signaling pathway and subsequently translocated glucose transporter 4 (GLUT4) to the plasma membrane⁸.

Although many aspects remain to be clarified in the pathophysiology of IR, its long-term complications have generated great interest in the determination of ideal methods that allow the promotion of an early and accurate diagnosis in risky populations⁹. In this sense, the Euglycemic-Hyperinsulinemic Clamp is considered the gold standard for the determination of IR¹⁰, but the high cost and impracticability of this method has promoted the development of new techniques for the estimation of insulin sensitivity. Many mathematical models have been proposed in recent years with the objective of simplifying the measurement of IR¹¹, highlighting the Homeostasis Model Assessment (HOMA-IR), a validated method to measure IR from serum glucose and fasting serum insulin¹². This index has been studied in our population, which was conducted in 2026 subjects and evaluated the factors related to insulin resistance (defined as $HOMA2-IR=2$)¹³. However, one of the biggest limitations is the lack of accessibility to populations with lower incomes, since all the individuals require insulin blood tests.

Simental-Mendia *et al.*^{14,15} have proposed and validated a new formula to evaluate IR from the levels of serum triglycerides (TAG) and fasting glucose (FG), which is known as 'Triglyceride/Glucose Index' (TGI), this formula is a potential diagnostic tool when other standard methods are not available. Based on the information above, the objective of the study is to determine an optimal cut point of the TGI to determine IR and to evaluate the behavior according to the main sociodemographic characteristics in an adult population from Maracaibo, Zulia, Venezuela.

Methods

Sample selection and study design

The Maracaibo City Metabolic Syndrome Prevalence Study (MMSPS) study is a descriptive, cross-sectional, randomized, multi-stage sampling study that was carried out in Maracaibo-Venezuela; the second most populated city in the country with an approximate population of 2,500,000, in order to evaluate the cardiovascular and metabolic risk factors of this locality, during the period May 2007 - December 2009¹⁶. The only inclusion criteria was individuals older than 18 years. The sample (1,986 individuals) was calculated based on the estimates of the population in the city given by the National Institute of Statistics (1,428,043 inhabitants for 2007). A total of 244 individuals (12%) were added through oversampling, in order to increase the accuracy of the estimates obtained from the smaller subgroups of the sample, representing a total of 2230 individuals of both genders. Sampling details have been previously published¹⁶. The study was approved by the Bioethics Committee of the Endocrine and Metabolic Research Center – University of Zulia (approval number: BEC-006-0305). This ethical approval included all future studies that used the data from the MMSPS. All participants signed written consent before being questioned and physically examined by a trained team.

For this analysis, 2004 individuals were selected according to the availability of baseline insulin, in addition to the exclusion of individuals with TAG $\geq$ 500 mg/dl. Based on this sub-sample, a reference population was selected based on all the following criteria: abdominal obesity, total cholesterol, high blood pressure and personal

history of DM2, coronary artery diseases, cardiac arrhythmias, acute cerebrovascular disease, and polycystic ovaries; with the purpose of establishing a subsample of healthy and unhealthy subjects without using definitions or diagnostic criteria that include TAG and glycaemia values to avoid variables correlation *a priori* (Figure 1).

Individual evaluation

The data was gathered through a complete medical history performed by the trained team: the past medical and family history for cardiovascular and endocrine-metabolic diseases was assessed.

Blood pressure and anthropometric evaluation

The auscultation method was used for the measurement of blood pressure, using stethoscopes and calibrated sphygmomanometers adequately validated. The procedure was performed with the individual at rest (at least 15 minutes) and sitting with both feet on the floor; three measurements were taken, with 15 minutes of separation between one measurement and the other. Systolic blood pressure was determined by auscultation of the first Korotkoff noise, while diastolic blood pressure was determined on auscultation of the fifth Korotkoff noise.

Weight was determined through a dielectric balance (Tanita, TBF-310 GS Body Composition Analyzer, Tokyo, Japan), and height was obtained by using vertical tape calibrated in centimeters and millimeters. Individuals were standing and barefoot, with light

clothing throughout the evaluation, maintaining a straight posture and head up. Body Mass Index was calculated using the weight/height formula and classified according to the criteria proposed by the World Health Organisation (low weight, normal weight, overweight, Obesity type I, II and III)¹⁷. The measurement of the abdominal circumference was taken with a plastic metric tape in centimeters at equidistant points between the costal ridge and the iliac crest, according to the protocol proposed by the National Salute Institute of the United States¹⁸. Abdominal obesity was defined according to specific cutoff points for our previously determined population, ≥ 90 cm in women and ≥ 95 cm in men¹⁹.

Laboratory analysis

Blood samples were collected between 8:00 A.M. and 10:00 A.M. after an overnight fast (~10 hours). Determination of glucose, total cholesterol, triglycerides, and HDL-C was done with an automated analyzer (Human Gesellschaft für Biochemica und Diagnostica mbH, Germany). The intra-assay coefficient of variation for total cholesterol, TAG, glucose and insulin was 3%, 5%, 3%, and <10%, respectively. Insulin was determined using an ultrasensitive ELISA double-sandwich method (DRG Instruments GmbH, Germany, Inc.), with a Detection Limit <1 mU/L. The HOMA2IR index was calculated using the software administered by the Oxford Diabetes Center, Endocrinology and Metabolism (available at <http://www.dtu.ox.ac.uk/homacalculator/index.php>). A cutoff value of ≥ 2 was used to determine IR²⁰.

†Abdominal obesity defined as: abdominal circumference > 90 cm in the female sex and > 95 cm in the male sex

Figure 1. Diagram of reference population selection for cut-off points determination of the triglyceride-glucose index in Maracaibo city, Venezuela. ROC Curves: Receiver operating characteristic curves.

Calculation of TGI

The calculation of the TGI was done using the equation: $\ln [\text{Fasting TAG (mg/dl)} \times \text{FG (mg/dl)}] / 2$; thus being expressed on a logarithmic scale^{14,21}.

Statistical analysis

Qualitative variables were expressed in absolute and relative frequencies. While the quantitative variables were expressed as arithmetic mean $\pm$ SD, with the previous analysis of normality through means of a Geary test. Significant differences between groups were assessed using the Student's *t*-test, while an ANOVA was used for comparisons between three or more groups. Data were analyzed through the SPSS v.21 for Windows (IBM Chicago, IL), considering statistically significant results when $p < 0.05$.

ROC curves were plotted in the reference population (Figure 1) to analyze the predictive capacity and to determine an optimal cutoff point for the TGI. ROC curves were gender-specific using R version 3.4.1. Several indices were calculated to evaluate the optimum cutoff point in the curve. The area under the curve (AUC) is used to establish the ability of the test to obtain an appropriate cutoff where an AUC of 1.00 is considered a perfect diagnostic test²². Comparisons between AUC were performed using Delong's test²³. The Youden Index (J) was calculated using the formula $[J = \text{sensitivity} + \text{specificity} - 1 = S - (1 - Es)]^{23}$, obtaining the value

of true positives (sensitivity) and false positives (1-specificity) when $J > 1$. The minimum cutoff point was calculated using the nearest point to 0.1 in the ROC curves formula: square root of $[(1 - \text{sensitivity})^2 + (1 - \text{specificity})^2]^{24}$. In addition, probability radius positive [sensitivity/1-specificity] and negative [1-sensitivity/specificity] were calculated to aid in the selection of the cutoff point together with the Youden index. Likelihood values > 1 indicate association with the disease, while those < 1 indicate association with the absence of the disease²⁵.

Results

General characteristics of the sample

A total of 2004 individuals were studied, 53.4% ($n = 1050$) were female, the mean age of the population was 39.6 ± 15.3 . The general characteristics of the population are shown in Table 1. The mean TGI in the general population was 4.6 ± 0.3 , with higher values in males (males: 4.66 ± 0.34 vs. females: 4.56 ± 0.33 , $p = 8.93 \times 10^{-10}$). The epidemiological behavior of the TGI according to age and ethnicity is shown in Table 2, which shows an increase of the index as age increases. In regards to ethnicity, no statistically significant differences were found between means ($p = 0.326$).

When assessing the reference population ($n = 351$); healthy: $n = 207$ – unhealthy: $n = 144$, a similar behavior of the TGI was found according to age and ethnicity (Table 3). In addition, there

Table 1. General characteristics of the sample studied, Maracaibo city, Venezuela.

	Female (n= 1050)		Male (n= 954)		Total (n=2004)	
	n	%	n	%	n	%
Age (years)						
<30	308	29.3	363	38.1	671	33.5
30–49	420	40.0	350	36.7	770	38.4
≥ 50	322	30.7	241	25.3	563	28.1
Ethnicity						
Mixed	794	75.6	740	77.6	1534	76.5
White Hispanic	171	16.3	145	15.2	316	15.8
Afro-Venezuelan	27	2.6	32	3.4	59	2.9
American Indian	47	4.5	36	3.8	83	4.1
Other*	11	1.0	1	0.1	12	0.6
Triglycerides (mg/dl)						
<150	815	77.6	644	67.5	1459	72.8
≥ 150	235	22.4	310	32.5	545	27.2
Glycaemic status[†]						
Euglycemic	774	73.9	662	69.4	1436	71.7
Impaired fasting glucose	186	17.7	212	22.2	398	19.9
Type 2 diabetes mellitus	88	8.4	80	8.4	168	8.4

*Asian and Arabic descent.

[†]Criteria according to the ADA 2016 consensus

were no differences in TGI mean between the general and reference population.

Cutoff points for the TGI in the reference population by gender

For the determination of cutoff points of the TGI, ROC curves were plotted for the reference population by gender (Figure 2). An AUC of 0.889 (95% CI: 0.854-0.924) was obtained for the general population with a proposed cutoff of 4.49 (82.6% sensitivity, and 82.1% specificity), while the AUC calculated for males was 0.903 (95% CI: 0.856-0.950) and for females was 0.871 (95% CI: 0.818-0.925); Delong's test: $p=0.37$. The cutoff points and index calculated according to the ROC curves are shown in Table 4.

Cutoff points for the TGI in the reference population by age

When assessing the predictive capacity of the TGI according to age, a higher AUC was obtained in individuals between the ages of 30 and 50 years old (AUC=0.876; 95% CI: 0.812-0.939); however, when comparing AUCs among age groups, no significant differences were found. The cutoff points obtained according to

age are shown in Table 5, with means similar to that proposed for the reference population (<30 years: 4.48 [65.0% sensitivity, and 84.8% specificity]; 30–50 years: 4.51 [84.7% sensitivity, and 78.3% specificity], ≥50 years, 4.51 [86.5% sensitivity, and 66.7% specificity]).

HOMA2IR according to cutoff points of the TGI

Finally, when assessing HOMA2-IR levels according to the proposed cutoff point of the TGI for the general population (Figure 3), individuals with $TGI \geq 4.5$ exhibited higher levels of HOMA2IR than those who had a $TGI < 4.5$ (2.48 vs 1.74, respectively, $p < 0.001$), with similar behavior by gender.

Table 2. Epidemiological behavior of the triglyceride-glucose index in the general population according to sociodemographic variables, in Maracaibo city, Venezuela.

	n	TGI (n=2004)	
		Mean±SD	p*
Gender			8.93×10 ⁻¹⁰
Female	1050	4.56±0.33	
Male	954	4.66±0.34	
Age (years)			6.21×10 ⁻⁷⁷
<30	671	4.43±0.28	
30–49	770	4.65±0.33	
≥50	563	4.77±0.31	
Ethnicity			0.326
Mixed	1534	4.60±0.33	
White Hispanic	316	4.62±0.35	
Afro-Venezuelan	59	4.68±0.32	
American Indian	83	4.60±0.31	
Other [†]	12	4.51±0.28	

*Student's t-test (for more than two groups one-way ANOVA was used).

[†]Asian and Arabic descent.

SD: standard deviation

Post-hoc Tukey: <30 years vs 30–49 years, $p=5.09 \times 10^{-9}$; <30 years vs ≥50 years, $p=5.09 \times 10^{-9}$; 30–49 years vs ≥50 years, $p=5.10 \times 10^{-9}$.

Table 3. Epidemiological behavior of the triglyceride-glucose index in the reference population according to sociodemographic variables, in Maracaibo city, Venezuela.

	TGI (n=351)		
	n	Mean±SD	p*
Gender			5.83·10 ⁻⁶
Female	190	4.41±0.29	
Male	161	4.56±0.32	
Age (years)			8.42×10 ⁻²⁸
<30	178	4.31±0.29	
30–49	118	4.61±0.31	
≥50	55	4.72±0.23	
Ethnicity			0.413
Mixed	268	4.48±0.32	
White Hispanic	52	4.48±0.34	
Afro-Venezuelan	8	4.59±0.26	
American Indian	20	4.39±0.23	
Other [†]	3	4.25±0.26	

*Student's t-test (for more than two groups one way-ANOVA was used).

[†]Asian and Arabic descent.

SD: Standard deviation.

Post-hoc Tukey: <30 years vs 30–49 years, $p=5.09 \times 10^{-9}$; <30 years vs ≥50 years, $p=5.09 \times 10^{-9}$; 30–49 years vs ≥50 years, $p=0.03$.

Figure 2. Receiver Operating Characteristic curves for the triglyceride-glucose index in the reference population by gender, in Maracaibo city, Venezuela. DtR, Distance to ROC; J, Youden Index; AUC, are under the curve; sen, sensitivity; spe, specificity.

Table 4. Cut-off points for triglyceride-glucose index (TGI) selected in the reference population and by gender, in Maracaibo city, Venezuela.

Gender	TGI [†]	Sensitivity (%)	Specificity (%)	Youden index	ROC distance	LR+	AUC (95%CI)
Female	4.45	80.3	80.6	0.61	0.276	4.14	0.871 (0.818–0.925)
Male	4.51	87.2	83.1	0.70	0.212	5.15	0.903 (0.856–0.950)
Total	4.49	82.6	82.1	0.65	0.249	4.61	0.889 (0.854–0.924)

[†]Cut-off points selected according to the best combination of indices. Delong's test= 0.37

Table 5. Cutoff points for triglyceride-glucose index (TGI) selected in the reference population according to age groups, in Maracaibo city, Venezuela.

Age (years)	TGI [†]	Sensitivity (%)	Specificity (%)	Youden index	ROC distance	LR+	AUC (95%CI)
<30	4.49	65.0	84.8	0.50	0.381	4.27	0.789 (0.684–0.895)
30–50	4.51	84.7	78.3	0.63	0.265	3.89	0.876 (0.812–0.939)
≥50	4.51	86.5	66.7	0.53	0.359	2.59	0.776 (0.602–0.949)

[†]Cut-off points selected according to the best combination of indices. Delong's test: <30 vs 30–50 years; $p=0.171$. <30 vs ≥50 years; $p=0.885$. 30–50 vs ≥50 years; $p=0.34$.

Figure 3. HOMA2-IR levels according to the specific cutoff point for the triglyceride-glucose index in the reference population, in Maracaibo city, Venezuela.

Dataset 1. Data for the study ‘Optimal cutoff for the evaluation of insulin resistance through triglyceride-glucose index: A cross-sectional study in a Venezuelan population’

<http://dx.doi.org/10.5256/f1000research.12170.d171840>

This data is available in both .SAV and .xls forms. BMI: Body Mass Index; BP: Blood Pressure.

Discussion

The evaluation of IR is an objective that continues to acquire relevance in clinical and epidemiological research, due to the potential role of this disorder in the pathophysiology of MS²⁶, the consequent risk of developing DM2²⁷ and CVD²⁸. In developing countries with economic difficulties in health systems, such as Venezuela, routine measurements of plasma insulin are not easily accessible, which forces the use of other indices based on the role of glucolipotoxicity as a key element in the development of IR²⁹. In this way, the TGI has recently been proposed and validated as a useful alternative in clinical settings, and has been well accepted due to the high predictive power observed with respect to other indices³⁰.

Several studies have shown that the TGI better predicts HOMA-IR levels than variables, such as the TAG/HDL index, visceral adiposity index, leptin, Apo-B/Apo-AI, and lipid parameters^{29,30}, representing a very good correspondence with this index³¹, which constitutes an important tool with high validity for the clinician when facing limited access to lab work-ups. However, in view of the wide variability of TAG levels according to the ethnic, sociocultural and genetic characteristics of each population, the need arises to evaluate their epidemiological behavior and establish reference values specific to each region.

In regards to our study, the mean TGI was higher in men compared to women. Several studies have reported similar findings^{14,32}; however, when plotted ROC curves for the TGI for each gender, significant differences in the AUC values and effect size by gender were not observed. We propose the use of a single cutoff point of 4.5 for the identification of unhealthy individuals in the clinical practice as an easy tool for the clinician.

These findings corresponded to those originally reported by the index's authors in 2010¹⁵, when evaluating the discriminative ability of this index to determine IR against the euglycemic-hyperinsulinemic clamp in a population of 99 individuals (11 healthy, 34 obese, 22 with prediabetes and 32 with DM2) suggesting a cutoff point of 4.68 with high sensitivity (96.5%) and specificity (85.0%). Based on these findings, several studies have attempted to establish specific cutoff points for their populations, assessing the clinical utility of this index. In 2011, in a Brazilian population, Vasques *et al.*³³ stipulated that the TGI had a slightly better performance to diagnose IR compared to HOMA2-IR (AUC=0.79 vs AUC=0.77, respectively). Although these investigators did not perform any statistical tests to compare the diagnostic capacity between both indices, the similar behavior allows the use of TGI in clinical settings routinely.

On the other hand, Unger *et al.*²¹ conducted a cross-sectional study in an Argentinean population with the aim of evaluating the discriminative capacity of the TGI for the diagnosis of MS, taking into account that the development of this syndrome is related to IR. They set a cutoff point of 8.8 (sensitivity = 79%, specificity = 86%) to diagnose MS in their population, a value that differs markedly from the values in our region and those originally proposed. These differences have been observed in other studies^{34–36}, so considering that the means of TAG do not vary significantly between these and our study, this discrepancy could be attributable to errors in the calculation of the original formula³⁷.

In regards to age, an increase in the average TGI was observed as age increased, similar to that found by Navarro-González *et al.*³⁸ and Cuda *et al.*³⁹. This can be associated with the increase in oxidative stress inherent to aging, which would favor the development of IR, as well as the elevation of plasma levels of TAG⁴⁰. In this sense, Guerrero-Romero *et al.*³¹ recently evaluated the performance of the TGI to determine IR against the euglycemic-hyperinsulinemic clamp in a young Mexican adult population (mean age: 19.2±1.4), with the goal of establishing a specific cutoff point in this population. However, the cutoff points of 4.55 in men and 4.68 in women were similar to the value proposed previously for the general population, and similar to the one proposed in our study. Also, when we plotted ROC curves for each age group, we did not observe significant differences between AUC neither effect size in Cohen's *d* analysis, ruling out the need to establish age-specific cutoff points.

In relation to ethnicity, the small sample in each of the categories precludes the generalization of these results, but it shows that it is necessary to evaluate the behavior of the TGI in the different ethnic groups on a large scale and to determine if there are significant differences between them because there is no data reported on this diagnostic method among various ethnicities.

Based on previous analyzes, the usefulness of the TGI has been extended in several regions of the world, being also used in the screening of other pathological metabolic states in which the IR underlies the fundamental pathophysiological mechanism, such as DM2⁴¹ and atypical metabolic phenotypes^{42,43}. In fact, the TGI appears to be a better predictor of incidence of DM2 than TAG³⁸, weight gain⁴³ and other IR indices, such as TAG/HDL and HOMA-IR⁴¹. Moreover, Lee *et al.* performed a retrospective study involving 2900 Korean non-diabetic adults determinate as a cutoff point 8, (AUC=0.751, 95% CI 0.704-0.799) to predict the incidence of DM2 in their population. In addition, only individuals with a TGI >8.8 were associated with a significant risk of DM2 incidence, regardless of the presence of obesity⁴⁴.

Additionally, the index can better predict the patient's metabolic status, as it has been shown in discriminating atypical metabolic phenotypes, such as metabolically obese normal weight and healthy obese, as well as their progression throughout time³⁵. Cutoff points have been determined to discriminate these

pathological metabolic status in two populations of Korea^{34,35}, which could facilitate the definition of these phenotypes. Future studies in our population should evaluate the utility of this index in the identification of these abnormal pathological statuses. It is important to mention that within the limitations of this study it is necessary to consider the transversal design and the lack of gold standard during the process of selection of the reference population.

Conclusions

The TGI is an instrument of interest when it comes to identifying IR in the general population. We propose a single cutoff point of 4.5 to identify patients with IR, as we identify the need for standardization of the formula calculation in order to be able to adequately compare the differences observed in various studies. Despite this, due to the easier application in clinical practice, future studies in our population should evaluate the predictive capacity of this index to determine atypical metabolic phenotypes, DM2 and CVD risk.

Data availability

Dataset 1: Data for the study 'Optimal cutoff for the evaluation of insulin resistance through triglyceride-glucose index: A cross-sectional study in a Venezuelan population'. This data is available in both .SAV and .xls forms. BMI: Body Mass Index; BP: Blood Pressure. doi, [10.5256/f1000research.12170.d171840](https://doi.org/10.5256/f1000research.12170.d171840)⁴⁵

Competing interests

No competing interests were disclosed.

Grant information

This work was supported by the Technological, Humanistic, and Scientific Development Council (Consejo de Desarrollo Científico, Humanístico y Tecnológico; CONDES), University of Zulia (grant n° CC-0437-10-21-09-10).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- Wilcox G: **Insulin and insulin resistance**. *Clin Biochem Rev*. 2005; **26**(2): 19–39. [PubMed Abstract](#) | [Free Full Text](#)
- Morigny P, Houssier M, Mouiel E, *et al.*: **Adipocyte lipolysis and insulin resistance**. *Biochimie*. 2016; **125**: 259–66. [PubMed Abstract](#) | [Publisher Full Text](#)
- Patel TP, Rawal K, Bagchi AK, *et al.*: **Insulin resistance: an additional risk factor in the pathogenesis of cardiovascular disease in type 2 diabetes**. *Heart Fail Rev*. 2016; **21**(1): 11–23. [PubMed Abstract](#) | [Publisher Full Text](#)
- Luchsinger JA: **Insulin resistance, type 2 diabetes, and AD: cerebrovascular disease or neurodegeneration?** *Neurology*. 2010; **75**(9): 758–759. [PubMed Abstract](#) | [Publisher Full Text](#)
- Rojas J, Bermúdez V, Leal E, *et al.*: **Insulinorresistencia E Hiperinsulinemia Como Factores De Riesgo Para Enfermedad Cardiovascular**. *AVTF*. 2008; **27**(1): 30–40. [Reference Source](#)
- Gastaldelli A: **Role of beta-cell dysfunction, ectopic fat accumulation and insulin resistance in the pathogenesis of type 2 diabetes mellitus**. *Diabetes Res Clin Pract*. 2011; **93**(suppl 1): S60–S65. [PubMed Abstract](#) | [Publisher Full Text](#)
- Randle PJ, Garland PB, Hales CN, *et al.*: **The glucose fatty-acid cycle. Its role in insulin sensitivity and the metabolic disturbances of diabetes mellitus**. *Lancet*. 1963; **1**(7285): 785–789. [PubMed Abstract](#) | [Publisher Full Text](#)
- Shulman GI: **Cellular mechanisms of insulin resistance**. *J Clin Invest*. 2000; **106**(2): 171–6. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Borai A, Livingstone C, Kaddam I, *et al.*: **Selection of the appropriate method for the assessment of insulin resistance**. *BMC Med Res Methodol*. 2011; **11**(1): 158. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- DeFronzo RA, Tobin JD, Andres R: **Glucose clamp technique: a method for quantifying insulin secretion and resistance**. *Am J Physiol*. 1979; **237**(3): E214–223. [PubMed Abstract](#)
- Singh B, Saxena A: **Surrogate markers of insulin resistance: A review**. *World J Diabetes*. 2010; **1**(2): 36–47. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Matthews DR, Hosker JP, Rudenski AS, *et al.*: **Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man**. *Diabetologia*. 1985; **28**(7): 412–9. [PubMed Abstract](#) | [Publisher Full Text](#)
- Bermúdez V, Salazar J, Martínez MS, *et al.*: **Prevalence and Associated Factors of Insulin Resistance in Adults from Maracaibo City, Venezuela**. *Adv Prev Med*. 2016; **2016**: 9405105. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Simental-Mendía LE, Rodríguez-Morán M, Guerrero-Romero F: **The product of fasting glucose and triglycerides as surrogate for identifying insulin resistance in apparently healthy subjects**. *Metab Syndr Relat Disord*. 2008; **6**(4): 299–304. [PubMed Abstract](#) | [Publisher Full Text](#)
- Guerrero-Romero F, Simental-Mendía LE, González-Ortiz M, *et al.*: **The product of triglycerides and glucose, a simple measure of insulin sensitivity. Comparison with the euglycemic-hyperinsulinemic clamp**. *J Clin Endocrinol Metab*. 2010; **95**(7): 3347–51. [PubMed Abstract](#) | [Publisher Full Text](#)
- Bermúdez V, Marciano RP, Cano C, *et al.*: **The Maracaibo city metabolic syndrome prevalence study: design and scope**. *Am J Ther*. 2010; **17**(3): 288–294. [PubMed Abstract](#) | [Publisher Full Text](#)
- World Health Organization: **Obesity: preventing and managing the global epidemic. Report of a WHO consultation**. *World Health Organ Tech Rep Ser*. Geneva: The Organization; 2000; **894**: i–xii, 1–253. [PubMed Abstract](#)
- Health Statistics: **NHANES III reference manuals and reports (CDROM)**. Hyattsville, MD: Centers for Disease Control and Prevention, 1996. [Reference Source](#)
- Bermúdez V, Rojas J, Salazar J, *et al.*: **Optimal Waist Circumference Cut-Off Point for Multiple Risk Factor Aggregation: Results from the Maracaibo City Metabolic Syndrome Prevalence Study**. *Epidemiol Res Int*. 2014; **2014**: 718571. [Publisher Full Text](#)
- Bermúdez V, Rojas J, Martínez MS, *et al.*: **Epidemiologic Behavior and Estimation of an Optimal Cut-Off Point for Homeostasis Model Assessment-2 Insulin Resistance: A Report from a Venezuelan Population**. *Int Sch Res Notices*. 2014; **2014**: 616271. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Unger G, Benozzi SF, Perruzza F, *et al.*: **Triglycerides and glucose index: a useful indicator of insulin resistance**. *Endocrinol Nutr*. 2014; **61**(10): 533–40. [PubMed Abstract](#) | [Publisher Full Text](#)
- Akobeng AK: **Understanding diagnostic tests 3: Receiver operating characteristic curves**. *Acta Paediatr*. 2007; **96**(5): 644–7. [PubMed Abstract](#) | [Publisher Full Text](#)
- Demler OV, Pencina MJ, D'Agostino RB Sr: **Misuse of DeLong test to compare AUCs for nested models**. *Stat Med*. 2012; **31**(23): 2577–87. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Böhning D, Böhning W, Holling H: **Revisiting Youden's index as a useful measure of the misclassification error in meta-analysis of diagnostic studies**. *Stat*

- Methods Med Res.* 2008; **17**(6): 543–54.
[PubMed Abstract](#) | [Publisher Full Text](#)
25. Perkins NJ, Schisterman EF: **The inconsistency of “optimal” cutpoints obtained using two criteria based on the receiver operating characteristic curve.** *Am J Epidemiol.* 2006; **163**(7): 670–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 26. Samson SL, Garber AJ: **Metabolic syndrome.** *Endocrinol Metab Clin North Am.* 2014; **43**(1): 1–23.
[PubMed Abstract](#) | [Publisher Full Text](#)
 27. Lyssenko V, Jonsson A, Almgren P, *et al.*: **Clinical risk factors, DNA variants, and the development of type 2 diabetes.** *N Engl J Med.* 2008; **359**(21): 2220–2232.
[PubMed Abstract](#) | [Publisher Full Text](#)
 28. Miller M, Stone NJ, Ballantyne C, *et al.*: **Triglycerides and cardiovascular disease: a scientific statement from the American Heart Association.** *Circulation.* 2011; **123**(20): 2292–2333.
[PubMed Abstract](#) | [Publisher Full Text](#)
 29. Er LK, Wu S, Chou HH, *et al.*: **Triglyceride Glucose-Body Mass Index Is a Simple and Clinically Useful Surrogate Marker for Insulin Resistance in Nondiabetic Individuals.** *PLoS One.* 2016; **11**(3): e0149731.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 30. Du T, Yuan G, Zhang M, *et al.*: **Clinical usefulness of lipid ratios, visceral adiposity indicators, and the triglycerides and glucose index as risk markers of insulin resistance.** *Cardiovasc Diabetol.* 2014; **13**(1): 146.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 31. Guerrero-Romero F, Villalobos-Molina R, Jiménez-Flores JR, *et al.*: **Fasting triglycerides and glucose index as a diagnostic test for insulin resistance in Young adults.** *Arch Med Res.* 2016; **47**(5): 382–387.
[PubMed Abstract](#) | [Publisher Full Text](#)
 32. Irace C, Carallo C, Scavelli FB, *et al.*: **Markers of insulin resistance and carotid atherosclerosis. A comparison of the homeostasis model assessment and triglyceride glucose index.** *Int J Clin Pract.* 2013; **67**(7): 665–672.
[PubMed Abstract](#) | [Publisher Full Text](#)
 33. Vasques AC, Novaes FS, de Oliveira Mda S, *et al.*: **TyG index performs better than HOMA in a Brazilian population: a hyperglycemic clamp validated study.** *Diabetes Res Clin Pract.* 2011; **93**(3): e98–e100.
[PubMed Abstract](#) | [Publisher Full Text](#)
 34. Lee SH, Han K, Yang HK, *et al.*: **A novel criterion for identifying metabolically obese but normal weight individuals using the product of triglycerides and glucose.** *Nutr Diabetes.* 2015; **5**(4): e149.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 35. Lee SH, Yang HK, Ha HS, *et al.*: **Changes in Metabolic Health Status Over Time and Risk of Developing Type 2 Diabetes: A Prospective Cohort Study.** *Medicine (Baltimore).* 2015; **94**(40): e1705.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 36. Abbasi F, Reaven GM: **Comparison of two methods using plasma triglyceride concentration as a surrogate estimate of insulin action in nondiabetic subjects: triglycerides × glucose versus triglyceride/high-density lipoprotein cholesterol.** *Metabolism.* 2011; **60**(12): 1673–1676.
[PubMed Abstract](#) | [Publisher Full Text](#)
 37. Hosseini SM: **Triglyceride-Glucose (TyG) Index Computation and Cut-Off.** *Acta Endo (Buc).* 2015; **11**(1): 130–131.
[Publisher Full Text](#)
 38. Navarro-González D, Sánchez-Íñigo L, Pastrana-Delgado J, *et al.*: **Triglyceride-glucose index (TyG index) in comparison with fasting plasma glucose improved diabetes prediction in patients with normal fasting glucose: The Vascular-Metabolic CUN cohort.** *Prev Med.* 2016; **86**: 99–105.
[PubMed Abstract](#) | [Publisher Full Text](#)
 39. Cuda G, Lentini M, Gallo L, *et al.*: **Fasting triglycerides and glucose index in an unselected consecutive Italian population of outpatients.** *Riv Ital Med Lab.* 2011; **7**(4): 226–227.
[Publisher Full Text](#)
 40. Monickaraj F, Aravind S, Nandhini P, *et al.*: **Accelerated fat cell aging links oxidative stress and insulin resistance in adipocytes.** *J Biosci.* 2013; **38**(1): 113–122.
[PubMed Abstract](#) | [Publisher Full Text](#)
 41. Lee SH, Kwon HS, Park YM, *et al.*: **Predicting the Development of Diabetes Using the Product of Triglycerides and Glucose: The Chungju Metabolic Disease Cohort (CMC) Study.** *PLoS One.* 2014; **9**(2): e90430.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 42. Lee SH, Han K, Yang HK, *et al.*: **Identifying subgroups of obesity using the product of triglycerides and glucose: the Korea National Health and Nutrition Examination Survey, 2008-2010.** *Clin Endocrinol (Oxf).* 2014; **82**(2): 213–220.
[PubMed Abstract](#) | [Publisher Full Text](#)
 43. Navarro-González D, Sánchez-Íñigo L, Fernández-Montero A, *et al.*: **TyG Index Change Is More Determinant for Forecasting Type 2 Diabetes Onset Than Weight Gain.** *Medicine (Baltimore).* 2016; **95**(19): e3646.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 44. Lee DY, Lee ES, Kim JH, *et al.*: **Predictive Value of Triglyceride Glucose Index for the Risk of Incident Diabetes: A 4-Year Retrospective Longitudinal Study.** *PLoS One.* 2016; **11**(9): e0163465.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 45. Salazar J, Bermúdez V, Calvo M, *et al.*: **Dataset 1 in: Optimal cutoff for the evaluation of insulin resistance through triglyceride-glucose index: A cross-sectional study in a Venezuelan population.** *F1000Research.* 2017.
[Data Source](#)

Open Peer Review

Current Peer Review Status:

Version 2

Reviewer Report 05 January 2018

<https://doi.org/10.5256/f1000research.14039.r28438>

© 2018 Balasubramanyam M. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Muthuswamy Balasubramanyam

Department of Cell and Molecular Biology, Madras Diabetes Research Foundation (MDRF), Chennai, India

Insulin resistance (IR) is the root cause for several major metabolic disorders and identifying individuals with IR is right now a public health priority as this will pave way for appropriate prevention and management of type 2 diabetes and associated non-communicable diseases. While Euglycemic-Hyperinsulinemic Clamp which is considered as the gold standard for the determination of IR is an invasive and impractical procedure for risk assessment in the general population, the measurement of IR by Homeostasis Model Assessment (HOMA-IR) also requires costly tests like insulin assays which are difficult in the low-resource settings. In this context, assessing IR and metabolic risk via triglyceride-glucose index (TGI) as detailed in this manuscript receives some attention and importance. The main advantages of the TGI is that both the triglyceride and glucose measurements are cost-effective and hence affordable in epidemiological and screening programs to assess the relative cardiometabolic risk in the general populations and in the clinics.

However, the manuscript should delineate several limitations of the study (as detailed below):

- As the study nature is of cross-sectional observation study design, it is difficult to evaluate a direct causal relationship.
- TGI may have limited inference when used in subjects with extremely high triglyceride levels.
- Extrapolating the findings of this study to other races or ethnicities should be interpreted cautiously; there might be a need for ethnic / race specific TGI cut-offs.
- TGI may suffer from confounding factors such as medications that lower mainly triglycerides. Authors should discuss these medication (statins, fenofibrate, omega-3 etc) effects.
- Also, TGI values could be confounded by diet and physical activity among the groups. Therefore, authors should discuss the role of diet (if FFQ is available) and physical activity in relation to cut-offs of TGI. These should be discussed.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Partly

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Disease-biology; biochemical and molecular medicine aspects of metabolic disorders with special reference to diabetes and its associated complications

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 25 August 2017

<https://doi.org/10.5256/f1000research.13172.r25043>

© 2017 Benozzi S et al. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Silvia Fabiana Benozzi

Universidad Nacional del Sur, Bahía Blanca, Argentina

Gisela Unger

Universidad Nacional del Sur, Bahía Blanca, Argentina

This research article deals with an interesting subject related to insulin resistance, its evaluation without the need of an insulin blood test. We believe that it can contribute to identify patients with metabolic risk. Authors have clearly elaborated the background of the problem and current literature is cited. The study is properly conducted and described. But, there are some shortcomings that should be corrected:

- Abstract: Line 1, there are two dots after the word *evaluated*.
- Keywords are absent.
- Abbreviations: Page 3, we considered unnecessary this subtitle because the abbreviations are spelled out in the text.
- Sample selection and study design: We have interpreted that the presence of all the cited criteria was necessary to classify an individual in the reference population, but it is not clear in the text. Authors should explain how many of the cited criteria were considered to classify an individual as healthy or sick.
- Figure 1: Authors should consider the previous correction also in this figure. Authors should replace the explanation [†]*Abdominal circumference > 90 cm in the female sex and > 95 cm in the male sex* with [†]*Abdominal obesity defined as: abdominal circumference > 90 cm in the female sex and > 95 cm in the male sex*. Authors should highlight in the respective box that the population of 2004 individuals is the General population.
- Laboratory analysis: If possible, authors should provide the fasting time in hours owing the lack of harmonization in this preanalytical aspect worldwide. Glucose and insulin analytical coefficients of variation should be provided. It is not necessary HDL-C description as this analyte was not used in the present research.
- Statistical analysis: Authors should verify the ROC curve formula described to calculate the minimum cutoff point.
- Results: Authors should correct the second subtitle because the cutoff points were calculated in the reference population (n=351) – as stated in Methods – and not in the general population (n=2004); also in the second line of the respective text. Authors should correct the third subtitle because the cutoff points were calculated in the reference population (n=351) – as stated in Methods – and not in the general population (n= 2004); authors should clarify the comparison made in line 7 of the respective text, we understand that they are comparing the cutoff values obtained by age with the cutoff value obtained for the total reference population.
- Table 2: Authors should replace the word *TOTAL* with *TGI* so as to allow a better interpretation of the values shown in the table. Also the number of individuals should be described by variable, as in Table 3.
- Table 3: Authors should replace the word *TOTAL* with *TGI* so as to allow a better interpretation of the values shown in the table.
- Table 4: Authors should correct the table title because the cutoff points were calculated in the reference population (n=351) – as stated in Methods – and not in the general population (n= 2004).
- Table 5: Authors should indicate in the title that the cutoff points were calculated in the reference population (n= 351), as in the other tables.

- Figure 2: Authors should correct the title because the cutoff points were calculated in the reference population (n=351) – as stated in Methods – and not in the general population (n=2004).
- Figures and tables in general: Authors should verify that each figure or table has a concise title of no more than 15 words. Authors should provide a legend for each figure and table that briefly describes the key points and explain any symbols and abbreviations used. Also the *n* value should be provided in each table and figure to help reader's interpretation.
- Discussion: Authors states that TGI was higher in men compared to women, and that an increase in its average value was observed as age increased. It would be interesting to evaluate the effect size of these statistical differences so as to interpret the clinical importance of these increases;Cohen's*d*—<https://www.uccs.edu/~lbecker/>— can be used for this purpose if the variable has a parametric distribution.
- General considerations: Authors should express *p* values with three decimals in all the text, tables and figures. The word *cutoff* is written as *cut-off* in same parts of the text, figures and tables, the same happens with the word *indexes* and *indices*. They should be written in the same way in all the text. It would be advisable to replace in the introduction: *blood work to calculate the serum insulin* with *insulinblood test*; in all the text: *sick* with *unhealthy*; in the conclusions: *generalization* with *harmonization* or *standardization*.

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

We confirm that we have read this submission and believe that we have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reader Comment 20 Oct 2017

F1000 Research, UK

The Editorial Team can confirm that keywords are included in the article, but only in the article's metadata rather than being displayed on the Article Page or part of the PDF.

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research