RESEARCH ARTICLE

REVISED RNAi targeting *Caenorhabditis elegans* α -arrestins has little effect on lifespan [version 3; peer review: 2 approved, 1 approved with reservations]

Previously titled: RNAi targeting *Caenorhabditis elegans* α -arrestins has small or no effects on lifespan

Sangsoon Park ¹, Yoonji Jung¹, Seon Woo A. An¹, Heehwa G. Son¹,
Woosoon Hwang¹, Dongyeop Lee¹, Murat Artan¹, Hae-Eun H. Park¹,
Dae-Eun Jeong¹, Yujin Lee¹, Seung-Jae V. Lee ^{1,2}

¹Department of Life Sciences, Pohang University of Science and Technology, Pohang, Gyeongbuk, 37673, South Korea

²School of Interdisciplinary Bioscience and Bioengineering, Pohang University of Science and Technology, Pohang, Gyeongbuk, 37673, South Korea

V3 First published: 18 Aug 2017, 6:1515
<https://doi.org/10.12688/f1000research.12337.1>
 Second version: 19 Oct 2017, 6:1515
<https://doi.org/10.12688/f1000research.12337.2>
 Third version: 14 Nov 2017, 6:1515
<https://doi.org/10.12688/f1000research.12337.3>
 Latest published: 08 Dec 2017, 6:1515
<https://doi.org/10.12688/f1000research.12337.4>

Abstract

Background: α -arrestins are a family of proteins that are implicated in multiple biological processes, including metabolism and receptor desensitization.

Methods: Here, we sought to examine the roles of α -arrestins in the longevity of *Caenorhabditis elegans* through an RNA interference screen.

Results: We found that knocking down each of 24 out of total 29 *C. elegans* α -arrestins had little effect on lifespan. Thus, individual *C. elegans* α -arrestins may have minor effects on longevity.

Conclusions: This study will provide useful information for future research on the functional role of α -arrestins in aging and longevity.

Keywords

C. elegans, α -arrestin, insulin/IGF-1 signaling, aging, lifespan

Open Peer Review

Approval Status

	1	2	3
version 4			
(revision)
08 Dec 2017	view		
version 3			
(revision)
14 Nov 2017	view		
version 2			
(revision)
19 Oct 2017	view	view	view
version 1			
18 Aug 2017
	view	view	

1. **Alessandro Bitto**, University of Washington, Seattle, Seattle, USA

Jason Pitt, University of Washington, Seattle, USA

2. **Roger Pocock** , Monash University, Melbourne, Australia

3. **Meng-Qiu Dong** , National Institute of Biological Sciences (NCBS), Beijing, China

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Seung-Jae V. Lee (seungjaelee@postech.ac.kr)

Author roles: **Park S:** Conceptualization, Formal Analysis, Investigation, Project Administration, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Jung Y:** Investigation, Validation, Writing – Review & Editing; **An SWA:** Investigation; **Son HG:** Investigation; **Hwang W:** Investigation; **Lee D:** Investigation; **Artan M:** Investigation; **Park HEH:** Investigation; **Jeong DE:** Investigation; **Lee Y:** Investigation; **Lee SJV:** Conceptualization, Funding Acquisition, Project Administration, Supervision, Validation, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by the Korean Government (MSIP) through the National Research Foundation of Korea (NRF) [NRF-2016R1E1A1A01941152] funded to S-J.V.L.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2017 Park S *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. Data associated with the article are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

How to cite this article: Park S, Jung Y, An SWA *et al.* **RNAi targeting *Caenorhabditis elegans* α -arrestins has little effect on lifespan [version 3; peer review: 2 approved, 1 approved with reservations]** F1000Research 2017, 6:1515
<https://doi.org/10.12688/f1000research.12337.3>

First published: 18 Aug 2017, 6:1515 <https://doi.org/10.12688/f1000research.12337.1>

REVISED Amendments from Version 2

In response to the reviewers' comments, we revised our paper and further changes are listed below:

1. We changed the title of our article
2. We specified the technical problems for RNAi clone generation in the Methods section
3. We revised misleading sentences throughout the article

See referee reports

Introduction

α -arrestins are a family of proteins that contain arrestin domains whose sequences and structures have similarities with those of classical visual and β -arrestins¹⁻³. α -arrestins are considered as ancestral forms of arrestins because their orthologs exist in fungi, including yeast, which do not have visual or β -arrestins^{1,4}. Several mammalian α -arrestins modulate metabolism and receptor desensitization^{5,6}, but much remains to be elucidated concerning the functions of α -arrestins in many organisms. The human genome encodes 6 α -arrestins and 4 visual or β -arrestins. Interestingly, the simple roundworm *Caenorhabditis elegans* genome contains 29 α -arrestin and 1 β -arrestin genes^{1,7}. Therefore, the *C. elegans* system provides opportunities for the genetic analysis of α -arrestins in various aspects of physiology both individually and combinatorially. However, information regarding the functions of *C. elegans* α -arrestins is limited^{8,9}.

C. elegans is an excellent genetic model organism that has been exploited for studying conserved biological processes, including apoptosis, behavior, development and aging. In particular, its short lifespan in combination with genetic amenability has made *C. elegans* one of the most popular models for research on aging and longevity^{10,11}. Many factors, including components in insulin/insulin-like growth factor 1 (IGF-1) signaling (IIS), have been identified as lifespan and aging regulators in *C. elegans*¹¹⁻¹³. For example, genetic inhibition of IIS components, such as DAF-2/insulin/IGF-1 receptor, robustly extends lifespan and delays physiological aging through up-regulating transcription factors, including DAF-16/FOXO¹¹⁻¹³. Importantly, the roles of these aging-regulatory factors in *C. elegans* have been shown to be conserved in other species, including *Drosophila* and mammals^{12,13}. One of the powerful ways to identify novel factors that influence aging is by employing genetic screens, such as an RNA interference (RNAi) screen. We previously identified several genetic factors, including RNA helicases, that modulate longevity in *C. elegans*, through targeted or genome-wide RNAi screens¹⁴⁻¹⁷. Because of its robust longevity phenotype that confers sensitivity to changes in lifespan, *daf-2*/insulin/IGF-1 receptor mutants serve as an excellent platform for the identification of novel lifespan-regulating factors^{17,18}.

In this study, we aimed to determine whether any α -arrestins played a role in the lifespan regulation of wild-type or *daf-2* mutants. We performed a lifespan assay-based RNAi screen targeting 24 out of 29 *C. elegans* α -arrestins. We found that knocking down each of the α -arrestins had little effect on the

lifespan of wild-type and *daf-2* mutants. Thus, *C. elegans* α -arrestins may play minor or modulatory roles in lifespan regulation. Based on our results, it will be important to test the roles of α -arrestins in combinatorial manners and/or by using strong loss-of-function mutations in future research.

Methods

Caenorhabditis elegans strains

All strains were maintained as previously described¹⁹. The following strains were used in this study: N2 wild-type, CF1041 *daf-2(e1370)* III outcrossed six times to N2.

Phylogenetic analysis

The protein sequences of 27 α -arrestins, except *arnd* (arrestin domain protein)-20 and *arnd*-21, were obtained from Wormbase (www.wormbase.org, version WS259). The protein sequences of *arnd*-20 and *arnd*-21 were obtained from Ensembl (<http://www.ensembl.org>, release 89). The phylogenetic tree of 29 α -arrestins in *C. elegans* was generated using Clustal Omega (<http://www.ebi.ac.uk/Tools/msa/clustalo/>)²⁰ and re-visualized using the Dendroscope 3 (version 3.5.9)²¹. For the α -arrestins that have multiple isoforms, isoform a was used for the analysis.

RNAi clones

Twenty one RNAi clones that target *C. elegans* α -arrestin genes were used from two commercial *C. elegans* RNAi feeding libraries. Specifically, RNAi clones targeting *arnd*-2, *arnd*-6, *arnd*-7, *arnd*-8, *arnd*-9, *arnd*-10, *arnd*-13, *arnd*-16, *arnd*-18, *arnd*-23, *arnd*-24, *arnd*-25, *arnd*-28 and *ttm*-2 (toxin-regulated targets of MAPK 2) were obtained from Ahringer laboratory library (Geneservice Ltd., Cambridge, UK), while those against *arnd*-1, *arnd*-3, *arnd*-4, *arnd*-5, *arnd*-14, *arnd*-15 and *arnd*-19 were from Vidal laboratory library (Source BioScience, Nottingham, UK). Three RNAi clones targeting *arnd*-11, *arnd*-17, and *arnd*-26 were generated by molecular cloning using infusion recombinase (EZ-Fusion™ Cloning Kit, Enzygnomics, Daejeon, South Korea). The N2 genomic DNA was obtained through the lysis of worms using proteinase K (Invitrogen, Carlsbad, CA, USA) and N2 complementary DNA was obtained from RNA extraction using RNAiso Plus (Takara, Shiga, Japan) followed by reverse transcription using ImProm-II reverse transcriptase (Promega, Madison, WI, USA, USA). The infusion reaction between PCR products and pL4440 plasmids (Fire lab *C. elegans* vector kit, 1999) digested with HindIII (New England Biolabs, Ipswich, MA, USA) and Acc65I (New England Biolabs) was followed by transformation of in-house competent *E. coli* HT115 cells and by selection of positive clones on ampicillin (USB, Santa Clara, CA, USA)-containing LB plates. Primers (CosmoGenetech, Seoul, South Korea) that were used to amplify coding regions of *arnd*-11 from N2 genomic DNA, and those of *arnd*-17 and *arnd*-26 from N2 complementary DNA are as follows: forward primer 5'-GAATTCGATATCAAGCTCCCTCGT-GCAAATTAGGAAA-3' and reverse primer 5'-CTATAG-GGCGAATTGGGGTTCCTCCCACTCCATACA-3' for *arnd*-11; forward primer 5'-GAATTCGATATCAAGCTATGGTGCAGTTA-GATCGTTTTG-3' and reverse primer 5'-CTATAGGGCGAATT-GGTTAATCGGTATAAAATGG-3' for *arnd*-17; forward primer

5'-GAATTCGATATCAAGCTATGAAGGTCGATTACTTCG-3' and reverse primer 5'-CTATAGGGCGAATTGGCTACTTCTCG-GAGCCATTG-3' for *arrd-26*. The sequences of all these 24 α -arrestin RNAi clones were confirmed using DNA sequencing (Solgent, Daejeon, South Korea) before lifespan assays. RNAi clone for *arrd-12*, *arrd-22*, or *arrd-27* was not generated because we were not able to obtain transformed *E. coli* (HT115) colonies using *arrd-12* or *arrd-22* infusion reaction products, or the PCR product for *arrd-27* genomic or complementary DNA. *arrd-20* and *arrd-21* are predicted to be pseudogenes (Wormbase, version WS259) and were excluded from our screen. All RNAi clones used in our screen were examined for their potential off-target effects by using Clone Mapper (<http://bioinformatics.lif.univ-mrs.fr/RNAiMap>)²², and no significant off-target was predicted except *arrd-19* RNAi, which may additionally target a predicted pseudogene *arrd-21* (Table S1). Experimental validation by qRT-PCR will be necessary to completely exclude the possible off-target effects caused by RNAi clones.

RNAi screen using lifespan assay

The RNAi screen employing lifespan assay was performed as previously described¹⁷. Briefly, *E. coli* HT115 bacteria that expressed specific double-stranded RNAs (dsRNAs) were cultured overnight at 37°C in LB media containing 50 µg/ml ampicillin (USB). The cultured bacteria were seeded onto nematode growth media plates containing 50 µg/ml ampicillin and incubated overnight at 37°C. The seeded bacteria were treated with 1 mM isopropyl β-D-1-thiogalactopyranoside (Gold Biotechnology, St. Louis, MO, USA) and incubated at room temperature for approximately 24 h to induce dsRNAs. Age-synchronized wild-type N2 and *daf-2(e1370)* mutant animals were grown on RNAi plates from embryo to L4 stage at 20°C. Worms were then transferred onto RNAi plates containing 5 µM 5-fluoro-2'-deoxyuridine (FUDR; Sigma, St. Louis, MO, USA), which prevents eggs from hatching, at young (day 1) adult stage, and transferred again to new FUDR-containing RNAi plates after 1 or 2 days. Eggs laid by *tmm-2* RNAi-treated worms hatched more frequently than control eggs on FUDR-containing plates in two independent lifespan experiments. All lifespan assays were performed at 20°C as duplicates by two independent researchers. The survival of worms was determined by gently touching worms with a platinum wire. Worms that did not respond were counted as dead worms and removed from the plates. Worms that crawled off the plates, burrowed into the agar media, or displayed internal hatching or vulval rupture were censored, but included in the statistical analysis. Lifespan data from two independent lifespan experiments within the experimental sets were pooled and used for statistical analysis (see Dataset 1–Dataset 3). OASIS 2 (Online Application for Survival Analysis 2; <https://sbi.postech.ac.kr/oasis2>)²³ was used for the statistical analysis of lifespan results. *P* values were calculated using long-rank (Mantel-Cox method) test.

Results

Knocking down each of several *C. elegans* α -arrestins marginally influenced lifespan

We measured the lifespan of wild-type and long-lived *daf-2* mutant animals upon knocking down each of 24 of 29 genes encoding putative α -arrestin proteins (Figure 1A). We used *daf-16* RNAi that largely suppressed the longevity of *daf-2(-)* mutants as a positive control (Figure 1B, Figure S1A, Table S2)^{17,18}. We found that knockdown of specific α -arrestin genes caused a minor reduction of lifespan in wild-type or in *daf-2(-)* mutants (Figure 1B, Figure S1B–U, Table S2). Out of the 24 RNAi clones, RNAi targeting *arrd* (arrestin domain protein)-13, *arrd-16*, *arrd-23*, *arrd-24* or *arrd-25* in wild-type, and RNAi against *arrd-1*, *arrd-2*, *arrd-5*, *arrd-24* or *arrd-28* in *daf-2(-)* mutant animals decreased lifespan by more than 5% (Figure 1B–E, Figures S1B, D, K, Q, R, T, Table S2). Specifically, *arrd-16* RNAi decreased lifespan in wild-type by 9%, and *arrd-1* RNAi decreased lifespan in *daf-2(-)* mutants by 7% (Figure 1B and C). In addition, RNAi targeting *arrd-24* decreased the lifespan of wild-type and *daf-2* mutant animals by 11% and 6%, respectively (Figure 1E). In contrast, *arrd-3* RNAi increased the lifespan of wild-type animals by 10% (Figure 1F). Overall, genetic inhibition of individual α -arrestin genes using RNAi appears to have minor effects on *C. elegans* lifespan.

Dataset 1. Kaplan-Meier estimator of RNAi lifespan experiments

<http://dx.doi.org/10.5256/f1000research.12337.d173158>

Kaplan-Meier estimate values were calculated from pooled lifespan data of two independent lifespan experiments using OASIS2 (<https://sbi.postech.ac.kr/oasis2/>). 'At risk' indicates the number of individuals at risk just prior to a specific time point. 'S_hat' indicates Kaplan-Meier estimate of survival function, 'Var S_hat' indicates variance of 'S_hat', and 'SE(S)' indicates standard error of 'S_hat'.

Dataset 2. Mean lifespan and mortality rates of RNAi experiment results

<http://dx.doi.org/10.5256/f1000research.12337.d173159>

Mean lifespan, ages at different percent mortality, linear interpolation of mortality curve at specific mortality rate were calculated from pooled lifespan data of two independent lifespan experiments. Data were obtained using OASIS2 (<https://sbi.postech.ac.kr/oasis2/>).

Dataset 3. Statistical analysis of lifespan data

<http://dx.doi.org/10.5256/f1000research.12337.d173160>

Statistical test results (Chi square, *p* value, Bonferroni *p* value) were calculated between 'condition 1' and 'condition 2'. Test results were obtained using OASIS2 (<https://sbi.postech.ac.kr/oasis2/>).

Figure 1. RNAi targeting each of *Caenorhabditis elegans* α -arrestins had little effect on lifespan. (A) A phylogenetic tree showing *C. elegans* α -arrestin family members. Asterisks (*) indicate α -arrestins that were not examined in this study and number signs (#) indicate predicted pseudogenes. (B) Each circle represents percent mean lifespan change by treatment with individual α -arrestin gene RNAi in wild-type (WT) and *daf-2(e1370)* (*daf-2(-)*) mutants. Red triangles indicate lifespan changes by *daf-16* RNAi, which was used as a positive control. (C–F) Lifespan curves of WT and *daf-2(-)* animals treated with RNAi targeting *arrd-16* (C), *arrd-1* (D), *arrd-24* (E), and *arrd-3* (F). See Table S2 for statistical analysis.

Discussion

The lifespan-regulatory functions of α -arrestins remain largely unexplored at the organism level. Here we showed that RNAi knockdown of individual *C. elegans* α -arrestins had little effect on lifespan in wild-type or *daf-2* mutants. Our study has limitations that need to be considered for interpretation. First, RNAi targeting some α -arrestins might be insufficient for causing strong lifespan phenotypes. This may be because many of *C. elegans* α -arrestins are predicted to be expressed in neurons^{7,9,24–26}, which are refractory to RNAi^{27–29}. Lifespan assays using RNAi-hypersensitive mutants, including *rif-3(-)* and *eri-1(-)* animals^{29–31}, treated with α -arres-

tin RNAi, or using α -arrestin null mutants will help address this issue. In addition, as we did not test whether RNAi targeting each α -arrestin gene was effective by using quantitative RT-PCR, our negative data should be interpreted with caution. Second, *C. elegans* α -arrestins may have functional redundancy, considering the large number of the α -arrestin family members in *C. elegans* and their sequence similarity^{1,7}, which may obscure examining the functional importance of each α -arrestin. In addition, some α -arrestins may mostly function by modulating the action of their interacting proteins⁶. In this case, genetic inhibition of α -arrestins may rather subtly affect the functions of their interacting partners

that directly regulate physiology, such as aging and longevity, causing weak or no phenotypes. Thus, it will be interesting to test the effects of simultaneous inhibition of α -arrestins possibly through targeting the arrestin domain, and to identify and to functionally characterize proteins that bind *C. elegans* α -arrestins. Third, it is possible that three α -arrestins, *arrd-12*, *arrd-22* and *arrd-27*, which were not tested in our screen, may play crucial roles in lifespan regulation. Thus, it will be important to examine if genetic inhibition of each of these three α -arrestin genes affects lifespan in future studies.

In mammals, several α -arrestins are implicated in metabolic regulation⁵. TXNIP (thioredoxin-interacting protein), an inhibitor of thioredoxin in mammals^{32–34}, is a crucial negative regulator of glucose uptake^{35,36}. ARRDC4 inhibits glucose uptake in cultured mammalian cells as well³⁶, and ARRDC3 deficiency protects against obesity in male mice through increasing energy expenditure³⁷. Because metabolism is closely associated with aging³⁸, it will be interesting to test whether α -arrestins in complex metazoans play roles in aging via regulating metabolism.

Data availability

Dataset 1. Kaplan-Meier estimator of RNAi lifespan experiments. Kaplan-Meier estimate values were calculated from pooled lifespan data of two independent lifespan experiments using OASIS2 (<https://sbi.postech.ac.kr/oasis2/>). ‘At risk’ indicates the number of individuals at risk just prior to a specific time point. ‘S_hat’ indicates Kaplan-Meier estimate of survival function, ‘Var S_hat’ indicates variance of ‘S_hat’, and ‘SE(S)’ indicates standard error of ‘S_hat’. doi, [10.5256/f1000research.12337.d173158](https://doi.org/10.5256/f1000research.12337.d173158)³⁹

Dataset 2. Mean lifespan and mortality rates of RNAi experiment results. Mean lifespan, ages at different percent mortality, linear interpolation of mortality curve at specific mortality rate were calculated from pooled lifespan data of two independent lifespan experiments. Data were obtained using OASIS2 (<https://sbi.postech.ac.kr/oasis2/>). doi, [10.5256/f1000research.12337.d173159](https://doi.org/10.5256/f1000research.12337.d173159)⁴⁰

Dataset 3. Statistical analysis of lifespan data. Statistical test results (Chi square, *p* value, Bonferroni *p* value) were calculated between ‘condition 1’ and ‘condition 2’. Test results were obtained using OASIS2 (<https://sbi.postech.ac.kr/oasis2/>). doi, [10.5256/f1000research.12337.d173160](https://doi.org/10.5256/f1000research.12337.d173160)⁴¹

Competing interests

No competing interests were disclosed.

Grant information

This work was supported by the Korean Government (MSIP) through the National Research Foundation of Korea (NRF) [NRF-2016R1E1A1A01941152] funded to S-J.V.L.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgments

We thank the Lee laboratory members and Dr. Young Kwon for critical comments on the manuscript.

Supplementary material

Figure S1. RNAi knockdown of each *Caenorhabditis elegans* α -arrestin had minor effects on the lifespan of wild-type or *daf-2(-)* mutants. (A–U). Lifespan curves of wild-type (WT) and *daf-2(e1370)* (*daf-2(-)*) mutants treated with RNAi targeting *daf-16* (A), which was used as a positive control, *arrd-2* (B), *arrd-4* (C), *arrd-5* (D), *arrd-6* (E), *arrd-7* (F), *arrd-8* (G), *arrd-9* (H), *arrd-10* (I), *arrd-11* (J), *arrd-13* (K), *arrd-14* (L), *arrd-15* (M), *arrd-17* (N), *arrd-18* (O), *arrd-19* (P), *arrd-23* (Q), *arrd-25* (R), *arrd-26* (S), *arrd-28* (T) and *ttn-2* (U). See Table S2 for statistical analysis and for additional repeats of lifespan data using *daf-16* RNAi.

[Click here to access the data.](#)

Table S1. Possible RNAi target prediction. RNAi clones used in the screen were examined for their possible off-targets by Clone Mapper (<http://bioinformatics.lif.univ-mrs.fr/RNAiMap>)²². For RNAi clone names, ‘mv’ indicates an RNAi clone from Vidal library and ‘sjj’ indicates an RNAi clone from Ahringer library. RNAi clones generated by us for this study are labeled as ‘Lee lab’. Scores for potential target transcripts were calculated as described previously²² and were rounded off to the second digit after the decimal point.

[Click here to access the data.](#)

Table S2. Lifespan data of *Caenorhabditis elegans* α -arrestin RNAi clones. Lifespan data within the solid lines indicate the pooled experimental datasets that are performed at the same time by two independent researchers. *P* values were calculated within the sets using log-rank (Mantel-Cox) method²³. Percent (%) changes and *p* values for the lifespan data of specific α -arrestin RNAi-treated wild-type (WT) and *daf-2(e1370)* worms were calculated against those of control RNAi-treated WT and *daf-2(e1370)* worms within the sets, respectively. Percent (%) changes and *p* values for the lifespan data of control RNAi-treated *daf-2(e1370)* mutants were calculated against those of control RNAi-treated WT worms. SEM represents standard error of the mean, and 75th percentile indicates the age (days) at 75% mortality.

[Click here to access the data.](#)

References

1. Alvarez CE: **On the origins of arrestin and rhodopsin.** *BMC Evol Biol.* 2008; **8**: 222.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
2. Aubry L, Guetta D, Klein G: **The arrestin fold: variations on a theme.** *Curr Genomics.* 2009; **10**(2): 133–142.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
3. Polekhina G, Ascher DB, Kok SF, *et al.*: **Structure of the N-terminal domain of human thioredoxin-interacting protein.** *Acta Crystallogr D Biol Crystallogr.* 2013; **69**(Pt 3): 333–344.
[PubMed Abstract](#) | [Publisher Full Text](#)
4. Becuwe, M, Herrador A, Haguener-Tsapir R, *et al.*: **Ubiquitin-mediated regulation of endocytosis by proteins of the arrestin family.** *Biochem Res Int.* 2012; **2012**: 242764.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
5. Patwari P, Lee RT: **An expanded family of arrestins regulate metabolism.** *Trends Endocrinol Metab.* 2012; **23**(5): 216–222.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
6. Puca L, Brou C: **A-arrestins - new players in Notch and GPCR signaling pathways in mammals.** *J Cell Sci.* 2014; **127**(Pt 7): 1359–1367.
[PubMed Abstract](#) | [Publisher Full Text](#)
7. Hobert O: **The neuronal genome of *Caenorhabditis elegans*.** *WormBook.* 2013: 1–106.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
8. Huffman DL, Abrami L, Sasik R, *et al.*: **Mitogen-activated protein kinase pathways defend against bacterial pore-forming toxins.** *Proc Natl Acad Sci USA.* 2004; **101**(30): 10995–11000.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
9. Jee C, Choi TW, Kalichamy K, *et al.*: **CNP-1 (ARRD-17), a novel substrate of calcineurin, is critical for modulation of egg-laying and locomotion in response to food and lysine sensation in *Caenorhabditis elegans*.** *J Mol Biol.* 2012; **417**(3): 165–178.
[PubMed Abstract](#) | [Publisher Full Text](#)
10. Tissenbaum HA: **Using *C. elegans* for aging research.** *Invertebr Reprod Dev.* 2015; **59**(sup1): 59–63.
[Publisher Full Text](#) | [Free Full Text](#)
11. An SWA, Artan M, Park S, *et al.*: **Longevity Regulation by Insulin/IGF-1 Signalling.** In *Ageing: Lessons from C. elegans*. A. Olsen and M.S. Gill, Editors. Springer International Publishing, 2017; 63–81.
[Publisher Full Text](#)
12. Altintas O, Park S, Lee SJ: **The role of insulin/IGF-1 signaling in the longevity of model invertebrates, *C. elegans* and *D. melanogaster*.** *BMB Rep.* 2016; **49**(2): 81–92.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
13. Kenyon CJ: **The genetics of ageing.** *Nature.* 2010; **464**(7288): 504–512.
[PubMed Abstract](#) | [Publisher Full Text](#)
14. Lee SJ, Hwang AB, Kenyon C: **Inhibition of respiration extends *C. elegans* life span via reactive oxygen species that increase HIF-1 activity.** *Curr Biol.* 2010; **20**(23): 2131–2136.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
15. Hwang W, Artan M, Seo M, *et al.*: **Inhibition of elongin C promotes longevity and protein homeostasis via HIF-1 in *C. elegans*.** *Aging Cell.* 2015; **14**(6): 995–1002.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
16. Lee D, Jeong DE, Son HG, *et al.*: **SREBP and MDT-15 protect *C. elegans* from glucose-induced accelerated aging by preventing accumulation of saturated fat.** *Genes Dev.* 2015; **29**(23): 2490–2503.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
17. Seo M, Seo K, Hwang W, *et al.*: **RNA helicase HEL-1 promotes longevity by specifically activating DAF-16/FOXO transcription factor signaling in *Caenorhabditis elegans*.** *Proc Natl Acad Sci U S A.* 2015; **112**(31): E4246–E4255.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
18. Samuelson AV, Carr CE, Ruvkun G: **Gene activities that mediate increased life span of *C. elegans* insulin-like signaling mutants.** *Genes Dev.* 2007; **21**(22): 2976–2994.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
19. Stiernagle T: **Maintenance of *C. elegans*.** *WormBook.* 2006; 1–11.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
20. Sievers F, Higgins DG: **Clustal omega.** *Curr Protoc Bioinformatics.* 2014; **48**: 3.13.1–3.13.16.
[PubMed Abstract](#) | [Publisher Full Text](#)
21. Huson DH, Scornavacca C: **Dendroscope 3: an interactive tool for rooted phylogenetic trees and networks.** *Syst Biol.* 2012; **61**(6): 1061–1067.
[PubMed Abstract](#) | [Publisher Full Text](#)
22. Thakur N, Pujol N, Tichit L, *et al.*: **Clone mapper: an online suite of tools for RNAi experiments in *Caenorhabditis elegans*.** *G3 (Bethesda).* 2014; **4**(11): 2137–2145.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
23. Han, SK, Lee D, Lee H, *et al.*: **OASIS 2: online application for survival analysis 2 with features for the analysis of maximal lifespan and healthspan in aging research.** *Oncotarget.* 2016; **7**(35): 56147–56152.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
24. Kunitomo H, Uesugi H, Kohara Y, *et al.*: **Identification of ciliated sensory neuron-expressed genes in *Caenorhabditis elegans* using targeted pull-down of poly(A) tails.** *Genome Biol.* 2005; **6**(2): R17.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
25. Lynch AS, Briggs D, Hope IA: **Developmental expression pattern screen for genes predicted in the *C. elegans* genome sequencing project.** *Nat Genet.* 1995; **11**(3): 309–313.
[PubMed Abstract](#) | [Publisher Full Text](#)
26. Mounsey A, Bauer P, Hope IA: **Evidence suggesting that a fifth of annotated *Caenorhabditis elegans* genes may be pseudogenes.** *Genome Res.* 2002; **12**(5): 770–775.
[PubMed Abstract](#) | [Free Full Text](#)
27. Timmons L, Court DL, Fire A: **Ingestion of bacterially expressed dsRNAs can produce specific and potent genetic interference in *Caenorhabditis elegans*.** *Gene.* 2001; **263**(1–2): 103–112.
[PubMed Abstract](#) | [Publisher Full Text](#)
28. Kamath RS, Martinez-Campos M, Zipperlen P, *et al.*: **Effectiveness of specific RNA-mediated interference through ingested double-stranded RNA in *Caenorhabditis elegans*.** *Genome Biol.* 2001; **2**(1): RESEARCH0002.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
29. Kennedy S, Wang D, Ruvkun G: **A conserved siRNA-degrading RNase negatively regulates RNA interference in *C. elegans*.** *Nature.* 2004; **427**(6975): 645–649.
[PubMed Abstract](#) | [Publisher Full Text](#)
30. Simmer F, Tijsterman M, Parrish S, *et al.*: **Loss of the putative RNA-directed RNA polymerase RRF-3 makes *C. elegans* hypersensitive to RNAi.** *Curr Biol.* 2002; **12**(15): 1317–1319.
[PubMed Abstract](#) | [Publisher Full Text](#)
31. Sijen T, Fleener J, Simmer F, *et al.*: **On the role of RNA amplification in dsRNA-triggered gene silencing.** *Cell.* 2001; **107**(4): 465–476.
[PubMed Abstract](#) | [Publisher Full Text](#)
32. Junn E, Han SH, Im JY, *et al.*: **Vitamin D₃ up-regulated protein 1 mediates oxidative stress via suppressing the thioredoxin function.** *J Immunol.* 2000; **164**(12): 6287–6295.
[PubMed Abstract](#) | [Publisher Full Text](#)
33. Nishiyama A, Matsui M, Iwata S, *et al.*: **Identification of thioredoxin-binding protein-2/vitamin D₃ up-regulated protein 1 as a negative regulator of thioredoxin function and expression.** *J Biol Chem.* 1999; **274**(31): 21645–21650.
[PubMed Abstract](#) | [Publisher Full Text](#)
34. Yamanaka H, Maehira F, Oshiro M, *et al.*: **A possible interaction of thioredoxin with VDUP1 in HeLa cells detected in a yeast two-hybrid system.** *Biochem Biophys Res Commun.* 2000; **271**(3): 796–800.
[PubMed Abstract](#) | [Publisher Full Text](#)
35. Parikh H, Carlsson E, Chutkow WA, *et al.*: **TXNIP regulates peripheral glucose metabolism in humans.** *PLoS Med.* 2007; **4**(5): e158.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
36. Patwari P, Chutkow WA, Cummings K, *et al.*: **Thioredoxin-independent regulation of metabolism by the alpha-arrestin proteins.** *J Biol Chem.* 2009; **284**(37): 24996–25003.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
37. Patwari P, Emilsson V, Schadt EE, *et al.*: **The arrestin domain-containing 3 protein regulates body mass and energy expenditure.** *Cell Metab.* 2011; **14**(5): 671–683.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
38. Finkel T: **The metabolic regulation of aging.** *Nat Med.* 2015; **21**(12): 1416–1423.
[PubMed Abstract](#) | [Publisher Full Text](#)
39. Park S, Jung Y, An SWA, *et al.*: **Dataset 1 in: RNAi targeting *Caenorhabditis elegans*. α -arrestins marginally affects lifespan.** *F1000Research.* 2017.
[Data Source](#)
40. Park S, Jung Y, An SWA, *et al.*: **Dataset 2 in: RNAi targeting *Caenorhabditis elegans*. α -arrestins marginally affects lifespan.** *F1000Research.* 2017.
[Data Source](#)
41. Park S, Jung Y, An SWA, *et al.*: **Dataset 3 in: RNAi targeting *Caenorhabditis elegans*. α -arrestins marginally affects lifespan.** *F1000Research.* 2017.
[Data Source](#)

Open Peer Review

Current Peer Review Status:

Version 3

Reviewer Report 15 November 2017

<https://doi.org/10.5256/f1000research.14215.r27946>

© 2017 Bitto A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Alessandro Bitto

Department of Pathology, University of Washington, Seattle, Seattle, WA, USA

I appreciate the wish for the authors to move past this set of experiments and communicate their results to the field. Nevertheless, I must reiterate that, without testing the efficacy of their RNAi, their results cannot be interpreted, rather than be "interpreted with caution" as the authors suggest in the discussion.

Indeed, the authors themselves agree in their response to my previous review that "providing evidence for RNAi targeting *arrd* genes is a key for concluding that knockdown of *arrd* genes does not affect lifespan."

The authors suggest that performing qPCR on a new set of RNAi experiments would not demonstrate whether the knockdown worked on the worms followed for lifespan curves. While technically correct, I think it is fair to assume that their RNAi results are reproducible enough to warrant the assumption that whatever happens in a new set of RNAi experiments has happened in a previous one.

The authors bring forth a third reason for not testing the efficiency of their RNAi by claiming that their results are still useful to the field as they are. In fact, as I have suggested in my previous review, without qPCR testing of the efficacy of their RNAi, their results cannot be distinguished from the natural variation in lifespan of a control population of N2 worms, and are thus potentially meaningless.

In all honesty, I do not think that performing 24 qPCR reactions is an unreasonable request, especially when it is agreed that it would be "key for concluding that knockdown of *arrd* genes does not affect lifespan", which is the main point of the manuscript.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Biology of Aging

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response (F1000Research Advisory Board Member) 06 Dec 2017

Seung-Jae V Lee, Department of Biological Sciences, Korea Advanced Institute of Science and Technology, South Korea

Dear Dr. Alessandro Bitto,

We greatly appreciate your time and efforts for further valuable comments. We totally agree with your concern that we do not know the degree of RNAi knockdown efficiency in worms except *daf-16* RNAi, which was used as a positive control and showed expected lifespan phenotypes. Therefore, to precisely describe what we did, we changed "RNAi knockdown" to "feeding bacteria expressing dsRNA" throughout the manuscript. We sincerely thank you for fixing our misleading interpretation.

Competing Interests: No competing interests were disclosed.

Version 2

Reviewer Report 26 October 2017

<https://doi.org/10.5256/f1000research.13946.r27143>

© 2017 Bitto A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Alessandro Bitto

Department of Pathology, University of Washington, Seattle, Seattle, WA, USA

The authors took effort to address most of the concerns we raised in our initial review, but did not address at all our major concern, apart from a note in the discussion about the limitations of their results.

Thus our major concern stands: the authors must provide evidence that RNAi knockdown is functioning for each of the genes tested before their lifespan results can be interpreted correctly. Until such evidence is provided, there is no way of determining whether knockdown of individual arrestins really has limited effect on lifespan.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Biology of Aging

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response (F1000Research Advisory Board Member) 02 Nov 2017

Seung-Jae V Lee, Department of Biological Sciences, Korea Advanced Institute of Science and Technology, South Korea

Dear Dr. Alessandro Bitto,

We greatly appreciate your comments and respect your concerns on our paper. We totally agree with your point that providing evidence for RNAi targeting *arrd* genes is a key for concluding that knockdown of *arrd* genes does not affect lifespan. Nevertheless, we think it is more beneficial for the research field and for our lab to publish our data as they are, instead of not publishing a paper or performing all the qRT-PCR experiments retrospectively because of the following reasons.

First, we emphasized the limitation and the pitfall of our data thoroughly and therefore the readers will be fully aware of that.

Second, even if we perform the *arrd* RNAi experiments again and measure mRNA levels now, we will not be able to know whether the RNAi worked for the ones that we used for our lifespan screen. In addition, we do not have a plan on working on *arrd* genes any more. Third, with all these issues and limitations in mind, we think our negative data are still useful for the field. If some researchers want to do lifespan experiments for *arrd* genes in the future, they would rather use RNAi-sensitive strains and/or mutants by knowing our negative results. We think publishing these results is better for the field than keeping the data in our lab unpublished.

In any case, we sincerely appreciate your efforts for evaluating our manuscript.

Competing Interests: No competing interests were disclosed.

Reviewer Report 24 October 2017

<https://doi.org/10.5256/f1000research.13946.r25721>

© 2017 Dong M. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Meng-Qiu Dong

National Institute of Biological Sciences (NCBS), Beijing, China

In this paper, Park et al. knocked down 24 out of 29 alpha-arrestin genes in *C. elegans* and found

that none of them produced a sizable effect on the lifespans of the wild type or the long-lived daf-2 worms. The methods used in this study were sound, and the data quality were up to standard with hundreds of worms assayed for RNAi of each alpha-arrestin gene. I think the work is honest, and the negative results in this paper are reliable.

Concerns:

- (1) I agree with the previous 2 reviewers that it is strange to leave out 5 members of the C. elegans alpha-arrestin genes.
- (2) I think it is fair to say that none of the 24 genes extended lifespan when knocked down. The statistically significant 10% increase of wild-type lifespan by arrd-3 RNAi is negligible, given the variation of the lifespan assay. The relatively small p-value (0.0045) is blessed by the large sample size (n = 240). I wonder what the p-values would be if the two independent lifespan experiments had not been pooled.
- (3) I would like the authors to report how many worms were censored.
- (4) I wonder whether the authors had observed any noticeable phenotypes from their RNAi experiments, in particular, egg-laying defects. Strong egg-laying defects are easy to detect. G-protein signaling and beta-arrestin regulate the egg-laying behavior.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

I cannot comment. A qualified statistician is required.

Are all the source data underlying the results available to ensure full reproducibility?

Partly

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: C. elegans aging

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response (F1000Research Advisory Board Member) 02 Nov 2017

Seung-Jae V Lee, Department of Biological Sciences, Korea Advanced Institute of Science

and Technology, South Korea

Dear Dr. Meng-Qiu Dong,

We sincerely thank you for your valuable comments on our manuscript. We admit the limitations of our research and revised our manuscript to downplay our results so as not to confuse the readers. We addressed your concerns and revised our manuscript as described below.

In this paper, Park et al. knocked down 24 out of 29 alpha-arrestin genes in *C. elegans* and found that none of them produced a sizable effect on the lifespans of the wild type or the long-lived *daf-2* worms. The methods used in this study were sound, and the data quality were up to standard with hundreds of worms assayed for RNAi of each alpha-arrestin gene. I think the work is honest, and the negative results in this paper are reliable.

Concerns:

(1) I agree with the previous 2 reviewers that it is strange to leave out 5 members of the *C. elegans* alpha-arrestin genes.

> We appreciate your comments. We leave out *arrd-20* and *arrd-21* because they are predicted to be pseudogenes (Wormbase, version WS259) that are expected not to encode functional proteins. In addition, we did not test *arrd-12*, *arrd-22* or *arrd-27* because we were not able to obtain the corresponding RNAi clones during our cloning. We addressed this issue in Methods and discussed potential lifespan-regulatory roles of those genes in Discussion as follows,

In Methods, RNAi clones: "RNAi clone for *arrd-12*, *arrd-22*, or *arrd-27* was not generated because we were not able to obtain transformed *E. coli* (HT115) colonies using *arrd-12* or *arrd-22* infusion reaction products, or the PCR product for *arrd-27* genomic or complementary DNA."

In Discussion: "Third, it is possible that three α -arrestins, *arrd-12*, *arrd-22* and *arrd-27*, which were not tested in our screen, may play crucial roles in lifespan regulation. Thus, it will be important to examine if genetic inhibition of each of these three α -arrestin genes affects lifespan in future studies."

(2) I think it is fair to say that none of the 24 genes extended lifespan when knocked down. The statistically significant 10% increase of wild-type lifespan by *arrd-3* RNAi is negligible, given the variation of the lifespan assay. The relatively small p-value (0.0045) is blessed by the large sample size ($n = 240$). I wonder what the p-values would be if the two independent lifespan experiments had not been pooled.

> We appreciate your valuable comments. We agree with you that the extent of lifespan changes by several *arrd* RNAi clones was small but statistically significant because of the large sample size. Even if the lifespan data are not pooled, many independent lifespan results for *arrd* RNAi clones were again statistically significant. In either case, pooled or not pooled, we think the main point of our data indicates little effects of *arrd* RNAi on lifespan.

(3) I would like the authors to report how many worms were censored.

> We appreciate your comments. We provided raw lifespan data that contain the number of censored worms on each day of counting in Dataset1. One can also easily know the total number of censored worms by the difference between the number of animals that died and the total number of worms in Supplementary Table S2.

(4) I wonder whether the authors had observed any noticeable phenotypes from their RNAi experiments, in particular, egg-laying defects. Strong egg-laying defects are easy to detect. G-protein signaling and beta-arrestin regulate the egg-laying behavior.

> We appreciate your comments. Because we focused on the lifespan phenotype of *arrd* RNAi, unfortunately we did not pay attention to other physiological phenotypes and might have missed egg-laying phenotypes caused by some *arrd* RNAi clones. It will be an important point for future research.

Competing Interests: No competing interests were disclosed.

Reviewer Report 20 October 2017

<https://doi.org/10.5256/f1000research.13946.r27142>

© 2017 Pocock R. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Roger Pocock

Development and Stem Cells Program, Monash Biomedicine Discovery Institute, Department of Anatomy and Developmental Biology, Monash University, Melbourne, Vic, Australia

The authors have partially addressed my concerns. Below I suggest further changes.

1. Change title from 'RNAi targeting *Caenorhabditis elegans* α -arrestins has small or no effects on lifespan' to 'RNAi targeting *Caenorhabditis elegans* α -arrestins has little effect on lifespan'
2. Also change from 'small or no effects' to 'little effect' throughout the text.
3. The authors state that they did not study *arrd*-12, 22 and 27 because of 'unknown technical problems'. It would be helpful for the field to be more specific here. Could they not be amplified?
4. In the results paragraph the authors state that 'knockdown of each of individual α -arrestin genes tended to slightly reduce lifespan in wild-type or in *daf-2(-)* mutants' This is a little misleading. Change to 'knockdown of specific α -arrestin genes caused a minor reduction of lifespan in wild-type or in *daf-2(-)* mutants'
5. Change from 'In addition, as we did not test whether RNAi targeting each of α -arrestin

genes was effective by using quantitative RT-PCR' to 'In addition, as we did not test whether RNAi targeting of each α -arrestin gene was effective by using quantitative RT-PCR'

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response (F1000Research Advisory Board Member) 02 Nov 2017

Seung-Jae V Lee, Department of Biological Sciences, Korea Advanced Institute of Science and Technology, South Korea

Dear Dr. Roger Pocock,

We greatly appreciate your comments on our manuscript. We further addressed your concerns and revised our manuscript as described below.

The authors have partially addressed my concerns. Below I suggest further changes.

1. Change title from 'RNAi targeting *Caenorhabditis elegans* α -arrestins has small or no effects on lifespan' to 'RNAi targeting *Caenorhabditis elegans* α -arrestins has little effect on lifespan'

> We appreciate your comment on the title. We changed the title as follows,
Title: RNAi targeting *Caenorhabditis elegans* α -arrestins has little effect on lifespan

2. Also change from 'small or no effects' to 'little effect' throughout the text.

> We also changed the 'small or no effects' to 'little effect' in the main text as follows,
In Abstract: "We found that knocking down each of 24 out of total 29 *C. elegans* α -arrestins

had little effect on lifespan."

In Introduction: "We found that knocking down each of the α -arrestins had little effect on the lifespan of wild-type and *daf-2* mutants."

In Discussion: "Here we showed that RNAi knockdown of individual *C. elegans* α -arrestins had little effect on lifespan in wild-type or *daf-2* mutants."

In Figure legend: "Figure 1. RNAi targeting each of *Caenorhabditis elegans* α -arrestins had little effect on lifespan."

3. The authors state that they did not study *arrd-12*, 22 and 27 because of 'unknown technical problems'. It would be helpful for the field to be more specific here. Could they not be amplified?

> We appreciate your comments. We were not able to obtain transformed *E. coli* HT115 with infusion reaction products for *arrd-12* or *arrd-22* RNAi clones, or the PCR product for *arrd-27* genomic or complementary DNA. We specified the 'unknown technical problems' by mentioning these details in the main text as follows,

In Methods, RNAi clones: "RNAi clone for *arrd-12*, *arrd-22*, or *arrd-27* was not generated because we were not able to obtain transformed *E. coli* (HT115) colonies using *arrd-12* or *arrd-22* infusion reaction products, or the PCR product for *arrd-27* genomic or complementary DNA."

4. In the results paragraph the authors state that 'knockdown of each of individual α -arrestin genes tended to slightly reduce lifespan in wild-type or in *daf-2(-)* mutants'

This is a little misleading. Change to 'knockdown of specific α -arrestin genes caused a minor reduction of lifespan in wild-type or in *daf-2(-)* mutants'

> We appreciate your comments. We changed the sentence as follows,

In Results: "We found that knockdown of specific α -arrestin genes caused a minor reduction of lifespan in wild-type or in *daf-2(-)* mutants (Figure 1B, Figure S1B-U, Table S2)."

5. Change from 'In addition, as we did not test whether RNAi targeting each of α -arrestin genes was effective by using quantitative RT-PCR' to 'In addition, as we did not test whether RNAi targeting of each α -arrestin gene was effective by using quantitative RT-PCR'

> We appreciate your comments. We changed the sentence as suggested.

In Discussion: "In addition, as we did not test whether RNAi targeting each α -arrestin gene was effective by using quantitative RT-PCR"

Competing Interests: No competing interests were disclosed.

Version 1

Reviewer Report 06 September 2017

<https://doi.org/10.5256/f1000research.13357.r25722>

© 2017 Pocock R. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Roger Pocock

Development and Stem Cells Program, Monash Biomedicine Discovery Institute, Department of Anatomy and Developmental Biology, Monash University, Melbourne, Vic, Australia

In this manuscript Park *et al.* performed RNAi knockdown of 24 of the 29 alpha-arrestin genes in *C. elegans* to ask whether they are important for lifespan regulation in wild type and daf-2 mutant animals. Their data shows that, in their experimental conditions, most alpha-arrestin genes are individually not important for regulating lifespan.

Concerns:

One major issue with the study is whether RNAi knockdown of the alpha-arrestin genes is working. As some alpha-arrestins are expressed in the nervous system, which is refractory to RNAi, this is a concern. To determine whether the RNAi knockdown is working then Q-RTPCR should be performed. However, this is not a small task for 24 genes, therefore, further experiments using sensitized genetic backgrounds for RNAi or elevated temperatures may tease out phenotypes for some alpha-arrestin genes. This could particularly be performed for the alpha-arrestin genes that when knocked down engender a slight effect on lifespan.

In parallel to this, the authors could align the alpha-arrestin gene sequences to ask if there is a common sequence that could be used by RNAi targeting to knockdown multiple alpha-arrestin genes simultaneously.

The authors knockdown of 24 of the 29 alpha-arrestin genes. I wonder why they didn't generate RNAi clones for the remaining 5, especially since they did generate 3 other constructs for their analysis. Potentially, the important alpha-arrestin genes is amongst those not examined.

The authors did not report any other behavioural or morphological phenotypes observed by alpha-arrestin gene knockdown. Any information on observed phenotypes would be of interest to the research community.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Partly

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Partly

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response (F1000Research Advisory Board Member) 16 Oct 2017

Seung-Jae V Lee, Department of Biological Sciences, Korea Advanced Institute of Science and Technology, South Korea

Dear Dr. Roger Pocock,

We sincerely appreciate your valuable comments on our manuscript. We agree with all your points and revised our manuscript to describe the limitations of this study. We would like to mention that the main motivation for publishing this manuscript is to provide our negative results to researchers who are interested in characterizing α -arrestins in *C. elegans*. Our point-by-point responses are described below. We hope you find our revised manuscript is suitable for publication in *F1000Research*.

In this manuscript Park et al. performed RNAi knockdown of 24 of the 29 alpha-arrestin genes in *C. elegans* to ask whether they are important for lifespan regulation in wild type and *daf-2* mutant animals. Their data shows that, in their experimental conditions, most alpha-arrestin genes are individually not important for regulating lifespan.

Concerns:

One major issue with the study is whether RNAi knockdown of the alpha-arrestin genes is working. As some alpha-arrestins are expressed in the nervous system, which is refractory to RNAi, this is a concern. To determine whether the RNAi knockdown is working then Q-RT-PCR should be performed. However, this is not a small task for 24 genes, therefore, further experiments using sensitized genetic backgrounds for RNAi or elevated temperatures may tease out phenotypes for some alpha-arrestin genes. This could particularly be performed for the alpha-arrestin genes that when knocked down engender a slight effect on lifespan.

> We agree with the reviewer's comment that the RNAi knockdown efficiency is a major concern in interpreting our screen results. It would have been great if we had performed qRT-PCR analysis. However, as the reviewers pointed out, this is not a small task. We currently do not have any plan for following up this project and want to inform other researchers of our screen data. Therefore, we further emphasized the pitfall of our results in the Discussion as follows,
In Discussion: "The lifespan-regulatory functions of α -arrestins remain largely unexplored at the organism level. Here we showed that RNAi knockdown of individual *C. elegans* α -

arrestins had small or no effects on lifespan in wild-type or *daf-2* mutants. Our study has limitations that need to be considered for interpretation. First, RNAi targeting some α -arrestins might be insufficient for causing strong lifespan phenotypes. This may be because many of *C. elegans* α -arrestins are predicted to be expressed in neurons^{7, 9, 24-26}, which are refractory to RNAi²⁷⁻²⁹. Lifespan assays using RNAi-hypersensitive mutants, including *rrf-3(-)* and *eri-1(-)* animals²⁹⁻³¹, treated with α -arrestin RNAi, or using α -arrestin null mutants will help address this issue. In addition, as we did not test whether RNAi targeting each of α -arrestin genes was effective by using quantitative RT-PCR, our negative data should be interpreted with caution."

In parallel to this, the authors could align the alpha-arrestin gene sequences to ask if there is a common sequence that could be used by RNAi targeting to knockdown multiple alpha-arrestin genes simultaneously.

> We appreciate the reviewer's comment. The α -arrestin family members share arrestin-domains (O. Hobert, *Wormbook*, 2013). Thus, it will be intriguing if we could examine whether targeting arrestin domain for RNAi to knock down multiple *arrd* genes simultaneously and if multiple *arrd* gene knock down has effects on lifespan. We modified our manuscript to address this issue in Discussion as follows, In Discussion: "Second, *C. elegans* α -arrestins may have functional redundancy considering the large number of the α -arrestin family members in *C. elegans* and their sequence similarity^{1, 7}, which may obscure examining the functional importance of each α -arrestin. In addition, some α -arrestins may mostly function by modulating their interacting proteins⁶. In this case, genetic inhibition of α -arrestins may rather subtly affect the functions of their interacting partners that directly regulate physiology, such as aging and longevity, causing weak or no phenotypes. Thus, it will be interesting to test the effects of simultaneous inhibition of α -arrestins possibly through targeting the arrestin domain, and to identify and to functionally characterize proteins that bind *C. elegans* α -arrestins."

The authors knockdown of 24 of the 29 alpha-arrestin genes. I wonder why they didn't generate RNAi clones for the remaining 5, especially since they did generate 3 other constructs for their analysis. Potentially, the important alpha-arrestin genes is amongst those not examined.

> We appreciate the reviewers' comment. We aimed at obtaining all the α arrestin RNAi clones that are not commercially available, except two predicted pseudogenes, *arrd-20* and *arrd-21* (Wormbase, version WS259). We successfully constructed three RNAi clones targeting *arrd-11*, *arrd-17* and *arrd-26*, but could not obtain the other three RNAi clones, targeting *arrd-12*, *arrd-22* and *arrd-27*. We addressed this issue in Methods for RNAi clones and in Discussion as follows,

In Methods, RNAi clones: "RNAi clones for *arrd-12*, *arrd-22*, and *arrd-27* were not generated because of unknown technical problems and were not included in the lifespan screen. *arrd-20* and *arrd-21* are predicted to be pseudogenes (Wormbase, version WS259) and were excluded from our screen."

In Discussion: "Third, it is possible that three α -arrestins, *arrd-12*, *arrd-22* and *arrd-27*, which were not tested in our screen, may play crucial roles in lifespan regulation. Thus, it will be important to examine if genetic inhibition of each of these three α -arrestin genes affects

lifespan in future studies.”

The authors did not report any other behavioural or morphological phenotypes observed by alpha-arrestin gene knockdown. Any information on observed phenotypes would be of interest to the research community.

> We thank the reviewer for the valuable comment. One noticeable phenotype we observed was that the eggs laid by worms fed with *ttn-2* RNAi bacteria hatched more frequently than control eggs on FUDR-treated plates, although we did not quantify the results. We described this issue in Methods as follows,

In Methods, RNAi screen using lifespan assays: “Eggs laid by *ttn-2* RNAi-treated worms hatched more frequently than control eggs on FUDR-containing plates in two independent lifespan experiments.”

Competing Interests: No competing interests were disclosed.

Reviewer Report 04 September 2017

<https://doi.org/10.5256/f1000research.13357.r25290>

© 2017 Bitto A et al. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Alessandro Bitto

Department of Pathology, University of Washington, Seattle, Seattle, WA, USA

Jason Pitt

Department of Pathology, University of Washington, Seattle, WA, USA

In the manuscript *RNAi targeting Caenorhabditis elegans alpha-arrestins marginally affects lifespan* Park and co-workers provide largely negative results from lifespan experiments following RNAi knockdown of most of the worm alpha-arrestins. While we welcome the effort to publish negative results, there is one major concern with the experimental approach and several minor concerns with the interpretation of the results that should be addressed before the manuscript is fully accepted for indexing in *F1000Research*.

Major concern:

The authors data is almost entirely negative: there were only 4 clones which had a statistically significant effect. The authors attempt to address this concern by providing RNAi data for *daf-16*, which is known to strongly suppress the lifespan extension of *daf-2(e1370)* as well as decreasing N2 lifespan. However, the authors must demonstrate that alpha-arrestin RNAi knockdown is functioning. As suggested by the authors, there is some evidence that alpha-arrestins are expressed in neurons which are notoriously refractory to RNAi. The authors must show that they can use RNAi to knockdown alpha-arrestins by QRT-PCR analysis. The authors could also address

this concern by performing the RNAi in one of the mutant strains in which RNAi works in neurons and report if the lifespan still does not change, as they suggest in the discussion. We believe that providing these data is necessary to correctly interpret the results from the lifespan analyses.

Minor concerns:

1. In the text the authors speak broadly of individual arrestins having no role in longevity however the authors only tested 24 of the 29 arrestins in the worm. If the authors had limited themselves to only those clones available in commercial RNAi libraries this would be understandable, however the authors did construct 3 new RNAi feeding constructs. This brings up the question to why the authors did not prepare the remaining 5 constructs. If the authors have reason to exclude those genes from their analysis (ie. they are pseudogenes, not expressed, etc) they should discuss these reasons in the text. Besides, there is the possibility that one of the missing constructs may indeed have major effects on lifespan in *C. elegans*.
2. The title of the manuscript implies there are more than one result than was observed. The simpler conclusion from the results provided is that individual alpha-arrestin knockdown has no effect of lifespan. This is the fundamental observation of the paper considering that most of them (20 out of 24) had no effect and that 4 of them had such an extremely small effect that it is arguable that the sample size is insufficient to support any conclusion¹. We understand the authors' motivation to make the manuscript sound more appealing, however, their interpretation is disingenuous, given their results.
3. Because alpha-arrestins share significant sequence homology the authors should make some mention as to the likelihood that RNAi of any one arrestin would have off target effects on other arrestins. The authors could easily address this question experimentally with QRTPCR analysis. This is particularly relevant as one of the main conclusions of the manuscript is that future alpha-arrestin studies should knockdown multiple alpha-arrestins simultaneously.
4. Given that the authors observed the RNAi treated animals for weeks or months (in daf-2) it is surprising that no other behavioral or physical phenotypes from the alpha-arrestin RNAi were observed during the course of the analysis. If the authors truly did not observe any other phenotypes they should state it briefly in the text.

References

1. Petrascheck M, Miller DL: Computational Analysis of Lifespan Experiment Reproducibility. *Front Genet.* 2017; **8**: 92 [PubMed Abstract](#) | [Publisher Full Text](#)

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Partly

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Partly

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

No

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Biology of Aging

We confirm that we have read this submission and believe that we have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however we have significant reservations, as outlined above.

Author Response (F1000Research Advisory Board Member) 16 Oct 2017

Seung-Jae V Lee, Department of Biological Sciences, Korea Advanced Institute of Science and Technology, South Korea

Dear Dr. Alessandro Bitto and Dr. Jason Pitt,

We greatly appreciate the reviewers' valuable comments on our manuscript. We agree with all the reviewers' comments and modified our manuscript to describe the limitations of our studies raised by the reviewers. We would like to mention that the main purpose of publishing our current manuscript is to provide our data to other researchers who are interested in the characterization of *C. elegans arrd* genes, instead of keeping our current data unpublished. Our point-by-point responses are described below. We hope you find our revised manuscript is suitable for publication in *F1000Research*.

Reviewers' comments:

In the manuscript RNAi targeting *Caenorhabditis elegans* alpha-arrestins marginally affects lifespan Park and co-workers provide largely negative results from lifespan experiments following RNAi knockdown of most of the worm alpha-arrestins. While we welcome the effort to publish negative results, there is one major concern with the experimental approach and several minor concerns with the interpretation of the results that should be addressed before the manuscript is fully accepted for indexing in *F1000Research*.

Major concern:

The authors data is almost entirely negative: there were only 4 clones which had a statistically significant effect. The authors attempt to address this concern by providing RNAi data for *daf-16*, which is known to strongly suppress the lifespan extension of *daf-2(e1370)* as well as decreasing N2 lifespan. However, the authors must demonstrate that alpha-arrestin RNAi knockdown is functioning. As suggested by the authors, there is some evidence that alpha-arrestins are expressed in neurons which are notoriously refractory to RNAi. The authors must show that they can use RNAi to knockdown alpha-arrestins by

QRTPCR analysis. The authors could also address this concern by performing the RNAi in one of the mutant strains in which RNAi works in neurons and report if the lifespan still does not change, as they suggest in the discussion. We believe that providing these data is necessary to correctly interpret the results from the lifespan analyses.

> We completely agree with the reviewers for the comment. It would have been great if we had performed qRTPCR for validating the RNAi efficiency, while we were doing lifespan assays. As our experiments were done as an RNAi screen, our initial plan was validating RNAi efficiency and using RNAi-sensitive mutants if we had identified some positive RNAi clones. However, as the reviewers pointed out, our screen results were negative. We currently do not have any plan for following up this project and just want to inform other researchers of our screen data. Therefore, instead of putting efforts on this project retrospectively, we emphasized the limitations of our data raised by the reviewers to inform the readers of pitfalls by revising the manuscript as follows,

In Discussion: "The lifespan-regulatory functions of α -arrestins remain largely unexplored at the organism level. Here we showed that RNAi knockdown of individual *C. elegans* α -arrestins had small or no effects on lifespan in wild-type or *daf-2* mutants. Our study has limitations that need to be considered for interpretation. First, RNAi targeting some α -arrestins might be insufficient for causing strong lifespan phenotypes. This may be because many of *C. elegans* α -arrestins are predicted to be expressed in neurons^{7, 9, 24-26}, which are refractory to RNAi²⁷⁻²⁹. Lifespan assays using RNAi-hypersensitive mutants, including *rrf-3*(-) and *eri-1*(-) animals²⁹⁻³¹, treated with α -arrestin RNAi, or using α -arrestin null mutants will help address this issue. In addition, as we did not test whether RNAi targeting each of α -arrestin genes was effective by using quantitative RT-PCR, our negative data should be interpreted with caution."

Minor concerns:

1. In the text the authors speak broadly of individual arrestins having no role in longevity however the authors only tested 24 of the 29 arrestins in the worm. If the authors had limited themselves to only those clones available in commercial RNAi libraries this would be understandable, however the authors did construct 3 new RNAi feeding constructs. This brings up the question to why the authors did not prepare the remaining 5 constructs. If the authors have reason to exclude those genes from their analysis (ie. they are pseudogenes, not expressed, etc) they should discuss these reasons in the text. Besides, there is the possibility that one of the missing constructs may indeed have major effects on lifespan in *C. elegans*.

> We appreciate the reviewers' comment. We initially aimed at obtaining all the α arrestin RNAi clones that are not commercially available except two predicted pseudogenes, *arrd-20* and *arrd-21* (Wormbase, version WS259). We successfully constructed three RNAi clones targeting *arrd-11*, *arrd-17* and *arrd-26*, but could not obtain the other three RNAi clones for *arrd-12*, *arrd-22* and *arrd-27* because of unknown technical problems. We agree with the reviewers' point that these three genes that were not tested may have major effects on lifespan. We addressed this issue in Methods for RNAi clones as well as in Discussion as follows,

In Methods, RNAi clones: "RNAi clones for *arrd-12*, *arrd-22*, and *arrd-27* were not generated because of unknown technical problems and were not included in the lifespan screen. *arrd-*

20 and *arrd-21* are predicted to be pseudogenes (Wormbase, version WS259) and were excluded from our screen."

In Discussion: "Third, it is possible that three α -arrestins, *arrd-12*, *arrd-22* and *arrd-27*, which were not tested in our screen, may play crucial roles in lifespan regulation. Thus, it will be important to examine if genetic inhibition of each of these three α -arrestin genes affects lifespan in future studies."

2. The title of the manuscript implies there are more than one result than was observed. The simpler conclusion from the results provided is that individual alpha-arrestin knockdown has no effect of lifespan. This is the fundamental observation of the paper considering that most of them (20 out of 24) had no effect and that 4 of them had such an extremely small effect that it is arguable that the sample size is insufficient to support any conclusion¹. We understand the authors' motivation to make the manuscript sound more appealing, however, their interpretation is disingenuous, given their results.

> We thank the reviewers' comment on the title and changed the title as follows,
Title: "RNAi targeting *Caenorhabditis elegans* α -arrestins has small or no effects on lifespan"

3. Because alpha-arrestins share significant sequence homology the authors should make some mention as to the likelihood that RNAi of any one arrestin would have off target effects on other arrestins. The authors could easily address this question experimentally with QRT-PCR analysis. This is particularly relevant as one of the main conclusions of the manuscript is that future alpha-arrestin studies should knockdown multiple alpha-arrestins simultaneously.

> We agree with the reviewers that the RNAi targeting each of α -arrestin genes may have caused off-target effects on other α -arrestin genes because of sequence similarities. We performed off-target prediction using Clone Mapper (<http://bioinformatics.lif.univ-mrs.fr/RNAiMap>), and at least with the prediction, the RNAi clones used in our screen do not appear to have major off-target effects on any other genes including α -arrestin genes, except *arrd-19* RNAi which may also target *arrd-21* which is a predicted pseudogene. Although qRT-PCR will be a straightforward way as the reviewers suggested, again currently we do not have any plan on following up this project and we therefore instead acknowledged the limitation. We added the prediction results in the Supplementary Table S1 and described this issue in the Methods, RNAi clones part as follows,
In Methods, RNAi clones: "All RNAi clones used in our screen were examined for their potential off-target effects by using Clone Mapper (<http://bioinformatics.lif.univ-mrs.fr/RNAiMap>)²², and no significant off-target was predicted except *arrd-19* RNAi, which may additionally target a predicted pseudogene *arrd-21* (Table S1). Experimental validation by qRT-PCR will be necessary to completely exclude the possible off-target effects by RNAi clones."

In Supplementary material: "Table S1. Possible RNAi target prediction. RNAi clones used in the screen were examined for their possible off-targets by Clone Mapper (<http://bioinformatics.lif.univ-mrs.fr/RNAiMap>)²². For RNAi clone names, 'mv' indicates a RNAi clone from Vidal library and 'sjj' indicates a RNAi clone from Ahinger library. RNAi clones generated by us for this study are labeled as 'Lee lab'. Scores for potential target transcripts were calculated as described previously²² and were rounded off to the second

digit after the decimal point.”

In Reference: 22. Thakur, N., et al., Clone Mapper: An Online Suite of Tools for RNAi Experiments in *Caenorhabditis elegans*. G3 (Bethesda), 2014. 4(11): p. 2137-2145.23.

4. Given that the authors observed the RNAi treated animals for weeks or months (in *daf-2*) it is surprising that no other behavioral or physical phenotypes from the alpha-arrestin RNAi were observed during the course of the analysis. If the authors truly did not observe any other phenotypes they should state it briefly in the text.

> We appreciate the reviewers' comment. As our main interests were lifespan changes from a large scale screen, we did not pay attention to other phenotypes. Nevertheless, one noticeable phenotype we observed was that the eggs laid by worms fed with *ttm-2* RNAi bacteria hatched more frequently than control eggs on FUDR-treated plates, although we did not quantitate the results. We described this issue in Methods as follows, In Methods, RNAi screen using lifespan assays: “Eggs laid by *ttm-2* RNAi-treated worms hatched more frequently than control eggs on FUDR-containing plates in two independent lifespan experiments.”

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research