Check for updates

CASE REPORT

Case Report: Spontaneous cholecystocutaneous fistula, a rare choledocholithiasis complication [version 1; peer review: 3 approved]

Nunzio Maria Angelo Rinzivillo , Riccardo Danna, Vito Leanza, Melissa Lodato, Salvatore Marchese, Francesco Basile, Guido Nicola Zanghì

Department of Surgery, Policlinico Vittorio Emanuele University Hospital, University of Catania, Catania, Sicily, 95100, Italy

v1 First published: 27 Sep 2017, 6:1768
<https://doi.org/10.12688/f1000research.12235.1>
 Latest published: 27 Sep 2017, 6:1768
<https://doi.org/10.12688/f1000research.12235.1>

Abstract

One of the most unusual complications in choledocholithiasis is spontaneous cholecystocutaneous fistula, which has only been reported a few times in the literature. We report the case of a 76 year old man who presented with a right hypochondrium subcutaneous abscess, with pain evoked through palpation. No comorbidity in the patient's medical history were noted. Confirmation of cholecystocutaneous fistula was made using the proper diagnostic process, which is computed tomography with contrast media, followed by hepatobiliary MRI. This confirmed the presence of a fistulous pathway between the gallbladder and the skin. The patient underwent cholecystectomy surgery and open laparotomy with *en block* aponeurotic muscle, skin and fistula orifice excision.

Keywords

Cholecystocutaneous, Fistula, choledocholithiasis, abdominal fistula, gallbladder stones, gallbladder cancer, Bile Ducts, Laparotomy

Open Peer Review

Approval Status

	1	2	3
version 1			
27 Sep 2017	view	view	view
1. Matteo Novello , Cleveland Clinic, Cleveland, USA			
Salomone Di Saverio, Ospedale Maggiore, Bologna, Italy			
2. Antonino Agrusa , University of Palermo, Palermo, Italy			
3. Giuseppe Navarra , University Hospital of Messina, Messina, Italy			

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding authors: Melissa Lodato (melissalodato@gmail.com), Guido Nicola Zanghì (gzanghi@unict.it)

Author roles: Rinzivillo NMA: Writing – Original Draft Preparation; Danna R: Conceptualization; Leanza V: Supervision; Lodato M: Writing – Review & Editing; Marchese S: Resources; Zanghì GN: Investigation, Methodology, Supervision, Validation

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2017 Rinzivillo NMA *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Rinzivillo NMA, Danna R, Leanza V *et al.* Case Report: Spontaneous cholecystocutaneous fistula, a rare choledocholithiasis complication [version 1; peer review: 3 approved] F1000Research 2017, 6:1768
<https://doi.org/10.12688/f1000research.12235.1>

First published: 27 Sep 2017, 6:1768 <https://doi.org/10.12688/f1000research.12235.1>

Introduction

Spontaneous cholecystocutaneous abscess or fistula is an extremely uncommon complication of gallbladder disease. Less than 100 cases have been described in the literature. The first descriptions of cholecystocutaneous fistula was made by Thelusus in 1670. Later Courvoisier reported 169 cases of biliary fistula in the 19th century¹. The natural history of the disease has changed from suppurative cholecystitis with spontaneous rupture to operative external drainage of an abscess². Early and effective medical and surgical management of biliary tract disease can prevent this rare condition. Stone obstruction of the biliary tree plays a crucial role in the pathophysiology of the development of this condition; intra-gallbladder pressure can increase dramatically due to the obstruction of the cystic duct. The unresolved obstruction of the bile outflow compromise gallbladder wall blood circulation, as well as lymphatic drainage, resulting in necrosis of the gallbladder wall with the fistula formation. Once pierced, the gallbladder may drain into the peritoneal cavity, causing peritoneal localized abscess or the abscess can lead into an external fistula due to its adherence to the abdominal wall¹⁻³.

Case report

A 76 year old man was admitted to our University Hospital “Ospedale Vittorio Emenuele” and seen in our surgical department. He presented with a 3 cm tumefaction of the right hypochondrium, surrounded by an erythematous skin area, with small secretion of a yellowing-green material, attributable to a bile leaking (Figure 1). The patient’s medical history was clear from previous medical disease and surgery; he only referred to previous upper right quadrant pain and nonspecific dyspeptic disorder.

An abdominal ultrasound examination revealed the presence of a lesion in the aponeurotic muscle wall, but the possible underlying pathology was unknown. No other signs of pathology were observed. Routine blood work was normal.

Abdominal computed tomography scan with contrast media showed the gallbladder walls had diffuse thickened and blurred edges, and the right and transverse abdominal muscles were almost covered and embedded with minute hypo-dense ailments compatible with relapsing phlogistic processes (Figure 2). Hepatobiliary MRI detected that the gallbladder had anteriorly shifted and adhered

to the right abdominal muscles. The side wall showed a break through with consensual purulent collection, which extruded through the thick abdominal wall (Figure 3). Eventually, several different-sized stones were revealed inside the cholecyst. Consequently, a diagnosis of spontaneous cholecystocutaneous fistula was made.

The patient underwent cholecystectomy surgery 10 days after diagnosis, with an open laparotomy with *en block* aponeurotic muscle, skin and fistula orifice excision. In order to have a good abdominal wall reconstruction, a properly shaped Prolene Prosthesis was placed using fibrin glue⁴⁻⁷. The patient received broad-spectrum antibiotics after surgery (Piperacillin/Tazobactam 4.5 g, 3 times a day for 5 days via IV).

Considering the patient’s good general condition and good post-operative course, discharge was on the seventh day post-surgery. Surgical wound re-dressing was made one week after discharge in our facility, where the surgical stitches were removed. Scar appearance was good and without concern. No other dressing was needed and the scar was covered with a bandage. The first follow-up was scheduled 15 days after discharge, second one 60 days from discharge. For both follow-ups, routine blood work and surgical scar checking were performed, the results of which were normal and the scar was healing normally.

A histological examination confirmed the diagnosis of chronic cholecystitis with gallstones and cholecystocutaneous fistula.

Figure 2. Abdominal computed tomography images. Left image, * indicates the gallbladder; right image, the arrows indicates the fistula.

Figure 1. Cholecystocutaneous fistula, macroscopic appearance.

Figure 3. Hepatobiliary MRI. The white arrows show the fistula. From left to right: coronal and sagittal fistula view.

Discussion

Thanks to the progress made with medical imaging and surgical techniques, biliary fistula is today a very rare pathology⁸⁻¹¹. Fistulas often represent the result of post-surgical¹² or post-traumatic¹⁵ complications that generally involve the duodenum (77%) and colon (15%)¹⁶.

Spontaneous cholecystocutaneous fistula represents a truly exceptional event, as confirmed by the analysis of the literature, which revealed only 28 cases published over the last 10 years (Table 1). This disease mainly affects female subjects over the age of 60. Etiology is generally due to an acute inflammatory process as a consequence of a cholecystitis or chronic gallstones disease¹⁷⁻²⁰, although there are described cases of spontaneous cholecystocutaneous

fistula in the absence of gallstones²¹. Rarely does cholecystocutaneous fistula evolve into a neoplastic process. Instead, fistula can be a sign of gallbladder cancer^{19,20}. According to Sibakoti, polyarteritis nodosa with gallbladder vasculitis and prolonged use of high dose steroids can be considered predisposing factors²¹. Fistula *primum movens* is by cystic duct obstruction, which increases the pressure within the gallbladder, with wall distension and impaired vascularization, resulting in the formation of focal necrosis of the wall with perforation evolution and abscess formation to the surrounding area that will rupture in to the continuous structures. In the present case, the abscess drained through the abdominal wall and the fistulose pathway originated from the bottom of the gallbladder. This area is the most distant from the cystic artery and physiologically the least vascularized and therefore more susceptible to ischemia¹⁷.

Table 1. Publications within the last 10 years concerning cholecystocutaneous fistula.

Author(s)	Year published	Number of cases	Country	Age	Gender	Treatment technique
Maynard <i>et al.</i> ³⁵	2016	1	United Kingdom	68	F	Open
Jayasinghe <i>et al.</i> ³⁶	2016	1	United Kingdom	>70	F	Open
Guardado-B <i>et al.</i> ³⁷	2015	1	Mexico	30	F	Open
Álvarez <i>et al.</i> ³⁸	2014	1	Argentina	79	F	Open
Dixon <i>et al.</i> ³⁹	2014	1	United Kingdom	94	F	Open
Pripotnev and Petrakos ⁴⁰	2014	1	Canada	85	F	Open
Kim <i>et al.</i> ⁴¹	2013	1	Australia			Open
Jayant <i>et al.</i> ⁴²	2013	1	India	42	F	Open
Sodhi <i>et al.</i> ⁴³	2012	1	India	66	F	Open
Kapoor <i>et al.</i> ⁴⁴	2013	1	India	45	M	Open
Ozdemir <i>et al.</i> ⁴⁵	2012	2	India	45 & 65	M	Open
Ugalde Serrano <i>et al.</i> ⁴⁶	2012	1	Spain	83	M	Open
Andersen and Friis-Andersen ^{24,47}	2012	1	Denmark	89	F	Open
Ioannidis <i>et al.</i> ⁴⁸	2012	1	Greece	71	M	Open
Cheng <i>et al.</i> ⁴⁹	2011	1	China			Open
Gordon ⁵⁰	2011	1	U.S.A.	83	F	Open
Sayed <i>et al.</i> ⁵¹	2010	1	United Kingdom	85	F	Open
Pezzilli <i>et al.</i> ⁵²	2010	1	Italy	90	F	Open
Metsemakers <i>et al.</i> ⁵³	2010	1	Belgium	69	M	Open
Tallón Aguilar <i>et al.</i> ⁵⁴	2010	1	Spain	83	F	Open
Kahn <i>et al.</i> ⁵⁵	2010	1	Ireland	76	M	Open
Hawari <i>et al.</i> ⁵⁶	2010	1	United Kingdom	84	M	Open
Murphy <i>et al.</i> ⁵⁷	2008	1	United Kingdom	80	M	Open
Ijaz <i>et al.</i> ⁵⁸	2008	1	United Kingdom	80	F	Open
Chatterjee <i>et al.</i> ⁵⁹	2007	1	India	45	F	Open
Malik <i>et al.</i> ⁶⁰	2007	1	United Kingdom	76	F	Laparoscopic

The external fistular orifice is usually on the right upper quadrant, but other locations have been described, including the left hypocondrium, umbilical scar, right lumbar, and right iliac fossa, and rarely the right gluteus and breast region^{19,24,47}.

The diagnostic process always begins with upper abdomen ultrasound and ends with **hepatobiliary** MRI to visualize the biliary tree. Considering that 11% of cholecystitis have concomitant presence of gallstones in the main bile duct, it is advisable to perform endoscopic retrograde cholangiopancreatography (ERCP)^{2–29}. In our case CT with CM and hepatobiliary MRI confirmed the fistula presence and it was not necessary to execute the ERCP before surgery. Although an intraoperative cholangiogram was performed to check that the bile ducts were clear from gallstones. Cholecystocutaneous fistula has always been treated by two different strategies. The first includes a two-step approach: percutaneous drainage and antibiotic therapy, and subsequently cholecystectomy. The second directly involves laparotomy cholecystectomy execution with *en block* aponeurotic muscles, as well as skin and fistula orifice excision.

The second strategy is the most commonly used since the two-step approach treatment is reserved for patients with sepsis and poor general condition^{12,15,29}.

In 1998, Kumar described the first case of gynecological fistula treated with laparoscopic technique, proposing to the scientific community the feasibility of this innovative approach^{27,30}.

Conclusion

Rarity of this pathology confirms the great quality of progress made by early diagnostic techniques and medical treatment to prevent complication of cholethiasis. Although cholecystocutaneous spontaneous fistula is not common, it can lead to a serious condition. If not quickly treated, it can rapidly evolve into a generalized septic state with severe impairment prognosis. In our case, the patient was in good health arguably because the fistula was draining the most of the abscess outside the body and not in the peritoneum space. Surgical treatment was, however, essential to restore the physiologic bile flow and adequate broad-spectrum antibiotic prophylaxis lowered the risk of post-operative infections. Although laparoscopic approaches have been described since 1998, this pathology is, in most cases, continuing to be treated with open technique, most likely because it is easier and with fewer risks of post-surgical complications^{31–34}.

Consent

Written informed consent was obtained from the patient for the publication of the patient's clinical details and related images.

Competing interests

No competing interests were disclosed.

Grant information

The author(s) declared that no grants were involved in supporting this work.

References

- Flora HS, Bhattacharya S: **Spontaneous cholecystocutaneous fistula**. *HPB (Oxford)*. 2001; **3**(4): 279–280.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Gupta V, Benerjee S, Garg H, *et al.*: **Spontaneous cholecysto-antral-cutaneous fistula: a consequence of neglected calculus cholecystitis**. *Singapore Med J*. 2012; **53**(10): e201–3.
[PubMed Abstract](#)
- Mathew G, Bhimji SS: **Fistula, Cholecystocutaneous**. University of Pelita Harapan, 2017.
[PubMed Abstract](#)
- Zanghi G, Rinzivillo NM, Caponnetto AM, *et al.*: **Sentinel lymph node biopsy in breast cancer New indications and our experience**. *Ann Ital Chir*. 2015; **86**: 508–12.
[PubMed Abstract](#)
- Biondi A, Tropea A, Monaco G, *et al.*: **[Complications in the laparoscopic treatment of primary and secondary hernias of the abdominal wall]**. *Ann Ital Chir*. 2010; **81**(3): 193–198.
[PubMed Abstract](#)
- Zanghi G, Di Stefano G, Leanza V, *et al.*: **Incisional hernia in day surgery: our personal experience**. *G Chir*. 2012; **33**(6–7): 218–220.
[PubMed Abstract](#)
- Leanza V, Zanghi G, Vecchio R, *et al.*: **How to prevent mesh erosion in transobturator tension-free incontinence cystocoele treatment (TICT): a comparative survey**. *G Chir*. 2015; **36**(1): 21–25.
[PubMed Abstract](#) | [Free Full Text](#)
- Courvoisier L: **Pathologie und Chirurgie der Gallenwege**. Leipzig, Germany: FCW Vogel; 1890.
- Naunyn B: **Ulcerative affections of the biliary passage and fistula formation**. In: *A Treatise on Cholelithiasis*. New Sydenham Society; (English version 1896). New Sydenham Society; 1892; 138–151.
[Reference Source](#)
- Bonnet: **Fistule biliaire cutanee**. *Lyon Med*. 1897; **85**.
- Henry CL, Ort TG Jr: **Spontaneous external biliary fistulas**. *Surgery*. 1949; **26**(4): 641–646.
[PubMed Abstract](#)
- Czerniak A, Blumgart LH: **External biliary fistula. (External Biliary Fistula, Surgery of the Liver and Biliary Tract)**. Churchill Livingstone; Edinburgh, 1994; **1**.
- Smith EE, Bowley N, Allison DJ: **The management of post-traumatic intrahepatic cutaneous biliary fistulas**. *Br J Surg*. 1982; **69**(6): 317–318.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Glenn F, Reed C, Grafe WR: **Biliary enteric fistula**. *Surg Gynecol Obstet*. 1981; **153**(4): 527–531.
[PubMed Abstract](#)
- Cruz RJ Jr, Nahas J, de Figueiredo LF: **Spontaneous cholecystocutaneous fistula: a rare complication of gallbladder disease**. *Sao Paulo Med J*. 2006; **124**(4): 234–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Flora HS, Bhattacharya S: **Spontaneous cholecystocutaneous fistula**. *HPB (Oxford)*. 2001; **3**(4): 279–280.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ugalde Serrano P, Solar García L, Miyar de León A, *et al.*: **[Cholecystocutaneous fistula as a first sign of presentation of a gallbladder adenocarcinoma]**. *Cir Esp*. 2013; **91**(6): 396–397.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Zanghi G, Leanza V, Vecchio R, *et al.*: **Neoplastic sigmoid-uterine fistula. An exceptional complication of large intestine cancer**. *G Chir*. 2017; **38**(1): 37–40.
[PubMed Abstract](#)
- Kuo YC, Wu CS: **Spontaneous cutaneous biliary fistula: a rare complication of cholangiocarcinoma**. *J Clin Gastroenterol*. 1990; **12**(4): 451–453.
[PubMed Abstract](#)
- Birch BR, Cox SJ: **Spontaneous external biliary fistula uncomplicated by gallstones**. *Postgrad Med J*. 1991; **67**(786): 391–2.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Sibakoti YC, Basnet RB, Poudel R: **Spontaneous Cholecystocutaneous Fistula:**

A Case Report. *PMJM*. 2012; 12(1): 51–54.

Reference Source

22. Gibson TC, Howat JM: **Cholecystocutaneous fistula. *Br J Clin Pract*. 1987; 41(10): 980–2.**
PubMed Abstract
23. Nicholson T, Born MW, Garber E: **Spontaneous cholecystocutaneous fistula presenting in the gluteal region. *J Clin Gastroenterol*. 1999; 28(3): 276–7.**
PubMed Abstract | Publisher Full Text
24. Andersen P, Friis-Andersen H: **[Spontaneous cholecystocutaneous fistula presenting in the right breast]. *Ugeskr Laeger*. 2012; 174(18): 1235–1236.**
PubMed Abstract
25. Costi R, Gnocchi A, Di Mario F, et al.: **Diagnosis and management of choledocholithiasis in the golden age of imaging, endoscopy and laparoscopy. *World J Gastroenterol*. 2014; 20(37): 13382–13401.**
PubMed Abstract | Publisher Full Text | Free Full Text
26. Cacopardo C, Pinzone M, Berretta S, et al.: **Localized and systemic bacterial infections in necrotizing pancreatitis submitted to surgical necrosectomy or percutaneous drainage of necrotic secretions. *BMC Surg*. 2013; 13(Suppl 2): S50.**
PubMed Abstract | Publisher Full Text | Free Full Text
27. Bentivegna F, Cortese V, Bentivegna S, et al.: **Treatment of the pancreatic stump after DCP. *Eur Rev Med Pharmacol Sci*. 2014; 18(Suppl 2): 36–39.**
PubMed Abstract
28. Zanghi G, Leanza V, Rinzivillo NM, et al.: **Radio-guided surgery in breast cancer. *Euromediterranean Biomedical Journal*. 2016; 11(13): 101–106.**
Publisher Full Text
29. Vasanth A, Siddiqui A, O'Donnell K: **Spontaneous cholecystocutaneous fistula. *South Med J*. 2004; 97(2): 183–5.**
PubMed Abstract | Publisher Full Text
30. Kumar SS: **Laparoscopic management of a cholecystocutaneous abscess. *Am Surg*. 1998; 64(12): 1192–4.**
PubMed Abstract
31. Leanza V, Zanghi G, Leanza G, et al.: **A specific application of two psychological measures on pelvic organ prolapse patients. *Giornale Italiano di Ostetricia e Ginecologia*. 2015; 37(2): 82–86.**
Reference Source
32. Zanghi G, Leanza V, Vecchio R, et al.: **Single-Incision Laparoscopic Cholecystectomy: our experience and review of literature. *G Chir*. 2015; 36(6): 243–246.**
PubMed Abstract | Publisher Full Text | Free Full Text
33. Vecchio R, Marchese S, Famoso S, et al.: **Colorectal cancer in aged patients. Toward the routine treatment through laparoscopic surgical approach. *G Chir*. 2015; 36(1): 9–14.**
PubMed Abstract | Publisher Full Text | Free Full Text
34. Zanghi G, Di Stefano G, Caponnetto A, et al.: **Breast cancer and sentinel lymph node micrometastases: indications for lymphadenectomy and literature review. *G Chir*. 2014; 35(11–12): 260–65.**
PubMed Abstract | Free Full Text
35. Maynard W, McGlone ER, Deguarra J: **Unusual aetiology of abdominal wall abscess: cholecystocutaneous fistula presenting 20 years after open subtotal cholecystectomy. *BMJ Case Rep*. 2016; 2016: pii: bcr2015213326.**
PubMed Abstract | Publisher Full Text
36. Jayasinghe G, Adam J, Abdul-Aal Y: **Unusual presentation of gallbladder perforation. *Int J Surg Case Rep*. 2016; 18: 42–44.**
PubMed Abstract | Publisher Full Text | Free Full Text
37. Guardado-Bermúdez F, Aguilar-Jaimes A, Ardisson-Zamora FJ, et al.: **[Spontaneous cholecystocutaneous fistula]. *Cir Cir*. (English Edition). 2015; 83(1): 61–64.**
PubMed Abstract | Publisher Full Text
38. Álvarez Florencia, Meraldi A, Emery NC, et al.: **[Spontaneous cholecystocutaneous fistula in an elderly woman]. *Rev Med Chil*. 2014; 142(8): 1076–7.**
PubMed Abstract | Publisher Full Text
39. Dixon S, Sharma M, Holtham S: **Cholecystocutaneous fistula: an unusual complication of a para-umbilical hernia repair. *BMJ Case Rep*. 2014; 2014: pii: bcr2013202417.**
PubMed Abstract | Publisher Full Text | Free Full Text
40. Pripotnev S, Petrakos A: **Cholecystocutaneous fistula after percutaneous gallbladder drainage. *Case Rep Gastroenterol*. 2014; 8(1): 119–22.**
PubMed Abstract | Publisher Full Text | Free Full Text
41. Kim M, Beenen E, Fergusson J: **Spontaneous cholecystocutaneous fistula by gallstone erosion into abdominal wall. *ANZ J Surg*. 2014; 84(11): 888–89.**
PubMed Abstract | Publisher Full Text
42. Jayant M, Kaushik R, Attri AK: **Spontaneous cholecystocutaneous abscess. *Indian J Gastroenterol*. 2014; 33(5): 498.**
PubMed Abstract | Publisher Full Text
43. Sodhi K, Athar M, Kumar V, et al.: **Spontaneous cholecysto-cutaneous fistula complicating carcinoma of the gall bladder: a case report. *Indian J Surg*. 2012; 74(2): 191–93.**
PubMed Abstract | Publisher Full Text | Free Full Text
44. Kapoor Y, Singh G, Khokhar M: **Spontaneous cholecystocutaneous fistula-not an old time story. *Indian J Surg*. 2013; 75(Suppl 1): 188–91.**
PubMed Abstract | Publisher Full Text | Free Full Text
45. Ozdemir Y, Yucel E, Sucullu I, et al.: **Spontaneous cholecystocutaneous fistula as a rare complication of gallstones. *Bratisl Lek Listy*. 2012; 113(7): 445–47.**
PubMed Abstract | Publisher Full Text
46. Ugalde Serrano P, Solar García L, Miyar de León A, et al.: **[Cholecystocutaneous fistula as a first sign of presentation of a gallbladder adenocarcinoma]. *Cir Esp*. 2013; 91(6): 396–97.**
PubMed Abstract | Publisher Full Text
47. Andersen P, Friis-Andersen H: **[Spontaneous cholecystocutaneous fistula presenting in the right breast]. *Ugeskr Laeger*. 2012; 174(18): 1235–36.**
PubMed Abstract
48. Ioannidis O, Paraskevas G, Kotronis A, et al.: **Spontaneous cholecystocutaneous fistula draining from an abdominal scar from previous surgical drainage. *Ann Ital Chir*. 2012; 83(1): 67–69.**
PubMed Abstract
49. Cheng HT, Wu CI, Hsu YC: **Spontaneous cholecystocutaneous fistula managed with percutaneous transhepatic gallbladder drainage. *Am Surg*. 2011; 77(12): E285–286.**
PubMed Abstract
50. Gordon PE, Miller DL, Rattner DW, et al.: **Image of the Month—Quiz Case. *Arch Surg*. 2011; 146(4): 487–8.**
PubMed Abstract | Publisher Full Text
51. Sayed L, Sangal S, Finch G: **Spontaneous cholecystocutaneous fistula: a rare presentation of gallstones. *J Surg Case Rep*. 2010; 2010(5): 5.**
PubMed Abstract | Publisher Full Text | Free Full Text
52. Pezzilli R, Barakat B, Corinaldesi R, et al.: **Spontaneous Cholecystocutaneous Fistula. *Case Rep Gastroenterol*. 2010; 4(3): 356–60.**
PubMed Abstract | Publisher Full Text | Free Full Text
53. Metsemakers WJ, Quanten I, Vanhoenacker F, et al.: **Spontaneous Cholecystocutaneous Abscess. *JBR-BTR*. 2010; 93(4): 198–200.**
PubMed Abstract | Publisher Full Text
54. Tallón Aguilar L, López Porras M, Molina García D, et al.: **Fístula colecistocutánea. Una rara complicación de la coledocolitiasis. *Gastroenterol Hepatol*. 2010; 33(7): 553–54.**
Publisher Full Text
55. Khan AA, Azhar MZ, Khan AA, et al.: **Spontaneous Cholecystocutaneous Fistula. *J Coll Physicians Surg Pak*. 2005; 15(11): 726–27.**
PubMed Abstract
56. Hawari M, Wemyss-Holden S, Parry GW: **Recurrent Chest Wall Abscesses Overlying a Pneumonectomy Scar: An Unusual Presentation of a Cholecystocutaneous Fistula. *Interact Cardiovasc Thorac Surg*. 2010; 10(5): 828–29.**
PubMed Abstract | Publisher Full Text
57. Murphy JA, Vimalachandran CD, Howes N, et al.: **Anterior abdominal wall abscess secondary to subcutaneous gallstones. *Case Rep Gastroenterol*. 2008; 2(2): 219–23.**
PubMed Abstract | Publisher Full Text | Free Full Text
58. Ijaz S, Lidder S, Mohamid W, et al.: **Cholecystocutaneous fistula secondary to chronic calculous cholecystitis. *Case Rep Gastroenterol*. 2008; 2(1): 71–75.**
PubMed Abstract | Publisher Full Text | Free Full Text
59. Chatterjee S, Choudhuri T, Ghosh G, et al.: **Spontaneous Cholecystocutaneous Fistula in a Case of Chronic Coliculous Cholecystitis—a Case Report. *J Indian Med Assoc*. 2007; 105(11): 644, 646, 656.**
PubMed Abstract
60. Malik AH, Nadeem M, Ockrim J: **Complete laparoscopic management of cholecystocutaneous fistula. *Ulster Med J*. 2007; 76(3): 166–7.**
PubMed Abstract | Free Full Text

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 24 November 2017

<https://doi.org/10.5256/f1000research.13246.r28008>

© 2017 Navarra G. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Giuseppe Navarra

Surgical Oncology Unit, Department of Human Pathology of Adult and Evolutive Age, University Hospital of Messina, Messina, Italy

The article Case Report: Spontaneous cholecystocutaneous fistula, a rare cholelithiasis complication covers a subject that nowadays is quite uncommon. It goes through the background of the case's history, it provides details of diagnostic tests, treatment given and outcomes reached. It could be useful for other practitioners. It can be accepted for indexing with minor English revision.

Is the background of the case's history and progression described in sufficient detail?

Yes

Are enough details provided of any physical examination and diagnostic tests, treatment given and outcomes?

Yes

Is sufficient discussion included of the importance of the findings and their relevance to future understanding of disease processes, diagnosis or treatment?

Yes

Is the case presented with sufficient detail to be useful for other practitioners?

Yes

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 21 November 2017

<https://doi.org/10.5256/f1000research.13246.r28162>

© 2017 Agrusa A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Antonino Agrusa

Department of General Surgery, Urgency and Organ Transplantation, University of Palermo, Palermo, Italy

My compliments to the authors for the interesting and rare case report. I approve it for publication in F1000 Research.

Please give more explanation for the 10 days delay between patient admission and operative surgical management. On the basis of rare case report the authors should clarify the operative technique with more details.

Is the background of the case's history and progression described in sufficient detail?

Yes

Are enough details provided of any physical examination and diagnostic tests, treatment given and outcomes?

Yes

Is sufficient discussion included of the importance of the findings and their relevance to future understanding of disease processes, diagnosis or treatment?

Yes

Is the case presented with sufficient detail to be useful for other practitioners?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: General and emergency surgery

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 20 November 2017

<https://doi.org/10.5256/f1000research.13246.r27856>

© 2017 Novello M et al. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Matteo Novello

Colorectal Surgery Department, Cleveland Clinic, Cleveland, OH, USA

Salomone Di Saverio

General Surgery Department, Ospedale Maggiore, Bologna, Italy

The authors should clarify the reasons for the 10 days delay in the operative management of their patient.

More insight should be provided on the choice of aggressive operative treatment as compared to a percutaneous drainage. Perhaps the drainage would have been more readily available and also less expensive as compared to elective surgery. Keeping in mind the age of the patient, it might have been a reasonable option proceeding with conservative management and percutaneous drainage trial.

Did the authors investigate the use of somatostatin (provided in the form of Octreotide) in such case? It is currently a first line approach for uncomplicated pancreatic-cutaneous fistulas and might as well work with the Gallbladder considering the physiological functions of the hormone.

All in all the case is interesting and, considering the rarity of the event, it deserves publication.

Is the background of the case's history and progression described in sufficient detail?

Yes

Are enough details provided of any physical examination and diagnostic tests, treatment given and outcomes?

Yes

Is sufficient discussion included of the importance of the findings and their relevance to future understanding of disease processes, diagnosis or treatment?

Yes

Is the case presented with sufficient detail to be useful for other practitioners?

Yes

Competing Interests: No competing interests were disclosed.

We confirm that we have read this submission and believe that we have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research