REVIEW

Recent advances in eosinophilic esophagitis [version 1; peer review: 3 approved]

Sandy Durrani , Marc Rothenberg

Division of Allergy and Immunology, Department of Pediatrics, Cincinnati Children's Hospital Medical Center, University of Cincinnati, Cincinnati OH, 45529, USA

V1 First published: 28 Sep 2017, 6(F1000 Faculty Rev):1775
<https://doi.org/10.12688/f1000research.11798.1>

Latest published: 28 Sep 2017, 6(F1000 Faculty Rev):1775
<https://doi.org/10.12688/f1000research.11798.1>

Abstract

Eosinophilic esophagitis is a chronic, antigen-driven, eosinophil-predominant inflammatory disease of the esophagus and affects both children and adults. Cutting-edge technologies, such as genome-wide association studies, have advanced our understanding of the disease pathogenesis at a remarkable rate. Recent insights from genetic and mechanistic studies have concluded that a complex interplay between genetic and environmental risk factors, allergic sensitization, and esophageal-specific pathways leads to disease pathogenesis. Importantly, recent epidemiologic studies have found that the incidence and prevalence of eosinophilic esophagitis continue to rise. New guidelines have advocated the elimination of the term proton pump inhibitor (PPI)-responsive esophageal eosinophilia and have recommended using PPIs as a first-line treatment modality. Systemic reviews and meta-analyses confirm the efficacy of PPIs, topical corticosteroids, and empiric food elimination diets. Unmet needs include the development of birth cohort studies, validated diagnostic scoring systems, minimally invasive disease-monitoring methods, and the development of new therapies.

Keywords

Eosinophilic esophagitis, Proton-Pump inhibitor, Topical Corticosteroid

Open Peer Review

Approval Status

	1	2	3
version 1 28 Sep 2017

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Michael F Vaezi**, Vanderbilt University Medical Center, Nashville, USA
2. **Edaire Cheng**, University of Texas Southwestern Medical Center, Dallas, Texas, USA
Diana Montoya Melo, University of Texas Southwestern Medical Center, Dallas, Texas, USA
3. **Yoshikazu Kinoshita**, Shimane University School of Medicine, Shimane, Japan

Any comments on the article can be found at the end of the article.

Corresponding author: Marc Rothenberg (marc.rothenberg@cchmc.org)

Author roles: Durrani S: Writing – Original Draft Preparation, Writing – Review & Editing; Rothenberg M: Writing – Review & Editing

Competing interests: MER is a consultant for NKT Therapeutics, Pulm One, Spoon Guru, Celgene, Shire, Astra Zeneca, GlaxoSmithKline, and Novartis and has an equity interest in the first three listed and Immune Pharmaceuticals and royalties from reslizumab (Teva Pharmaceuticals). He is an inventor of patents, owned by Cincinnati Children's Hospital Medical Center. SRD declares that he has no competing interests.

Grant information: This work was supported in part by National Institutes of Health U19 AI070235, National Institutes of Health R01 AI124355, and R37 R37 A1045898; the Campaign Urging Research for Eosinophilic Disease Foundation; the Buckeye Foundation; the American Partnership for Eosinophilic Disorders; and the Sunshine Charitable Foundation and its supporters, Denise A. Bunning and David G. Bunning.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2017 Durrani S and Rothenberg M. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Durrani S and Rothenberg M. **Recent advances in eosinophilic esophagitis [version 1; peer review: 3 approved]** F1000Research 2017, 6(F1000 Faculty Rev):1775 <https://doi.org/10.12688/f1000research.11798.1>

First published: 28 Sep 2017, 6(F1000 Faculty Rev):1775 <https://doi.org/10.12688/f1000research.11798.1>

Introduction

Eosinophilic esophagitis (EoE) is a fascinating case study in modern medicine today as it was only recently recognized in the 1990s as a clinicopathologic disorder—a metaphorical infant when compared with other allergic diseases such as asthma and allergic rhinitis. Importantly, owing to the development and use of cutting-edge molecular diagnostic technology, the scientific breakthroughs in our understanding of EoE rival those of diseases recognized centuries ago. Nonetheless, much work remains to be done to understand this enigmatic food antigen-induced disorder of chronic esophageal inflammation of both children and adults. If left untreated, EoE can progress over time to a fibrostenotic disorder characterized by dysphagia and esophageal strictures. As a result of the overwhelming number of genetic, mechanistic, and clinical breakthroughs in EoE over the last two decades, the research and medical communities are in need of frequent and comprehensive reviews of the literature by international experts. Thus, the following review highlights and summarizes important epidemiologic, genetic, pathophysiologic, and clinical findings over the last three years.

Epidemiology

The incidence and prevalence of EoE have been rising sharply over the last 20 years in Western countries¹. Indeed, numerous studies seem to confirm a rapid rise in the incidence and prevalence of EoE². Recently, a systematic review of 13 population-based studies confirmed a higher incidence (7.2 per 100,000 person-years) and pooled prevalence (28.1 cases per 100,000 inhabitant-years) comparing studies before and after 2008. Moreover, the incidence and prevalence of EoE were higher in adults than in children and the incidence of EoE was higher in the US than in Europe³. Finally, there are also now reports of EoE cases in Africa—a heretofore undocumented finding⁴.

EoE has been classically observed in studies to be a disease of Caucasian males²; however, two recent studies have found that African-American male children have the expected burden of EoE seen in Caucasians^{5,6}. Weiler *et al.*⁶ found that African-American children presented at younger ages with higher rates of failure to thrive, vomiting, and atopic dermatitis. Moreover, an older retrospective study of 208 EoE cases in North Carolina (US) found no differences in clinical, endoscopic, and histologic features by either race or gender⁷. Furthermore, African-Americans were more likely to present earlier with failure to thrive whereas Caucasians were more likely to present with dysphagia and esophageal rings⁷. Taken together, these studies suggest that African-Americans may develop EoE more frequently and more aggressively than previously reported⁵⁻⁷. However, a recent population-based study of an electronic medical database in the US demonstrated that Caucasians are affected at higher rates than African-Americans⁸. Regardless, an important area of future research is to confirm racial EoE phenotypes and determine whether racially and ethnically unique mechanisms underpin disease presentation and treatment response and influence health-care disparities similar to asthma.

EoE has been found to predominantly affect males in numerous studies with estimates as high as 70–85%^{2,9}. A recent meta-analysis found that the risk for EoE was significantly higher for

males³. Moreover, a retrospective, multicenter, cross-sectional analysis of 793 patients with EoE found that the disease predominantly affected male patients (72%)¹⁰. There were no differences in clinical, endoscopic, and histologic characteristics except that a higher percentage of males had strictures¹⁰. Importantly, a smaller recent retrospective case control study found that adult women with EoE were more likely to report chest pain (as well as a non-significant trend toward more heartburn symptoms) but that adult males with EoE were more likely to report dysphagia and food impactions⁹. It is possible that women may be underdiagnosed due to displaying more inflammatory symptoms such as chest pain and heartburn rather than the classic symptoms of dysphagia and food impactions of EoE. Indeed, males may have more fibrostenotic symptoms and thus are more easily diagnosed. Finally, the authors astutely point out that their study is one of the few focused on gender differences in EoE suggesting gender health-care disparities⁹.

Pathophysiology

Elucidating the heritability and pathophysiology of EoE via genetic risk variants and molecular pathways has involved the use of cutting-edge technologies such as genome-wide association studies (GWASs), candidate gene studies, epigenomics, and DNA methylation profiling and chromatin immunoprecipitation sequencing technologies¹¹. The factors that contribute to the risk of EoE are genetics as well as early-life environmental exposures (for example, antibiotic use in infancy, caesarean delivery, preterm birth, season of birth, breastfeeding, and exposures affecting the microbiome)¹²⁻¹⁴. Disease inception involves a complex interplay between epithelial inflammatory pathways, impaired barrier function, and dysregulated transforming growth factor beta (TGF- β) activity/production and activity and induction of allergic TH₂ inflammation mediated by eosinophils and mast cells¹². Although a detailed discussion on disease pathophysiology is beyond the scope of this review, we will highlight two recent discoveries involving thymic stromal lymphopoietin (TSLP) and calpain 14 (CAPN14).

An important reason why our understanding of the EoE disease inception and pathophysiology has progressed rapidly is the use of GWASs. These studies allow disease risk variants to be identified in a more unbiased fashion. A previous GWAS has identified a strong association between EoE and the 5q22 locus, which spans the TSLP domain¹⁵. Importantly, TSLP is a potent mediator of TH₂ allergic responses and basophil activation¹⁶. Furthermore, a separate candidate gene approach found a coding variant in the gene encoding the receptor for TSLP (cytokine receptor-like factor 2, or CRLF2), which is (fascinatingly) associated with EoE risk in men only¹⁷. This represents a possible explanation for why risk of EoE is significantly higher in men (approximately 3:1 male:female)³.

An independently confirmed¹⁸, more recent GWAS was performed on 2.5 million genetic variants of 726 EoE cases and 9,246 controls. In addition to reproducing EoE risk association at the 5q22/TSLP locus, the authors found a strong association at 2p23, which encodes CAPN14¹⁹. CAPN14 is an esophagus-specific intracellular epithelial protease that is induced by interleukin-13 (IL-13) (and genetic variants are risk factors for EoE). Although

its role is still being defined, CAPN14 is thought to directly and indirectly (through desmoglein 1) impair epithelial barrier function in EoE (Figure 1)²⁰. Importantly, *CAPN14* is expressed exclusively in the esophagus when compared with 130 other tissues—a potentially remarkable breakthrough. The *CAPN14* risk variant represents evidence of esophageal-specific pathways in EoE disease pathophysiology and has potential in future, targeted therapeutic applications in EoE. As a result of the above findings, a two-hit mechanism of EoE susceptibility has been proposed¹². The first hit occurs at 5q22, leading to TSLP-induced allergic sensitization. The second hit occurs at 2p23, leading to activation of CAPN14—a potent esophageal-specific protease now shown to regulate epithelial cell barrier function^{12,19,20}.

Diagnosis

The diagnostic criteria for EoE have evolved but mainly required symptoms of esophageal dysfunction with histologic evidence of eosinophil-predominant inflammation consisting of a peak value of at least 15 eosinophils per high-power field. Prior to 2011, the mucosal eosinophilia had to be isolated to the esophagus and persist after 8 weeks of treatment with a proton pump inhibitor (PPI) trial to meet diagnostic criteria^{21,22}. Since 2011, PPI-responsive esophageal eosinophilia (PPI-REE) has been recognized as a specific disease phenotype in which patients respond clinically and histologically to PPI²¹. However, there is strong evidence that PPI-REE and EoE share overlapping genetic, molecular, histologic, and

endoscopic features to suggest that the two are of the same disease spectrum^{2,23}. Therefore, the recently published European guidelines recommended retraction of the term PPI-REE and further recommended that PPI be considered a first-line treatment for EoE rather than a diagnostic criterion².

An important unmet need has been the development of validated scoring systems for additional histologic features besides peak eosinophil counts. Recently, Collins *et al.*²⁴ developed and validated an EoE histologic scoring system (EoEHSS) taking into account eight EoE-associated features, including eosinophil density and basal zone hyperplasia. Interestingly, the EoEHSS better discriminated treated versus untreated patients compared with peak eosinophil counts²⁴. Importantly, this study underscores the observed discordance between histology and symptoms clinically and meets an important need to develop prospectively validated instruments to better characterize whether symptoms associate with histology/disease activity. Indeed, it has been difficult to evaluate symptom-histology associations without a standardized scoring system to compare and confirm findings from separate studies²⁵. Interestingly, a recent multicenter prospective study found only modest predictive capacity of the EoE Activity Index in predicting remission²⁶.

Finally, and perhaps most importantly, there is still a glaring lack of minimally invasive methods to monitor EoE disease activity.

Figure 1. Interleukin-13 (IL-13) induces calpain 14 (CAPN14) effector and regulatory roles in genetically predisposed patients. An important function in eosinophilic esophagitis (EoE) disease pathogenesis involves IL-13 stimulating CAPN14 expression and desmoglein 1 (DSG1) downregulation (a). CAPN14 and IL-13 reduce DSG1 expression, which leads to decreased barrier function and likely increased TH₂ responses seen in EoE (b). Adapted from Davis *et al.*²⁰. EDC, epidermal differentiation complex; TSLP, thymic stromal lymphopoietin.

This unmet need remains of paramount importance as patients continue to be subjected to repeat endoscopies in order to diagnose, monitor, and effectively treat EoE. Moreover, repeat endoscopies significantly impact quality of life and health-care costs. Unfortunately, potential biomarkers of interest have not correlated with endoscopic and histologic findings thus far², although a transcriptome analysis of esophageal biopsies that has shown a sensitivity of 96% and a specificity of 98% for EoE is promising. Furthermore, the transcriptome analysis performs well with only one biopsy but still requires endoscopy to obtain tissue. There is emerging evidence that assessing mucosal integrity using endoscopic impedance measurements correlates with esophageal eosinophilia and may have value in disease management^{27,28}. Finally, the String Test²⁹ and the CytoSponge³⁰, in which surface esophageal samples are retrieved with these semi-invasive devices, have shown promise in smaller studies. Perhaps using either the String Test or the CytoSponge in combination with the transcriptome analysis is the future of EoE disease surveillance and management.

Treatment

The goal of therapy in EoE is to reduce inflammation and halt long-term progression to a fibrostenotic state. The established first-line treatments for EoE are PPIs, topical corticosteroids (CSs), and empiric food elimination diets. PPIs have recently been confirmed to establish histologic remission and symptom improvement in 50% and 60% of patients with EoE, respectively^{2,31}. However, this meta-analysis should be interpreted with caution, as many of the studies were of poor quality². Nonetheless, the new, recently published European guidelines developed

a treatment algorithm for EoE with PPIs considered as a first-line treatment for EoE; an evaluation and treatment algorithm is summarized in Figure 2². Interestingly, PPIs block signal transducer and activator of transcription 6 (STAT6) activation and IL-13-induced signaling, which reduces eotaxin-3 production and consequently may have an anti-eosinophil effect³². This mechanism, along with improvement of barrier function, may be the reason why PPIs are an effective form of treatment for some EoE cases^{12,32}. A new and potent potassium channel acid blocker called vonoprazan has also shown evidence of clinical and histologic responsiveness in EoE³³.

The numbers of individual patients in each EoE trial are quite low; however, the cumulative numbers across all studies are now more substantial and have allowed insightful systematic reviews and meta-analyses to be performed². Multiple systematic reviews and meta-analyses have confirmed the efficacy of topical CSs for histologic remission EoE^{34–37}; however, two of these reviews of the literature^{36,37} did not find evidence of symptom improvement with topical CSs. This disconnect may be due in large part to the use of different, non-validated, symptom-scoring tools for each study as well as overall small sample sizes². A recent prospective study in children found that 63% of children had sustained remission at 2 years of treatment with swallowed CSs, suggesting along with other studies that topical CSs can achieve long-term remission^{2,38}. Finally, although swallowed budesonide and fluticasone are almost always prescribed, no formulation of topical CSs has been approved for EoE. Two different budesonide formulations—an effervescent tablet for orodispersible use (BET) and a viscous suspension (BVS)—are

Figure 2. Treatment algorithm for patients with confirmed eosinophilic esophagitis (EoE). Adapted from European consensus guidelines². *Proton pump inhibitor (PPI) is first-line but consider swallowed corticosteroids (SCS) if there are severe symptoms, failure to thrive, or provider-patient preferences. *Can be done in addition to PPI or other drug or diet therapies. FED, food elimination diet.

currently in clinical trials (<https://clinicaltrials.gov/ct2/show/NCT02493335?recrs=a&cond=Eosinophilic+Esophagitis&draw=1&rank=9> and <https://clinicaltrials.gov/ct2/show/NCT02736409?recrs=a&cond=Eosinophilic+Esophagitis&draw=2&rank=11>). Both BET and BVS have been found to be highly safe and effective thus far³⁹.

In terms of safety, a recent meta-analysis observed that topical CSs were not associated with significant adverse events other than a risk for developing asymptomatic esophageal candidiasis^{2,37}. However, a study out of Cincinnati found that 10% of children whose EoE was treated with fluticasone (>440 µg daily) developed evidence of adrenal suppression⁴⁰. Moreover, another recent study found decreased cortisol stimulation with no evidence of adrenal insufficiency or growth delay in children on oral viscous budesonide⁴¹. Many patients with EoE are on other forms of steroids for comorbid allergic conditions; therefore, more studies are needed to characterize the risk of adrenal suppression with topical CSs.

Recently, a meta-analysis found a 72% histologic remission rate using an empiric six-food elimination diet (6FED)³. Importantly, there are trials ongoing to understand whether less restrictive elimination diets can be used. One-food (1FED; milk), two-food (2FED; milk and wheat), and four-food (4FED; milk, wheat, soy, and eggs) elimination diets are currently being investigated (<https://clinicaltrials.gov/ct2/show/NCT02610816>)². Specifically, a multicenter trial of a 4FED (milk, wheat, egg, and soy) was found to induce remission in 60% of children with EoE⁴². Furthermore, a recent study found that 61% of children had histologic remission with elimination of milk only⁴³. Initial empiric elimination diets that are less restrictive have the potential advantages of reducing the number of endoscopies while improving nutrition and quality of life.

Finally, the cumulative evidence is leading many to conclude that food allergy testing-directed elimination diets have little role in the management of EoE^{2,44}. Indeed, at Cincinnati Children's Hospital Medical Center, clinicians who routinely care for patients with EoE do not test for causative food triggers by using skin tests, *in vitro* IgE measures, or patch testing unless there is clear evidence of an IgE-mediated reaction (for example, hives and anaphylaxis). It is the authors' opinion that the totality of evidence is fairly conclusive that test-directed elimination approaches are inadequate to identify food triggers in EoE, lead to confusion, and consequently delay attaining histologic and clinical control and remission.

Conclusions

Our understanding of EoE is exponentially increasing. This is primarily due to cutting-edge technologies that have allowed researchers to discover disease mechanisms at an astonishing rate. A multi-hit mechanism of EoE pathogenesis involving a complex interplay between genetic and early-life environmental risk factors, allergic sensitization, and esophageal-specific pathways has been proposed¹². These pathways will offer new insights into mechanisms and potentially new therapeutic applications. The use of transcriptome analyses may further allow clinicians to "personalize" treatment depending on endotype¹². Unmet needs include developing and following high-risk birth or early-life cohorts prospectively over decades to understand disease inception, natural history, and sequelae. Finally, comprehensive standardized scoring systems for symptoms, quality of life, endoscopy, and histology need to be prospectively and universally validated. The recent formation of the Consortium of Eosinophilic Gastrointestinal Disease Researchers, which is part of the Rare Disease Clinical Research Network of the National Institutes of Health, provides an opportunity to unite clinicians, scientists, and a full spectrum of key stakeholders, including patients, to better understand and treat EoE (<https://www.rarediseasesnetwork.org/cms/cegir/>).

Competing interests

MER is a consultant for NKT Therapeutics, Pulm One, Spoon Guru, Celgene, Shire, Astra Zeneca, GlaxoSmithKline, and Novartis and has an equity interest in the first three listed and Immune Pharmaceuticals and royalties from reslizumab (Teva Pharmaceuticals). He is an inventor of patents, owned by Cincinnati Children's Hospital Medical Center. SRD declares that he has no competing interests.

Grant information

This work was supported in part by National Institutes of Health U19 AI070235, National Institutes of Health R01 AI124355, and R37 R37 A1045898; the Campaign Urging Research for Eosinophilic Disease Foundation; the Buckeye Foundation; the American Partnership for Eosinophilic Disorders; and the Sunshine Charitable Foundation and its supporters, Denise A. Bunning and David G. Bunning.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- Lucendo AJ, Arias A: **Eosinophilic gastroenteritis: an update.** *Expert Rev Gastroenterol Hepatol.* 2012; 6(5): 591–601.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lucendo AJ, Molina-Infante J, Arias Á, *et al.*: **Guidelines on eosinophilic esophagitis: evidence-based statements and recommendations for diagnosis and management in children and adults.** *United European Gastroenterol J.* 2017; 5(3): 335–58.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
- Arias Á, Pérez-Martínez I, Tenías JM, *et al.*: **Systematic review with meta-analysis: the incidence and prevalence of eosinophilic oesophagitis in children and adults in population-based studies.** *Aliment Pharmacol Ther.* 2016; 43(1): 3–15.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
- Hunter SS, Helmy DO, Zayed NA, *et al.*: **Eosinophilic esophagitis in Egyptian adult patients presenting with upper gastrointestinal symptoms.** *OJGAS.* 2014; 4(2): 88–95, 43157.
[Publisher Full Text](#)
- Gill RK, Al-Subu A, Elitsur Y, *et al.*: **Prevalence and characteristics of**

- eosinophilic esophagitis in 2 ethnically distinct pediatric populations. *J Allergy Clin Immunol*. 2014; **133**(2): 576–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
6. Weiler T, Mikhail I, Singal A, *et al.*: Racial differences in the clinical presentation of pediatric eosinophilic esophagitis. *J Allergy Clin Immunol Pract*. 2014; **2**(3): 320–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
 7. Sperry SL, Woosley JT, Shaheen NJ, *et al.*: Influence of race and gender on the presentation of eosinophilic esophagitis. *Am J Gastroenterol*. 2012; **107**(2): 215–21.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 8. **F** Mansoor E, Cooper GS: The 2010–2015 Prevalence of Eosinophilic Esophagitis in the USA: A Population-Based Study. *Dig Dis Sci*. 2016; **61**(10): 2928–34.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 9. **F** Lynch KL, Dhalla S, Chedid V, *et al.*: Gender is a determinative factor in the initial clinical presentation of eosinophilic esophagitis. *Dis Esophagus*. 2016; **29**(2): 174–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 10. **F** Moawad FJ, Dellon ES, Achem SR, *et al.*: Effects of Race and Sex on Features of Eosinophilic Esophagitis. *Clin Gastroenterol Hepatol*. 2016; **14**(1): 23–30.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 11. Davis BP, Rothenberg ME: Mechanisms of Disease of Eosinophilic Esophagitis. *Annu Rev Pathol*. 2016; **11**: 365–93.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 12. Rothenberg ME: Molecular, genetic, and cellular bases for treating eosinophilic esophagitis. *Gastroenterology*. 2015; **148**(6): 1143–57.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 13. Alexander ES, Martin LJ, Collins MH, *et al.*: Twin and family studies reveal strong environmental and weaker genetic cues explaining heritability of eosinophilic esophagitis. *J Allergy Clin Immunol*. 2014; **134**(5): 1084–1092.e1.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 14. Jensen ET, Kappelman MD, Kim HP, *et al.*: Early life exposures as risk factors for pediatric eosinophilic esophagitis. *J Pediatr Gastroenterol Nutr*. 2013; **57**(1): 67–71.
[PubMed Abstract](#) | [Publisher Full Text](#)
 15. Rothenberg ME, Spergel JM, Sherrill JD, *et al.*: Common variants at 5q22 associate with pediatric eosinophilic esophagitis. *Nat Genet*. 2010; **42**(4): 289–91.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 16. **F** Noti M, Wojno ED, Kim BS, *et al.*: Thymic stromal lymphopoietin-elicited basophil responses promote eosinophilic esophagitis. *Nat Med*. 2013; **19**(8): 1005–13.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 17. **F** Sherrill JD, Gao PS, Stucke EM, *et al.*: Variants of thymic stromal lymphopoietin and its receptor associate with eosinophilic esophagitis. *J Allergy Clin Immunol*. 2010; **126**(1): 160–5.e3.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 18. Sleiman PM, Wang ML, Cianferoni A, *et al.*: GWAS identifies four novel eosinophilic esophagitis loci. *Nat Commun*. 2014; **5**: 5593.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 19. **F** Kottyan LC, Davis BP, Sherrill JD, *et al.*: Genome-wide association analysis of eosinophilic esophagitis provides insight into the tissue specificity of this allergic disease. *Nat Genet*. 2014; **46**(8): 895–900.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 20. Davis BP, Stucke EM, Khorki ME, *et al.*: Eosinophilic esophagitis-linked calpain 14 is an IL-13-induced protease that mediates esophageal epithelial barrier impairment. *JCI Insight*. 2016; **1**(4): e86355.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 21. **F** Liacouras CA, Furuta GT, Hirano I, *et al.*: Eosinophilic esophagitis: updated consensus recommendations for children and adults. *J Allergy Clin Immunol*. 2011; **128**(1): 3–20.e6; quiz 21–2.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 22. **F** Dellon ES, Gonsalves N, Hirano I, *et al.*: ACG clinical guideline: Evidence-based approach to the diagnosis and management of esophageal eosinophilia and eosinophilic esophagitis (EoE). *Am J Gastroenterol*. 2013; **108**(5): 679–92; quiz 693.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 23. Molina-Infante J, Bredenoord AJ, Cheng E, *et al.*: Proton pump inhibitor-responsive oesophageal eosinophilia: an entity challenging current diagnostic criteria for eosinophilic oesophagitis. *Gut*. 2016; **65**(3): 524–31.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 24. Collins MH, Martin LJ, Alexander ES, *et al.*: Newly developed and validated eosinophilic esophagitis histology scoring system and evidence that it outperforms peak eosinophil count for disease diagnosis and monitoring. *Dis Esophagus*. 2017; **30**(3): 1–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 25. Lucendo AJ, Arias Á, Tenias JM: Systematic review: the association between eosinophilic oesophagitis and coeliac disease. *Aliment Pharmacol Ther*. 2014; **40**(5): 422–34.
[PubMed Abstract](#) | [Publisher Full Text](#)
 26. **F** Safroneeva E, Straumann A, Coslovsky M, *et al.*: Symptoms Have Modest Accuracy in Detecting Endoscopic and Histologic Remission in Adults With Eosinophilic Esophagitis. *Gastroenterology*. 2016; **150**(3): 581–590.e4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 27. **F** Warners MJ, van Rhijn BD, Verheij J, *et al.*: Disease activity in eosinophilic esophagitis is associated with impaired esophageal barrier integrity. *Am J Physiol Gastrointest Liver Physiol*. 2017; **313**(3): G230–G238.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 28. **F** Katzka DA, Ravi K, Geno DM, *et al.*: Endoscopic Mucosal Impedance Measurements Correlate With Eosinophilia and Dilatation of Intercellular Spaces in Patients With Eosinophilic Esophagitis. *Clin Gastroenterol Hepatol*. 2015; **13**(7): 1242–1248.e1.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 29. **F** Furuta GT, Kagalwalla AF, Lee JJ, *et al.*: The oesophageal string test: a novel, minimally invasive method measures mucosal inflammation in eosinophilic oesophagitis. *Gut*. 2013; **62**(10): 1395–405.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 30. **F** Katzka DA, Geno DM, Ravi A, *et al.*: Accuracy, safety, and tolerability of tissue collection by Cytosponge vs endoscopy for evaluation of eosinophilic esophagitis. *Clin Gastroenterol Hepatol*. 2015; **13**(1): 77–83.e2.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 31. **F** Lucendo AJ, Arias Á, Molina-Infante J: Efficacy of Proton Pump Inhibitor Drugs for Inducing Clinical and Histologic Remission in Patients With Symptomatic Esophageal Eosinophilia: A Systematic Review and Meta-Analysis. *Clin Gastroenterol Hepatol*. 2016; **14**(1): 13–22.e1.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 32. Zhang X, Cheng E, Huo X, *et al.*: Omeprazole blocks STAT6 binding to the eotaxin-3 promoter in eosinophilic esophagitis cells. *PLoS One*. 2012; **7**(11): e50037.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 33. **F** Ishimura N, Ishihara S, Kinoshita Y: Sustained Acid Suppression by Potassium-Competitive Acid Blocker (P-CAB) May Be An Attractive Treatment Candidate for Patients with Eosinophilic Esophagitis. *Am J Gastroenterol*. 2016; **111**(8): 1203–4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 34. **F** Tan ND, Xiao YL, Chen MH: Steroids therapy for eosinophilic esophagitis: Systematic review and meta-analysis. *J Dig Dis*. 2015; **16**(8): 431–42.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 35. **F** Sawas T, Dhalla S, Sayyar M, *et al.*: Systematic review with meta-analysis: pharmacological interventions for eosinophilic oesophagitis. *Aliment Pharmacol Ther*. 2015; **41**(9): 797–806.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 36. **F** Murali AR, Gupta A, Attar BM, *et al.*: Topical steroids in eosinophilic esophagitis: Systematic review and meta-analysis of placebo-controlled randomized clinical trials. *J Gastroenterol Hepatol*. 2016; **31**(6): 1111–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 37. **F** Chuang MY, Chinnarath MA, Hancock DG, *et al.*: Topical Steroid Therapy for the Treatment of Eosinophilic Esophagitis (EoE): A Systematic Review and Meta-Analysis. *Clin Transl Gastroenterol*. 2015; **6**(3): e82.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 38. **F** Gutiérrez-Junquera C, Fernández-Fernández S, Cilleruelo ML, *et al.*: High Prevalence of Response to Proton-pump Inhibitor Treatment in Children With Esophageal Eosinophilia. *J Pediatr Gastroenterol Nutr*. 2016; **62**(5): 704–10.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 39. **F** Miehlke S, Hruz P, Vieth M, *et al.*: A randomised, double-blind trial comparing budesonide formulations and dosages for short-term treatment of eosinophilic oesophagitis. *Gut*. 2016; **65**(3): 390–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 40. **F** Golekoh MC, Hornung LN, Mikkada VA, *et al.*: Adrenal Insufficiency after Chronic Swallowed Glucocorticoid Therapy for Eosinophilic Esophagitis. *J Pediatr*. 2016; **170**: 240–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 41. **F** Harel S, Hursh BE, Chan ES, *et al.*: Adrenal Suppression in Children Treated With Oral Viscous Budesonide for Eosinophilic Esophagitis. *J Pediatr Gastroenterol Nutr*. 2015; **61**(2): 190–3.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 42. **F** Kagalwalla AF, Wechsler JB, Amsden K, *et al.*: Efficacy of a 4-Food Elimination Diet for Children With Eosinophilic Esophagitis. *Clin Gastroenterol Hepatol*. 2017; pii: S1542-3565(17)30689-4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 43. **F** Kruszewski PG, Russo JM, Franciosi JP, *et al.*: Prospective, comparative effectiveness trial of cow's milk elimination and swallowed fluticasone for pediatric eosinophilic esophagitis. *Dis Esophagus*. 2016; **29**(4): 377–84.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 44. Simon D, Cianferoni A, Spergel JM, *et al.*: Eosinophilic esophagitis is characterized by a non-IgE-mediated food hypersensitivity. *Allergy*. 2016; **71**(5): 611–20.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. **Yoshikazu Kinoshita**

Department of Gastroenterology and Hepatology, Shimane University School of Medicine, Shimane, Japan

Competing Interests: No competing interests were disclosed.

2. **Edaire Cheng**

Pediatric Gastroenterology, Hepatology & Nutrition Division, University of Texas Southwestern Medical Center, Dallas, Texas, USA

Diana Montoya Melo

Pediatric Gastroenterology, Hepatology & Nutrition Division, University of Texas Southwestern Medical Center, Dallas, Texas, USA

Competing Interests: No competing interests were disclosed.

3. **Michael F Vaezi**

Division of Gastroenterology, Hepatology and Nutrition, Vanderbilt University Medical Center, Nashville, USA

Competing Interests: No competing interests were disclosed.

Comments on this article

Version 1

Author Response (F1000Research Advisory Board Member) 01 Oct 2017

Marc Rothenberg, Cincinnati Children's Hospital Medical Center, University of Cincinnati, Cincinnati OH, 45529, USA

This paper was published under Open Peer Review, which is a new format for publishing, where the paper is subjected to non-confidential review. Comments from the original reviewers and subsequent readers are open for dialog/comment. I am pleased to spread information about eosinophilic disorders through this new format. I am also pleased to have partnered with a

talented faculty member in my Division, Dr. Sandy Durrani, and welcome him to get more involved in the EoE and F1000 community.”

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research