Check for updates

REVIEW

Acute Coronary Syndromes in the Elderly [version 1; peer review: 3 approved]

Niels Engberding ¹, Nanette K. Wenger ^{2,3}¹Department of Medicine, Division of Cardiology, National Jewish Health, Denver, Colorado, USA²Emory Heart and Vascular Center, Atlanta, Georgia, USA³Department of Medicine, Division of Cardiology, Emory University School of Medicine, Atlanta, Georgia, USA**V1** First published: 02 Oct 2017, 6(F1000 Faculty Rev):1791
<https://doi.org/10.12688/f1000research.11064.1>Latest published: 02 Oct 2017, 6(F1000 Faculty Rev):1791
<https://doi.org/10.12688/f1000research.11064.1>

Abstract

The clinical evidence for treatment of acute coronary syndrome (ACS) in the elderly is less robust than in patients younger than 75 years. The elderly have the highest incidence of cardiovascular disease and frequently present with ACS. This number can be expected to increase over time because society is aging. Older adults often sustain unfavorable outcomes from ACS because of atypical presentation and delay in recognition. In addition, elderly patients commonly do not receive optimal guideline-directed ACS treatment. Owing to their high baseline risk of ischemic complications, the elderly also fare worse even with optimal ACS treatment as they frequently have more complex coronary disease, more comorbidities, less cardiovascular reserve, and a higher risk of treatment complications. They are also subjected to a broader range of pharmacologic treatment. Treatment complications can be mitigated to some extent by meticulous dose adjustment of antithrombotic and adjunctive therapies. While careful transitions of care and appropriate utilization of post-discharge secondary preventive measures are important in ACS patients of all ages, the elderly are more vulnerable to system errors and thus deserve special attention from the clinician.

Keywords

acute coronary, elderly, cardiovascular disease

Open Peer Review

Approval Status

	1	2	3
version 1 02 Oct 2017			

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Anna Sonia Petronio**, University of Pisa, Pisa, Italy
2. **Stefano Savonitto**, Manzoni Hospital, Lecco, Italy
3. **Sanjay Patel**, University of Sydney, Sydney, Australia

Any comments on the article can be found at the end of the article.

Corresponding author: Nanette K. Wenger (nwenger@emory.edu)

Competing interests: The authors declare that they have no competing interests.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2017 Engberding N and Wenger NK. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Engberding N and Wenger NK. **Acute Coronary Syndromes in the Elderly [version 1; peer review: 3 approved]** F1000Research 2017, 6(F1000 Faculty Rev):1791 <https://doi.org/10.12688/f1000research.11064.1>

First published: 02 Oct 2017, 6(F1000 Faculty Rev):1791 <https://doi.org/10.12688/f1000research.11064.1>

Epidemiologic data

Elderly patients (>75 years of age)¹ constitute a large proportion of those patients presenting with acute coronary syndrome (ACS), and temporal trends in the incidence of myocardial infarction document a shift toward older adults². The average ages at first ACS presentation in the US are 65 years for men and 72 years for women. About two thirds of myocardial infarctions occur in patients older than 65 years of age, and one third in patients older than 75 years of age. Randomized clinical trials, on the other hand, have included substantially fewer elderly patients than clinicians encounter in real life³. Thus, the basis of evidence forming the foundation of ACS treatment may not apply to a large number of patients, and clinicians need to extrapolate evidence to match their older patients' needs and preferences. Sixty percent of ACS hospitalizations occur in patients older than 65 years, and 85% of ACS mortality occurs in the Medicare population. Most deaths related to myocardial infarction occur in patients older than 65 years of age⁴.

Age is not only a powerful risk factor for cardiovascular disease; it is also an independent risk factor for adverse outcomes after cardiovascular events, for complications of cardiovascular procedures and interventions, and for side effects of pharmacotherapy, particularly from antithrombotic therapies. The mortality rate after a first non-ST segment elevation myocardial infarction (non-STEMI) in very elderly patients is very high: with respect to 1-year outcomes, among patients who were 65–79, 80–84, 85–89, and at least 90 years old, mortality increased progressively from 13.3% to 23.6%, 33.6%, and 45.5%, respectively⁵.

In addition, older patients generally have more complex cardiovascular disease, more comorbidities, and generally a more atypical clinical presentation. There is a greater prevalence of hypertension, congestive heart failure (CHF), atrial fibrillation, cerebrovascular disease, anemia, and renal insufficiency in older patients with ACS. Age also has important implications on pharmacokinetics and pharmacodynamics⁶. Challenges in taking care of elderly patients with ACS include timely recognition, not withholding lifesaving therapies on the basis of age alone, and respecting the patients' preferences and goals of care.

Atypical symptoms

There may be several explanations for why the elderly have worse outcomes with ACS. While chest pain remains the most common presentation for ACS, elderly patients frequently present with atypical symptoms (meaning, without chest pain)⁷. In patients who present without chest pain, the diagnosis of ACS is often missed or delayed, leading to worse outcomes. Notably, chest pain as a presenting symptom occurs in only 40% of patients older than 85 years but is present in nearly 80% of patients under 65 years. Common symptoms in the elderly presenting with ACS include dyspnea, diaphoresis, nausea and vomiting, and syncope. In patients at least 85 years old, an atypical presentation of myocardial infarction appears to be the standard and the clinician must be prepared to diagnose ACS in many acutely ill patients of this age⁸. Acute pulmonary edema is more commonly a presentation of the elderly patient with ACS. Increased arterial stiffness as manifested with increased arterial pulse pressure as

well as increased prevalence of multivessel coronary artery disease (CAD) may explain why older patients with ACS are more likely to present with signs and symptoms of CHF⁹.

Aside from atypical symptoms, the 12-lead electrocardiogram (ECG), a standard investigation in patients with suspected ACS, may be non-diagnostic and therefore serial ECGs are recommended to diagnose high-risk findings such as ST segment elevation. The diagnosis of a STEMI is more challenging in patients presenting with left bundle branch block (LBBB). Therefore, the higher prevalence of LBBB in the elderly may contribute to diagnostic uncertainty in the early phase of presentation, when rapid risk stratification and triage are most important.

Prehospital delays also contribute to prevent prompt treatment. Despite having more severe coronary disease than younger patients at coronary angiography, elderly patients are more likely to be treated medically and experience more adverse outcomes¹⁰. Additionally, the hemodynamic impact of a given infarct size may be more pronounced in the elderly because of reduced cardiac reserve. The age-related decline in cardiac reserve may be related to reduced beta-adrenergic responsiveness. There is also a higher likelihood of comorbid illnesses with advancing age. Not only do these comorbidities obscure the presentation of ACS, they also contribute to worse outcomes. Type 2 myocardial infarctions, which result from increased myocardial oxygen demand in the setting of severe fixed obstructive CAD, are commonly caused by comorbidities such as tachycardia, pneumonia with hypoxemia, chronic pulmonary disease, and bleeding episodes. These comorbidities frequently complicate hospital admissions of elderly patients and need to be recognized since they require treatment strategies different than those of type 1 myocardial infarctions. In general, elderly patients are more likely to experience complications of ACS, such as CHF, heart block, ventricular rupture, and atrial fibrillation. Furthermore, frailty and disability can complicate acute hospitalization as well as convalescence and rehabilitation. On the other hand, frailty is not considered in clinically accepted risk scores.

General considerations in management

The treatment of ACS has evolved significantly over the past 40 years. Classically, ACS is caused by thrombotic obstruction of an epicardial coronary artery. Thus, treatment focuses primarily on early coronary revascularization supported by the use of antithrombotic pharmacotherapy¹¹. In general, more aggressive use of invasive coronary procedures and antithrombotic medications is associated with lower risk of ischemic complications but higher risk of a bleeding complication. Outcomes research has revealed that the elderly have been treated less effectively¹². Presumably, practitioners consider the risk-benefit ratio of cardiac procedures to be less favorable in the elderly. While elderly patients did not share in the improved survival rates observed in younger patients in the early days of coronary interventions (1979–1994)², more recent data showed that mortality after hospital admission of elderly patients with acute myocardial infarction has substantially decreased over the past 15 years¹³. This improvement is likely mediated by increasing use of recommended management strategies; thus, the application of guidelines derived from trials

mostly including younger patients may benefit the elderly populations as well. However, adverse outcomes increase with age across the whole spectrum of ACS^{10,14–16}. Observational data suggest that outcomes of elderly patients improve when they receive guideline-directed cardiac procedures¹⁷. Registry data from ACS events in England and Wales between 2003 and 2010 show substantial reduction in in-hospital mortality in all age groups, including the old and very old, males and females, and STEMI and non-STEMI. Within the study time frame, the use of percutaneous coronary intervention (PCI) and evidence-based pharmacologic therapies increased significantly for all age groups. Although elderly patients still received fewer coronary revascularizations than younger patients, it is noteworthy that almost half of all STEMI patients older than 85 years received a PCI, and the adherence to evidence-based pharmacotherapy at discharge was about 90% in this group¹⁸. The Italian Elderly ACS study suggests that elderly patients with troponin elevation benefit from an early invasive approach¹⁹. Data from the same trial, when pooled with registry data, suggested that coronary revascularization in elderly women was associated with lower 1-year mortality when compared with an early conservative approach, without an increase in severe bleeding²⁰. The results from the After Eighty study support the use of an invasive strategy in patients at least 80 years old²¹. Even in nonagenarians and centenarians with ACS, there is evidence that adherence to guideline-recommended therapies is associated with decreased mortality²². Despite an increased risk for major bleeding in patients older than 75 years of age, a routine early invasive strategy significantly improved outcomes in elderly patients with ACS²³. Registry data suggest that, over the last 15 years, the progressive switch from a conservative to a more invasive approach in elderly patients may have contributed to mortality reduction across the ACS spectrum, irrespective of age and gender^{24,25}. Accordingly, the absolute benefit of early invasive therapies in the elderly appears to be greater than in younger patients because of their high mortality at baseline.

Pharmacologic considerations

It has been reported that the risk of adverse drug reaction increases with the number of medications taken concurrently. With two concurrent medications, there is a 13% risk of an adverse drug interaction, and the risk increases to 38% for four medications and 82% for seven or more medications prescribed simultaneously²⁶. In light of these data, it is crucial to balance the risks of polypharmacy with the benefit of not withholding guideline-directed medications, proven to be of benefit in the elderly.

Age-related decline in organ function, muscle mass, and volume of distribution and changes in pharmacokinetics require the treating physician to meticulously adjust medication dosing (Table 1). As bleeding risk increases with age, dose adjustments are particularly important when it comes to anticoagulant therapies. The creatinine clearance is preferably calculated by the Cockcroft-Gault equation and should form the basis for renally dosed medications rather than the serum creatinine level. Observational data revealed that patients with ACS often receive antithrombotic therapies at excess doses. Factors associated with excess dosing included older age as well as female sex, renal insufficiency,

Table 1. Acute coronary syndrome medications.

Antiplatelet
Aspirin
<i>P₂Y₁₂ antagonists</i>
Clopidogrel ^a
Prasugrel ^{a,b} (relatively contraindicated in patients who are at least 75 years old or whose body weight is less than 60 kg)
Ticagrelor
Cangrelor ^b
<i>Glycoprotein IIb/IIIa inhibitors</i>
Tirofiban ^{b,c}
Eptifibatide ^{b,c}
Abciximab ^b
Anticoagulant
Heparin ^b
Enoxaparin (low-molecular-weight heparin) ^{a,b,c}
Dalteparin (low-molecular-weight heparin) ^b
Fondaparinux ^c (in the US, use for acute coronary syndrome is off-label)
Argatroban ^b
Bivalirudin ^{b,c} (vascular access site bleeding more often in patients older than 65 years)
Anti-ischemic
Nitroglycerin
Beta blocker ^c (in the elderly, consider lower initial doses and titrate to response)
Adjunct
Statins ^c
Angiotensin-converting enzyme inhibitor ^c
Angiotensin receptor blocker ^c

^aNeed for age-based dose adjustment. ^bNeed for weight-based dose adjustment. ^cNeed for renal-based dose adjustment.

low body weight, diabetes mellitus, and CHF²⁷. Among patients who recently had an ACS, a high-dose statin regimen is known to provide greater protection against death or major cardiovascular events than a low- or moderate-dose statin regimen²⁸. A subgroup analysis from the PROVE-IT-TIMI 22 (Pravastatin or Atorvastatin Evaluation and Infection Therapy–Thrombolysis in Myocardial Infarction 22) trial including 634 elderly patients suggested that a high-dose statin regimen achieved a greater reduction in adverse events in the elderly than in the younger study subjects. Not only was the incidence of major statin side effects among the elderly similar to that in younger patients, it also did not differ with the intensity of the statin regimen²⁹. In similar fashion to the general ACS population, optimal medical therapy should be initiated before hospital discharge, as this strategy has been shown to improve long-term outpatient adherence. While randomized data are sparse in patients older than 75 years, an observational

study has shown that early beta-blocker therapy was not used for 51% of patients at least 65 years old who were hospitalized with an acute myocardial infarction, although they did not have a clear contraindication to this therapy. Yet the same study showed that patients who received beta blockers had a lower in-hospital mortality rate than patients who did not receive beta blockers³⁰. Results from the CRUSADE (Can Rapid Risk Stratification of Unstable Angina Patients Suppress Adverse Outcomes With Early Implementation of the American College of Cardiology/American Heart Association Guidelines) National Quality Improvement Initiative have shown that early in-hospital use of aspirin and beta blockers was less likely in patients past 65 years of age and that heparin was significantly less used past 85 years of age. The acute use of clopidogrel and platelet glycoprotein IIb/IIIa inhibitors was most affected by age. Only 30% of patients older than 85 years received clopidogrel, and only 12.8% received platelet glycoprotein IIb/IIIa inhibitors. For patients surviving the index hospitalization, use of many discharge medications was similar in young and old patients except that clopidogrel and lipid-lowering therapy remained less commonly prescribed in elderly patients. While in-hospital mortality and complication rates increased with advancing age, those receiving more recommended therapies had lower mortality than those who did not³¹.

Revascularization

Most myocardial infarctions in older adults present without ST elevations on the ECG and this has been attributed to the presence of more severe multivessel CAD that may have led to ischemic preconditioning or significant collateral growth³². Less than 30% of ACS presentations in patients older than 75 years are caused by STEMI⁹. While it is well established that coronary reperfusion is crucial within 12 hours of symptom onset, in the Global Registry of Acute Coronary Events (GRACE) registry, 30% of STEMI patients presenting within 12 hours of symptoms did not receive reperfusion therapy. Being 75 years of age or older was among the characteristics of those less likely to receive revascularization³³.

Current data support the use of reperfusion therapies, including fibrinolysis, up to the age of 85 years¹⁴. The selection between fibrinolytics or PCI is determined largely by factors other than age, such as time from presentation, travel time to cardiac catheterization laboratory, comorbidity, and signs of cardiogenic shock³⁴. The safety and efficacy of reperfusion, specifically fibrinolytic therapy, in the very elderly (≥ 85 years of age) have yet to be determined as the risks of intracranial bleeding and cardiac rupture increase with age.

When revascularization options for the elderly are being considered, it is reasonable to choose coronary artery bypass graft (CABG) surgery over PCI in older patients with non-ST segment elevation ACS who are appropriate candidates, particularly those with diabetes mellitus or complex multivessel CAD, to improve outcomes¹¹. CABG may become the only option in certain clinical settings, such as a coronary anatomy not amenable to PCI, unsuccessful PCI, or the need for surgical repair of mechanical complications or concomitant valve disease.

When the decision is made for an invasive approach, the transradial access may be particularly appealing in the elderly as it reduces the risk of access site bleeding complications. In a subgroup analysis of the RIVAL (Radial Versus Femoral Access for Coronary Intervention) trial, elderly patients undergoing cardiac catheterization had lower rates of major bleeding or access site complications, yet the elderly also had higher rates of access site crossover with radial access compared with femoral access³⁵. Operative mortality rates and risk of complications for CABG increase with age. In addition, duration of hospitalization and post-surgery convalescence may be significantly prolonged in older patients after CABG and therefore should be considered in counselling the patient. It is up to the clinician to individualize the patient assessment by acknowledging the wide heterogeneity that exists between chronological and biological age and determining the patient's preferences and goals for life. Another underutilized measure in the care of elderly patients with ACS is referral to cardiac rehabilitation. Given the overall reduced functional status in the older adult, the elderly are at an elevated risk of disability following ACS. There is evidence that the benefit of cardiac rehabilitation applies to both the elderly and younger coronary patients³⁶. Older adults may even benefit more, as cardiac rehabilitation in the elderly may extend beyond the cardiovascular system and include an improvement in physical fitness as well as enhancement in balance, stability, muscle strength, and tone.

Conclusions

As the population continues to age, physicians will be confronted with an increasing number of elderly and very elderly patients presenting with ACS. While care needs to be individualized, age alone should never be the reason to withhold potentially lifesaving procedures and interventions. Elderly patients are at high risk for bleeding complications, but they are also at the highest risk for ischemic complications if less aggressive treatment strategies are pursued. So clinicians are tasked with meticulous risk stratification for ischemic risk and bleeding risk while taking into account assessment of frailty, quality of life, goals of care, and individual preferences. Early invasive protocols seem to be just as feasible in the elderly as in the general population. In order to mitigate bleeding risk and adverse medication side effects, it is imperative to correct dosages of pharmacotherapy to age- and gender-adjusted renal function and volume of distribution.

Abbreviations

ACS, acute coronary syndrome; CABG, coronary artery bypass graft; CAD, coronary artery disease; CHF, congestive heart failure; ECG, electrocardiogram; LBBB, left bundle branch block; PCI, percutaneous coronary intervention; STEMI, ST segment elevation myocardial infarction.

Competing interests

The authors declare that they have no competing interests.

Grant information

The author(s) declared that no grants were involved in supporting this work.

References

1. Alexander KP, Newby LK, Cannon CP, *et al.*: **Acute coronary care in the elderly, part I: Non-ST-segment-elevation acute coronary syndromes: a scientific statement for healthcare professionals from the American Heart Association Council on Clinical Cardiology: in collaboration with the Society of Geriatric Cardiology.** *Circulation.* 2007; 115(19): 2549–69.
[PubMed Abstract](#) | [Publisher Full Text](#)
2. Roger VL, Jacobsen SJ, Weston SA, *et al.*: **Trends in the incidence and survival of patients with hospitalized myocardial infarction, Olmsted County, Minnesota, 1979 to 1994.** *Ann Intern Med.* 2002; 136(5): 341–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
3. Lee PY, Alexander KP, Hammill BG, *et al.*: **Representation of elderly persons and women in published randomized trials of acute coronary syndromes.** *JAMA.* 2001; 286(6): 708–13.
[PubMed Abstract](#) | [Publisher Full Text](#)
4. Goldberg RJ, McCormick D, Gurwitz JH, *et al.*: **Age-related trends in short- and long-term survival after acute myocardial infarction: a 20-year population-based perspective (1975–1995).** *Am J Cardiol.* 1998; 82(11): 1311–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
5. Lopes RD, Gharacholou SM, Holmes DN, *et al.*: **Cumulative incidence of death and rehospitalization among the elderly in the first year after NSTEMI.** *Am J Med.* 2015; 128(6): 582–90.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
6. Mangoni AA, Jackson SH: **Age-related changes in pharmacokinetics and pharmacodynamics: basic principles and practical applications.** *Br J Clin Pharmacol.* 2004; 57(1): 6–14.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
7. Brieger D, Eagle KA, Goodman SG, *et al.*: **Acute coronary syndromes without chest pain, an underdiagnosed and undertreated high-risk group: insights from the Global Registry of Acute Coronary Events.** *Chest.* 2004; 126(2): 461–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
8. Bayer AJ, Chadha JS, Farag RR, *et al.*: **Changing presentation of myocardial infarction with increasing old age.** *J Am Geriatr Soc.* 1986; 34(4): 263–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
9. Mehta RH, Rathore SS, Radford MJ, *et al.*: **Acute myocardial infarction in the elderly: differences by age.** *J Am Coll Cardiol.* 2001; 38(3): 736–41.
[PubMed Abstract](#) | [Publisher Full Text](#)
10. Stone PH, Thompson B, Anderson HV, *et al.*: **Influence of race, sex, and age on management of unstable angina and non-Q-wave myocardial infarction: The TIMI III registry.** *JAMA.* 1996; 275(14): 1104–12.
[PubMed Abstract](#) | [Publisher Full Text](#)
11. Amsterdam EA, Wenger NK, Brindis RG, *et al.*: **2014 AHA/ACC guideline for the management of patients with non-ST-elevation acute coronary syndromes: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines.** *Circulation.* 2014; 130(25): e344–426.
[PubMed Abstract](#) | [Publisher Full Text](#)
12. Avezum A, Makdisse M, Spencer F, *et al.*: **Impact of age on management and outcome of acute coronary syndrome: observations from the Global Registry of Acute Coronary Events (GRACE).** *Am Heart J.* 2005; 149(1): 67–73.
[PubMed Abstract](#) | [Publisher Full Text](#)
13. Puymirat E, Aissaoui N, Cayla G, *et al.*: **Changes in One-Year Mortality in Elderly Patients Admitted with Acute Myocardial Infarction in Relation with Early Management.** *Am J Med.* 2017; 130(5): 555–63.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
14. White HD, Barbash GI, Califf RM, *et al.*: **Age and outcome with contemporary thrombolytic therapy. Results from the GUSTO-I trial. Global Utilization of Streptokinase and TPA for Occluded coronary arteries trial.** *Circulation.* 1996; 94(8): 1826–33.
[PubMed Abstract](#) | [Publisher Full Text](#)
15. Maggioni AP, Maseri A, Fresco C, *et al.*: **Age-related increase in mortality among patients with first myocardial infarctions treated with thrombolysis. The Investigators of the Gruppo Italiano per lo Studio della Sopravvivenza nell'Infarto Miocardico (GISSI-2).** *N Engl J Med.* 1993; 329(20): 1442–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
16. DeGeare VS, Stone GW, Grines L, *et al.*: **Angiographic and clinical characteristics associated with increased in-hospital mortality in elderly patients with acute myocardial infarction undergoing percutaneous intervention (a pooled analysis of the primary angioplasty in myocardial infarction trials).** *Am J Cardiol.* 2000; 86(1): 30–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
17. Alter DA, Manuel DG, Gunraj N, *et al.*: **Age, risk-benefit trade-offs, and the projected effects of evidence-based therapies.** *Am J Med.* 2004; 116(8): 540–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
18. Gale CP, Cattle BA, Woolston A, *et al.*: **Resolving inequalities in care? Reduced mortality in the elderly after acute coronary syndromes. The Myocardial Ischaemia National Audit Project 2003–2010.** *Eur Heart J.* 2012; 33(5): 630–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
19. Savonitto S, Cavallini C, Petronio AS, *et al.*: **Early aggressive versus initially conservative treatment in elderly patients with non-ST-segment elevation acute coronary syndrome: a randomized controlled trial.** *JACC Cardiovasc Interv.* 2012; 5(9): 906–16.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
20. De Carlo M, Morici N, Savonitto S, *et al.*: **Sex-Related Outcomes in Elderly Patients Presenting With Non-ST-Segment Elevation Acute Coronary Syndrome: Insights From the Italian Elderly ACS Study.** *JACC Cardiovasc Interv.* 2015; 8(6): 791–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
21. Tegn N, Abdelnoor M, Aaberge L, *et al.*: **Invasive versus conservative strategy in patients aged 80 years or older with non-ST-elevation myocardial infarction or unstable angina pectoris (After Eighty study): an open-label randomised controlled trial.** *Lancet.* 2016; 387(10023): 1057–65.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
22. Skolnick AH, Alexander KP, Chen AY, *et al.*: **Characteristics, management, and outcomes of 5,557 patients age > or =90 years with acute coronary syndromes: results from the CRUSADE Initiative.** *J Am Coll Cardiol.* 2007; 49(17): 1790–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
23. Bach RG, Cannon CP, Weintraub WS, *et al.*: **The effect of routine, early invasive management on outcome for elderly patients with non-ST-segment elevation acute coronary syndromes.** *Ann Intern Med.* 2004; 141(3): 186–95.
[PubMed Abstract](#) | [Publisher Full Text](#)
24. De Luca L, Olivari Z, Bolognese L, *et al.*: **A decade of changes in clinical characteristics and management of elderly patients with non-ST elevation myocardial infarction admitted in Italian cardiac care units.** *Open Heart.* 2014; 1(1): e000148.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
25. De Luca L, Marini M, Gonzini L, *et al.*: **Contemporary Trends and Age-Specific Sex Differences in Management and Outcome for Patients With ST-Segment Elevation Myocardial Infarction.** *J Am Heart Assoc.* 2016; 5(12): pii: e004202.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
26. Goldberg RM, Mabee J, Chan L, *et al.*: **Drug-drug and drug-disease interactions in the ED: analysis of a high-risk population.** *Am J Emerg Med.* 1996; 14(5): 447–50.
[PubMed Abstract](#) | [Publisher Full Text](#)
27. Alexander KP, Chen AY, Roe MT, *et al.*: **Excess dosing of antiplatelet and antithrombin agents in the treatment of non-ST-segment elevation acute coronary syndromes.** *JAMA.* 2005; 294(24): 3108–16.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
28. Cannon CP, Braunwald E, McCabe CH, *et al.*: **Intensive versus moderate lipid lowering with statins after acute coronary syndromes.** *N Engl J Med.* 2004; 350(15): 1495–504.
[PubMed Abstract](#) | [Publisher Full Text](#)
29. Ray KK, Bach RG, Cannon CP, *et al.*: **Benefits of achieving the NCEP optional LDL-C goal among elderly patients with ACS.** *Eur Heart J.* 2006; 27(19): 2310–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
30. Krumholz HM, Radford MJ, Wang Y, *et al.*: **Early beta-blocker therapy for acute myocardial infarction in elderly patients.** *Ann Intern Med.* 1999; 131(9): 648–54.
[PubMed Abstract](#) | [Publisher Full Text](#)
31. Alexander KP, Roe MT, Chen AY, *et al.*: **Evolution in cardiovascular care for elderly patients with non-ST-segment elevation acute coronary syndromes: results from the CRUSADE National Quality Improvement Initiative.** *J Am Coll Cardiol.* 2005; 46(8): 1479–87.
[PubMed Abstract](#) | [Publisher Full Text](#)
32. Alexander KP, Newby LK, Armstrong PW, *et al.*: **Acute coronary care in the elderly, part II: ST-segment-elevation myocardial infarction: a scientific statement for healthcare professionals from the American Heart Association Council on Clinical Cardiology: in collaboration with the Society of Geriatric Cardiology.** *Circulation.* 2007; 115(19): 2570–89.
[PubMed Abstract](#) | [Publisher Full Text](#)
33. Eagle KA, Goodman SG, Avezum A, *et al.*: **Practice variation and missed opportunities for reperfusion in ST-segment-elevation myocardial infarction: findings from the Global Registry of Acute Coronary Events (GRACE).** *Lancet.* 2002; 359(9304): 373–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
34. American College of Emergency Physicians, Society for Cardiovascular Angiography and Interventions, O'Gara PT, *et al.*: **2013 ACCF/AHA guideline for the management of ST-elevation myocardial infarction: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines.** *J Am Coll Cardiol.* 2013; 61(4): e78–140.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
35. Cantor WJ, Mehta SR, Yuan F, *et al.*: **Radial versus femoral access for elderly patients with acute coronary syndrome undergoing coronary angiography and intervention: insights from the RIVAL trial.** *Am Heart J.* 2015; 170(5): 880–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
36. Frengley JD, Sansone GR, Alba A, *et al.*: **Influence of age on rehabilitation outcomes and survival in post-acute inpatient cardiac rehabilitation.** *J Cardiopulm Rehabil Prev.* 2011; 31(4): 230–8.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

- 1. Sanjay Patel**
Sydney Medical School, University of Sydney, Sydney, Australia
Competing Interests: No competing interests were disclosed.
- 2. Stefano Savonitto**
Division of Cardiology, Manzoni Hospital, Lecco, Italy
Competing Interests: No competing interests were disclosed.
- 3. Anna Sonia Petronio**
University of Pisa, Pisa, Italy
Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research