REVIEW

Novel therapeutic approaches to simultaneously target rhinovirus infection and asthma/COPD pathogenesis [version 1; peer review: 2 approved]

Carmen Mirabelli , Els Scheers, Johan Neyts

Laboratory of Virology and Chemotherapy, Rega Institute for Medical Research, Leuven, Belgium

V1 First published: 19 Oct 2017, 6(F1000 Faculty Rev):1860
<https://doi.org/10.12688/f1000research.11978.1>

Latest published: 19 Oct 2017, 6(F1000 Faculty Rev):1860
<https://doi.org/10.12688/f1000research.11978.1>

Abstract

Rhinoviruses are exclusive respiratory pathogens and the etiological agents of the common cold. These viruses are increasingly reported to cause exacerbations of asthma and chronic obstructive pulmonary disease (COPD). Here, we review the role of rhinovirus infections in the pathogenesis of asthma and COPD and we discuss the current and potential future treatments. We propose that, in order to prevent exacerbations, the design of novel therapeutics should focus on directly acting antivirals but also include the design of drugs that simultaneously inhibit viral replication and alleviate symptoms of asthma and COPD.

Keywords

rhinovirus, asthma, COPD, antiviral

Open Peer Review

Approval Status

	1	2
version 1
19 Oct 2017		

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **James Gern**, University of Wisconsin-Madison, Madison, USA

2. **Lena Uller**, Lund University, Lund, Sweden

Any comments on the article can be found at the end of the article.

Corresponding author: Carmen Mirabelli (carmen.mirabelli@kuleuven.be)

Author roles: **Mirabelli C:** Conceptualization, Project Administration, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Scheers E:** Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Neyts J:** Supervision, Writing – Review & Editing

Competing interests: The authors declare that they have no competing interests.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2017 Mirabelli C *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Mirabelli C, Scheers E and Neyts J. **Novel therapeutic approaches to simultaneously target rhinovirus infection and asthma/COPD pathogenesis [version 1; peer review: 2 approved]** F1000Research 2017, 6(F1000 Faculty Rev):1860 <https://doi.org/10.12688/f1000research.11978.1>

First published: 19 Oct 2017, 6(F1000 Faculty Rev):1860 <https://doi.org/10.12688/f1000research.11978.1>

General introduction of rhinoviruses

Rhinoviruses (RVs) belong to the large family of *Picornaviridae*¹ and are grouped in three species: RV-A, RV-B, and the relatively recently characterized RV-C. Species and type assignment relies on genetic classification, in particular 13% nucleotide divergence in VP1 or more than 10% divergence in VP4/VP2 or both². This classification does not correspond to the receptor usage of RVs, the routes of entry, or the interaction with the immune system of the host³. RV-A and RV-B can bind intercellular adhesion molecule 1 (ICAM-1, major group: the majority of RV-A types and all RV-B types) or low-density lipoprotein receptor (LDLR, minor group: 12 RV-A types). The receptor for RV-C15 was recently identified as the cadherin-related family member 3 (CDHR3)⁴ and is likely shared by all members of species C, although no empirical evidence has been produced for other RV-C types. The viral cycle of RVs is exclusively cytoplasmic. The genome, a single-stranded positive RNA strain, undergoes cap-independent translation into a long polypeptide, cleaved by viral proteases 2A and 3C. Replication and encapsidation of new viral particles occur in loosely organized membrane systems burgeoning from the Golgi apparatus⁵ and are assisted by a number of host factors—protein kinase D, oxysterol binding protein, phosphatidylinositol 4-kinase III beta, and glutathione^{6–8}—together with viral proteins: 3D viral polymerase and the multi-factorial 2C and 3A proteins. Lytic exit of RVs from cells may not be the sole mechanism by which viruses are released. Evidence was recently produced to support the non-lytic release of the newly produced virions within phosphatidylserine lipid-enriched vesicles⁹.

RVs are extremely host-specific respiratory tract pathogens, mainly associated with upper respiratory tract infections in humans. RVs have long been appointed as non-harmful pathogens, and the cure for the common cold has not been considered a high-priority medical need universally. Furthermore, the development of antiviral strategies was largely hampered by the unavailability of models of infection (particularly small animal models). The extreme antigenic diversity of RVs discouraged, and still discourages, the development of vaccines. In addition, the genetic variability within the species necessitates the development of broad-spectrum, pan-RV-active antiviral drugs.

Lately, infections with RVs have been increasingly reported in the bronchial epithelium of the lower respiratory tract and even in B lymphocytes¹⁰, and there are overwhelming data demonstrating that infection with these viruses causes exacerbations of asthma and chronic obstructive pulmonary disease (COPD). In the following paragraphs, we will discuss the involvement of RVs in the potential initiation or the development of these respiratory chronic syndromes (or both) and the possible therapeutic strategies to adopt for prevention and treatment.

Asthma and rhinovirus

Asthma is a disease of the airways characterized by airflow obstruction and airway hyper-responsiveness. Depending on the severity of the disease, asthma is clinically classified as intermittent asthma or persistent asthma (mild, moderate, or severe). However, the

biological definition of asthma is more complex given that more than 10 phenotypes have been described¹¹. Overall, initiation of asthma is associated with enhanced type 2 (T helper 2, Th2) or reduced type 1 immunologic responses (or both), particularly mediated by eosinophils (interleukin [IL]-5) or neutrophils (IL-8). Both eosinophils and neutrophils trigger sustained and chronic airway inflammation, which causes airway remodeling and results in decreased lung function. The majority of asthma cases have an allergic component (immunoglobulin E, IgE). Asthma attacks, clinically termed exacerbations, are mostly induced by allergen exposure and respiratory tract infections. The causes of asthma initiation are currently unknown, and both genetic and environmental factors may contribute to the establishment of the pathological condition.

Potential role of rhinovirus in asthma initiation

Early-life human RV infection has been associated with asthma development in infants. Clinical prospective and retrospective studies demonstrated that children with RV-related wheezing illnesses present an increased risk of developing childhood asthma (Childhood Allergy Study [CAS], Childhood Origins of Asthma [COAST], Copenhagen Prospective Studies on Asthma in Childhood [COPSAC], and Tucson Children's Respiratory Study [TCRS] studies). These findings have been extensively reviewed elsewhere^{12,13}. The role of RV infection in the initiation of asthma remains unclear. Recently, a genome-wide association study revealed a link between a polymorphism in the CDHR3 gene (C529Y) and early childhood asthma¹⁴. Interestingly, CDHR3 is the putative receptor of the RV-C type and the CDHR3-Y₅₂₉ polymorphism has been associated with a higher expression of the receptor at the cell surface than CDHR3-C₅₂₉. A consequent increased susceptibility to RV-C infection could cause illness early in life, airway damage and remodeling, lung function decline, and ultimately asthma development¹⁵. A convincing study by the Hershenov group recently described an increased expression of the type 2 cytokines IL-13, IL-4, and IL-5 and decreased expression of type 1 cytokine genes, interferon gamma (IFN γ) and IL12p40, immediately after RV infection in neonatal mice and not in adult mice¹⁶. In addition, in these neonatal mice, RV infection induced sustained IL-13 expression, mucous metaplasia, and airway hyper-responsiveness, which are typical pathophysiologic changes described in asthmatic humans¹⁷. In addition to having a potential role in asthma initiation, RV infection is known to be a major cause of asthma exacerbations.

In adults, a study of cohabiting partners, one of whom had asthma, found that although there was no difference in the frequency of RV infections, the partners with asthma exhibited greater lower respiratory symptoms and change in airway physiology¹⁸. The causal link between allergic inflammation and increased susceptibility to RVs was recently proved by prophylactic treatment with an anti-IgE, omalizumab. It was shown that prophylactic treatment with omalizumab decreases fall exacerbation rates¹⁹ and reduces the duration of RV infections, viral shedding, and the risk of RV illnesses²⁰ in asthmatic children and young adults. Little is known about the fundamental mechanisms responsible for the fact that

asthmatics are more susceptible to RV-induced airway disease. Recent evidence points to an abnormal innate antiviral immunity (IFN dysregulation in airways), exaggerated production of inflammatory molecules such as the thymic stromal lymphopoietin, which primes Th2 responses, and an altered antibacterial host defense during acute RV infection²¹.

Chronic obstructive pulmonary disease and rhinovirus

COPD is a leading cause of mortality with an estimated 3 million deaths globally per year (5% of all deaths). COPD is a group of respiratory disorders characterized by a progressive, non-reversible airflow limitation, associated with a chronic inflammatory response in the lung to inhaled environmental agents, mainly cigarette smoke and biomass fuels. In contrast to asthma, COPD develops slowly and usually becomes apparent after the age of 40 years with risk for mortality from age 50²². Although it is well known that COPD is triggered by environmental factors (and mainly cigarette smoking), the underlying mechanism of individual susceptibility is not yet fully understood²³. The principal feature of COPD is airflow limitation by thickening of the small-airway walls and destruction of lung tissue, mainly caused by enhanced pulmonary inflammation (recruitment of macrophages, neutrophils, and CD8⁺ T cells), oxidative stress (augmented activity of reactive oxygen species), and a protease/antiprotease imbalance^{24–26}. As the disease progresses, tertiary lymphoid aggregates have been shown to develop around the small airways^{24,27}.

Mechanisms of rhinovirus-induced exacerbation in chronic obstructive pulmonary disease

Much of the morbidity, mortality, and health-care costs of COPD is attributable to acute worsening of respiratory symptoms or exacerbations. Exacerbations are commonly caused by viral or bacterial airway infections or both. The viral pathogens responsible for 60% to 80% of all exacerbations are RV, the respiratory syncytial virus, and influenza virus²⁸. In particular, RV has been detected (by PCR) in 3.1% to 27% of COPD exacerbations²⁹. Recently, a human model to study RV-induced exacerbations of COPD formally proved a causative relationship between RV infection and COPD exacerbation²⁵. George and colleagues reported that the frequency of exacerbations correlated with RV viral loads in the sputum³⁰. The authors proposed that the mechanism of increased viral susceptibility in patients with COPD resides in the modulation of ICAM-1 on respiratory epithelial cells³⁰. Interestingly, increased expression of ICAM-1 at the surface of airway epithelial cells was also reported in response to tobacco smoke exposure³¹. Mallia and colleagues have provided evidence that RV infection in patients with COPD is associated with recruitment of circulating T lymphocytes to the lungs and that the T-cell numbers in bronchial alveolar lavage correlate with viral load³². Evidence of a direct effect of the viral protease 2A in the induction of Th1 and Th2 immune responses from CD4 T cells was also described *in vitro*³³. Furthermore, intranasal administration of recombinant RV protease 2A in mice resulted in an increased airway hyper-reactivity, lung inflammation, and IL-4 and IFN- γ production, suggesting a direct role of 2A in the Th1 and Th2 hyper-response during COPD exacerbations³³.

Antiviral strategies for rhinovirus infection

The genetic diversity of RVs has so far hampered the development of both prophylactic and therapeutic antiviral strategies. More than 150 serotypes of RV are known, and cross-protection appears to be limited. The slow and disappointing efforts to develop RV vaccines have been exhaustively reviewed elsewhere^{34,35}. For the development of small antiviral molecules, each step of RV viral cycle could theoretically represent a target, and several candidates have been described over the years. Comprehensive reviews on efforts to develop small-molecule inhibitors of RV/enterovirus replication have been published^{36,37}. The most-studied compounds are presented in Figure 1A.

Capsid binders are inhibitors of viral entry, and the most/best-studied series is composed of the so-called WIN compounds. These molecules bind a hydrophobic pocket located under the floor of the canyon, which is the depression on the viral capsid surface involved in cell receptor binding. The drug fits the pocket and stabilizes the virion by pushing the floor of the canyon upwards, thereby preventing capsid conformational changes necessary for virus uncoating³⁸. The potential efficacy of three WIN-capsid binders (that is, pirodnavir, pleconaril, and vapendavir) against RV infections in humans was assessed by the Janssen Research Foundation, Schering-Plough (nasal formulation by ViroPharma), and Biota, respectively (39, clinicaltrial.gov). Pirodnavir was the first small-molecule shown to prevent experimental RV infection in a human challenge model. However, the compound was active only when administered six times daily and when treatment was initiated within 10 minutes after RV challenge. When start of treatment was delayed until 24 or 48 hours after challenge, the antiviral effect was no longer observed⁴⁰. Pleconaril failed the first clinical trial with an oral formulation that was given three times per day when treatment started 24 hours after challenge (only 1 to 1.5 days of reduction in symptom alleviation time was demonstrated)⁴¹. Furthermore, drug-resistant RV strains were identified in 24% of enrolled patients: 13% of patients were naturally resistant at baseline, and 11% exhibited a reduced susceptibility by day 5 of treatment^{39,42}. A phase II study, which used an intranasal formulation of pleconaril, failed to show a statistically significant result for RV-positive participants either with or without asthma exacerbations³⁹. Recently, vapendavir in a phase IIb clinical trial (SPIRITUS) resulted in a statistically significant antiviral effect (RV PCR-negative) for patients who received the compound within 24 hours after onset of symptoms. However, no reduction in secondary endpoints (lung function and reduction in asthma exacerbations) was noted (Biota website). Overall, although capsid binders are attractive and potent early-stage inhibitors of RV replication *in vitro*, problems in pharmacodynamics, *in vivo* efficacy, and resistance development were reported. In addition, with the recent resolution of the capsid structure, it was concomitantly demonstrated that RV-C species lack the hydrophobic pocket, the binding target of capsid binders. Therefore, pan-RV coverage cannot be achieved with this specific class of compounds⁴³. In contrast to capsid binders, the pan-enterovirus 3C protease inhibitor rupintrivir, a Michaelis acceptor, exerts low nanomolar potency against a large panel of RV-A and RV-B serotypes *in vitro*. Moreover, activity was demonstrated against RV-C in a replicon

Figure 1. Rhinovirus inhibitors and asthma/COPD therapeutics. (A) Schematic representation of the rhinovirus replication cycle, including different classes of inhibitors and representative examples thereof (*italics*). **(B)** Cross-section of a bronchial tube with asthma/COPD pathogenesis and detail of the epithelial barrier illustrating the enhanced type 2 (Th2) immunologic response leading to airway remodeling. Relevant selections of therapeutics used for asthma/COPD treatment are indicated in *italics*. DC, dendritic cell; IFN γ , interferon gamma; IL, interleukin; LABA, long-acting B2-agonist bronchodilator; PDE4, phosphodiesterase 4; SABA, short-acting B2-agonist bronchodilator; ssRNA, single-stranded RNA; TGF β , transforming growth factor-beta; UTR, untranslated region.

system⁴⁴. Other protease inhibitors have been successfully developed as antiviral drugs for the treatment of infections with the human immunodeficiency virus and more recently the hepatitis C virus (HCV). However, the antiviral effect of the orally available analogue of rupintrivir, named “Compound 1”, was investigated in naturally infected RV patients and no reduction in disease severity or viral load was noted. Hence, further development was halted³⁹, and the effect of the treatment was not evaluated in the context of patients with asthma or COPD.

A safe and efficient antiviral for the treatment of RVs would be appreciated by many, including otherwise-healthy patients with common cold, to reduce the time of symptoms and, in turn, reduce the costs of school/work productivity loss. But mostly, a highly efficient therapy is needed for the treatment or prophylaxis of RV infections in those with underlying conditions such as asthma and COPD. Unfortunately, so far, there has been no proof of concept that a direct-targeting anti-RV treatment could prevent exacerbations and alleviate asthma or COPD symptoms. In the next paragraphs, we will discuss the potential design and development of combination therapies, which may combat RV infection/replication and asthma/COPD symptoms simultaneously and constitute

an interesting strategy in patients with compromising chronic conditions.

Current treatment for asthma and chronic obstructive pulmonary disease

Asthma is a progressive pathologic condition, which generally manifests before 12 years of age. For intermittent episodes of asthma, patients undergo short-acting B2-agonist bronchodilator (SABA) treatment. For persistent asthma, inhaled corticosteroids (ICSs), the most effective anti-inflammatory medication, are coupled with SABAs or long-acting B2-agonist bronchodilators (LABAs), according to the severity of the disease. In case of uncontrolled severe asthma, tiotropium (patients > 12 years old, exacerbated), anti-IgE (patients > 6 years old), or anti-IL-5 (patients > 12 years old, eosinophilic asthma) is added⁴⁵. Patients with COPD differ markedly from patients with asthma: the former are older and mostly current or ex-smokers and have impaired lung function and their airflow obstruction is not reversible. Current treatments for COPD exacerbations consist of supportive therapies: SABA or LABA in addition to ICS according to the severity of disease. In the case of severe COPD, oral phosphodiesterase 4 inhibitors and azithromycin are prescribed⁴⁶. Tiotropium is also used for

maintenance therapy during stable COPD to reduce symptoms and the frequency and severity of exacerbations^{46,47}. No new therapeutics have been developed over the past three decades, and the use of antibiotics for COPD is debated given the emergence and spread of antibiotic resistance. The main therapeutics for asthma and COPD are illustrated in [Figure 1B](#).

Rhinovirus infection interferes with the current treatment for asthma and chronic obstructive pulmonary disease

Glucocorticoids (GCs), belonging to the ICS family of drugs, are a highly effective group of anti-inflammatory agents, which are widely used in the treatment of chronic inflammatory airway diseases, including asthma and COPD. However, results from clinical trials suggest that respiratory viral infections affect GC therapy^{48–50}. In particular, RV-induced exacerbations of asthma and COPD have been shown to be refractory to the anti-inflammatory effect of GCs⁵⁰. Xia and colleagues suggested the mechanism of reduced sensitivity through an airway epithelium model. RV infection appeared to increase the expression and activity of transforming growth factor-beta, which in turn induced GC insensitivity⁵¹.

Interference of viral infection on current treatment was also reported in the context of budesonide or formoterol treatment (or both), two common LABAs. *In vitro* data on RV-stimulated peripheral blood mononuclear cells suggest that the combination of budesonide and formoterol inhibits RV-induced upregulation of CXCL10 and in turn has a negative effect on antiviral responses⁵². Indeed, IFN- α secretion and expression of the type I IFN-inducible genes were enhanced, suggesting a negative interference of drugs on RV infection (less protection and more infection). In contrast, when bronchial epithelial cells were infected with RV, budesonide/formoterol treatment did not induce a significant inhibitory effect on CXCL10 secretion or type I and III IFN gene induction, suggesting that the drugs could have a beneficial effect on RV-induced asthma exacerbations (more protection and less infection)⁵³. It remains to be elucidated which of the findings is relevant to the situation in patients.

Azithromycin is a macrolide antibiotic used for the treatment of COPD since it shows a reduction in exacerbation frequency together with an improvement in quality of life of patients with COPD⁵⁴. When primary bronchial epithelial cells from COPD donors or healthy individuals were treated with azithromycin 24 hours before infection with RV, azithromycin transiently induced increases of IFN β , IFN λ , and RIG-I expression. After infection, azithromycin augmented RV-induced expression of IFN- and RIG-I like-helicases in cells derived from patients with COPD and this decreased viral load. This effect was not observed in cultures obtained from healthy individuals. These data support azithromycin's emerging role in the prevention of exacerbations of COPD⁵⁵.

Conclusion and future perspective

The aforementioned examples point to the importance of studying the potential interference between RV infection and asthma/COPD treatment agents. New clinical trials take into consideration

this problem of interference (clinicaltrials.gov). Therefore, the efficacy of (a) OC459, an antagonist of the prostaglandin D2 receptor 2 (known to stimulate the *in vitro* chemotaxis of Th2 cells and leukocytes and to release the granule content of eosinophils and basophils), and (b) CNTO 3157, an antagonist of Toll-like receptor 3 (which mediates antiviral response and is involved in asthma exacerbation and development), is now being explored in the context of RV infection in humans. When human RV challenge experiments are not possible or not ethical (in the case of patients with severe asthma/COPD symptoms), physiologically relevant *in vitro* models should be used. In particular, airway epithelium culture derived from healthy patients or patients with underlying chronic respiratory syndromes should be collected and differentiated *in vitro*. Moreover, the strain of RV to infect such cultures should be clinically relevant. Although they are highly prevalent in RV-induced exacerbation of asthma and COPD, little is known about the effect of RV-C species. Mechanistic studies addressing the potential differences of these strains in the context of asthma/COPD are awaited and the results should be directly compared with those obtained with the better-studied RV-A strains.

In the development of novel therapeutic approaches, efforts should be addressed toward dual-target molecules, which inhibit RV replication and target the underlying pathogenesis of asthma and COPD at the same time. Dual-target therapies are often considered far-fetched and unrealistic. However, the interesting mechanism of action of azithromycin demonstrates that dual-target therapies can be achieved⁵⁵. IFN signaling could be an important target to alleviate asthma symptoms but also to prime the cells and subsequently prevent RV infection. Besides targeting IFN-associated pathways, other cellular targets may be used to develop synergistically acting therapeutics. We recently reported on a class of broad-spectrum enterovirus/RV inhibitors that are also correctors of the cystic fibrosis transmembrane conductance regulator (CFTR) folding⁵⁶. Deficiencies in CFTR are associated with more than 90% of cystic fibrosis cases. Similarly, cytokines such as the thymic stromal lymphopoietin or its receptor are pivotal in the pathophysiology of asthma (Th2-upstream activator) and are becoming an attractive cellular target for the treatment of asthma⁵⁷. Multi-target drug design could be employed to identify chemical scaffolds with dual activity as antiviral and anti-asthmatic compound.

The development of directly acting antivirals with an additional effect on exacerbations of asthma and COPD should also be explored. To that end, the RV 2A protease, which is an essential protein for viral replication, may be a potentially interesting target. Indeed, as previously mentioned, the expression of RV 2A was reported to induce Th1- and Th2-mediated inflammation *in vitro* and *in vivo*, suggesting a direct role of 2A in COPD exacerbations³³. Highly potent and directly acting antivirals have been developed in recent years for HCV, which, similarly to RV, is a positive single-stranded RNA virus. Today, treatment of chronic HCV is extremely successful and most patients are cured from the infection and are no longer at risk of death from chronic liver cirrhosis and hepatocellular carcinoma. Likewise, highly potent

inhibitors for RV, to be used either prophylactically or therapeutically⁵⁸, may be sufficient to cure the infection and block the progression of underlying chronic conditions such as asthma and COPD. We propose that, together with direct pan-RV antiviral strategies, novel multi-target or combination approaches that simultaneously interfere with viral replication and the pathogenesis (of exacerbations) of asthma and COPD should be rationally designed and developed.

Abbreviations

CDHR3, cadherin-related family member 3; CFTR, cystic fibrosis transmembrane conductance regulator; COPD, chronic obstructive pulmonary disease; GC, glucocorticoid; HCV, hepatitis C virus; IgE,

immunoglobulin E; ICAM-1, intercellular adhesion molecule 1; ICS, inhaled corticosteroid; IFN, interferon; IL, interleukin; LABA, long-acting B2-agonist bronchodilator; RV, rhinovirus; SABA, short-acting B2-agonist bronchodilator; Th2, T helper 2.

Competing interests

The authors declare that they have no competing interests.

Grant information

The author(s) declared that no grants were involved in supporting this work.

References

- Royston L, Tapparel C: **Rhinoviruses and Respiratory Enteroviruses: Not as Simple as ABC**. *Viruses*. 2016; **8**(1): pii: E16.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Bochkov YA, Grindle K, Vang F, *et al.*: **Improved molecular typing assay for rhinovirus species A, B, and C**. *J Clin Microbiol*. 2014; **52**(7): 2461–71.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Blaas D, Fuchs R: **Mechanism of human rhinovirus infections**. *Mol Cell Pediatr*. 2016; **3**(1): 21.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Bochkov YA, Watters K, Ashraf S, *et al.*: **Cadherin-related family member 3, a childhood asthma susceptibility gene product, mediates rhinovirus C binding and replication**. *Proc Natl Acad Sci U S A*. 2015; **112**(17): 5485–90.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Quiner CA, Jackson WT: **Fragmentation of the Golgi apparatus provides replication membranes for human rhinovirus 1A**. *Virology*. 2010; **407**(2): 185–95.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Guedán A, Swieboda D, Charles M, *et al.*: **Investigation of the Role of Protein Kinase D in Human Rhinovirus Replication**. *J Virol*. 2017; **91**(9): pii: e00217-17.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Spickler C, Lippens J, Laberge MK, *et al.*: **Phosphatidylinositol 4-kinase III beta is essential for replication of human rhinovirus and its inhibition causes a lethal phenotype in vivo**. *Antimicrob Agents Chemother*. 2013; **57**(7): 3358–68.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Thibaut HJ, van der Linden L, Jiang P, *et al.*: **Binding of glutathione to enterovirus capsids is essential for virion morphogenesis**. *PLoS Pathog*. 2014; **10**(4): e1004039.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Chen YH, Du W, Hagemeijer MC, *et al.*: **Phosphatidylserine vesicles enable efficient en bloc transmission of enteroviruses**. *Cell*. 2015; **160**(4): 619–30.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Aab A, Wirz O, van de Veen W, *et al.*: **Human rhinoviruses enter and induce proliferation of B lymphocytes**. *Allergy*. 2017; **72**(2): 232–43.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Edwards MR, Bartlett NW, Hussell T, *et al.*: **The microbiology of asthma**. *Nat Rev Microbiol*. 2012; **10**(7): 459–71.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Jackson DJ, Gern JE, Lemanske RF Jr: **The contributions of allergic sensitization and respiratory pathogens to asthma inception**. *J Allergy Clin Immunol*. 2016; **137**(3): 659–65; quiz 666.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Jackson DJ, Gern JE, Lemanske RF: **Lessons learned from birth cohort studies conducted in diverse environments**. *J Allergy Clin Immunol*. 2017; **139**(2): 379–86.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Bonnelykke K, Sleiman P, Nielsen K, *et al.*: **A genome-wide association study identifies CDHR3 as a susceptibility locus for early childhood asthma with severe exacerbations**. *Nat Genet*. 2014; **46**(1): 51–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Bochkov YA, Gern JE: **Rhinoviruses and Their Receptors: Implications for Allergic Disease**. *Curr Allergy Asthma Rep*. 2016; **16**(4): 30.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Hong JY, Bentley JK, Chung Y, *et al.*: **Neonatal rhinovirus induces mucous metaplasia and airways hyperresponsiveness through IL-25 and type 2 innate lymphoid cells**. *J Allergy Clin Immunol*. 2014; **134**(2): 429–39.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Al-Muhsen S, Johnson JR, Hamid Q: **Remodeling in asthma**. *J Allergy Clin Immunol*. 2011; **128**(3): 451–62; quiz 463–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Corne JM, Marshall C, Smith S, *et al.*: **Frequency, severity, and duration of rhinovirus infections in asthmatic and non-asthmatic individuals: a longitudinal cohort study**. *Lancet*. 2002; **359**(9309): 831–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Teach SJ, Gill MA, Togias A, *et al.*: **Preseasonal treatment with either omalizumab or an inhaled corticosteroid boost to prevent fall asthma exacerbations**. *J Allergy Clin Immunol*. 2015; **136**(6): 1476–85.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Esquivel A, Busse WW, Calatroni A, *et al.*: **Effects of Omalizumab on Rhinovirus Infections, Illnesses and Exacerbations of Asthma**. *Am J Respir Crit Care Med*. 2017.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Perez GF, Rodriguez-Martinez CE, Nino G: **Rhinovirus-Induced Airway Disease: A Model to Understand the Antiviral and Th2 Epithelial Immune Dysregulation in Childhood Asthma**. *J Invest Med*. 2015; **63**(6): 792–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Lozano R, Naghavi M, Foreman K, *et al.*: **Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010**. *Lancet*. 2012; **380**(9859): 2095–128.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Vestbo J, Hurd SS, Agustí AG, *et al.*: **Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: GOLD executive summary**. *Am J Respir Crit Care Med*. 2013; **187**(4): 347–65.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Decramer M, Janssens W, Miravittles M: **Chronic obstructive pulmonary disease**. *Lancet*. 2012; **379**(9823): 1341–51.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Gunawardana N, Finney L, Johnston SL, *et al.*: **Experimental rhinovirus infection in COPD: implications for antiviral therapies**. *Antiviral Res*. 2014; **102**: 95–105.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Gao W, Li L, Wang Y, *et al.*: **Bronchial epithelial cells: The key effector cells in the pathogenesis of chronic obstructive pulmonary disease?** *Respirology*. 2015; **20**(5): 722–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Polverino F, Seys LJ, Bracke KR, *et al.*: **B cells in chronic obstructive pulmonary disease: moving to center stage**. *Am J Physiol Lung Cell Mol Physiol*. 2016; **311**(4): L687–L695.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Biancardi E, Fennell M, Rawlinson W, *et al.*: **Viruses are frequently present as the infecting agent in acute exacerbations of chronic obstructive pulmonary disease in patients presenting to hospital**. *Intern Med J*. 2016; **46**(10): 1160–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)

29. Kurai D, Saraya T, Ishii H, *et al.*: **Virus-induced exacerbations in asthma and COPD.** *Front Microbiol.* 2013; 4: 293.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
30. **F** George SN, Garcha DS, Mackay AJ, *et al.*: **Human rhinovirus infection during naturally occurring COPD exacerbations.** *Eur Respir J.* 2014; 44(1): 87–96.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
31. **F** Shukla SD, Mahmood MQ, Weston S, *et al.*: **The main rhinovirus respiratory tract adhesion site (ICAM-1) is upregulated in smokers and patients with chronic airflow limitation (CAL).** *Respir Res.* 2017; 18(1): 6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
32. **F** Mallia P, Message SD, Contoli M, *et al.*: **Lymphocyte subsets in experimental rhinovirus infection in chronic obstructive pulmonary disease.** *Respir Med.* 2014; 108(1): 78–85.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
33. Singh M, Lee SH, Porter P, *et al.*: **Human rhinovirus proteinase 2A induces TH1 and TH2 immunity in patients with chronic obstructive pulmonary disease.** *J Allergy Clin Immunol.* 2010; 125(6): 1369–1378.e2.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
34. McLean GR: **Developing a vaccine for human rhinoviruses.** *J Vaccines Immun.* 2014; 2(3): 16–20.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
35. Glanville N, Johnston SL: **Challenges in developing a cross-serotype rhinovirus vaccine.** *Curr Opin Virol.* 2015; 11: 83–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
36. Bauer L, Lyoo H, van der Schaar HM, *et al.*: **Direct-acting antivirals and host-targeting strategies to combat enterovirus infections.** *Curr Opin Virol.* 2017; 24: 1–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
37. Thibaut HJ, De Palma AM, Neyts J: **Combating enterovirus replication: state-of-the-art on antiviral research.** *Biochem Pharmacol.* 2012; 83(2): 185–92.
[PubMed Abstract](#) | [Publisher Full Text](#)
38. McKinlay MA, Pevear DC, Rossmann MG: **Treatment of the picornavirus common cold by inhibitors of viral uncoating and attachment.** *Annu Rev Microbiol.* 1992; 46: 635–54.
[PubMed Abstract](#) | [Publisher Full Text](#)
39. Patick AK: **Rhinovirus chemotherapy.** *Antiviral Res.* 2006; 71(2–3): 391–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
40. Hayden FG, Hipskind GJ, Woerner DH, *et al.*: **Intranasal pirodavir (R77,975) treatment of rhinovirus colds.** *Antimicrob Agents Chemother.* 1995; 39(2): 290–4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
41. Hayden FG, Coats T, Kim K, *et al.*: **Oral pleconaril treatment of picornavirus-associated viral respiratory illness in adults: efficacy and tolerability in phase II clinical trials.** *Antivir Ther.* 2002; 7(1): 53–65.
[PubMed Abstract](#)
42. Pevear DC, Hayden FG, Demenczuk TM, *et al.*: **Relationship of pleconaril susceptibility and clinical outcomes in treatment of common colds caused by rhinoviruses.** *Antimicrob Agents Chemother.* 2005; 49(11): 4492–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
43. **F** Liu Y, Hill MG, Klose T, *et al.*: **Atomic structure of a rhinovirus C, a virus species linked to severe childhood asthma.** *Proc Natl Acad Sci U S A.* 2016; 113(2): 8997–9002.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
44. Mello C, Aguayo E, Rodriguez M, *et al.*: **Multiple classes of antiviral agents exhibit *in vitro* activity against human rhinovirus type C.** *Antimicrob Agents Chemother.* 2014; 58(3): 1546–55.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
45. Becker AB, Abrams EM: **Asthma guidelines: the Global Initiative for Asthma in relation to national guidelines.** *Curr Opin Allergy Clin Immunol.* 2017; 17(2): 99–103.
[PubMed Abstract](#) | [Publisher Full Text](#)
46. Bostock-Cox B: **The 2017 GOLD guideline on COPD: implications for practice.** *Pract Nurse.* [Internet]. 2017; 47(1): 30–4.
[Reference Source](#)
47. **F** Vogelmeier CF, Asijee GM, Kupas K, *et al.*: **Tiotropium and Salmeterol in COPD Patients at Risk of Exacerbations: A Post Hoc Analysis from POET-COPD(®).** *Adv Ther.* 2015; 32(6): 537–47.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
48. Gustafson LM, Proud D, Hendley JO, *et al.*: **Oral prednisone therapy in experimental rhinovirus infections.** *J Allergy Clin Immunol.* 1996; 97(4): 1009–14.
[PubMed Abstract](#) | [Publisher Full Text](#)
49. Grünberg K, Sharon RF, Hiltermann TJ, *et al.*: **Experimental rhinovirus 16 infection increases intercellular adhesion molecule-1 expression in bronchial epithelium of asthmatics regardless of inhaled steroid treatment.** *Clin Exp Allergy.* 2000; 30(7): 1015–23.
[PubMed Abstract](#) | [Publisher Full Text](#)
50. **F** Hayward G, Thompson MJ, Perera R, *et al.*: **Corticosteroids for the common cold.** *Cochrane Database Syst Rev.* 2015; (10): CD008116.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
51. **F** Xia YC, Radwan A, Keenan CR, *et al.*: **Glucocorticoid Insensitivity in Virally Infected Airway Epithelial Cells Is Dependent on Transforming Growth Factor-β Activity.** *PLoS Pathog.* 2017; 13(1): e1006138.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
52. Davies JM, Carroll ML, Li H, *et al.*: **Budesonide and formoterol reduce early innate anti-viral immune responses *in vitro*.** *PLoS One.* 2011; 6(11): e27898.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
53. Bochkov YA, Busse WW, Brockman-Schneider RA, *et al.*: **Budesonide and formoterol effects on rhinovirus replication and epithelial cell cytokine responses.** *Respir Res.* 2013; 14(1): 98.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
54. **F** Uzun S, Djamin RS, Kluytmans JA, *et al.*: **Azithromycin maintenance treatment in patients with frequent exacerbations of chronic obstructive pulmonary disease (COLUMBUS): a randomised, double-blind, placebo-controlled trial.** *Lancet Respir Med.* 2014; 2(5): 361–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
55. **F** Menzel M, Akbarshahi H, Bjerrmer L, *et al.*: **Azithromycin induces anti-viral effects in cultured bronchial epithelial cells from COPD patients.** *Sci Rep.* 2016; 6: 28698.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
56. Tassini S, Sun L, Lanko K, *et al.*: **Discovery of Multitarget Agents Active as Broad-Spectrum Antivirals and Correctors of Cystic Fibrosis Transmembrane Conductance Regulator for Associated Pulmonary Diseases.** *J Med Chem.* 2017; 60(4): 1400–16.
[PubMed Abstract](#) | [Publisher Full Text](#)
57. Wang WL, Li HY, Zhang MS, *et al.*: **Thymic stromal lymphopoietin: a promising therapeutic target for allergic diseases.** *Int Arch Allergy Immunol.* 2013; 160(1): 18–26.
[PubMed Abstract](#) | [Publisher Full Text](#)
58. Thibaut HJ, Lacroix C, De Palma AM, *et al.*: **Toward antiviral therapy/prophylaxis for rhinovirus-induced exacerbations of chronic obstructive pulmonary disease: challenges, opportunities, and strategies.** *Rev Med Virol.* 2016; 26(1): 21–33.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. **Lena Uller**

Lund University, Lund, Sweden

Competing Interests: No competing interests were disclosed.

2. **James Gern**

University of Wisconsin-Madison, Madison, WI, USA

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research