REVIEW

Rehabilitation following hemorrhagic stroke: building the case for stroke-subtype specific recovery therapies [version 1; peer review: 2 approved]

Tomoko Kitago^{1,2}, Rajiv R. Ratan^{2,3}

¹Department of Neurology, Columbia University, New York, USA

²Burke Medical Research Institute, White Plains, New York, USA

³Departments of Neurology and Neuroscience, Weill Cornell Medicine, New York, USA

V1 First published: 23 Nov 2017, 6(F1000 Faculty Rev):2044
<https://doi.org/10.12688/f1000research.11913.1>

Latest published: 23 Nov 2017, 6(F1000 Faculty Rev):2044
<https://doi.org/10.12688/f1000research.11913.1>

Abstract

Intracerebral hemorrhage (ICH), a form of brain bleeding and minor subtype of stroke, leads to significant mortality and long-term disability. There are currently no validated approaches to promote functional recovery after ICH. Research in stroke recovery and rehabilitation has largely focused on ischemic stroke, but given the stark differences in the pathophysiology between ischemic and hemorrhagic stroke, it is possible that strategies to rehabilitate the brain in distinct stroke subtypes will be different. Here, we review our current understanding of recovery after primary intracerebral hemorrhage with the intent to provide a framework to promote novel, stroke-subtype specific approaches.

Keywords

Rehabilitation, Hemorrhagic stroke, Recovery

Open Peer Review

Approval Status

	1	2
version 1
23 Nov 2017		

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Craig S. Anderson**, Royal Prince Alfred Hospital, Sydney, Australia
Peking University Health Science Center, Beijing, China
University of New South Wales, Sydney, Australia
2. **Klaus Van Leyen**, Massachusetts General Hospital, Harvard Medical School, Boston, USA

Any comments on the article can be found at the

end of the article.

Corresponding author: Tomoko Kitago (tok7002@med.cornell.edu)

Author roles: **Kitago T:** Conceptualization, Writing – Original Draft Preparation; **Ratan RR:** Conceptualization, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This article was funded by an unrestricted gift from John K. Castle (TK).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2017 Kitago T and Ratan RR. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Kitago T and Ratan RR. **Rehabilitation following hemorrhagic stroke: building the case for stroke-subtype specific recovery therapies [version 1; peer review: 2 approved]** F1000Research 2017, 6(F1000 Faculty Rev):2044 <https://doi.org/10.12688/f1000research.11913.1>

First published: 23 Nov 2017, 6(F1000 Faculty Rev):2044 <https://doi.org/10.12688/f1000research.11913.1>

Introduction

Intracerebral hemorrhage (ICH) accounts for about 15% of all strokes in the USA and Europe and leads to significant mortality and long-term disability despite advances in acute medical and surgical treatment^{1,2}. In humans, primary ICH occurs because of a rupture of an intracerebral vessel, most often due to hypertension or cerebral amyloid angiopathy. Vessel rupture leads to mechanical tissue disruption (secondary to the jet of blood) and hematoma formation, followed by secondary injury processes, including disruption of the blood-brain barrier, expansion of perihematomal edema, inflammation, neuronal cell death, and chemical or cellular barriers to repair³. These secondary injury processes occur over hours to days after the primary injury and are attributable to toxins from lysed blood, including hemoglobin (Hgb), heme (oxidized form of heme), and thrombi⁴⁻⁸. These toxins could affect cell survival or repair via interactions with a host of cell types, not only neurons.

Although research in the field of stroke recovery and rehabilitation has grown, there have been very few studies specifically focusing on rehabilitation in ICH. In this review, we summarize the recent advances in our knowledge of the pathophysiology of primary ICH, how rehabilitation affects functional outcomes after ICH, and the mechanisms that may underlie the gains seen with rehabilitation in preclinical models and patients.

Pathophysiology of intracerebral hemorrhage

Preclinical models have provided important insights into the pathophysiological processes that occur after ICH, which differ from those seen after ischemic stroke⁹. The two most commonly used rodent models of ICH are autologous blood injection and bacterial collagenase infusion¹⁰. In the whole-blood injection model, lysed blood is injected directly into the brain parenchyma, with almost no spontaneous bleeding¹¹. The bacterial collagenase model involves intraparenchymal infusion of bacterial collagenase, an enzyme that damages the basal lamina and causes bleeding that evolves and grows over 24 hours¹². Models of hypertensive rodents in which ICH occurs spontaneously, which may more closely mimic what occurs in patients, have been less widely used. In reviewing the literature, one must consider that these preclinical models have different pathophysiological processes which may affect the course of recovery and response to rehabilitation; thus, the use of multiple models has been recommended to better understand post-ICH recovery in preclinical studies¹³.

After ICH, there is both immediate and ongoing cell death in both whole-blood and collagenase models¹⁴. In the collagenase model, there is markedly greater delayed cell death, leading to worse functional deficits even when the hematoma size is matched¹⁴. Historically, it was believed that Hgb or heme induced toxicity following ICH by loading neurons with redox-active free iron leading to the formation of hydroxyl radicals via Fenton chemistry⁹. Indirect support for this model came from studies, not replicated by all groups, that iron chelators such as deferoxamine (DFO) improve functional recovery in pig¹⁵ and rodent¹⁶ models. More recent studies suggest that DFO and a more selective metal chelator, adaptaquin, act not by inhibiting Fenton chemistry

but by inhibiting a specific family of iron-, 2-oxoglutarate-, and oxygen-dependent dioxygenases, the hypoxia-inducible factor prolyl hydroxylases (HIF PHDs)¹⁷. HIF PHDs are oxygen sensors and accordingly, with drugs such as adaptaquin, the brain can be fooled into thinking it is hypoxic when it is not. The consequence of turning on hypoxic adaptation in ICH with adaptaquin is to inhibit the synthesis of a cassette of genes (Chac1, Trib3, and Xc-transporter) that are involved in mediating neuronal death. Accordingly iron, likely via its ability to act as a cofactor for the HIF PHDs, promotes the synthesis of ATF4-dependent genes that mediate neuronal death.

Of note, ATF4 regulates the synthesis of a pseudokinase inhibitor of AKT, called tribbles homolog 3 (Trib3). Studies in non-neural cells have shown that Trib3 and ATF4 can dimerize to inhibit transcription by the pro-plasticity transcription factor, CREB (cAMP-responsive element-binding protein). Activators of CREB have been shown to improve learning and memory as well as functional recovery after stroke. Accordingly, iron's ability to load the HIF PHDs could lead to the transcriptional induction of genes (for example, Trib3) that constrain neuronal survival and plasticity following hemorrhagic stroke. The model is consistent with data showing that adaptaquin, a small-molecule inhibitor of the HIF PHDs, improves functional recovery out to a month in collagenase and blood infusion models in mice and rats, respectively. Future studies will clarify the role of ATF4 in regenerative failure following ICH.

How do cells die following ICH in cellular and rodent models? Recent studies from two groups suggest that cell death in ICH is completely distinct from that found following ischemic stroke. In ischemic stroke, cell death occurs via a poly(ADP-ribose) polymerase-1 (PARP)-dependent pathway leading to AIF release from the mitochondria and *Parthanatos*^{18,19}. By contrast, recent data suggest that cell death following ICH occurs via a ferroptotic pathway that also involves necroptosis^{7,20}. Of note, inhibitors of *Parthanatos* have no effect on ICH-induced death *in vitro*⁷, suggesting that the pathways of cell damage and, by extension, regenerative failure might be different.

Other studies have focused on the role of the immune system in secondary injury and repair. After ICH, there is a marked inflammatory reaction with activation of microglia and recruitment of leukocytes to the perihematomal region, leading to the production of inflammatory mediators, including cytokines, chemokines, and matrix metalloproteinases^{9,21}. Interestingly, activated microglia can take on pro-inflammatory (via classic activation) or anti-inflammatory (via alternative activation) phenotypes, the latter associated with wound healing and repair^{22,23}. A recent study demonstrated that, after ICH, microglia have dynamic changes in gene expression that result in a transition from an early, pro-inflammatory phenotype to an anti-inflammatory phenotype during the resolution phase of ICH²⁴. Transforming growth factor-beta 1 (TGF-β1) was identified as the most likely mediator of this effect, and treatment with TGF-β1 improved functional outcomes in a murine model. Furthermore, early increases in TGF-β1 levels were also shown to be highly associated with improved 90-day outcomes in patients with ICH.

Another area of research involves the transcription factor peroxisome proliferator-activated receptor gamma (PPAR γ), which plays a significant role in phagocytosis and modulating inflammation and neuroprotective mechanisms³. In a preclinical study, treatment with a PPAR γ agonist, rosiglitazone, promoted hematoma resolution, reduced neuronal damage, downregulated the expression of pro-inflammatory genes (tumor necrosis factor- α , interleukin-1 β , matrix metalloproteinase-9, and inducible nitric oxide synthase mRNA), reduced oxidative stress, and was associated with improved neurological function²⁵.

Recovery after intracerebral hemorrhage

Post-stroke recovery has been widely studied, but the majority of research has focused only on ischemic stroke^{26–29}. Given the differences in pathophysiology between ischemic stroke and ICH, one could assume that the recovery outcomes or mechanisms would be dissimilar between the two stroke subtypes. Comparisons between recovery outcomes in patients with ischemic stroke and ICH have yielded mixed results; some have found comparable activity limitation and recovery³⁰, whereas others have found greater recovery after ICH³¹.

The majority of recovery after ICH occurs early, within the first few months post-stroke³². A recent longitudinal study of patients with ICH characterized the time course of recovery of motor and sensory impairment and ambulation in 11 patients up to six months post-stroke³³. Sensory, truncal, and lower limb motor impairment reached a plateau after three months, whereas there was continued improvement of upper limb movement and ambulation to six months. The course of motor recovery has been shown to depend on the integrity of the corticospinal tract (CST), as measured by transcranial magnetic stimulation³⁴ or diffusion tensor tractography³². In a study of 36 patients with putaminal hemorrhage, motor function recovered in all patients up to four months post-stroke, but patients who had preserved CST integrity at baseline had greater recovery of motor function³². More studies are needed to better elucidate the natural history of recovery after ICH, as has been done for ischemic stroke.

Preclinical studies of rehabilitation after intracerebral hemorrhage

Rehabilitation has been shown to lead to functional improvements in rodent models using both collagenase-induced and whole blood-induced ICH^{13,35,36}. Different types of rehabilitation have been tested in preclinical models, including environmental enrichment^{37,38}, skilled reach training³⁷, constraint-induced movement therapy^{39,40}, and aerobic training⁴¹. Enriched rehabilitation (ER) combines environmental enrichment (group housing and access to tunnels, ramps, and various toys) and task-specific training (most commonly skilled reach training) and was first used as a successful rehabilitative intervention in preclinical models of ischemic stroke^{42,43}. Rehabilitation has been shown to be associated with improved behavioral recovery and enhanced neuroplasticity, including dendritic reorganization⁴⁴, astrocytic plasticity⁴⁵, and synaptogenesis in ipsilesional motor cortex and striatum⁴⁶.

In a collagenase-induced rodent model, rats that participated in a two-week course of enriched rehabilitation beginning one week after ICH improved their skilled reaching and walking ability³⁶. ER also significantly reduced the amount of perihematomal neuronal death even after one week of treatment, suggesting a possible neuroprotective effect of rehabilitation. This study also examined whether rehabilitation influenced iron toxicity and inflammation post-ICH; rehabilitation did not have an effect on levels of iron-binding proteins (ferritin and transferrin) or number of inflammatory cells in perihematomal tissue. The results of a follow-up study also support a neuroprotective role of rehabilitation; ER beginning one week after ICH was demonstrated to augment the clearance of toxic blood components, Hgb and iron, and reduce oxidative stress at the hematoma/perihematomal interface³⁵. The authors speculate that rehabilitation upregulates pathways for Hgb clearance (for example, Nrf2⁴⁷) and downregulates anticlearance pathways (for example, CD47⁴⁸) to expedite hematoma resolution and limit secondary injury. Further investigation is needed into a possible neuroprotective role of rehabilitation in preclinical models, as well as the extent to which secondary injury processes contribute to the functional deficits after ICH in humans, and the ability to reduce secondary injury with rehabilitation in humans.

In a whole-blood injection model, ER beginning one week after ICH also led to significant improvements in reaching ability¹³. Interestingly, the behavioral improvements in the whole-blood model, unlike those in the collagenase model, were not accompanied by a decrease in lesion volume or increase in dendritic length¹³. The authors propose that the lack of delayed injury in the whole-blood model compared with the collagenase model may account for these differences. It is still unknown which of these models better exemplifies what occurs after ICH in humans.

Taken together, evidence from preclinical studies shows a benefit of early rehabilitation after ICH, but the mechanisms underlying the behavioral gains are incompletely understood and vary according to model. Further refinement of neurorehabilitative interventions in preclinical models, which focuses on factors such as the timing, intensity, schedule, and total dose of therapy, is also needed to guide the development of optimal rehabilitation paradigms for patients⁴⁹.

Rehabilitation may be more effective after hemorrhagic stroke than after ischemic stroke, when matched for baseline clinical severity. In a preclinical study in size- and location-matched ischemic or hemorrhagic stroke, rodents with ICH were shown to have greater recovery of skilled walking ability than those with ischemic stroke⁵⁰. Similar results were seen in ICH patients who received the same type and dose of inpatient rehabilitation and had greater recovery of neurological impairment and reduced activity limitation by time of discharge, relative to patients with ischemic stroke³¹.

Clinical rehabilitation after intracerebral hemorrhage

There have been very few rehabilitation studies undertaken in patients with ICH. Many of the larger neurorehabilitation trials

have included both hemorrhagic and ischemic stroke patients^{51,52} or excluded ICH patients altogether from study⁵³. In animal models of ischemic stroke, there is an early 'sensitive period' after stroke, during which rehabilitation leads to larger behavioral improvements than when rehabilitation is delayed, even by one week^{42,54}. However, owing to the paucity of clinical trials, evidence is lacking for a 'sensitive period' in humans for ischemic or hemorrhagic stroke.

The effects of early rehabilitation after ICH were recently investigated in a multicenter, prospective, randomized controlled study of 243 patients with ICH in China⁵⁵. Patients with moderate motor impairment were randomly assigned to either very early rehabilitation (within 48 hours) or standard care (rehabilitation starting after seven days). The very early rehabilitation group had shorter lengths of stay, improved quality of life, greater independence in activities of daily living, and lower six-month mortality when compared with the standard-care group. Because this study was conducted in China, where rehabilitation is delivered by family members and is less standardized, it is unclear whether the results are generalizable to the general ICH population. Furthermore, information on known predictors of outcome, such as hematoma volume, was not available, and the outcomes were self-reported and subject to responder bias. Nevertheless, it is a promising result in favor of very early rehabilitation, which has been a topic of recent debate⁵⁶.

Most clinical guidelines support early initiation of rehabilitation after stroke⁵⁷, and multiple studies have shown that early therapy is safe and feasible^{58,59}. However, the results of the recently published A Very Early Rehabilitation Trial (AVERT) indicate that very early, intensive mobilization may be detrimental after stroke⁵². In this single-blinded, randomized controlled trial of 2,104 patients with ischemic or hemorrhagic stroke, patients who received very early mobilization (< 24 hours post-stroke with more frequent sessions) were more likely to have an unfavorable

outcome than patients who received usual care⁵². In a pre-specified subgroup analysis, this effect was stronger in patients with severe stroke and ICH. Furthermore, providing early rehabilitation for patients with ICH is particularly challenging because these patients are often admitted to intensive care units and require close neurological and hemodynamic monitoring. One must ensure that rehabilitation interventions do not cause detrimental fluctuations in blood pressure and intracranial pressure that can lead to hematoma expansion, which is associated with worsened functional outcomes after ICH^{9,60}.

Conclusions and future directions

Reducing impairment and disability after ICH requires a multifaceted approach with advances in the acute medical and surgical treatment and rehabilitation of patients with ICH. Rehabilitation leads to significant behavioral improvements in animal models of ICH and is associated with enhanced neuroplasticity and reduced neuronal degeneration, although the mechanisms contributing to behavioral recovery are not yet understood. Clinical rehabilitation studies in patients with ICH have been limited; more studies are needed to characterize the natural history of recovery after ICH and to examine the effect of rehabilitative interventions specifically in this patient population. Therapies to enhance neuroprotection and neuroplasticity are also being developed and may be used in conjunction with rehabilitation to promote recovery.

Competing interests

The authors declare that they have no competing interests.

Grant information

This article was funded by an unrestricted gift from John K. Castle (TK).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- Sacco S, Marini C, Toni D, *et al.*: **Incidence and 10-year survival of intracerebral hemorrhage in a population-based registry.** *Stroke*. 2009; **40**(2): 394–9. [PubMed Abstract](#) | [Publisher Full Text](#)
- Adeoye O, Broderick JP: **Advances in the management of intracerebral hemorrhage.** *Nat Rev Neurol*. 2010; **6**(11): 593–601. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Sonni S, Lioutas VA, Selim MH: **New avenues for treatment of intracranial hemorrhage.** *Curr Treat Options Cardiovasc Med*. 2014; **16**(1): 277. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Wu J, Hua Y, Keep RF, *et al.*: **Oxidative brain injury from extravasated erythrocytes after intracerebral hemorrhage.** *Brain Res*. 2002; **953**(1–2): 45–52. [PubMed Abstract](#) | [Publisher Full Text](#)
- Wang YC, Zhou Y, Fang H, *et al.*: **Toll-like receptor 2/4 heterodimer mediates inflammatory injury in intracerebral hemorrhage.** *Ann Neurol*. 2014; **75**(6): 876–89. [PubMed Abstract](#) | [Publisher Full Text](#)
- Huang FP, Xi G, Keep RF, *et al.*: **Brain edema after experimental intracerebral hemorrhage: role of hemoglobin degradation products.** *J Neurosurg*. 2002; **96**(2): 287–93. [PubMed Abstract](#) | [Publisher Full Text](#)
- Zille M, Karuppagounder SS, Chen Y, *et al.*: **Neuronal Death After Hemorrhagic Stroke In Vitro and In Vivo Shares Features of Ferroptosis and Necroptosis.** *Stroke*. 2017; **48**(4): 1033–43. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Xi G, Reiser G, Keep RF: **The role of thrombin and thrombin receptors in ischemic, hemorrhagic and traumatic brain injury: deleterious or protective?** *J Neurochem*. 2003; **84**(1): 3–9. [PubMed Abstract](#) | [Publisher Full Text](#)
- Keep RF, Hua Y, Xi G: **Intracerebral haemorrhage: mechanisms of injury and therapeutic targets.** *Lancet Neurol*. 2012; **11**(8): 720–31. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- MacLellan CL, Silasi G, Auriat AM, *et al.*: **Rodent models of intracerebral hemorrhage.** *Stroke*. 2010; **41**(10 Suppl): S95–8. [PubMed Abstract](#) | [Publisher Full Text](#)
- Bullock R, Mendelow AD, Teasdale GM, *et al.*: **Intracranial haemorrhage induced at arterial pressure in the rat. Part 1: Description of technique, ICP changes and neuropathological findings.** *Neurol Res*. 1984; **6**(4): 184–8. [PubMed Abstract](#) | [Publisher Full Text](#)
- Rosenberg GA, Mun-Bryce S, Wesley M, *et al.*: **Collagenase-induced intracerebral hemorrhage in rats.** *Stroke*. 1990; **21**(5): 801–7. [PubMed Abstract](#) | [Publisher Full Text](#)

13. MacLellan CL, Plummer N, Silasi G, *et al.*: **Rehabilitation promotes recovery after whole blood-induced intracerebral hemorrhage in rats.** *Neurorehabil Neural Repair.* 2011; 25(5): 477–83.
[PubMed Abstract](#) | [Publisher Full Text](#)
14. MacLellan CL, Silasi G, Poon CC, *et al.*: **Intracerebral hemorrhage models in rat: comparing collagenase to blood infusion.** *J Cereb Blood Flow Metab.* 2008; 28(3): 516–25.
[PubMed Abstract](#) | [Publisher Full Text](#)
15. Gu Y, Hua Y, Keep RF, *et al.*: **Deferoxamine reduces intracerebral hematoma-induced iron accumulation and neuronal death in piglets.** *Stroke.* 2009; 40(6): 2241–3.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
16. Okauchi M, Hua Y, Keep RF, *et al.*: **Deferoxamine treatment for intracerebral hemorrhage in aged rats: therapeutic time window and optimal duration.** *Stroke.* 2010; 41(2): 375–82.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
17. Karuppagounder SS, Alim I, Khim SJ, *et al.*: **Therapeutic targeting of oxygen-sensing prolyl hydroxylases abrogates ATF4-dependent neuronal death and improves outcomes after brain hemorrhage in several rodent models.** *Sci Transl Med.* 2016; 8(328): 328ra29.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
18. Andrabi SA, Dawson TM, Dawson VL: **Mitochondrial and nuclear cross talk in cell death: parthanatos.** *Ann N Y Acad Sci.* 2008; 1147: 233–41.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
19. Endres M, Wang ZQ, Namura S, *et al.*: **Ischemic brain injury is mediated by the activation of poly(ADP-ribose)polymerase.** *J Cereb Blood Flow Metab.* 1997; 17(11): 1143–51.
[PubMed Abstract](#) | [Publisher Full Text](#)
20. **F** Li Q, Han X, Lan X, *et al.*: **Inhibition of neuronal ferroptosis protects hemorrhagic brain.** *JCI Insight.* 2017; 2(7): e90777.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
21. Xue M, Del Bigio MR: **Intracerebral injection of autologous whole blood in rats: time course of inflammation and cell death.** *Neurosci Lett.* 2000; 283(3): 230–2.
[PubMed Abstract](#) | [Publisher Full Text](#)
22. Cherry JD, Olschowka JA, O'Banion MK: **Neuroinflammation and M2 microglia: the good, the bad, and the inflamed.** *J Neuroinflammation.* 2014; 11: 98.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
23. Gordon S: **Alternative activation of macrophages.** *Nat Rev Immunol.* 2003; 3(1): 23–35.
[PubMed Abstract](#) | [Publisher Full Text](#)
24. **F** Taylor RA, Chang CF, Goods BA, *et al.*: **TGF- β 1 modulates microglial phenotype and promotes recovery after intracerebral hemorrhage.** *J Clin Invest.* 2017; 127(1): 280–92.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
25. Zhao X, Sun G, Zhang J, *et al.*: **Hematoma resolution as a target for intracerebral hemorrhage treatment: role for peroxisome proliferator-activated receptor gamma in microglia/macrophages.** *Ann Neurol.* 2007; 61(4): 352–62.
[PubMed Abstract](#) | [Publisher Full Text](#)
26. **F** Cortes JC, Goldsmith J, Harran MD, *et al.*: **A Short and Distinct Time Window for Recovery of Arm Motor Control Early After Stroke Revealed With a Global Measure of Trajectory Kinematics.** *Neurorehabil Neural Repair.* 2017; 31(6): 552–60.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
27. Kwakkel G, Kollen B, Twisk J: **Impact of time on improvement of outcome after stroke.** *Stroke.* 2006; 37(9): 2348–53.
[PubMed Abstract](#) | [Publisher Full Text](#)
28. **F** Winters C, van Wegen EE, Daffertshofer A, *et al.*: **Generalizability of the Proportional Recovery Model for the Upper Extremity After an Ischemic Stroke.** *Neurorehabil Neural Repair.* 2015; 29(7): 614–22.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
29. **F** Byblow WD, Stinear CM, Barber PA, *et al.*: **Proportional recovery after stroke depends on corticomotor integrity.** *Ann Neurol.* 2015; 78(6): 848–59.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
30. Jørgensen HS, Nakayama H, Raaschou HO, *et al.*: **Intracerebral hemorrhage versus infarction: stroke severity, risk factors, and prognosis.** *Ann Neurol.* 1995; 38(1): 45–50.
[PubMed Abstract](#) | [Publisher Full Text](#)
31. Paolucci S, Antonucci G, Grasso MG, *et al.*: **Functional outcome of ischemic and hemorrhagic stroke patients after inpatient rehabilitation: a matched comparison.** *Stroke.* 2003; 34(12): 2861–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
32. **F** Jang SH, Park JW, Choi BY, *et al.*: **Difference of recovery course of motor weakness according to state of corticospinal tract in putaminal hemorrhage.** *Neurosci Lett.* 2017; 653: 163–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
33. **F** Lee KB, Kim JS, Hong BY, *et al.*: **The motor recovery related with brain lesion in patients with intracranial hemorrhage.** *Behav Neurol.* 2015; 2015: 258161.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
34. **F** Stinear CM, Byblow WD, Ackerley SJ, *et al.*: **Proportional Motor Recovery After Stroke: Implications for Trial Design.** *Stroke.* 2017; 48(3): 795–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
35. **F** Williamson MR, Dietrich K, Hackett MJ, *et al.*: **Rehabilitation Augments Hematoma Clearance and Attenuates Oxidative Injury and Ion Dyshomeostasis After Brain Hemorrhage.** *Stroke.* 2017; 48(1): 195–203.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
36. Caliaperumal J, Colbourne F: **Rehabilitation improves behavioral recovery and lessens cell death without affecting iron, ferritin, transferrin, or inflammation after intracerebral hemorrhage in rats.** *Neurorehabil Neural Repair.* 2014; 28(4): 395–404.
[PubMed Abstract](#) | [Publisher Full Text](#)
37. Auriat AM, Colbourne F: **Delayed rehabilitation lessens brain injury and improves recovery after intracerebral hemorrhage in rats.** *Brain Res.* 2009; 1251: 262–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
38. Auriat AM, Wowk S, Colbourne F: **Rehabilitation after intracerebral hemorrhage in rats improves recovery with enhanced dendritic complexity but no effect on cell proliferation.** *Behav Brain Res.* 2010; 214(1): 42–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
39. Ishida A, Tamakoshi K, Hamakawa M, *et al.*: **Early onset of forced impaired forelimb use causes recovery of forelimb skilled motor function but no effect on gross sensory-motor function after capsular hemorrhage in rats.** *Behav Brain Res.* 2011; 225(1): 126–34.
[PubMed Abstract](#) | [Publisher Full Text](#)
40. DeBow SB, Davies ML, Clarke HL, *et al.*: **Constraint-induced movement therapy and rehabilitation exercises lessen motor deficits and volume of brain injury after striatal hemorrhagic stroke in rats.** *Stroke.* 2003; 34(4): 1021–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
41. Takamatsu Y, Ishida A, Hamakawa M, *et al.*: **Treadmill running improves motor function and alters dendritic morphology in the striatum after collagenase-induced intracerebral hemorrhage in rats.** *Brain Res.* 2010; 1355: 165–73.
[PubMed Abstract](#) | [Publisher Full Text](#)
42. **F** Biernaskie J, Chernenko G, Corbett D: **Efficacy of rehabilitative experience declines with time after focal ischemic brain injury.** *J Neurosci.* 2004; 24(5): 1245–54.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
43. Biernaskie J, Corbett D: **Enriched rehabilitative training promotes improved forelimb motor function and enhanced dendritic growth after focal ischemic injury.** *J Neurosci.* 2001; 21(14): 5272–80.
[PubMed Abstract](#)
44. Santos MV, Pagnussat AS, Mestriner RG, *et al.*: **Motor Skill Training Promotes Sensorimotor Recovery and Increases Microtubule-Associated Protein-2 (MAP-2) Immunoreactivity in the Motor Cortex after Intracerebral Hemorrhage in the Rat.** *ISRN Neurol.* 2013; 2013: 159184.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
45. Mestriner RG, Pagnussat AS, Boisserand LS, *et al.*: **Skilled reaching training promotes astroglial changes and facilitated sensorimotor recovery after collagenase-induced intracerebral hemorrhage.** *Exp Neurol.* 2011; 227(1): 53–61.
[PubMed Abstract](#) | [Publisher Full Text](#)
46. Tamakoshi K, Ishida A, Takamatsu Y, *et al.*: **Motor skills training promotes motor functional recovery and induces synaptogenesis in the motor cortex and striatum after intracerebral hemorrhage in rats.** *Behav Brain Res.* 2014; 260: 34–43.
[PubMed Abstract](#) | [Publisher Full Text](#)
47. Zhao X, Sun G, Ting SM, *et al.*: **Cleaning up after ICH: the role of Nrf2 in modulating microglia function and hematoma clearance.** *J Neurochem.* 2015; 133(1): 144–52.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
48. Cao S, Zheng M, Hua Y, *et al.*: **Hematoma Changes During Clot Resolution After Experimental Intracerebral Hemorrhage.** *Stroke.* 2016; 47(6): 1626–31.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
49. **F** Saulle MF, Schambra HM: **Recovery and Rehabilitation after Intracerebral Hemorrhage.** *Semin Neurol.* 2016; 36(3): 306–12.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
50. Mestriner RG, Miguel PM, Bagatini PB, *et al.*: **Behavior outcome after ischemic and hemorrhagic stroke, with similar brain damage, in rats.** *Behav Brain Res.* 2013; 244: 82–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
51. **F** Lo AC, Guarino PD, Richards LG, *et al.*: **Robot-assisted therapy for long-term upper-limb impairment after stroke.** *N Engl J Med.* 2010; 362(19): 1772–83.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
52. **F** AVERT Trial Collaboration group: **Efficacy and safety of very early mobilisation within 24 h of stroke onset (AVERT): a randomised controlled trial.** *Lancet.* 2015; 386(9988): 46–55.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
53. Kwakkel G, Winters C, van Wegen EE, *et al.*: **Effects of Unilateral Upper Limb Training in Two Distinct Prognostic Groups Early After Stroke: The EXPLICIT-Stroke Randomized Clinical Trial.** *Neurorehabil Neural Repair.* 2016; 30(9): 804–16.
[PubMed Abstract](#) | [Publisher Full Text](#)
54. **F** Zeiler SR, Gibson EM, Hoesch RE, *et al.*: **Medial premotor cortex shows a**

- reduction in inhibitory markers and mediates recovery in a mouse model of focal stroke. *Stroke*. 2013; 44(2): 483–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
55. Liu N, Cadilhac DA, Andrew NE, *et al.*: **Randomized controlled trial of early rehabilitation after intracerebral hemorrhage stroke: difference in outcomes within 6 months of stroke.** *Stroke*. 2014; 45(12): 3502–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
56. Luft AR, Kesselring J: **Critique of A Very Early Rehabilitation Trial (AVERT).** *Stroke*. 2016; 47(1): 291–2.
[PubMed Abstract](#) | [Publisher Full Text](#)
57. Hemphill JC 3rd, Greenberg SM, Anderson CS, *et al.*: **Guidelines for the Management of Spontaneous Intracerebral Hemorrhage: A Guideline for Healthcare Professionals From the American Heart Association/American Stroke Association.** *Stroke*. 2015; 46(7): 2032–60.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
58. Bernhardt J, Godecke E, Johnson L, *et al.*: **Early rehabilitation after stroke.** *Curr Opin Neurol*. 2017; 30(1): 48–54.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
59. Yelnik AP, Quintaine V, Andriantsifanetra C, *et al.*: **AMOBES (Active Mobility Very Early After Stroke): A Randomized Controlled Trial.** *Stroke*. 2017; 48(2): 400–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
60. Dowlathshahi D, Demchuk AM, Flaherty ML, *et al.*: **Defining hematoma expansion in intracerebral hemorrhage: relationship with patient outcomes.** *Neurology*. 2011; 76(14): 1238–44.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. Klaus Van Leyen

Neuroprotection Research Laboratory, Massachusetts General Hospital, Harvard Medical School, Boston, Massachusetts, USA

Competing Interests: No competing interests were disclosed.

2. Craig S. Anderson

¹ Neurology Department, Royal Prince Alfred Hospital, Sydney, New South Wales, Australia

² The George Institute for Global Health, Peking University Health Science Center, Beijing, China

³ Department of Neurology and Epidemiology, Faculty of Medicine, University of New South Wales, Sydney, Australia

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research