REVIEW

Update on ventilator-associated pneumonia [version 1; peer review: 2 approved]

Jean-Francois Timsit ^{1,2}, Wafa Esaied¹, Mathilde Neuville², Lila Bouadma^{1,2}, Bruno Mourvillier^{1,2}

¹IAME, Inserm U1137, Paris Diderot University, Paris, F75018, France

²Medical and Infectious Diseases Intensive Care Unit, AP-HP, Bichat University Hospital, Paris, France

V1 First published: 29 Nov 2017, 6(F1000 Faculty Rev):2061
<https://doi.org/10.12688/f1000research.12222.1>

Latest published: 29 Nov 2017, 6(F1000 Faculty Rev):2061
<https://doi.org/10.12688/f1000research.12222.1>

Abstract

Ventilator-associated pneumonia (VAP) is the most frequent life-threatening nosocomial infection in intensive care units. The diagnostic is difficult because radiological and clinical signs are inaccurate and could be associated with various respiratory diseases. The concept of infection-related ventilator-associated complication has been proposed as a surrogate of VAP to be used as a benchmark indicator of quality of care. Indeed, bundles of prevention measures are effective in decreasing the VAP rate. In case of VAP suspicion, respiratory secretions must be collected for bacteriological secretions before any new antimicrobials. Quantitative distal bacteriological exams may be preferable for a more reliable diagnosis and therefore a more appropriate use antimicrobials. To improve the prognosis, the treatment should be adequate as soon as possible but should avoid unnecessary broad-spectrum antimicrobials to limit antibiotic selection pressure. For empiric treatments, the selection of antimicrobials should consider the local prevalence of microorganisms along with their associated susceptibility profiles. Critically ill patients require high dosages of antimicrobials and more specifically continuous or prolonged infusions for beta-lactams. After patient stabilization, antimicrobials should be maintained for 7–8 days. The evaluation of VAP treatment based on 28-day mortality is being challenged by regulatory agencies, which are working on alternative surrogate endpoints and on trial design optimization.

Keywords

ventilator-associated pneumonia, VAP, nosocomial infection, antimicrobials

Open Peer Review

Approval Status

	1	2
version 1		
29 Nov 2017		

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Saad Nseir**, Lille Univ Hosp, Lille, France
2. **Michael Klompas**, Harvard Medical School and Harvard Pilgrim Health Care Institute, Boston, USA
Brigham and Women's Hospital, Boston, USA

Any comments on the article can be found at the end of the article.

Corresponding author: Jean-Francois Timsit (jean-francois.timsit@bch.aphp.fr)

Author roles: **Timsit JF:** Conceptualization, Supervision, Validation, Writing – Original Draft Preparation, Writing – Review & Editing; **Esaied W:** Resources, Writing – Original Draft Preparation; **Neuville M:** Writing – Original Draft Preparation, Writing – Review & Editing; **Bouadma L:** Writing – Original Draft Preparation, Writing – Review & Editing; **Mourvillier B:** Supervision, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: J-FT gave lectures for Merck, Pfizer, and Gilead on antimicrobial therapy and participated in scientific boards of Paratek, Bayer, Merck, and 3M. The other authors declare that they have no competing interests.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2017 Timsit JF *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Timsit JF, Esaied W, Neuville M *et al.* **Update on ventilator-associated pneumonia [version 1; peer review: 2 approved]** F1000Research 2017, 6(F1000 Faculty Rev):2061 <https://doi.org/10.12688/f1000research.12222.1>

First published: 29 Nov 2017, 6(F1000 Faculty Rev):2061 <https://doi.org/10.12688/f1000research.12222.1>

Introduction

Hospital-acquired pneumonia (HAP) is defined by an infection of the lung parenchyma that occurred at least 48 hours after hospital admission. Ventilator-associated pneumonia (VAP) develops in intensive care unit (ICU) patients mechanically ventilated for at least 48 hours^{1,2}. In contrast, ventilator-associated tracheobronchitis (VAT) is characterized by signs of respiratory infection without new radiographic infiltrates in a patient mechanically ventilated for at least 48 hours^{3–5}.

In the past 10 years, a great deal of progress has been made in understanding VAP. New concepts of infection-related ventilator-associated complications (IVACs) and ventilator-associated events (VAEs) have been proposed as outcome indicators for prevention strategies⁶. In diagnostic strategies, criteria used to suspect a VAP have been challenged, as have optimal diagnostic tests used to confirm it⁷. Traditional risk factors of VAP due to multidrug-resistant (MDR) bacteria (based on early-onset occurrence and previous antimicrobial therapy) are no longer sufficient. Proposed empirical therapy has been modified accordingly. The optimization of pharmacokinetic/pharmacodynamic parameters is now considered a key factor to ensure adequate and successful therapy. The use of adjunctive aerosolized therapy is also more and more debated. In addition, regulatory agencies are trying to find surrogate endpoints to replace 28-day mortality and to improve the design of randomized clinical trials in this field of investigation⁸.

For VAP prevention, the concept of bundle of care was defined. It enabled great successes in VAP prevention; however, the insufficient compliance observed in clinical practice needs to be addressed in order to define easier-to-apply procedures.

This review aims to summarize the available knowledge on VAP, taking profit from the recent publication of North American⁷ and European guidelines on VAP management and highlighting recent advances and remaining controversies of the new concepts.

Epidemiology

VAP is the second most common nosocomial infection and the leading cause of death from nosocomial infections in critically ill patients⁹. Its incidence ranges from 5% to 67% depending on case mix and the diagnostic criteria used¹⁰, and the highest rates are in immunocompromised, surgical, and elderly patients. In the US, the incidence of VAP ranges from 2 to 16 episodes per 1,000 ventilator-days¹¹. The estimated risk of VAP is 1.5% per day and decreases to less than 0.5% per day after the 14th day of mechanical ventilation¹². VAP increases the duration of hospitalization by 7 days and health-care costs by approximately \$40,000 USD¹³.

In published studies, the crude mortality of patients with VAP is highly variable according to case mix and definitions used^{14–16}. The definition of attributable VAP mortality is the percentage of deaths that would not have occurred in the absence of the infection. Recent studies have reappraised the impact of VAP on mortality^{17–19}. Specifically, given that the risk of VAP is time-dependent, this could potentially result in a significant time-dependent bias because mortality and ICU discharge both act as competing endpoints. Indeed, the most recent studies reported an

attributable mortality below 10% with surgical patients¹⁸ whereas those with mid-range illness severity presented the highest associated risk^{18,19}.

Late-onset VAP is often reported to be associated with higher mortality rates than early-onset VAP^{20–22}. Using a multistate model, we confirmed that the attributable mortality for early-onset VAP (5.8%) was considerably lower than for late-onset VAP (10.6%)¹⁸.

Most studies showed that VAP is usually due to aerobic *Enterobacteriaceae* (25%), *Staphylococcus aureus* (20%), *Pseudomonas aeruginosa* (20%), *Haemophilus influenza* (10%), and streptococci²³. MDR pathogens are more common among late-onset cases. Trouillet *et al.*²⁴ found that prior use of broad-spectrum antibiotics and mechanical ventilation of more than 7 days were independent risk factors of infection caused by MDR pathogens. However, more recent reports^{25–29} have identified similar rates of etiologies in patients with early- versus late-onset VAP. This may be related to the worldwide rise in MDR pathogens; it emphasizes that the local ICU ecology³⁰ is the most important risk factor for acquiring MDR pathogens, irrespective of the length of intubation. In early-onset pneumonia, the initial VAP severity—that is, the presence of sepsis or septic shock (odds ratio [OR] = 3.7)—and pneumonia that developed in a center with a prevalence of resistant pathogens greater than 25% were independently associated with the presence of resistant pathogens (OR = 11.3)²⁶.

Risk factors of ventilator-associated pneumonia

VAP results from the microbial invasion of the normally sterile lower respiratory tract, which subsequently can overwhelm the host's defense and establish infection. The major route for microbial invasion is microaspiration of oropharyngeal secretions contaminated by endogenous flora around the endotracheal tube cuff³¹. VAP may also occur by other means^{2,32}. In terms of potential reservoirs, it has been suggested that the stomach hosts bacteria that colonize the oropharynx. It has been postulated by some researchers that embolization into the alveoli during suctioning or bronchoscopy is caused by the colonization of the endotracheal tube with bacteria encased in a biofilm³³. Inhalation of pathogens from contaminated aerosols and direct inoculation are less common, and hematogenous spread from either infected intravascular catheters or bacterial translocation of the gastrointestinal tract lumen are rarer in occurrence.

Consequently, two groups of risk factors for VAP have been identified—namely ventilation-related factors (instrumentation of the airway with an endotracheal tube and subsequent microaspirations) and, less frequently, patient-related factors (for example, pre-existing pulmonary disease)—and only the former is accessible to prevention (Table 1). As a result, VAP, unlike many other nosocomial infections, is difficult to prevent³⁴.

Prevention

First of all, reducing the exposure to risk factors for VAP is the most efficient way to prevent VAP onset (Figure 1). Therefore, intubation should be avoided whenever possible, and strategies

Table 1. Risk factors of ventilator-associated pneumonia.

Host-related risk factors	Intervention-related risk factors
Medical history and underlying illness	Peri-operative transfusion of blood products
Male gender	Duration of the mechanical ventilation
Extreme age	Reintubation
Prior central nervous system disorder	Supine head position in patients receiving enteral nutrition
Immunocompromised	Antibiotic therapy ^a
Acute underlying diseases	Enteral nutrition
Emergent surgery	Absence of subglottic secretion drainage ^b
Neurosurgery	Intra-hospital transports
Thoracic surgery	Continuous sedation, use of paralytic agents
Cardiac surgery	Nasogastric tubes
Burns	Tracheostomy
Re-intervention	Frequent ventilator circuit changes
Acute severity factors	Intracuff pressure of less than 20 cm H ₂ O
Organ system failure index of at least 3	
Acute renal failure	
Acute respiratory distress syndrome	
ECMO, intra-aortic support	
Ulcer disease	

Adapted from 2,35–38. ^aAntibiotic therapy protects from early-onset pneumonia due to susceptible bacteria but is a risk factor for late-onset pneumonia due to more resistant organisms. ^bProtective impact of subglottic secretion drainage is mainly demonstrated for cardiac surgery patients. ECMO, extra-corporeal membrane oxygenation.

Figure 1. Preventive measures of ventilator-associated pneumonia. Adapted from 37,39. QOE, quality of evidence.

such as non-invasive positive-pressure ventilation, sedation, and weaning protocols should be used to replace or shorten mechanical ventilation. In contrast, recent data suggest that the timing of the tracheotomy does not significantly change VAP incidence^{40–43}.

Patients at risk of VAP must be managed with a “bundle of preventive measures” (Figure 1). Indeed, no single preventive strategy will efficiently prevent VAP. Bundles group together a small straightforward set of key interventions that are from evidence-based guidelines—generally three to five—and that are expected to result in a better outcome when performed collectively and reliably instead of individually. However, the ideal set of key preventive measures is unknown⁴⁴. Importantly, although studies demonstrated great success in reducing VAP rates using bundle of care in recent years^{44–48}, meta-analyses showed that most of the preventive measures failed to demonstrate a sustained effect⁴⁹. This conclusion is in line with the absence of substantial improvement of VAP rates in the past decade⁵⁰. Figure 1 lists recommended preventive measures according to their level of evidence.

The sole preventive measures that positively impacted mortality are selective digestive decontamination (SDD) and selective oropharyngeal decontamination (SOD)⁵¹. Compared with SOD, SDD was associated with a lower mortality, reduced length of stay, lower rates of ICU-acquired bacteremia and candidemia, and lower prevalence of rectal carriage of antibiotic-resistant Gram-negative bacteria but with a pronounced gradual increase in aminoglycoside-resistant Gram-negative bacteria⁵². The main remaining question is the reproducibility of these results out of the Netherlands. Indeed, the antibiotic selection pressure induced by SOD or SDD may outweigh their benefits in countries with high levels of bacterial resistance.

Oral care with chlorhexidine is also debated. An updated meta-analysis focusing on double-blind studies in non-cardiac surgery patients showed that it had no impact on VAP rates or duration of mechanical ventilation or duration of ICU stay⁵³.

Ecological Effects of Decolonization Strategies in Intensive Care (RGNOSIS), a cluster-randomized study (ClinicalTrials.gov identifier NCT02208154) conducted in six European countries, is currently enrolling 10,800 patients into four arms: control, oral care with chlorhexidine, SOD, and SDD. The study’s new insights into these ongoing debates are awaited.

Many possible factors may explain why prevention measures did not result in reductions in mortality, duration of stay, or antibiotic consumption. First of all, the VAP definition may not be sufficiently accurate, especially when tested intervention could not be blinded. Second, in recent studies using modern statistics, the attributable mortality of VAP is only 3–4%, considerably smaller than previously reported¹⁹. Both factors may induce a dramatic decrease of the power of the studies available.

Even if there is convincing evidence that specific interventions might prevent VAP, translating research into practice remains a challenge (Figure 1). Two European surveys found that 37.0% of ICU physicians⁵⁴ and 22.3% of nurses⁵⁵ did not comply with

the published recommendations for VAP prevention. Beyond the theoretical frame, a great deal of attention must be given to the factors that might facilitate a bundle implementation and allow a sustained compliance. An educational session alone, without an associated behavioral strategy, is unlikely to induce profound behavioral changes. It should be kept in mind that, to engage an individual in a particular behavior and improve compliance, we need to act on predisposing factors (knowledge, perceptions, and beliefs) to favor the access to new processes or technologies and to continually reinforce the behavior by feedback^{56,57}.

Diagnosis

VAP, VAE, IVAC, and VAT: what do these abbreviations mean?

The diagnosis of VAP is traditionally based on clinical symptoms and radiographic criteria that require further bacteriological confirmation. However, it has been demonstrated that these criteria are inaccurate^{14,58,59}. Of note, VAP is now considered an indicator of performance in the US and some other countries. The National Healthcare Society Network reported a considerable decrease in the VAP incidence rate attributed to a multifaceted infection control program and its effective implementation⁶⁰. A 70% decrease of the incidence between 2006 and 2012 was reported by the Centers for Disease Control and Prevention (CDC). But in the same period, the Medicare Patient Safety Monitoring System reported an adjusted average annual change of 0% (95% confidence interval (CI) –0.05 to 0.07) in patients 65 years old or older⁵⁰. These findings emphasize a true discrepancy between rates reported in a quality monitoring program and rates observed in patients’ care program.

The discrepancies between results prompted the CDC to promote new objectives of the surveillance based on VAEs. Another important motivation of the CDC was to expand the purview of quality and safety surveillance to encompass multiple complications in mechanically ventilated patients instead of just pneumonia alone.

The VAE surveillance definition algorithm uses three new indicators: ventilator-associated complications (VACs), IVACs, and possible and probable VAP⁶¹. VAC is the first step of VAE surveillance, with the aim of identifying any complication occurring in mechanically ventilated patients, regardless of the origin or mechanism. To meet the definition of VAC, a mechanically ventilated patient must have at least 2 days of stability or improvement of respiratory parameters—such as a stable or decreasing daily minimum positive end-expiratory pressure (PEEP) or fraction of inspired oxygen (FiO₂)—followed by at least 2 days of worsened oxygenation (diagnosed by an increase of the daily minimum PEEP (at least 3 cm H₂O) or FiO₂ (at least 20%)). The concept of IVAC aims to identify the subgroup of VACs that are potentially related to infection. A VAC associated with an abnormal white blood cell count or a modified temperature becomes an IVAC if the initiation of a new antimicrobial agent is maintained for at least 4 days. With evidence of purulent respiratory secretions or positive results of microbiological tests performed on respiratory tract specimens or both, an IVAC becomes a possible VAP. All of these definitions are summarized in Figure 2.

Figure 2. Ventilator-associated events, definitions, and nosology. Ventilator-associated conditions (VACs): at least 2 calendar days of stable or decreasing daily minimum positive end-expiratory pressure (PEEP) or fraction of inspired oxygen (FiO_2) followed by rise in PEEP of at least 3 cm H_2O or rise in FiO_2 of at least 20 points sustained for at least 2 days. Infection-related ventilator-associated complications (IVACs): VAC plus: temperature of less than 36°C or more than 38°C OR white blood cell (WBC) count of not more than 4 or at least 12×10^3 cells/ mm^3 AND at least one new antibiotics continued for at least 4 days WITHIN 2 days of VAC onset EXCLUDING first 2 days on the ventilator. Possible ventilator-associated pneumonia (VAP) (Centers for Disease Control and Prevention [CDC] definitions): IVAC plus: criterion 1: Positive culture meeting specific quantitative or semi-quantitative threshold; criterion 2: Purulent respiratory secretions AND identification of organisms NOT meeting the quantitative or semi-quantitative thresholds; criterion 3: Organisms identified from pleural fluid specimen, positive lung histopathology, and positive diagnostic test for Legionella species or selected respiratory viruses WITHIN 2 days of VAC onset EXCLUDING first 2 days on the ventilator. (The updated January 2017 definitions and comprehensive examples are detailed in the CDC National Healthcare Society Network website; https://www.cdc.gov/nhsn/pdfs/pscmanual/10-vae_final.pdf; accessed 23 October 2017.) VAP: radiographic criteria (new or progressive and persistent infiltrates or consolidation or cavitation); systemic criteria (temperature of less than 36°C or more than 38°C OR WBC count of not more than 4 or at least 12×10^3 cells/ mm^3); pulmonary criteria (at least one of the following: (1) new onset or increase of purulent aspirates and (2) worsening gas exchange). Ventilator-associated tracheobronchitis (VAT): criteria for VAP but without radiographic criteria.

The VAE concept uses objective criteria and their collection can be automated for systematic recording. The VAP definition is still widely discussed, and a recent study showed that applying the various diagnostic criteria to the same patient population resulted in large differences in the incidence of VAP (that is, from 4% to 42%)⁶². Furthermore, even distal quantitative samples are not 100% reproducible^{63,64}.

This new approach might overcome the inaccuracy of the VAP definition, facilitate its electronic assessment, and make inter-ICU comparisons more relevant. Second, the association between VAE and antibiotic consumption (considering VAC rates and not only IVAC) was a point in favor of using VAC rate as one indicator of ICU quality of care for antimicrobial stewardship programs¹². Of note, VAE has very low sensitivity and specificity in diagnosing VAP¹². In our experience, VAP accounted for only 14.5% of the VAC episodes and 27.6% of the IVAC episodes¹²; in addition, not all IVAC episodes were related to a nosocomial infection.

Furthermore, radiological criteria are not taken into account, so that the IVAC definition includes VAT and VAP (Figure 2). Although VAP and VAT are both associated with an increased duration of mechanical ventilation, VAP impact on ICU mortality is higher than that of VAT²⁷. Finally, embedding VAP in the larger definition

of IVAC may hamper the understanding of VAP pathophysiology and thus its prevention improvement.

Which bacteriological samples should be collected in case of suspicion of ventilator-associated pneumonia?

Great controversies persist about the bacteriological samples that should be used for diagnosing VAP. Of note, when bacteriological analyses are not immediately available, processing of a bacteriological specimen refrigerated after collection is a reliable alternative⁶⁵. Invasive techniques, such as bronchoalveolar lavage or protected specimen brush with quantitative culture, require qualified clinicians. Randomized studies that have evaluated their value as compared with proximal qualitative samples yielded contradictory results^{66–68}. In one study with 413 patients⁶⁷, the invasive distal quantitative strategy was combined with an algorithm for treatment de-escalation and led to a significant increase in the number of antibiotic-free days at day 14 (5.0 ± 5.1 versus 2.2 ± 3.5) and day 28 (11.5 ± 9.0 versus 7.5 ± 7.6) in comparison with the strategy with non-invasive methods using qualitative cultures. In contrast, the Canadian Critical Care Trial Group reported no impact of distal quantitative samples on the day-28 antibiotic-free days or on survival⁶⁶. However, in that study, the research protocol may have facilitated appropriate discontinuation of antibiotics or targeted therapy in the two groups,

thus minimizing the differences between them. Cohort studies confirmed the potential advantages of distal quantitative samples in narrowing antimicrobial therapy and limiting antibiotic selection pressure without adverse effects on mortality or length of stay^{69–71}. Finally, an observational study in 89 patients with clinically suspected VAP and a negative quantitative bronchoalveolar lavage compared patients with early (within one day) and late antibiotic discontinuation. Early discontinuation was associated with a non-significant decrease in mortality and significantly lower risks of overall superinfections (22.5% versus 43%), respiratory superinfections (10% versus 29%), and superinfections due to MDR pathogens (7.5% versus 36%)⁷².

Considering available literature, recent US guidelines recommend non-invasive sampling with semi-quantitative culture⁷, whereas the European guidelines suggest obtaining distal samples with quantitative cultures to improve the accuracy of results⁹. Despite

this discrepancy, the two guidelines agreed that a bacteriological sample should be performed before any antibiotic treatment in order to reduce antibiotic exposure.

Treatment of ventilator-associated pneumonia

The initial treatment of VAP is based on empirical choices; however, an inappropriate initial antibiotic choice is associated with increased mortality^{21,73}. In addition, the recovery of MDR bacteria is clearly associated with an increased risk of inappropriate therapy⁷⁴. As discussed earlier, the risk of MDR is conditioned by the local ecological data, previous colonization, and previous antibiotic therapy received by the patients. The increase in the risk of MDR in late-onset infections is challenged by recent studies. Regimens proposed by the North American guidelines are listed in [Table 2](#)⁷. An algorithm for an empirical therapy strategy combining guidelines and practical rules is proposed in [Figure 3](#).

Table 2. Empirical treatment of hospital-acquired pneumonia/ventilator-associated pneumonia.

Not at high risk of mortality and no risk factors ^a	Not at high risk of mortality but with factors increasing the likelihood of Gram-negative bacteria	High risk of mortality or receipt of intravenous antibiotics during the prior 90 days
One of the following: Piperacillin-tazobactam 4.5 g IV q6h OR Cefepime 2 g IV q8h Levofloxacin 750 mg IV daily	Piperacillin-tazobactam 4.5 g IV q6h OR Cefepime or ceftazidime 2 g IV q8h OR Levofloxacin 750 mg IV daily Ciprofloxacin 400 mg IV q8h OR Imipenem 1g IV q8h Meropenem 1 g IV q6h	Piperacillin-tazobactam 4.5 g IV q6h OR Cefepime or ceftazidime 2 g IV q8h OR Levofloxacin 750 mg IV daily Ciprofloxacin 400 mg IV q8h OR Imipenem 1g IV q8h Meropenem 1 g IV q6h AND Amikacin 25 (30) mg/kg IV daily OR Gentamicin 5–7 mg/kg IV daily OR Tobramycin 5–7 mg/kg IV daily
	Vancomycin 15 mg/kg IV q8–12h with goal to target 15–20 mg/mL trough level (consider a loading dose of 25–30 mg/kg × 1 for severe illness) OR Linezolid 600 mg IV q12h	Vancomycin 15 mg/kg IV q8–12h with goal to target 15–20 mg/mL trough level (consider a loading dose of 25–30 mg/kg × 1 for severe illness) OR Linezolid 600 mg IV q12h

Adapted from Infectious Diseases Society of America/American Thoracic Society guidelines⁷. ^aRisk factors of multidrug-resistant ventilator-associated pneumonia (VAP) are prior intravenous use within 90 days, septic shock at VAP onset, acute respiratory distress syndrome preceding VAP, five or more days of hospitalization prior to VAP onset, and acute renal replacement therapy prior to VAP onset. IV, intravenous; q, every.

Figure 3. Proposed strategy for empirical therapy. *In areas with a risk of multidrug-resistant and carbapenemase-producing bacteria, the empirical choice should be decided on the basis of local ecology. 3rd GC, third-generation cephalosporin; ARDS, acute respiratory distress syndrome; ATB, antibiotics; GNB, Gram-negative bacteria; MDR, multidrug-resistant; MRSA, methicillin-resistant *Staphylococcus aureus*; PA, *Pseudomonas aeruginosa*; PIP/TAZ, piperacillin-tazobactam; R, Resistant; VAP, ventilator-associated pneumonia.

The challenge for the intensivist is to start an antimicrobial therapy that will be immediately effective while avoiding any overuse of extended-spectrum antimicrobials. New rapid diagnostic tests have been developed but their performances for VAP diagnosis remain to be evaluated^{75,76}. Rapid nucleic acid amplification or mass spectrometry-based techniques provide rapid identification of targeted microorganisms. Some of these new tests are also able to detect resistance genes. However, the presence of genes detected by these techniques does not mean that the pathogens are alive or dead, nor does it provide information regarding phenotypic antimicrobial susceptibility. Rapid culture with semi-automated rapid antibiotic susceptibility tests are also in development. Fluorescence *in situ* hybridization-based microscopy identification and antibiotic susceptibility test (ID/AST) systems can evaluate antibiotic susceptibility from respiratory secretions on a previously defined panel of pathogens. A recent pilot study reported promising results: the technique was able to detect pathogens in bronchoalveolar lavage after 5 ± 7 hours of culture and 5 hours of analysis, and sensitivity and specificity were 100% and 97%, respectively⁷⁷. Technical developments with a better selection and quantification of pathogens and resistance patterns are warranted.

Beta-lactams remain a cornerstone antibiotic for the treatment of VAP. Critical care patients exhibit high clearance and distribution volume, which contribute to low blood levels of antimicrobials⁷⁸. Therefore, the doses that should be used to treat the most severe patients are frequently higher than the ones

approved by regulatory agencies⁷⁹. For β -lactams, the best results seem to be associated with β -lactam plasma levels up to four times the minimal inhibitory concentration (MIC) of the involved pathogen and during 100% of the interval between each dose⁸⁰. For these agents, a loading dose followed by a continuous infusion may be a relevant method of administration to increase the antibiotic concentration in the blood and the lung lining fluid with a lower risk of neurological⁸¹ or renal⁸² toxicity. It is of special importance when bacteria are not fully susceptible or when MICs are high, as for *P. aeruginosa* and MDR Gram-negative bacteria⁸³.

Combination therapy with aminoglycosides increases the likelihood to immediately achieve an adequate therapy, especially for infection due to MDR Gram-negative bacteria⁸⁴. It is associated with an improved prognosis in the most severe patients⁸⁵. A dose as high as 25 mg/kg of amikacin is required to reach the optimal 60 mg/L peak concentration, even in the case of renal failure. Indeed, the distribution volume of aminoglycosides is not affected by renal dysfunction⁸⁶. However, renal impairment, present in almost 30% of ICU patients, will lead to prolonged intervals between doses, reducing the actual number of peak levels, thus possibly affecting the treatment efficiency.

Indeed, controversies still exist about advantages and disadvantages of aminoglycosides. Ong *et al.* compared empirical therapy of septic shock with or without gentamicin in two Dutch ICUs⁸⁷. One of the ICUs preferentially used aminoglycosides, whereas the other preferentially avoided them. After careful adjustment

on patients' characteristics, they found that gentamicin is an independent predictor of renal failure without affecting mortality. The main limitations of this study include (1) the absence of random allocation, (2) the absence of control on center effect, (3) the setting in two ICUs where the rate of MDR Gram-negative organisms is very low with a minimal risk of inadequate therapy, and (4) the absence of therapeutic drug monitoring for gentamicin. Nevertheless, the study by Ong *et al.*⁸⁷ emphasizes the importance of new well-conducted studies comparing antimicrobial therapy with or without aminoglycosides, given with careful therapeutic drug monitoring, for the most severe ICU patients.

Combination therapy with fluoroquinolones was associated with a decrease in the number of days alive without relapse or reinfection in a cohort of patients with VAP due to *P. aeruginosa* or *Enterobacteriaceae* and without any prior treatment with fluoroquinolones⁸⁸. However, the fluoroquinolones are inconsistently active against MDR Gram-negative bacteria and their use is associated with an important risk of emergence of MDR bacteria in the lung and gut microbiota⁸⁹.

Concerning extended-spectrum β -lactamase-producing *Enterobacteriaceae* (ESBL-PE), carbapenems remain the first-line agents despite leading to the risk of emergence and spread of carbapenemase-producing *Enterobacteriaceae*⁹⁰. Other options such as piperacillin/tazobactam or a high dose of third-generation cephalosporins administered by continuous infusion could be considered, especially as step-down therapy for ESBL-E with low MICs³⁰. New compounds such as ceftolozane/tazobactam and ceftazidime/avibactam have recently been released; however, against HAP/VAP, only the latter was non-inferior to meropenem in a recent randomized trial (REPROVE NCT01808092; results available on ClinicalTrials.gov, not yet published). Temocillin, a ticarcillin derivative that resists ESBL, can be used but only as a step-down therapy for pathogens with MICs below 8 mg/L⁹¹. For carbapenem-resistant Gram-negative bacteria, colistin is a cornerstone of the treatment, although ceftazidime/avibactam association might be effective. A new association of meropenem and vaborbactam (M-V), recently approved for severe urinary tract infections in the US, has been compared with best available therapy (BAT) in severe infection presumably due to carbapenemase R *Enterobacteriaceae*, including nosocomial pneumonia (TANGO-2 NCT 02168946). The study was presented at the IDWeek convention (in San Diego, CA, in October 2017; abstract 1867) and enrolled 43 patients (more than 80% with *Klebsiella pneumoniae carbapenemase*). It has been stopped prematurely for significant superiority in terms of clinical failure, nephrotoxicity, and non-significant improvement of day-28 mortality of patients with HAP/VAP (M-V 4/16 versus BAT 4/9). Further studies are awaited to confirm these encouraging results and make formal recommendations on its use. An intravenous colistin regimen should be considered with a loading dose of 9 MU and with caution regarding its potential nephrotoxicity⁹².

Inhaled antimicrobial therapy may be considered, as this route of administration enables very high concentrations of antimicrobials to be locally delivered^{93,94}. However, there are no solutions

specifically formulated for inhalation, and a limited number of devices are designed for the nebulization of antibiotics^{95,96}. Of note, despite the possible advantages in terms of microbiological eradication and emergence of resistance⁹⁷, no impact on patient prognosis has been demonstrated^{94,97-99}.

In units with rates of methicillin-resistant *Staphylococcus aureus* (MRSA) around 10–20%, include vancomycin or linezolid in the empirical therapy⁷. When the MIC to vancomycin is higher than 1.5 mg/L, the mortality of MRSA pneumonia is higher¹⁰⁰. Moreover, it is very difficult to reach pharmacokinetic targets using vancomycin¹⁰¹ without any increase in renal toxicity. Consequently, linezolid should be preferred, particularly in patients with renal impairment or if the MRSA MIC to vancomycin is over 1.5 mg/L¹⁰².

An 8-day antibiotic course appears safe in VAP. This duration can be shortened when a procalcitonin-guided algorithm is used¹⁰³ or when ventilator settings (PEEP ≤ 5 cm H₂O and FiO₂ $\leq 40\%$) are stable for 48 hours after antibiotic initiation¹⁰⁴. As procalcitonin levels above 1.5 ng/mL after three days of treatment seemed strongly associated with a poor outcome¹⁰⁵, re-evaluation of the accuracy of diagnosis and a search for drainable collections (for example, lung abscess or empyema) and revision of therapeutic antimicrobial regimens should be promptly revisited when procalcitonin levels remain high. However, definite data are lacking where *Pseudomonas*, *Acinetobacter*, *Stenotrophomonas*, and MRSA are concerned.

Abbreviations

BAT, best available therapy; CDC, Centers for Disease Control and Prevention; ESBL-PE, extended-spectrum beta-lactamase-producing *Enterobacteriaceae*; FiO₂, fraction of inspired oxygen; HAP, hospital-acquired pneumonia; ICU, intensive care unit; IVAC, infection-related ventilator-associated complication; MDR, multidrug-resistant; MIC, minimal inhibitory concentration; MRSA, methicillin-resistant *Staphylococcus aureus*; M-V, meropenem and vaborbactam; OR, odds ratio; PEEP, positive end expiratory pressure; SDD, selective digestive decontamination; SOD, selective oropharyngeal decontamination; VAC, ventilator-associated complication; VAE, ventilator-associated event; VAP, ventilator-associated pneumonia; VAT, ventilator-associated tracheobronchitis.

Competing interests

J-FT gave lectures for Merck, Pfizer, and Gilead on antimicrobial therapy and participated in scientific boards of Paratek, Bayer, Merck, and 3M. The other authors declare that they have no competing interests.

Grant information

The author(s) declared that no grants were involved in supporting this work.

Acknowledgments

The authors thank Celine Feger (EMIBiotech) for her editorial assistance.

References

1. American Thoracic Society, Infectious Diseases Society of America: **Guidelines for the management of adults with hospital-acquired, ventilator-associated, and healthcare-associated pneumonia.** *Am J Respir Crit Care Med.* 2005; 171(4): 388–416.
[PubMed Abstract](#) | [Publisher Full Text](#)
2. Chastre J, Fagon JY: **Ventilator-associated pneumonia.** *Am J Respir Crit Care Med.* 2002; 165(7): 867–903.
[PubMed Abstract](#) | [Publisher Full Text](#)
3. Craven DE, Hudcova J, Lei Y: **Diagnosis of ventilator-associated respiratory infections (VARI): microbiologic clues for tracheobronchitis (VAT) and pneumonia (VAP).** *Clin Chest Med.* 2011; 32(3): 547–57.
[PubMed Abstract](#) | [Publisher Full Text](#)
4. Martin-Loeches I, Papiol E, Almansa R, et al.: **Intubated patients developing tracheobronchitis or pneumonia have distinctive complement system gene expression signatures in the pre-infection period: a pilot study.** *Med Intensiva.* 2012; 36(4): 257–63.
[PubMed Abstract](#) | [Publisher Full Text](#)
5. Rello J, Riera J, Serrano R: **What's new in ventilator-associated pneumonia?** *Intensive Care Med.* 2015; 41(11): 1954–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
6. Magill SS, Klompas M, Balk R, et al.: **Developing a new, national approach to surveillance for ventilator-associated events: executive summary.** *Clin Infect Dis.* 2013; 57(12): 1742–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
7. Kalil AC, Metersky ML, Klompas M, et al.: **Management of Adults With Hospital-acquired and Ventilator-associated Pneumonia: 2016 Clinical Practice Guidelines by the Infectious Diseases Society of America and the American Thoracic Society.** *Clin Infect Dis.* 2016; 63(5): e61–e111.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
8. Timsit JF, de Kraker MEA, Sommer H, et al.: **Appropriate endpoints for evaluation of new antibiotic therapies for severe infections: a perspective from COMBACTE's STAT-Net.** *Intensive Care Med.* 2017; 43(7): 1002–1012.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
9. Torres A, Niederman MS, Chastre J, et al.: **International ERS/ESICM/ESCMID/ALAT guidelines for the management of hospital-acquired pneumonia and ventilator-associated pneumonia: Guidelines for the management of hospital-acquired pneumonia (HAP)/ventilator-associated pneumonia (VAP) of the European Respiratory Society (ERS), European Society of Intensive Care Medicine (ESICM), European Society of Clinical Microbiology and Infectious Diseases (ESCMID) and Asociación Latinoamericana del Tórax (ALAT).** *Eur Respir J.* 2017; 50(3): pii: 1700582.
[PubMed Abstract](#) | [Publisher Full Text](#)
10. Barbier F, Andremon A, Wolff M, et al.: **Hospital-acquired pneumonia and ventilator-associated pneumonia: recent advances in epidemiology and management.** *Curr Opin Pulm Med.* 2013; 19(3): 216–28.
[PubMed Abstract](#) | [Publisher Full Text](#)
11. Rosenthal VD, Bijie H, Maki DG, et al.: **International Nosocomial Infection Control Consortium (INICC) report, data summary of 36 countries, for 2004–2009.** *Am J Infect Control.* 2012; 40(5): 396–407.
[PubMed Abstract](#) | [Publisher Full Text](#)
12. Bouadma L, Sonnevile R, Garrouste-Orgeas M, et al.: **Ventilator-Associated Events: Prevalence, Outcome, and Relationship With Ventilator-Associated Pneumonia.** *Crit Care Med.* 2015; 43(9): 1798–806.
[PubMed Abstract](#) | [Publisher Full Text](#)
13. Warren DK, Shukla SJ, Olsen MA, et al.: **Outcome and attributable cost of ventilator-associated pneumonia among intensive care unit patients in a suburban medical center.** *Crit Care Med.* 2003; 31(5): 1312–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
14. Klompas M: **Does this patient have ventilator-associated pneumonia?** *JAMA.* 2007; 297(14): 1583–93.
[PubMed Abstract](#) | [Publisher Full Text](#)
15. Safdar N, Dezfoulian C, Collard HR, et al.: **Clinical and economic consequences of ventilator-associated pneumonia: a systematic review.** *Crit Care Med.* 2005; 33(10): 2184–93.
[PubMed Abstract](#) | [Publisher Full Text](#)
16. Timsit J, Zahar J, Chevret S: **Attributable mortality of ventilator-associated pneumonia.** *Curr Opin Crit Care.* 2011; 17(5): 464–71.
[PubMed Abstract](#) | [Publisher Full Text](#)
17. Melsen WG, Rovers MM, Groenwold RHH, et al.: **Attributable mortality of ventilator-associated pneumonia: a meta-analysis of individual patient data from randomised prevention studies.** *Lancet Infect Dis.* 2013; 13(8): 665–71.
[PubMed Abstract](#) | [Publisher Full Text](#)
18. Nguile-Makao M, Zahar JR, François A, et al.: **Attributable mortality of ventilator-associated pneumonia: respective impact of main characteristics at ICU admission and VAP onset using conditional logistic regression and multi-state models.** *Intensive Care Med.* 2010; 36(5): 781–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
19. Bekaert M, Timsit JF, Vansteelandt S, et al.: **Attributable mortality of ventilator-associated pneumonia: a reappraisal using causal analysis.** *Am J Respir Crit Care Med.* 2011; 184(10): 1133–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
20. Ibrahim EH, Ward S, Sherman G, et al.: **A comparative analysis of patients with early-onset vs late-onset nosocomial pneumonia in the ICU setting.** *Chest.* 2000; 117(5): 1434–42.
[PubMed Abstract](#) | [Publisher Full Text](#)
21. Moine P, Timsit J, De Lassence A, et al.: **Mortality associated with late-onset pneumonia in the intensive care unit: results of a multi-center cohort study.** *Intensive Care Med.* 2002; 28(2): 154–63.
[PubMed Abstract](#) | [Publisher Full Text](#)
22. Vallés J, Pobo A, García-Esquirol O, et al.: **Excess ICU mortality attributable to ventilator-associated pneumonia: the role of early vs late onset.** *Intensive Care Med.* 2007; 33(8): 1363–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
23. Park DR: **The microbiology of ventilator-associated pneumonia.** *Respir Care.* 2005; 50(6): 742–63; discussion 763–5.
[PubMed Abstract](#)
24. Trouillet JL, Chastre J, Vuagnat A, et al.: **Ventilator-associated pneumonia caused by potentially drug-resistant bacteria.** *Am J Respir Crit Care Med.* 1998; 157(2): 531–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
25. Ferrer M, Liapikou A, Valencia M, et al.: **Validation of the American Thoracic Society-Infectious Diseases Society of America guidelines for hospital-acquired pneumonia in the intensive care unit.** *Clin Infect Dis.* 2010; 50(7): 945–52.
[PubMed Abstract](#) | [Publisher Full Text](#)
26. Martin-Loeches I, Deja M, Koulenti D, et al.: **Potentially resistant microorganisms in intubated patients with hospital-acquired pneumonia: the interaction of ecology, shock and risk factors.** *Intensive Care Med.* 2013; 39(4): 672–81.
[PubMed Abstract](#) | [Publisher Full Text](#)
27. Martin-Loeches I, Povoa P, Rodríguez A, et al.: **Incidence and prognosis of ventilator-associated tracheobronchitis (TAVeM): a multicentre, prospective, observational study.** *Lancet Respir Med.* 2015; 3(11): 859–68.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
28. Gastmeier P, Sohr D, Geffers C, et al.: **Early- and late-onset pneumonia: is this still a useful classification?** *Antimicrob Agents Chemother.* 2009; 53(7): 2714–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
29. Restrepo MI, Peterson J, Fernandez JF, et al.: **Comparison of the bacterial etiology of early-onset and late-onset ventilator-associated pneumonia in subjects enrolled in 2 large clinical studies.** *Respir Care.* 2013; 58(7): 1220–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
30. Timsit JF, Pilmis B, Zahar JR: **How Should We Treat Hospital-Acquired and Ventilator-Associated Pneumonia Caused by Extended-Spectrum β -Lactamase-Producing Enterobacteriaceae?** *Semin Respir Crit Care Med.* 2017; 38(3): 287–300.
[PubMed Abstract](#) | [Publisher Full Text](#)
31. Lacherade JC, de Jonghe B, Guezennec P, et al.: **Intermittent subglottic secretion drainage and ventilator-associated pneumonia: a multicenter trial.** *Am J Respir Crit Care Med.* 2010; 182(7): 910–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
32. Bouadma L, Wolff M, Lucet JC: **Ventilator-associated pneumonia and its prevention.** *Curr Opin Infect Dis.* 2012; 25(4): 395–404.
[PubMed Abstract](#) | [Publisher Full Text](#)
33. Adair CG, Gorman SP, Feron BM, et al.: **Implications of endotracheal tube biofilm for ventilator-associated pneumonia.** *Intensive Care Med.* 1999; 25(10): 1072–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
34. Pronovost P, Needham D, Berenholtz S, et al.: **An intervention to decrease catheter-related bloodstream infections in the ICU.** *N Engl J Med.* 2006; 355(26): 2725–32.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
35. Reignier J, Darmon M, Sonnevile R, et al.: **Impact of early nutrition and feeding route on outcomes of mechanically ventilated patients with shock: a post hoc marginal structural model study.** *Intensive Care Med.* 2015; 41(5): 875–86.
[PubMed Abstract](#) | [Publisher Full Text](#)
36. Fitch ZW, Whitman GJ: **Incidence, risk, and prevention of ventilator-associated pneumonia in adult cardiac surgical patients: a systematic review.** *J Card Surg.* 2014; 29(2): 196–203.
[PubMed Abstract](#) | [Publisher Full Text](#)
37. Schwebel C, Clec'h C, Magne S, et al.: **Safety of intrahospital transport in ventilated critically ill patients: a multicenter cohort study*.** *Crit Care Med.* 2013; 41(8): 1919–28.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
38. Bornstain C, Azoulay E, De Lassence A, et al.: **Sedation, sucralfate, and antibiotic use are potential means for protection against early-onset ventilator-associated pneumonia.** *Clin Infect Dis.* 2004; 38(10): 1401–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
39. Klompas M, Branson R, Eichenwald EC, et al.: **Strategies to prevent ventilator-associated pneumonia in acute care hospitals: 2014 update.** *Infect Control Hosp Epidemiol.* 2014; 35 Suppl 2: S133–54.
[PubMed Abstract](#) | [Publisher Full Text](#)

40. Huang H, Li Y, Ariani F, *et al.*: **Timing of tracheostomy in critically ill patients: a meta-analysis.** *PLoS One*. 2014; 9(3): e92981.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
41. **F** Meng L, Wang C, Li J, *et al.*: **Early vs late tracheostomy in critically ill patients: a systematic review and meta-analysis.** *Clin Respir J*. 2016; 10(6): 684–92.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
42. **F** Szakmany T, Russell P, Wilkes AR, *et al.*: **Effect of early tracheostomy on resource utilization and clinical outcomes in critically ill patients: meta-analysis of randomized controlled trials.** *Br J Anaesth*. 2015; 114(3): 396–405.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
43. Clech C, Alberti C, Vincent F, *et al.*: **Tracheostomy does not improve the outcome of patients requiring prolonged mechanical ventilation: a propensity analysis.** *Crit Care Med*. 2007; 35(1): 132–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
44. Rello J, Lode H, Cornaglia G, *et al.*: **A European care bundle for prevention of ventilator-associated pneumonia.** *Intensive Care Med*. 2010; 36(5): 773–80.
[PubMed Abstract](#) | [Publisher Full Text](#)
45. **F** Bouadma L, Deslandes E, Lolom I, *et al.*: **Long-term impact of a multifaceted prevention program on ventilator-associated pneumonia in a medical intensive care unit.** *Clin Infect Dis*. 2010; 51(10): 1115–22.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
46. **F** Muscedere J, Sinuff T, Heyland DK, *et al.*: **The clinical impact and preventability of ventilator-associated conditions in critically ill patients who are mechanically ventilated.** *Chest*. 2013; 144(5): 1453–60.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
47. **F** Morris AC, Hay AW, Swann DG, *et al.*: **Reducing ventilator-associated pneumonia in intensive care: impact of implementing a care bundle.** *Crit Care Med*. 2011; 39(10): 2218–24.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
48. Speck K, Rawat N, Weiner NC, *et al.*: **A systematic approach for developing a ventilator-associated pneumonia prevention bundle.** *Am J Infect Control*. 2016; 44(6): 652–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
49. **F** Roquilly A, Marret E, Abraham E, *et al.*: **Pneumonia prevention to decrease mortality in intensive care unit: a systematic review and meta-analysis.** *Clin Infect Dis*. 2015; 60(1): 64–75.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
50. **F** Metersky ML, Wang Y, Klompas M, *et al.*: **Trend in Ventilator-Associated Pneumonia Rates Between 2005 and 2013.** *JAMA*. 2016; 316(22): 2427–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
51. **F** de Smet AM, Kluytmans JA, Cooper BS, *et al.*: **Decontamination of the digestive tract and oropharynx in ICU patients.** *N Engl J Med*. 2009; 360(1): 20–31.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
52. Oostdijk EAN, Kesecioglu J, Schultz MJ, *et al.*: **Effects of decontamination of the oropharynx and intestinal tract on antibiotic resistance in ICUs: a randomized clinical trial.** *JAMA*. 2014; 312(14): 1429–37.
[PubMed Abstract](#) | [Publisher Full Text](#)
53. **F** Klompas M: **Oropharyngeal Decontamination with Antiseptics to Prevent Ventilator-Associated Pneumonia: Rethinking the Benefits of Chlorhexidine.** *Semin Respir Crit Care Med*. 2017; 38(3): 381–90.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
54. Rello J, Lorente C, Bodí M, *et al.*: **Why do physicians not follow evidence-based guidelines for preventing ventilator-associated pneumonia?: a survey based on the opinions of an international panel of intensivists.** *Chest*. 2002; 122(2): 656–61.
[PubMed Abstract](#) | [Publisher Full Text](#)
55. Ricart M, Lorente C, Diaz E, *et al.*: **Nursing adherence with evidence-based guidelines for preventing ventilator-associated pneumonia.** *Crit Care Med*. 2003; 31(11): 2693–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
56. Pittet D: **The Lowbury lecture: behaviour in infection control.** *J Hosp Infect*. 2004; 58(1): 1–13.
[PubMed Abstract](#) | [Publisher Full Text](#)
57. Bouadma L, Mourvillier B, Deiler V, *et al.*: **Changes in knowledge, beliefs, and perceptions throughout a multifaceted behavioral program aimed at preventing ventilator-associated pneumonia.** *Intensive Care Med*. 2010; 36(8): 1341–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
58. **F** Nair GB, Niederman MS: **Ventilator-associated pneumonia: present understanding and ongoing debates.** *Intensive Care Med*. 2015; 41(1): 34–48.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
58. Tejerina E, Esteban A, Fernández-Segoviano P, *et al.*: **Accuracy of clinical definitions of ventilator-associated pneumonia: comparison with autopsy findings.** *J Crit Care*. 2010; 25(1): 62–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
60. Dudeck MA, Horan TC, Peterson KD, *et al.*: **National Healthcare Safety Network (NHSN) Report, data summary for 2010, device-associated module.** *Am J Infect Control*. 2011; 39(10): 798–816.
[PubMed Abstract](#) | [Publisher Full Text](#)
61. Magill SS, Klompas M, Balk R, *et al.*: **Developing a new, national approach to surveillance for ventilator-associated events*.** *Crit Care Med*. 2013; 41(11): 2467–75.
[PubMed Abstract](#) | [Publisher Full Text](#)
62. **F** Ego A, Preiser JC, Vincent JL: **Impact of diagnostic criteria on the incidence of ventilator-associated pneumonia.** *Chest*. 2015; 147(2): 347–55.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
63. Timsit JF, Misset B, Francoal S, *et al.*: **Is protected specimen brush a reproducible method to diagnose ICU-acquired pneumonia?** *Chest*. 1993; 104(1): 104–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
64. Gerbeaux P, Ledoray V, Boussuges A, *et al.*: **Diagnosis of nosocomial pneumonia in mechanically ventilated patients: repeatability of the bronchoalveolar lavage.** *Am J Respir Crit Care Med*. 1998; 157(1): 76–80.
[PubMed Abstract](#) | [Publisher Full Text](#)
65. de Lasseuse A, Joly-Guillou ML, Salah A, *et al.*: **Accuracy of delayed (24 hours) processing of bronchoalveolar lavage for diagnosing bacterial pneumonia.** *Crit Care Med*. 2004; 32(3): 680–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
66. **F** Canadian Critical Care Trials Group: **A randomized trial of diagnostic techniques for ventilator-associated pneumonia.** *N Engl J Med*. 2006; 355(25): 2619–30.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
67. Fagon JY, Chastre J, Wolff M, *et al.*: **Invasive and noninvasive strategies for management of suspected ventilator-associated pneumonia. A randomized trial.** *Ann Intern Med*. 2000; 132(8): 621–30.
[PubMed Abstract](#) | [Publisher Full Text](#)
68. Solé Violán J, Fernández JA, Benítez AB, *et al.*: **Impact of quantitative invasive diagnostic techniques in the management and outcome of mechanically ventilated patients with suspected pneumonia.** *Crit Care Med*. 2000; 28(8): 2737–41.
[PubMed Abstract](#) | [Publisher Full Text](#)
69. Bonten MJ, Bergmans DC, Stobbering EE, *et al.*: **Implementation of bronchoscopic techniques in the diagnosis of ventilator-associated pneumonia to reduce antibiotic use.** *Am J Respir Crit Care Med*. 1997; 156(6): 1820–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
70. Timsit JF, Cheval C, Gachot B, *et al.*: **Usefulness of a strategy based on bronchoscopy with direct examination of bronchoalveolar lavage fluid in the initial antibiotic therapy of suspected ventilator-associated pneumonia.** *Intensive Care Med*. 2001; 27(4): 640–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
71. Leone M, Bourgoin A, Cambon S, *et al.*: **Empirical antimicrobial therapy of septic shock patients: adequacy and impact on the outcome.** *Crit Care Med*. 2003; 31(2): 462–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
72. Raman K, Nailor MD, Nicolau DP, *et al.*: **Early antibiotic discontinuation in patients with clinically suspected ventilator-associated pneumonia and negative quantitative bronchoscopy cultures.** *Crit Care Med*. 2013; 41(7): 1656–63.
[PubMed Abstract](#) | [Publisher Full Text](#)
73. Zahar JR, Timsit JF, Garrouste-Orgeas M, *et al.*: **Outcomes in severe sepsis and patients with septic shock: pathogen species and infection sites are not associated with mortality.** *Crit Care Med*. 2011; 39(8): 1886–95.
[PubMed Abstract](#) | [Publisher Full Text](#)
74. Tabah A, Koultenti D, Laupland K, *et al.*: **Characteristics and determinants of outcome of hospital-acquired bloodstream infections in intensive care units: the EUROBACT International Cohort Study.** *Intensive Care Med*. 2012; 38(12): 1930–45.
[PubMed Abstract](#) | [Publisher Full Text](#)
75. **F** Decousser JW, Poirer L, Nordmann P: **Recent advances in biochemical and molecular diagnostics for the rapid detection of antibiotic-resistant *Enterobacteriaceae*: a focus on β -lactam resistance.** *Expert Rev Mol Diagn*. 2017; 17(4): 327–50.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
76. Laupland KB, Paiva JA, Timsit JF: **Focus on severe infections.** *Intensive Care Med*. 2017; 43(7): 1033–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
77. **F** Douglas IS, Price CS, Overdier KH, *et al.*: **Rapid automated microscopy for microbiological surveillance of ventilator-associated pneumonia.** *Am J Respir Crit Care Med*. 2015; 191(5): 566–73.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
78. Roberts JA, Taccone FS, Lipman J: **Understanding PK/PD.** *Intensive Care Med*. 2016; 42(11): 1797–1800.
[PubMed Abstract](#) | [Publisher Full Text](#)
79. Weiss E, Essaïed W, Adrie C, *et al.*: **Treatment of severe hospital-acquired and ventilator-associated pneumonia: a systematic review of inclusion and judgment criteria used in randomized controlled trials.** *Crit Care*. 2017; 21(1): 162.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
80. McKinnon PS, Paladino JA, Schentag JJ: **Evaluation of area under the inhibitory**

- curve (AUC) and time above the minimum inhibitory concentration ($T > MIC$) as predictors of outcome for cefepime and ceftazidime in serious bacterial infections. *Int J Antimicrob Agents*. 2008; 31(4): 345–51.
[PubMed Abstract](#) | [Publisher Full Text](#)
81. Chow KM, Hui AC, Szeto CC: Neurotoxicity induced by beta-lactam antibiotics: from bench to bedside. *Eur J Clin Microbiol Infect Dis*. 2005; 24(10): 649–53.
[PubMed Abstract](#) | [Publisher Full Text](#)
 82. Roberts JA, Lefrant JY, Lipman J: What's new in pharmacokinetics of antimicrobials in AKI and RRT? *Intensive Care Med*. 2017; 43(6): 904–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
 83. **F** Abdul-Aziz MH, Sulaiman H, Mat-Nor MB, *et al.*: Beta-Lactam Infusion in Severe Sepsis (BLISS): a prospective, two-centre, open-labelled randomised controlled trial of continuous versus intermittent beta-lactam infusion in critically ill patients with severe sepsis. *Intensive Care Med*. 2016; 42(10): 1535–45.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 84. Martínez JA, Cobos-Trigueros N, Soriano A, *et al.*: Influence of empiric therapy with a beta-lactam alone or combined with an aminoglycoside on prognosis of bacteremia due to gram-negative microorganisms. *Antimicrob Agents Chemother*. 2010; 54(9): 3590–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 85. **F** Kumar A, Zarychanski R, Light B, *et al.*: Early combination antibiotic therapy yields improved survival compared with monotherapy in septic shock: a propensity-matched analysis. *Crit Care Med*. 2010; 38(9): 1773–85.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 86. Boyer A, Gruson D, Bouchet S, *et al.*: Aminoglycosides in septic shock: an overview, with specific consideration given to their nephrotoxic risk. *Drug Saf*. 2013; 36(4): 217–30.
[PubMed Abstract](#) | [Publisher Full Text](#)
 87. **F** Ong DSY, Frencen JF, Klein Klouwenberg PMC, *et al.*: Short-Course Adjunctive Gentamicin as Empirical Therapy in Patients With Severe Sepsis and Septic Shock: A Prospective Observational Cohort Study. *Clin Infect Dis*. 2017; 64(12): 1731–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 88. Planquette B, Timsit JF, Misset BY, *et al.*: *Pseudomonas aeruginosa* ventilator-associated pneumonia: predictive factors of treatment failure. *Am J Respir Crit Care Med*. 2013; 188(1): 69–76.
[PubMed Abstract](#) | [Publisher Full Text](#)
 89. de Lastours V, Chau F, Roy C, *et al.*: Emergence of quinolone resistance in the microbiota of hospitalized patients treated or not with a fluoroquinolone. *J Antimicrob Chemother*. 2014; 69(12): 3393–400.
[PubMed Abstract](#) | [Publisher Full Text](#)
 90. Armand-Lefèvre L, Angebault C, Barbier F, *et al.*: Emergence of imipenem-resistant gram-negative bacilli in intestinal flora of intensive care patients. *Antimicrob Agents Chemother*. 2013; 57(3): 1488–95.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 91. Balakrishnan I, Awad-El-Kariem FM, Aali A, *et al.*: Temocillin use in England: clinical and microbiological efficacies in infections caused by extended-spectrum and/or derepressed AmpC β -lactamase-producing Enterobacteriaceae. *J Antimicrob Chemother*. 2011; 66(11): 2628–31.
[PubMed Abstract](#) | [Publisher Full Text](#)
 92. Vardakas KZ, Rellos K, Triarides NA, *et al.*: Colistin loading dose: evaluation of the published pharmacokinetic and clinical data. *Int J Antimicrob Agents*. 2016; 48(5): 475–84.
[PubMed Abstract](#) | [Publisher Full Text](#)
 93. Niederman MS, Chastre J, Corkery K, *et al.*: BAY41-6551 achieves bactericidal tracheal aspirate amikacin concentrations in mechanically ventilated patients with Gram-negative pneumonia. *Intensive Care Med*. 2012; 38(2): 263–71.
[PubMed Abstract](#) | [Publisher Full Text](#)
 94. **F** Kollef MH, Ricard JD, Roux D, *et al.*: A Randomized Trial of the Amikacin Fosfomycin Inhalation System for the Adjunctive Therapy of Gram-Negative Ventilator-Associated Pneumonia: IASIS Trial. *Chest*. 2017; 151(6): 1239–46.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 95. Kollef MH: COUNTERPOINT: Should Inhaled Antibiotic Therapy Be Used Routinely for the Treatment of Bacterial Lower Respiratory Tract Infections in the ICU Setting? No. *Chest*. 2017; 151(4): 740–3.
[PubMed Abstract](#) | [Publisher Full Text](#)
 96. **F** Bassetti M, Luyt CE, Nicolau DP, *et al.*: Characteristics of an ideal nebulized antibiotic for the treatment of pneumonia in the intubated patient. *Ann Intensive Care*. 2016; 6(1): 35.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 97. **F** Solé-Lleonart C, Rouby JJ, Blot S, *et al.*: Nebulization of Antiinfective Agents in Invasively Mechanically Ventilated Adults: A Systematic Review and Meta-analysis. *Anesthesiology*. 2017; 126(5): 890–908.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 98. **F** Zampieri FG, Nassar AP Jr, Gusmao-Flores D, *et al.*: Nebulized antibiotics for ventilator-associated pneumonia: a systematic review and meta-analysis. *Crit Care*. 2015; 19(1): 150.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 99. Rattanaumpawan P, Lorsutthitham J, Ungprasert P, *et al.*: Randomized controlled trial of nebulized colistimethate sodium as adjunctive therapy of ventilator-associated pneumonia caused by Gram-negative bacteria. *J Antimicrob Chemother*. 2010; 65(12): 2645–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
 100. **F** Choi EY, Huh JW, Lim CM, *et al.*: Relationship between the MIC of vancomycin and clinical outcome in patients with MRSA nosocomial pneumonia. *Intensive Care Med*. 2011; 37(4): 639–47.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 101. Patel N, Pai MP, Rodvold KA, *et al.*: Vancomycin: we can't get there from here. *Clin Infect Dis*. 2011; 52(8): 969–74.
[PubMed Abstract](#) | [Publisher Full Text](#)
 102. **F** Wunderink RG, Niederman MS, Kollef MH, *et al.*: Linezolid in methicillin-resistant *Staphylococcus aureus* nosocomial pneumonia: a randomized, controlled study. *Clin Infect Dis*. 2012; 54(5): 621–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 103. **F** Bouadma L, Luyt CE, Tubach F, *et al.*: Use of procalcitonin to reduce patients' exposure to antibiotics in intensive care units (PRORATA trial): a multicentre randomised controlled trial. *Lancet*. 2010; 375(9713): 463–74.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 104. **F** Klompas M, Li L, Menchaca JT, *et al.*: Ultra-Short-Course Antibiotics for Patients With Suspected Ventilator-Associated Pneumonia but Minimal and Stable Ventilator Settings. *Clin Infect Dis*. 2017; 64(7): 870–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 105. Luyt CE, Guérin V, Combes A, *et al.*: Procalcitonin kinetics as a prognostic marker of ventilator-associated pneumonia. *Am J Respir Crit Care Med*. 2005; 171(1): 48–53.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. Michael Klompas

¹ Department of Population Medicine, Harvard Medical School and Harvard Pilgrim Health Care Institute, Boston, MA, USA

² Department of Medicine, Brigham and Women's Hospital, Boston, MA, USA

Competing Interests: No competing interests were disclosed.

2. Saad Nseir

ICU, Lille Univ Hosp, Lille, France

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research