REVIEW

An update on phloem transport: a simple bulk flow under complex regulation [version 1; peer review: 5 approved]

Johannes Liesche^{1,2}, John Patrick³

¹Biomass Energy Center for Arid and Semi-arid lands, Northwest A&F University, Yangling, China

²College of Life Science, Northwest A&F University, Yangling, China

³Department of Biological Sciences, School of Environmental and Life Sciences, The University of Newcastle, Callaghan, Australia

V1 First published: 06 Dec 2017, 6(F1000 Faculty Rev):2096
<https://doi.org/10.12688/f1000research.12577.1>

Latest published: 06 Dec 2017, 6(F1000 Faculty Rev):2096
<https://doi.org/10.12688/f1000research.12577.1>

Abstract

The phloem plays a central role in transporting resources and signalling molecules from fully expanded leaves to provide precursors for, and to direct development of, heterotrophic organs located throughout the plant body. We review recent advances in understanding mechanisms regulating loading and unloading of resources into, and from, the phloem network; highlight unresolved questions regarding the physiological significance of the vast array of proteins and RNAs found in phloem saps; and evaluate proposed structure/function relationships considered to account for bulk flow of sap, sustained at high rates and over long distances, through the transport phloem.

Keywords

Phloem transport, signalling, plants, water

Open Peer Review

Approval Status

	1	2	3	4	5
version 1
06 Dec 2017					

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Antia Rodriguez-Villalon**, Swiss Federal Institute of Technology (ETH-Z), Zurich, Switzerland
2. **Friedrich Kragler**, Max Planck Institute of Molecular Plant Physiology, Potsdam, Germany
3. **Robert Turgeon**, Cornell University, New York, USA
4. **Brian Ayre**, University of North Texas, Texas, USA
5. **William W Adams**, University of Colorado, Boulder, USA

Any comments on the article can be found at the end of the article.

Corresponding authors: Johannes Liesche (liesche@nwafu.edu.cn), John Patrick (john.patrick@newcastle.edu.au)

Author roles: **Liesche J:** Writing – Original Draft Preparation, Writing – Review & Editing; **Patrick J:** Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: Published work authored by JWP was funded by the Australian Research Council.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2017 Liesche J and Patrick J. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Liesche J and Patrick J. **An update on phloem transport: a simple bulk flow under complex regulation** [version 1; peer review: 5 approved] F1000Research 2017, 6(F1000 Faculty Rev):2096 <https://doi.org/10.12688/f1000research.12577.1>

First published: 06 Dec 2017, 6(F1000 Faculty Rev):2096 <https://doi.org/10.12688/f1000research.12577.1>

Introduction

Successful functioning of land plants depends upon specialized transport systems. With some exceptions such as remobilized reduced carbon in winter-deciduous trees during the winter-spring transition, xylem conducts water and essential mineral elements from roots to transpiring leaves. Phloem transports water, mineral elements, amino nitrogen compounds, and sugars (resources), together with signalling molecules, from fully expanded leaves (sources) to meet the nutrient requirements of heterotrophic growth or storage organs (sinks) and to direct their development, respectively. Thus, phloem transport is at the heart of plant growth and development and, as such, is a primary factor determining crop yield potential. (For recent reviews, see Braun *et al.*¹, Ham and Lucas², and Yadav *et al.*³.)

Phloem transport occurs as a pressure-driven bulk flow through a longitudinally arrayed subset of transport-specialized cells termed sieve elements (SEs). During their development, SEs undergo partial autophagy that leaves a parietal enucleate cytoplasm enclosed by a plasma membrane (PM). (For more details of recent progress in understanding phloem development, see reviews by Rodriguez-Villalon⁴ and de Rybel *et al.*⁵.) In contrast to SEs, their adjoining companion cells (CCs) contain a dense cytoplasm and are connected to their corresponding SE through an extensive network of plasmodesmata (PD) to form a metabolic and genetic unit, the sieve element-companion cell complex (SECCC). Loading SECCCs in the collection phloem of source leaves generates an osmotically derived hydrostatic pressure potential that drives bulk flow through the SE arrays of the transport phloem to reach the release phloem where resources and signals are unloaded to enter pathways leading to plant growth or storage^{6,7}.

In this commentary, we review new insights into key aspects of phloem transport and highlight unresolved questions that need to be addressed to gain a fuller understanding of phloem transport.

Phloem loading

Sugar uptake into the phloem is a principal contributor to establishing source-sink hydrostatic pressure differentials driving phloem transport^{6,7}. Different phloem loading types have been identified, and abundant insight into phloem loading mechanisms has become available. However, two key aspects remain largely unexplored: the ecology and regulation of phloem loading.

Three major types of phloem loading

The three main types of phloem loading currently recognized are (i) active apoplastic, (ii) active symplasmic, and (iii) passive symplasmic (Figure 1). The mechanism of active apoplastic loading (Figure 1) is well understood with proton-coupled sucrose transporters (SUTs), also referred to as SUCs, concentrating sucrose into SECCCs^{8–10} released from surrounding cells by sucrose uniporters, termed sugars will eventually be exported transporters (SWEETs)^{11,12}. In contrast, the function of the two symplasmic loading types has been established only recently.

A remaining major question for active symplasmic loading is how PD enable steric filtering to allow sucrose movement into CCs while preventing the synthesized sugar oligomers from leaking back to the mesophyll cells (Figure 1). No method exists to resolve PD canal sub-structure to the degree required to answer this question, and current insights rest solely on mathematical modelling. For instance, Liesche and Schulz¹³ showed that a PD structure that is highly restrictive to solute movement, with cytosolic channels of 0.65 nm in diameter compared with diameters of 2.5 nm in PD

Figure 1. Major phloem loading types. In active apoplastic loading, the sieve element-companion cell complex (SECCC) is symplasmically isolated. Sucrose produced in the mesophyll cells (MCs) diffuses into phloem parenchyma cells (PPCs), where it is released into the apoplast by efflux carriers (SWEETs) before being taken up into the SECCC by plasma membrane-localized sucrose transporters. In active symplasmic loading, sucrose can diffuse or convect into the companion cell (CC) through the abundant plasmodesmata (PD). Sucrose in the CC is converted to sugar oligomers, which are hindered from diffusing back into the phloem parenchyma but instead enter the SE through the larger PD at the SE-CC interface. In passive symplasmic loading, sucrose can diffuse or convect along the whole phloem loading pathway from mesophyll cells to SEs following the sugar concentration gradient. Aquaporins facilitate the osmotic uptake of water from the phloem apoplast into SEs in all loading types. In symplasmic loaders, water also enters the SECCC through PD. SWEET, sugar will eventually be exported transporters.

between leaf mesophyll cells¹⁴, could enable steric sugar filtering while accommodating observed phloem loading rates. However, this extreme PD anatomy might not be necessary as differences in intracellular sugar concentrations along the loading pathway from mesophyll to phloem could be sufficient to drive a combination of diffusion and mass flow of molecules (convective flow) into SECCCs. As a consequence, diffusion of sugar oligomers out of the SECCC, against the direction of the convective flow, would be limited^{15,16}.

New evidence regarding passive symplasmic loading (Figure 1) was provided by the finding that sucrose export from poplar leaves was not compromised by blocking apoplasmic loading of sucrose¹⁷. This was achieved through expressing a cell wall invertase to cleave sucrose into hexoses, thereby making the sugar unavailable for uptake by SUTs. Furthermore, microfluidic experiments showed that passive phloem loading is feasible with convective flow through PD to the SECCCs¹⁸. However, it is unclear whether this concept is universally applicable across all species exhibiting features of passive symplasmic loading. For example, in several designated passive loaders, higher sugar concentrations were detected in phloem compared with mesophyll cells^{19,20}. In addition to methodological controversies surrounding measurements of intracellular sugar concentrations, a switch between loading types or even their concomitant operation could explain the conflicting conclusions²¹.

Parallel and sequential operation of different phloem loading types

Coexistence of phloem loading types has become apparent for many plant species²². For instance, two anatomical types of CCs have been detected in minor veins in 35% of 320 Asteridae species exhibiting PD characteristics consistent with active symplasmic or apoplasmic loading²³ as well as other species of various growth forms^{24,25}. Moreover, switching from active symplasmic to apoplasmic phloem loading, putatively by modifying PD conductance in the same CC, has been described for virus-infected melon plants²⁶. A switch between loading types in the same CC is thought to be impossible without adaptation of the relevant PD^{21,27}. Consequently, elucidating this potential mechanism of switching between loading types is a major task for future research on phloem loading.

In addition to switching between loading types, concomitant operation of passive and active loading by the same CC has been proposed²⁰ and shown to be theoretically possible though attenuated by a low efficiency²¹. Passive loading could function as a fallback mechanism as demonstrated by continued growth, at a reduced rate, of plants in which active loading is blocked through introduction of mutations that render either the relevant SUT or enzymes involved in sucrose oligomerization non-functional^{22,28,29}.

Ecology of phloem loading types

No satisfactory answer has been provided to account for why there are different phloem loading types. Slewinski *et al.*²² theorized that some plants load certain molecules symplasmically in addition to those transported across the PM of SECCCs, but no candidates have been identified so far. Analysis of growth form

concluded that herbaceous plants generally are not passive symplasmic loaders³⁰. Presumably, this is because active loading allows the plant to keep carbon content low in leaves, which provides a substantial, positive effect on growth rate³¹. However, this does not mean that the slower-growing trees are all passive loaders. Indeed, half of the families dominated by woody plant species exhibit an active loading mechanism^{21,30}. Evolutionary and distribution analyses similarly do not provide a clear explanation for why certain families use a particular phloem loading type³⁰. A closer examination of a plant's ecology might reveal ecophysiological adaptations that different loading types offer. For example, a link between loading type and a plant's adaptive environmental response has been demonstrated in a comparative study of minor vein structure developed under different temperature and light conditions³².

Regulation of phloem loading

While sugar export rates are a function of sugar availability and sink demand^{33,34} as well as phloem loading capacity^{35,36}, the underlying molecular mechanisms are poorly understood. SUT activities are expected to be principal regulators of sugar export in apoplasmic loaders⁵ as demonstrated by SUT overexpression³⁷. In this context, transcriptional SUT regulation by phytohormones and corresponding changes in sugar export rates were observed in potato³⁸. Several studies suggest that post-translational regulation of SUTs contributes to determining export rates (for example, Sakr *et al.*³⁹). How post-translational regulation of SUTs occurs *in vivo* is unclear but could involve differential intracellular localization⁴⁰, dimerization⁴¹, and protein-protein interactions⁴². However, other factors should be considered. For instance, simultaneous upregulated expression of SUTs and SWEETs, in response to water deficit, increased phloem loading⁴³. Furthermore, constitutive overexpression of a proton-pumping pyrophosphatase in *Arabidopsis thaliana* leaf CCs increased phloem loading possibly by enhancing the proton motive force driving sucrose flux through SECCC localized SUTs^{44,45}. Regulation of sucrose exchange between cytosol and vacuole of cells in the pre-phloem pathway influences phloem loading rates even in apoplasmic loaders^{46,47}. So far not demonstrated is the short-term regulation of phloem loading through modification of PD permeability, which would be especially relevant in symplasmic loaders²¹. After elucidation of the molecular mechanisms regulating phloem loading, the important question will be how these are linked to sink demand⁴⁸.

Phloem transport

This section combines a description of the transport phloem, especially how its anatomy facilitates sap flow, with a more general discussion on the mechanism of phloem transport and the molecules transported within.

The mechanism of phloem transport in trees

Osmotically driven pressure flow has been widely accepted as the mechanism of phloem transport in herbaceous plants^{49,50}. However, in regard to trees, where distances between source and sink can extend up to 100 m, there are doubts about whether a hydrostatic pressure potential sufficient to drive flow could be generated^{51,52}. A variety of approaches have been employed to

answer this question. Simple theoretical models of Münch-type pressure flow agree with measurements of stem diameter variations⁵³ or SE anatomy^{54,55}. Importantly, the main prediction of pressure flow, that SE conductivity scales with transport distance⁵⁶, could be confirmed. The scaling of SE conductivity with tree height was shown within a single tree (Figure 2A;^{57,58}), within a species⁵⁹, and across species (Figure 2B;^{55,60}), confirming that resistance decreases to accommodate mass flow in larger trees. Furthermore, it was recently shown in mature, field-grown Scots pine trees that there is an osmotic pressure gradient along the phloem pathway from leaves to the stem base (Figure 2C;⁶¹). The osmotic pressure gradient, supported by gravity, was calculated to be large enough to overcome the xylem water pressure potential and establish a phloem turgor pressure gradient that drives mass flow according to the Münch mechanism at all times across the diel cycle⁶¹. Taken together, these results confirm that Münch-type pressure flow works even in the tallest angiosperm and gymnosperm trees, although transport speed might be up to 10 times lower than in herbaceous plants⁶². However, this is in agreement with equally lower rates of photosynthesis and growth⁶³.

Open questions regarding the functional anatomy of the phloem

Despite major progress in determining how phloem anatomy facilitates osmotically driven pressure flow, not all questions regarding its functional anatomy have been resolved. For example, sieve plates forming the axial connection between adjoining SEs have been assumed to evolve toward a simple form associated with lower hydraulic resistance—a paradigm overturned by an anatomical analysis of 447 species⁶⁰. Rather, the distinct organizational patterns of sieve plates might relate to general vascular anatomy (for example, SE length or xylem cell anatomy) instead of differences in conductivity⁶⁰. Another aspect that is relevant for phloem transport is the symplasmic coupling along the

transport path. Even in apoplasmically loading herbaceous plants, unloading along the stem can switch between apoplasmic and symplasmic unloading pathways, depending upon the source/sink ratio, with high ratios favouring symplasmic unloading⁶⁴. In many tree species, especially conifers, the presence of PD connecting the SECCC with surrounding cells⁶⁵ indicates symplasmic coupling, presumably to efficiently supply the cambium with the large amounts of sugars needed for wood formation. The question is in how far symplasmic coupling along the path compromises whole-plant phloem transport. A recent theoretical analysis showed that unloading along the transport path considerably influences flow if it is not balanced by reloading⁶⁶. The most likely explanation is that unloading is symplasmic only in case of high sugar concentrations in the ground tissue, which then could also have a function in homeostatically maintaining SE turgor⁶⁷.

Another central issue is the relationship between fibrous SE protein (p-protein) agglomerations, considered to occlude sieve pores to prevent sap loss in the event of damage⁶⁸. Careful preparation of material for electron microscope examination reveals unobstructed SE lumens bounded by a parietal cytoplasm. However, Froelich *et al.*⁶⁹ showed that in *A. thaliana* phloem, the p-protein, AtSEOR1, forms a meshwork at the margins and clots in the lumen of intact SEs and their presence does not impede longitudinal flow. Together with discrepancies between theoretical flow speeds calculated according to SE conductivity and observed flow speeds⁷⁰, these observations have led to the conclusion that phloem transport in herbaceous plants and grasses is not limited by SE conductivity but only by sink strength⁷¹.

Phloem sap composition

Knowledge of phloem sap composition is relevant to reach a quantitative understanding of resource allocation and inter-organ signalling. Sugar concentrations are typically around 20%, but

Figure 2. Scaling of sieve element (SE) conductivity and osmotic pressure potential enable Münch-type pressure flow in trees. (A) Axial widening of phloem cells along the stem of individual trees of Norway spruce (blue), European ash (orange), and bitter willow (gray). Average values of cell diameter contain up to 30% non-SE cells. Axial widening of SEs toward the bottom of a tree translates to decreased hydraulic resistance, and the scaling relationship was found to enable optimal xylem-phloem water exchange that is critical to drive flow in tall trees. (B) SE resistance of 44 different tree species with sampled individuals grown to their typical maximum height. Negative scaling of resistance with transport distance illustrates the anatomical optimization of SEs for phloem transport in trees. (C) Measurements of osmolality indicated the presence of an osmotic pressure gradient along the phloem of 18 m high Scots pine trees, which changed in accordance with the xylem water potential. The differences in osmotic potential were calculated to be large enough to overcome the xylem water pressure potential and establish a phloem turgor pressure gradient that drives mass flow according to the Münch mechanism across the diel cycle⁵⁶. Frames (A–C) are based on data from Petit and Crivellaro⁵⁷, Liesche *et al.*⁶⁰, and Paljakka *et al.*⁶¹, respectively.

the sap contains other molecules (for example, amino nitrogen compounds, proteins, and RNAs) and mineral ions⁷². Determining the precise abundance of the different sap components remains one of the biggest challenges in phloem research as all methods available so far are prone to artifacts^{73–75}. Recently, laser-capture microdissection was used to uncover the link between seasonal differences in phloem metabolites and phloem formation in Norway spruce. However, since sampling could not be restricted to SEs, results were not representative of phloem sap⁷⁶. In the future, insight on relative sap composition might be gained from using rapid freezing and nanoscale secondary ion mass spectrometry (NanoSIMS), which enables semi-quantitative imaging of ions as well as isotope-labeled nitrogen and carbon compounds within SEs⁷⁷. Gaining a complete picture of the spatiotemporal dynamics of phloem sap composition will be instrumental in refining multi-compartmental metabolic models describing source-to-sink metabolite flows (for example, Zakhartsev *et al.*⁷⁸).

Controversy persists surrounding the function of phloem-mobile proteins and RNA⁷⁹. Recently, a meta-analysis comparing proteomes of xylem, phloem, and leaf apoplastic saps highlighted phloem sap as being the most enriched in signalling proteins of which 13 were conserved across experiments and that included the ubiquitous phloem-mobile protein FLOWERING LOCUS T⁸⁰. Grafting studies suggest phloem mobility of a multitude of additional proteins⁸¹. However, without further validation of their signalling function, many of these proteins could have entered the translocation stream “by accident” rather than by a control mechanism located at PD interconnecting CCs with SEs⁸². Of the proteins identified in phloem sap by proteomics, those associated with redox regulation and plant defense consistently have been found across species⁸⁰, suggesting that many phloem proteins function in the response to biotic and abiotic stress. Prominent examples of phloem-mobile RNAs include the transcription factors *BEL5*, which influences potato tuber induction⁸³, and the Cucurbit *NACP* regulating apical meristem development⁸⁴. Recently 2,006 *A. thaliana* phloem-mobile mRNAs were identified⁸⁵ with a conserved tRNA-derived sequence conferring mobility⁸⁶. However, since most of the mRNAs are expressed in mesophyll cells, it is unclear how they could enter the phloem stream in the apoplastic loader *A. thaliana*. In addition to those many phloem-mobile proteins and mRNAs, the role of microRNAs⁸⁷ and lipids⁸⁸ as systemic signals needs to be further explored.

Phloem unloading

Phloem unloading describes the movement of phloem sap constituents from SECCC lumens (SECCC unloading) and their subsequent cell-to-cell transport to final destinations in non-SECCC vascular or ground tissues. Cellular pathways of phloem unloading are apoplastic (Figure 3A) or symplasmic with or without an intervening apoplastic step located in the post-SECCC pathway (Figure 3B, C;⁸⁹). As outlined below, phloem unloading mechanisms are only partially resolved.

Apoplastic phloem unloading mechanisms

Possible mechanisms contributing to efflux across any cell PM, including SECCC PMs, are passive diffusion, transporter-mediated facilitated diffusion, and energy-coupled movement.

Passive diffusive fluxes of molecules and ions are the product of their membrane permeability coefficients and *trans*-membrane gradients in chemical potential for non-electrolytes or electrochemical potential for electrolytes⁹⁰. To illustrate the magnitude that these passive fluxes could reach for apoplastic SECCC unloading of sucrose along the entire phloem pathway from source to sink, a plausible maximum of 1 M for the SECCC *trans*-PM sucrose concentration difference and a membrane permeability coefficient of $10^{-10} \text{ m s}^{-191}$ predict a sucrose efflux of $1 \times 10^{-7} \text{ mol m}^{-2} \text{ s}^{-1}$. This matches the maximal membrane flux for loading sucrose into SECCCs ($2.3 \times 10^{-7} \text{ mol m}^{-2} \text{ s}^{-1}$; Giaquinta⁹²). Thus, it is imperative to experimentally determine rates of diffusive leak from SECCCs and include this component in theoretical models of phloem transport.

For facilitated membrane efflux of sugars by uniporters, SWEETs^{11,12} have been detected in transport phloem parenchyma cells⁹³, but whether these function in efflux to, or retrieval from, the phloem apoplast remains to be determined. However, sucrose/proton symporters occur in SECCCs of root and stem transport phloem (for example, 94,95). Depolarized membrane potentials (−55 mV) of root SECCCs predict that sucrose/proton symport could reverse to an efflux mode driven by a *trans*-membrane sucrose concentration difference of 85.5 mM (for more information, see Carpaneto *et al.*⁹⁴). In contrast, symporter reversal in stem phloem is highly unlikely as their SE/CC membrane potential of −110 mV⁹⁶ would require a sucrose concentration difference of 7,200 mM, a concentration difference that far exceeds physiological limits.

During sugar accumulation in fleshy fruits, phloem unloading follows an apoplastic route^{64,97–100}. Whether the sucrose leak from fruit SECCCs is augmented by reversal of sucrose/proton symporters^{101,102} cannot be evaluated, as their membrane potentials are unknown. However, pharmacological studies suggest that sucrose efflux from vascular bundles of apple fruit and grape berries is mediated by energy-coupled carriers^{97,103}. Resolving the identity (or identities) of the putative effluxer (or effluxers) is central to acquiring a full understanding of phloem unloading in fleshy fruits. What is clearer is that monosaccharide/proton symporters retrieve inverted disaccharides from the fruit apoplast into storage parenchyma cells^{104–107} and, together with cell wall invertases, function to co-regulate phloem unloading^{106,108,109}.

In all cases, unloading of osmotic solutes must be accompanied by a proportionate loss of phloem water to maintain water potential equilibrium. The exit of water likely occurs through aquaporins localized in PMs of the unloading SEs (Figure 3A and, for example, 110,111).

Symplasmic unloading mechanisms with or without an intervening apoplastic step located in the post-SECCC pathway

In recent years, it is becoming increasingly clear that a major component of symplasmic unloading from SECCCs is contributed by bulk flow. For instance, experimental manipulations of hydrostatic pressure gradients in root tips and stems cause changes in phloem unloading rates consistent with bulk flow (reviewed by Patrick¹¹²). In developing wheat seeds, hydrostatic pressure

Figure 3. Cellular pathways and mechanisms of phloem unloading. (A) Apoplastic sieve element-companion cell complex (SECCC) unloading of sucrose mediated by sucrose transporter reversal and/or possible by sugars will eventually be exported transporters (SWEETs) with energy coupled transporters retrieving sugars from the sink apoplasm while accompanying water transport across the plasma membranes facilitated by aquaporins. (B) Symplasmic phloem unloading by bulk flow. (C) Symplasmic phloem unloading by bulk flow with an intervening apoplastic step in the post-phloem pathway of developing seeds. Sucrose release to the seed apoplasm from maternal seed coats is mediated by SWEETs, sucrose facilitators and possibly a yet-to-be-cloned sucrose/proton antiporter. SWEETs and sucrose transporters recover released sucrose from the seed apoplasm into the endosperm/embryo. Membrane transport of water facilitated by aquaporins. GT, ground tissue; PPC, phloem parenchyma cell.

differences of up to 1.0 MPa between SEs and vascular parenchyma cells account for observed PD volume flow rates¹¹³. In contrast, modelling flows through large-diameter funnel-shaped PD, interconnecting SEs, and adjacent pericycle cells in *A. thaliana* root tips predicted that hydrostatic pressure differences of only 0.05 to 0.2 MPa were required¹¹⁴. Whether these model-based hydrostatic pressure differences can be reconciled with measured hydrostatic or osmotic pressures of 1.3 and 0.6 MPa, respectively, between SEs and surrounding cells in barley and maize root tips^{115,116} awaits determination.

For the apoplastic step in the mandatory phloem-unloading pathway of developing seeds (Figure 3B), sucrose release from maternal seed coats occurs by a combination of facilitated diffusion and sucrose/proton antiport while retrieval by embryo/endosperm is mediated by sucrose symporters¹¹⁷ or SWEETs¹¹⁸ or both. Cloned sucrose facilitators (SUFs) from grain legume seed coats exhibited transport properties consistent with those found for facilitated diffusion of sucrose in native membranes of seed coats and were expressed in cells considered responsible for sucrose efflux^{119,120}. In developing *A. thaliana* seeds, SWEETs

are positioned to provide a cascade of sucrose transport from the outer (SWEET15) and inner (SWEET12) integuments and uptake into the filial tissues (SWEET11 and SWEET15¹²¹) while SWEET11 facilitates release from maternal tissues of developing rice seeds¹²². Consistent with relative contributions of facilitated diffusion to sucrose effluxed from grain legume seed coats¹¹⁷, seed weights, but not number, were depressed by 50% in the *A. thaliana* triple-mutant *sweet11*, *12* and *15*, and by 65% in the rice *sweet11* mutant^{121,122}. The challenge now is to identify the gene or (genes) encoding the membrane protein (or proteins) responsible for energy-coupled sucrose/proton antiport from maternal seed tissues¹²³ that may also operate in fleshy fruit (see previous section). Membrane transport of solutes is matched by water movement through aquaporins located in PMs of maternal and filial tissues of developing seeds (Figure 3C and, for example, 124,125).

Conclusions and future directions

Throughout the text, we have highlighted a series of unresolved questions of phloem transport biology required to move the understanding of phloem loading, axial transport, and unloading forward. Their resolution will inform profitable approaches to

address the key question of how the components of phloem transport are integrated into a functional whole⁷¹ and how phloem transport mechanistically intermeshes with photosynthesis and sink demand for resources^{63,71}.

Abbreviations

CC, companion cell; PD, plasmodesmata; PM, plasma membrane; SE, sieve element; SECCC, sieve element-companion cell complex; SUT, sucrose transporter; SWEET, sugars will eventually be exported transporters

Competing interests

The authors declare that they have no competing interests.

Grant information

Published work authored by JWP was funded by the Australian Research Council.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- Braun DM, Wang L, Ruan YL: **Understanding and manipulating sucrose phloem loading, unloading, metabolism, and signalling to enhance crop yield and food security.** *J Exp Bot.* 2014; **65**(7): 1713–35.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Ham BK, Lucas WJ: **The angiosperm phloem sieve tube system: a role in mediating traits important to modern agriculture.** *J Exp Bot.* 2014; **65**(7): 1799–816.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Yadav UP, Ayre BG, Bush DR: **Transgenic approaches to altering carbon and nitrogen partitioning in whole plants: assessing the potential to improve crop yields and nutritional quality.** *Front Plant Sci.* 2015; **6**: 275.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rodriguez-Villalon A: **Wiring a plant: genetic networks for phloem formation in Arabidopsis thaliana roots.** *New Phytol.* 2016; **210**(1): 45–50.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- De Rybel B, Mähönen AP, Helariutta Y, et al.: **Plant vascular development: from early specification to differentiation.** *Nat Rev Mol Cell Biol.* 2016; **17**(1): 30–40.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Ayre BG: **Membrane-transport systems for sucrose in relation to whole-plant carbon partitioning.** *Mol Plant.* 2011; **4**(3): 377–94.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Ainsworth EA, Bush DR: **Carbohydrate export from the leaf: a highly regulated process and target to enhance photosynthesis and productivity.** *Plant Physiol.* 2011; **155**(1): 64–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Riesmeier JW, Willmitzer L, Frommer WB: **Isolation and characterization of a sucrose carrier cDNA from spinach by functional expression in yeast.** *EMBO J.* 1992; **11**(13): 4705–13.
[PubMed Abstract](#) | [Free Full Text](#)
- Riesmeier JW, Willmitzer L, Frommer WB: **Evidence for an essential role of the sucrose transporter in phloem loading and assimilate partitioning.** *EMBO J.* 1994; **13**(1): 1–7.
[PubMed Abstract](#) | [Free Full Text](#)
- Gahrtz M, Stolz J, Sauer N: **A phloem-specific sucrose-H⁺ symporter from Plantago major L. supports the model of apoplastic phloem loading.** *Plant J.* 1994; **6**(5): 697–706.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Chen LQ: **SWEET sugar transporters for phloem transport and pathogen nutrition.** *New Phytol.* 2014; **201**(4): 1150–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Chen LQ, Qu XQ, Hou BH, et al.: **Sucrose efflux mediated by SWEET proteins as a key step for phloem transport.** *Science.* 2012; **335**(6065): 207–11.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Liesche J, Schulz A: **Modeling the parameters for plasmodesmal sugar filtering in active symplasmic phloem loaders.** *Front Plant Sci.* 2013; **4**: 207.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Ding B, Turgeon R, Parthasarathy MV: **Substructure of freeze-substituted plasmodesmata.** *Protoplasma.* 1992; **169**(1–2): 28–41.
[Publisher Full Text](#)
- Münch E: **Die Stoffbewegungen in der Pflanze.** *Protoplasma.* 1932; **15**(1): 488–9.
[Publisher Full Text](#)
- Comtet J, Turgeon R, Stroock AD: **Phloem loading through plasmodesmata: a biophysical analysis.** *Plant Physiol.* 2017; **175**(2): 904–15.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Zhang C, Han L, Siewinski TL, et al.: **Symplastic phloem loading in poplar.** *Plant Physiol.* 2014; **166**(1): 306–13.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Comtet J, Jensen KH, Turgeon R, et al.: **Passive phloem loading and long-distance transport in a synthetic tree-on-a-chip.** *Nat Plants.* 2017; **3**: 17032.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Nadwodnik J, Lohaus G: **Subcellular concentrations of sugar alcohols and sugars in relation to phloem translocation in Plantago major, Plantago maritima, Prunus persica, and Apium graveolens.** *Planta.* 2008; **227**(5): 1079–89.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Oner-Sieben S, Lohaus G: **Apoplastic and symplastic phloem loading in Quercus robur and Fraxinus excelsior.** *J Exp Bot.* 2014; **65**(7): 1905–16.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Liesche J: **Sucrose transporters and plasmodesmal regulation in passive phloem loading.** *J Integr Plant Biol.* 2017; **59**(5): 311–21.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Siewinski TL, Zhang C, Turgeon R: **Structural and functional heterogeneity in phloem loading and transport.** *Front Plant Sci.* 2013; **4**: 244.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Batashev DR, Pakhomova MV, Razumovskaya AV, et al.: **Cytology of the minor-vein phloem in 320 species from the subclass Asteridae suggests a high diversity of phloem-loading modes.** *Front Plant Sci.* 2013; **4**: 312.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Öner-Sieben S, Rapp C, Sauer N, et al.: **Characterization, localization, and seasonal changes of the sucrose transporter FeSUT1 in the phloem of**

- Fraxinus excelsior.** *J Exp Bot.* 2015; **66**(15): 4807–19.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
25. van Bel AJ, van Kesteren WJ, Papenhuijzen C: **Ultrastructural indications for coexistence of symplastic and apoplastic phloem loading in *Commelina benghalensis* leaves: Differences in ontogenic development, spatial arrangement and symplastic connections of the two sieve tubes in the minor vein.** *Planta.* 1988; **176**(2): 159–72.
[PubMed Abstract](#) | [Publisher Full Text](#)
 26. **F** Gil L, Yaron I, Shalitin D, *et al.*: **Sucrose transporter plays a role in phloem loading in CMV-infected melon plants that are defined as symplastic loaders.** *Plant J.* 2011; **66**(2): 366–74.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 27. Hannah MA, Zuther E, Buchel K, *et al.*: **Transport and metabolism of raffinose family oligosaccharides in transgenic potato.** *J Exp Bot.* 2006; **57**(14): 3801–11.
[PubMed Abstract](#) | [Publisher Full Text](#)
 28. **F** McCaskill A, Turgeon R: **Phloem loading in *Verbascum phoeniceum* L. depends on the synthesis of raffinose-family oligosaccharides.** *Proc Natl Acad Sci U S A.* 2007; **104**(49): 19619–24.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 29. Srivastava AC, Ganesan S, Ismail IO, *et al.*: **Functional characterization of the Arabidopsis AtSUC2 Sucrose/H⁺ symporter by tissue-specific complementation reveals an essential role in phloem loading but not in long-distance transport.** *Plant Physiol.* 2008; **148**(1): 200–11.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 30. Davidson A, Keller F, Turgeon R: **Phloem loading, plant growth form, and climate.** *Protoplasma.* 2011; **248**(1): 153–63.
[PubMed Abstract](#) | [Publisher Full Text](#)
 31. Turgeon R: **The role of phloem loading reconsidered.** *Plant Physiol.* 2010; **152**(4): 1817–23.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 32. Adams WW 3rd, Cohu CM, Muller O, *et al.*: **Foliar phloem infrastructure in support of photosynthesis.** *Front Plant Sci.* 2013; **4**: 194.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 33. Jiao J, Grodzinski B: **The effect of leaf temperature and photorespiratory conditions on export of sugars during steady-state photosynthesis in *Salvia splendens*.** *Plant Physiol.* 1996; **111**(1): 169–78.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 34. Grodzinski B, Jiao J, Leonardos ED: **Estimating photosynthesis and concurrent export rates in C₃ and C₄ species at ambient and elevated CO₂.** *Plant Physiol.* 1998; **117**(1): 207–15.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 35. **F** Amiard V, Mueh KE, Demmig-Adams B, *et al.*: **Anatomical and photosynthetic acclimation to the light environment in species with differing mechanisms of phloem loading.** *Proc Natl Acad Sci U S A.* 2005; **102**(36): 12968–73.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 36. Muller O, Cohu CM, Stewart JJ, *et al.*: **Association between photosynthesis and contrasting features of minor veins in leaves of summer annuals loading phloem via symplastic versus apoplastic routes.** *Physiol Plant.* 2014; **152**(1): 174–83.
[PubMed Abstract](#) | [Publisher Full Text](#)
 37. **F** Wang L, Lu Q, Wen X, *et al.*: **Enhanced sucrose loading improves rice yield by increasing grain size.** *Plant Physiol.* 2015; **169**(4): 2848–62.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 38. **F** Chincinska IA, Liesche J, Krügel U, *et al.*: **Sucrose transporter StSUT4 from potato affects flowering, tuberization, and shade avoidance response.** *Plant Physiol.* 2008; **146**(2): 515–28.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 39. Sakr S, Noubahni M, Bourbouloux A, *et al.*: **Cutting, ageing and expression of plant membrane transporters.** *Biochim Biophys Acta.* 1997; **1330**(2): 207–16.
[PubMed Abstract](#) | [Publisher Full Text](#)
 40. Liesche J, He HX, Grimm B, *et al.*: **Recycling of Solanum sucrose transporters expressed in yeast, tobacco, and in mature phloem sieve elements.** *Mol Plant.* 2010; **3**(6): 1064–74.
[PubMed Abstract](#) | [Publisher Full Text](#)
 41. **F** Krügel U, Veenhoff LM, Langbein J, *et al.*: **Transport and sorting of the solanum tuberosum sucrose transporter SUT1 is affected by posttranslational modification.** *Plant Cell.* 2008; **20**(9): 2497–513.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 42. Krügel U, Kühn C: **Post-translational regulation of sucrose transporters by direct protein-protein interactions.** *Front Plant Sci.* 2013; **4**: 237.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 43. **F** Durand M, Porcheron B, Hennen N, *et al.*: **Water deficit enhances C export to the roots in *Arabidopsis thaliana* plants with contribution of sucrose transporters in both shoot and roots.** *Plant Physiol.* 2016; **170**(3): 1460–79.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 44. **F** Khadilkar AS, Yadav UP, Salazar C, *et al.*: **Constitutive and companion cell-specific overexpression of *AVP1*, encoding a proton-pumping pyrophosphatase, enhances biomass accumulation, phloem loading, and long-distance transport.** *Plant Physiol.* 2016; **170**(1): 401–14.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 45. **F** Pizzio GA, Paez-Valencia J, Khadilkar AS, *et al.*: **Arabidopsis type I proton-pumping pyrophosphatase expresses strongly in phloem, where it is required for pyrophosphate metabolism and photosynthate partitioning.** *Plant Physiol.* 2015; **167**(4): 1541–53.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 46. Leach KA, Tran TM, Slewinski TL, *et al.*: **Sucrose transporter2 contributes to maize growth, development, and crop yield.** *J Integr Plant Biol.* 2017; **59**(6): 390–408.
[PubMed Abstract](#) | [Publisher Full Text](#)
 47. **F** Wingenter K, Schulz A, Wormit A, *et al.*: **Increased activity of the vacuolar monosaccharide transporter TMT1 alters cellular sugar partitioning, sugar signaling, and seed yield in Arabidopsis.** *Plant Physiol.* 2010; **154**(2): 665–77.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 48. Griffiths CA, Paul MJ, Foyer CH: **Metabolite transport and associated sugar signalling systems underpinning source/sink interactions.** *Biochim Biophys Acta.* 2016; **1857**(10): 1715–25.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 49. Crafts AS, Crisp CE: **Phloem transport in plants.** *Bioscience.* 1972; **22**: 236.
 50. Knoblauch M, Knoblauch J, Mullendore DL, *et al.*: **Testing the Münch hypothesis of long distance phloem transport in plants.** *eLife.* 2016; **5**: pii: e15341.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 51. Knoblauch M, Peters WS: **Münch, morphology, microfluidics - our structural problem with the phloem.** *Plant Cell Environ.* 2010; **33**(9): 1439–52.
[PubMed Abstract](#) | [Publisher Full Text](#)
 52. de Schepper V, de Swaef T, Bauweraerts I, *et al.*: **Phloem transport: a review of mechanisms and controls.** *J Exp Bot.* 2013; **64**(16): 4839–50.
[PubMed Abstract](#) | [Publisher Full Text](#)
 53. de Schepper V, Steppe K: **Development and verification of a water and sugar transport model using measured stem diameter variations.** *J Exp Bot.* 2010; **61**(8): 2083–99.
[PubMed Abstract](#) | [Publisher Full Text](#)
 54. Jensen KH, Liesche J, Bohr T, *et al.*: **Universality of phloem transport in seed plants.** *Plant Cell Environ.* 2012; **35**(6): 1065–76.
[PubMed Abstract](#) | [Publisher Full Text](#)
 55. Hölttä T, Kurppa M, Nikinmaa E: **Scaling of xylem and phloem transport capacity and resource usage with tree size.** *Front Plant Sci.* 2013; **4**: 496.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 56. **F** Mencuccini M, Hölttä T: **The significance of phloem transport for the speed with which canopy photosynthesis and belowground respiration are linked.** *New Phytol.* 2010; **185**(1): 189–203.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 57. Petit G, Crivellaro A: **Comparative axial widening of phloem and xylem conduits in small woody plants.** *Trees.* 2014; **28**(3): 915–21.
[Publisher Full Text](#)
 58. **F** Jyske T, Hölttä T: **Comparison of phloem and xylem hydraulic architecture in *Picea abies* stems.** *New Phytol.* 2015; **205**(1): 102–15.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 59. Woodruff DR: **The impacts of water stress on phloem transport in Douglas-fir trees.** *Tree Physiol.* 2014; **34**(1): 5–14.
[PubMed Abstract](#) | [Publisher Full Text](#)
 60. Liesche J, Pace MR, Xu Q, *et al.*: **Height-related scaling of phloem anatomy and the evolution of sieve element end wall types in woody plants.** *New Phytol.* 2017; **214**(1): 245–56.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 61. **F** Paljakka T, Jyske T, Lintunen A, *et al.*: **Gradients and dynamics of inner bark and needle osmotic potentials in Scots pine (*Pinus sylvestris* L.) and Norway spruce (*Picea abies* L. Karst).** *Plant Cell Environ.* 2017; **40**(10): 2160–73.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 62. Liesche J, Windt C, Bohr T, *et al.*: **Slower phloem transport in gymnosperm trees can be attributed to higher sieve element resistance.** *Tree Physiol.* 2015; **35**(4): 376–86.
[PubMed Abstract](#) | [Publisher Full Text](#)
 63. Demmig-Adams B, Stewart JJ, Adams WW Jr: **Environmental regulation of intrinsic photosynthetic capacity: an integrated view.** *Curr Opin Plant Biol.* 2017; **37**: 34–41.
[PubMed Abstract](#) | [Publisher Full Text](#)
 64. Patrick JW, Offler CE: **Post-sieve element transport of photoassimilates in sink regions.** *J Exp Bot.* 1996; **47** Spec No: 1165–77.
[PubMed Abstract](#) | [Publisher Full Text](#)
 65. den Outer RW: **Histological investigations of the secondary phloem of gymnosperms.** Meded Landbouwhogeschool Wageningen. H. Veenman & Zonen N.V.: Wageningen. 1967; 67–7: 1-119.
[Reference Source](#)
 66. **F** Minchin PEH, Lacointe A: **Consequences of phloem pathway unloading/reloading on equilibrium flows between source and sink: A modelling approach.** *Functional Plant Biol.* 2017; **44**: 507–514.
[Publisher Full Text](#) | [F1000 Recommendation](#)
 67. **F** Gould N, Minchin PEH, Thorpe MR: **Direct measurements of sieve element hydrostatic pressure reveal strong regulation after pathway blockage.** *Functional Plant Biol.* 2004; **31**: 987–993.
[Publisher Full Text](#) | [F1000 Recommendation](#)

68. Knoblauch M, Froelich DR, Pickard WF, *et al.*: **SEORious business: structural proteins in sieve tubes and their involvement in sieve element occlusion.** *J Exp Bot.* 2014; **65**(7): 1879–93.
[PubMed Abstract](#) | [Publisher Full Text](#)
69. **F** Froelich DR, Mullendore DL, Jensen KH, *et al.*: **Phloem ultrastructure and pressure flow: Sieve-Element-Occlusion-Related agglomerations do not affect translocation.** *Plant Cell.* 2011; **23**(12): 4428–45.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
70. **F** Mullendore DL, Windt CW, van As H, *et al.*: **Sieve tube geometry in relation to phloem flow.** *Plant Cell.* 2010; **22**(3): 579–93.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
71. Patrick JW, Botha FC, Birch RG: **Metabolic engineering of sugars and simple sugar derivatives in plants.** *Plant Biotechnol J.* 2013; **11**(2): 142–56.
[PubMed Abstract](#) | [Publisher Full Text](#)
72. Turgeon R, Wolf S: **Phloem transport: cellular pathways and molecular trafficking.** *Annu Rev Plant Biol.* 2009; **60**: 207–21.
[PubMed Abstract](#) | [Publisher Full Text](#)
73. **F** Will T, Tjallingii WF, Thönnessen A, *et al.*: **Molecular sabotage of plant defense by aphid saliva.** *Proc Natl Acad Sci U S A.* 2007; **104**(25): 10536–41.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
74. **F** Liu DD, Chao WM, Turgeon R: **Transport of sucrose, not hexose, in the phloem.** *J Exp Bot.* 2012; **63**(11): 4315–20.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
75. Dinant S, Kehr J: **Sampling and analysis of phloem sap.** *Methods Mol Biol.* 2013; **953**: 185–94.
[PubMed Abstract](#) | [Publisher Full Text](#)
76. Jyske TM, Suuronen JP, Pranovich AV, *et al.*: **Seasonal variation in formation, structure, and chemical properties of phloem in *Picea abies* as studied by novel microtechniques.** *Planta.* 2015; **242**(3): 613–29.
[PubMed Abstract](#) | [Publisher Full Text](#)
77. Kaiser C, Kilburn MR, Clode PL, *et al.*: **Exploring the transfer of recent plant photosynthates to soil microbes: mycorrhizal pathway vs direct root exudation.** *New Phytol.* 2015; **205**(4): 1537–51.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
78. Zakhartsev M, Medvedeva I, Orlov Y, *et al.*: **Metabolic model of central carbon and energy metabolisms of growing *Arabidopsis thaliana* in relation to sucrose translocation.** *BMC Plant Biol.* 2016; **16**(1): 262.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
79. Schulz A: **Long-distance trafficking: lost in transit or stopped at the gate?** *Plant Cell.* 2017; **29**(3): 426–30.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
80. Rodríguez-Celma J, Ceballos-Laita L, Grusak MA, *et al.*: **Plant fluid proteomics: Delving into the xylem sap, phloem sap and apoplastic fluid proteomes.** *Biochim Biophys Acta.* 2016; **1864**(8): 991–1002.
[PubMed Abstract](#) | [Publisher Full Text](#)
81. Guan D, Yan B, Thieme C, *et al.*: **PlaMoM: a comprehensive database compiles plant mobile macromolecules.** *Nucleic Acids Res.* 2017; **45**(D1): D1021–D1028.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
82. **F** Paultre DS, Gustin MP, Molnar A, *et al.*: **Lost in transit: long-distance trafficking and phloem unloading of protein signals in *Arabidopsis* homografts.** *Plant Cell.* 2016; **28**(9): 2016–2025, pii: tpc.00249.2016.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
83. Banerjee AK, Chatterjee M, Yu Y, *et al.*: **Dynamics of a mobile RNA of potato involved in a long-distance signaling pathway.** *Plant Cell.* 2006; **18**(12): 3443–57.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
84. Ruiz-Medrano R, Xoconostle-Cázares B, Lucas WJ: **Phloem long-distance transport of CmNACP mRNA: implications for supracellular regulation in plants.** *Development.* 1999; **126**(20): 4405–19.
[PubMed Abstract](#)
85. Thieme CJ, Rojas-Triana M, Stecyk E, *et al.*: **Endogenous *Arabidopsis* messenger RNAs transported to distant tissues.** *Nat Plants.* 2015; **1**(4): 15025.
[PubMed Abstract](#) | [Publisher Full Text](#)
86. Zhang W, Thieme CJ, Kollwig G, *et al.*: **tRNA-related sequences trigger systemic mRNA transport in plants.** *Plant Cell.* 2016; **28**(6): 1237–49.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
87. Huen AK, Rodríguez-Medina C, Ho AYY, *et al.*: **Long-distance movement of phosphate starvation-responsive microRNAs in *Arabidopsis*.** *Plant Biol (Stuttg).* 2017; **19**(4): 643–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
88. Barbaglia AM, Hoffmann-Benning S: **Long-distance lipid signaling and its role in plant development and stress response.** *Subcell Biochem.* 2016; **86**: 339–61.
[PubMed Abstract](#) | [Publisher Full Text](#)
89. Patrick JW: **PHLOEM UNLOADING: Sieve element unloading and post-sieve element transport.** *Annu Rev Plant Physiol Mol Biol.* 1997; **48**: 191–222.
[PubMed Abstract](#) | [Publisher Full Text](#)
90. Patrick JW, Tyeramn SD, van Bel AJE: **Long-distance transport.** In *Biochemistry & Molecular Biology of Plants*, eds Buchanan BB, Gruissem W, Jones RL, John Wiley and Sons, Oxford UK, 2015; 658–710.
[Reference Source](#)
91. Cram WJ: **Mannitol transport and suitability as an osmoticum in root cells.** *Physiol Plant.* 1984; **61**(3): 396–404.
[Publisher Full Text](#)
92. Giaquinta RT: **Phloem loading of sucrose.** *Annu Rev Plant Physiol.* 1983; **34**: 347–87.
[Publisher Full Text](#)
93. **F** Le Hir R, Spinner L, Klemens PA, *et al.*: **Disruption of the sugar transporters *AtSWEET11* and *AtSWEET12* affects vascular development and freezing tolerance in *Arabidopsis*.** *Mol Plant.* 2015; **8**(11): 1687–90.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
94. **F** Carpaneto A, Geiger D, Bamberg E, *et al.*: **Phloem-localized, proton-coupled sucrose carrier ZmSUT1 mediates sucrose efflux under the control of the sucrose gradient and the proton motive force.** *J Biol Chem.* 2005; **280**(22): 21437–43.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
95. Milne RJ, Perroux JM, Rae AL, *et al.*: **Sucrose transporter localization and function in phloem unloading in developing stems.** *Plant Physiol.* 2017; **173**(2): 1330–41.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
96. **F** Hafke JB, Höll S, Kühn C, *et al.*: **Electrophysiological approach to determine kinetic parameters of sucrose uptake by single sieve elements or phloem parenchyma cells in intact *Vicia faba* plants.** *Front Plant Sci.* 2013; **4**: 274.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
97. **F** Zhang LY, Peng YB, Pelleschi-Travier S, *et al.*: **Evidence for apoplastic phloem unloading in developing apple fruit.** *Plant Physiol.* 2004; **135**(1): 574–86.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
98. Zhang XY, Wang XL, Wang XF, *et al.*: **A shift of Phloem unloading from symplasmic to apoplastic pathway is involved in developmental onset of ripening in grape berry.** *Plant Physiol.* 2006; **142**(1): 220–32.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
99. Hu L, Sun H, Li R, *et al.*: **Phloem unloading follows an extensive apoplastic pathway in cucumber (*Cucumis sativus* L.) fruit from anthesis to marketable maturing stage.** *Plant Cell Environ.* 2011; **34**(11): 1835–48.
[PubMed Abstract](#) | [Publisher Full Text](#)
100. Chen C, Yuan Y, Zhang C, *et al.*: **Sucrose phloem unloading follows an apoplastic pathway with high sucrose synthase in *Actinidia* fruit.** *Plant Sci.* 2017; **255**: 40–50.
[PubMed Abstract](#) | [Publisher Full Text](#)
101. **F** Hackel A, Schauer N, Carrari F, *et al.*: **Sucrose transporter LeSUT1 and LeSUT2 inhibition affects tomato fruit development in different ways.** *Plant J.* 2006; **45**(2): 180–92.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
102. Zanon L, Falchi R, Santi S, *et al.*: **Sucrose transport and phloem unloading in peach fruit: potential role of two transporters localized in different cell types.** *Physiol Plant.* 2015; **154**(2): 179–93.
[PubMed Abstract](#) | [Publisher Full Text](#)
103. Wang ZP, Deloire A, Carbonneau A, *et al.*: **An *in vivo* experimental system to study sugar phloem unloading in ripening grape berries during water deficiency stress.** *Ann Bot.* 2003; **92**(4): 523–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
104. Ruan YL, Patrick JW, Brady C: **Protoplast hexose carrier activity is a determinate of genotypic difference in hexose storage in tomato fruit.** *Plant Cell Environ.* 1997; **20**(3): 341–9.
[Publisher Full Text](#)
105. Hayes MA, Davies C, Dry IB: **Isolation, functional characterization, and expression analysis of grapevine (*Vitis vinifera* L.) hexose transporters: differential roles in sink and source tissues.** *J Exp Bot.* 2007; **58**(8): 1985–97.
[PubMed Abstract](#) | [Publisher Full Text](#)
106. McCurdy DW, Dibley S, Cahyanegara R, *et al.*: **Functional characterization and RNAi-mediated suppression reveals roles for hexose transporters in sugar accumulation by tomato fruit.** *Mol Plant.* 2010; **3**(6): 1049–63.
[PubMed Abstract](#) | [Publisher Full Text](#)
107. Cheng JT, Li X, Yao FZ, *et al.*: **Functional characterization and expression analysis of cucumber (*Cucumis sativus* L.) hexose transporters, involving carbohydrate partitioning and phloem unloading in sink tissues.** *Plant Sci.* 2015; **237**: 46–56.
[PubMed Abstract](#) | [Publisher Full Text](#)
108. Jin Y, Ni DA, Ruan YL: **Posttranslational elevation of cell wall invertase activity by silencing its inhibitor in tomato delays leaf senescence and increases seed weight and fruit hexose level.** *Plant Cell.* 2009; **21**(7): 2072–89.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
109. **F** Zanon MI, Osorio S, Nunes-Nesi A, *et al.*: **RNA interference of *LIN5* in tomato confirms its role in controlling Brix content, uncovers the influence of sugars on the levels of fruit hormones, and demonstrates the importance of sucrose cleavage for normal fruit development and fertility.** *Plant Physiol.* 2009; **150**(3): 1204–18.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
110. Shi J, Wang J, Li R, *et al.*: **Expression patterns of genes encoding plasma membrane aquaporins during fruit development in cucumber (*Cucumis sativus* L.).** *Plant Physiol Biochem.* 2015; **96**: 329–36.
[PubMed Abstract](#) | [Publisher Full Text](#)

111. **F** Stanfield RC, Hacke UG, Laur J: **Are phloem sieve tubes leaky conduits supported by numerous aquaporins?** *Am J Bot.* 2017; **104**(5): 719–32.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
112. Patrick JW: **Does Don Fisher's high-pressure manifold model account for phloem transport and resource partitioning?** *Front Plant Sci.* 2013; **4**: 184.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
113. Fisher DB, Cash-Clark CE: **Gradients in water potential and turgor pressure along the translocation pathway during grain filling in normally watered and water-stressed wheat plants.** *Plant Physiol.* 2000; **123**(1): 139–48.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
114. **F** Ross-Elliott TJ, Jensen KH, Haaning KS, *et al.*: **Phloem unloading in Arabidopsis roots is convective and regulated by the phloem-pole pericycle.** *eLife.* 2017; **6**: pii: e24125.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
115. Gould N, Thorpe MR, Minchin PEH, *et al.*: **Solute is imported to elongating root cells of barley as a pressure driven-flow of solution.** *Functional Plant Biol.* 2004; **31**(4): 391–397.
[Publisher Full Text](#)
116. Warmbrodt RD: **Solute concentrations in the phloem and apex of the root of Zea mays.** *Am J Bot.* 1987; **74**(3): 394–402.
[Publisher Full Text](#)
117. Zhang W, Zhou Y, Dibley KE, *et al.*: **Review: Nutrient loading of developing seeds.** *Functional Plant Biol.* 2007; **34**(4): 314–331.
[Publisher Full Text](#)
118. **F** Sosso D, Luo D, Li QB, *et al.*: **Seed filling in domesticated maize and rice depends on SWEET-mediated hexose transport.** *Nat Genet.* 2015; **47**(12): 1489–93.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
119. Ritchie RJ, Fieuw-Makaroff S, Patrick JW: **Sugar retrieval by coats of developing seeds of Phaseolus vulgaris L. and Vicia faba L.** *Plant Cell Physiol.* 2003; **44**(2): 163–72.
[PubMed Abstract](#) | [Publisher Full Text](#)
120. **F** Zhou Y, Qu H, Dibley KE, *et al.*: **A suite of sucrose transporters expressed in coats of developing legume seeds includes novel pH-independent facilitators.** *Plant J.* 2007; **49**(4): 750–64.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
121. **F** Chen LQ, Lin IW, Qu X, *et al.*: **A cascade of sequentially expressed sucrose transporters in the seed coat and endosperm provides nutrition for the Arabidopsis embryo.** *Plant Cell.* 2015; **27**(3): 607–19.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
122. **F** Ma L, Zhang D, Miao Q, *et al.*: **Essential role of sugar transporter OsSWEET11 during the early stage of rice grain filling.** *Plant Cell Physiol.* 2017; **58**(5): 863–73.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
123. Walker NA, Patrick JW, Zhang W, *et al.*: **Efflux of photosynthate and acid from developing seed coats of Phaseolus vulgaris L: A chemiosmotic analysis of pump-driven efflux.** *J Exp Bot.* 1995; **46**(5): 539–49.
[Publisher Full Text](#)
124. Zhou Y, Setz N, Niemietz C, *et al.*: **Aquaporins and unloading of phloem-imported water in coats of developing bean seeds.** *Plant Cell Environ.* 2007; **30**(12): 1566–77.
[PubMed Abstract](#) | [Publisher Full Text](#)
125. **F** Hayashi H, Ishikawa-Sakurai J, Murai-Hatano M, *et al.*: **Aquaporins in developing rice grains.** *Biosci Biotechnol Biochem.* 2015; **79**(9): 1422–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

- 1. William W Adams**
Department of Ecology and Evolutionary Biology, University of Colorado, Boulder, USA
Competing Interests: No competing interests were disclosed.
- 2. Brian Ayre**
Department of Biological Sciences, University of North Texas, Texas, USA
Competing Interests: No competing interests were disclosed.
- 3. Robert Turgeon**
Plant Biology, Cornell University, New York, USA
Competing Interests: No competing interests were disclosed.
- 4. Friedrich Kragler**
Department of Intercellular Macromolecular Transport, Max Planck Institute of Molecular Plant Physiology, Potsdam, Germany
Competing Interests: No competing interests were disclosed.
- 5. Antia Rodriguez-Villalon**
Department of Biology, Swiss Federal Institute of Technology (ETH-Z), Zurich, Switzerland
Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research