CORRESPONDENCE

Can circulating tumor DNA be used for direct and early stage cancer detection? [version 1; peer review: 2 approved]

Eleftherios P. Diamandis ¹⁻³, Clare Fiala³¹ Department of Clinical Biochemistry, University Health Network, Toronto, Ontario, Canada² Department of Laboratory Medicine and Pathobiology, University of Toronto, Toronto, Ontario, Canada³ Department of Pathology and Laboratory Medicine, Mount Sinai Hospital, Toronto, Ontario, Canada

V1 First published: 13 Dec 2017, 6:2129
<https://doi.org/10.12688/f1000research.13440.1>
Latest published: 13 Dec 2017, 6:2129
<https://doi.org/10.12688/f1000research.13440.1>

Abstract

In the August 16th issue of Science Translational Medicine, Phallen et al propose a method for early cancer diagnosis by using circulating tumor DNA (1). One major advance of this paper includes optimized sequencing of cell-free/circulating tumor DNA (ctDNA) without knowledge of tumor mutations. Evaluation of 200 patients with colorectal, breast, lung and ovarian cancer revealed mutations in ctDNA in approx. 60-70% of all patients, including stage 1 and stage 2 disease. If this data can be reproduced in asymptomatic individuals, they will likely have a major impact on early cancer detection and patient outcomes. In this commentary, we examine the feasibility of this approach for detecting small, asymptomatic tumors, based on previously published empirical data.

Keywords

Circulating tumor, Cancer diagnosis, Cancer detection, Tumor measurement, Colorectal, breast, lung, ovarian cancer data

Open Peer Review

Approval Status

	1	2
version 1
13 Dec 2017	view	view

1. **Nicholas J. Wald** , Queen Mary University of London, London, UK

Robert W Old, Queen Mary University of London, London, UK

2. **Catherine Alix-Panabières**, University Medical Center of Montpellier, Montpellier, France

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Eleftherios P. Diamandis (Eleftherios.diamandis@sinaihhealthsystem.ca)

Author roles: **Diamandis EP:** Conceptualization, Data Curation, Formal Analysis, Funding Acquisition, Project Administration, Resources, Supervision, Validation, Writing – Original Draft Preparation, Writing – Review & Editing; **Fiala C:** Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This work was funded by a grant to EPD from the Canadian Cancer Society (Grant #703873)
The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2017 Diamandis EP and Fiala C. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Diamandis EP and Fiala C. **Can circulating tumor DNA be used for direct and early stage cancer detection? [version 1; peer review: 2 approved]** F1000Research 2017, 6:2129 <https://doi.org/10.12688/f1000research.13440.1>

First published: 13 Dec 2017, 6:2129 <https://doi.org/10.12688/f1000research.13440.1>

Correspondence

Phallen *et al.* recently proposed a method for early cancer diagnosis by using circulating tumor DNA¹. One major advance of this paper includes optimized sequencing of cell-free/circulating tumor DNA (ctDNA) without knowledge of tumor mutations. Evaluation of 200 patients with colorectal, breast, lung and ovarian cancer revealed mutations in ctDNA in approximately 60–70% of all patients, including stage 1 and stage 2 disease. If this data can be reproduced in asymptomatic individuals, they will likely have a major impact on early cancer detection and patient outcomes. In this correspondence, we examine if this approach is effective in detecting small, asymptomatic tumors, based on previously published empirical data.

An important question is the desirable size of a tumor to be detected with such methods. For this discussion, we will use data mostly from breast cancer, for which there is ample experience from screening programs. Some organizations set a tumor diameter of 1 mm as the goal of early detection² since tumors of this size are localized, less complex and likely to be cured by radical resection. The literature suggests that the chances of progression of these small, 1 mm diameter tumors, is less than 1%³. However, such small tumors could be found in many asymptomatic individuals and this may lead to over-diagnosis and over-treatment³. Consequently, we also include larger tumors in this discussion. It could be argued that 5 mm in diameter is an optimal tumor size for early and curable cancer detection. At 5 mm diameter, the chances of this tumor progressing are small (around 6%)³, and currently, these small lesions are only detectable by mammography (or other imaging modalities) in only 26% of cases⁴. We also suggest that detection of a 10 mm diameter tumor may not be an advance in breast cancer screening, since at this size, the chances of progression increase considerably (to about 50%), and most of these tumors (> 90%), are currently detected by mammographic screening.

We further assume that the circulating free DNA concentration (cfDNA) in individuals without cancer, or very small cancers (ctDNA), is about 5 ng/mL on average⁵, which is equivalent to about 6,000 whole haploid genomes per 4 mL of plasma (10 mL blood draw). For this discussion we also use well-established measurements of tumor volume and cellularity. A tumor of approximately 12.5 mm in diameter, weights approx. 1 gram, has a volume of 1 cm³ and contains approx. 100 million to 1 billion cells⁶. According to recent estimates of ctDNA and tumor volume, a 10 gram tumor has a volume of 10 cm³⁷. For such tumors, the average percent fraction of mutant DNA has been reported to be 0.1 %, or 1 mutant DNA molecule per 1,000 non-mutated DNA molecules in the circulation⁷.

Based on these published assumptions, we constructed [Table 1](#).

[Table 1](#) shows the minimum tumor measurements required for 4 mL of plasma to contain at least 1 mutated genome. The percent fraction of mutant DNA is approximately 1 in 10,000 (0.01%) which corresponds to a tumor volume of 1 cm³ or 12.5 mm in diameter (assuming a spherical nodule). Most, if not all, of these tumors are currently detectable by mammographic screening or other imaging modalities, which have an approximate limit of detection of about 4 mm diameter⁸. [Table 1](#) also shows that when the tumor diameter drops below 10 mm, the chances of this method working are minimal, since there will be not enough tumor DNA (at least one copy) in 4 mL of plasma to make detection possible. Consequently, [Table 1](#) predicts that 5 mm diameter tumors will not be detected due to this sampling error. These estimates are corroborated by the data presented by Phallen *et al.*¹. They used tumors of varying sizes, stages and types, but universally, the mutant fraction of ctDNA is never less than 0.01%, the approximate threshold for 1 genome copy in 4 mL of plasma. We do not know if the patients included in the study have been preselected to have a

Table 1. Tumor measurements and chances of progression or detection by mammographic screening (assuming spherical tumor).

Tumor diameter, mm	Tumor weight, mg	Tumor Volume, mL(cm ³)	Number of cells	%Fraction of mutant DNA	Number of genomes per 10 mL blood/4 mL plasma	Chance of progression ³	Mammographic screen sensitivity ⁴
27	10,000	10¹	1,000,000,000	1:1,000	6		
12.5	1,000	1²	100,000,000	1:10,000	0.6		
10	500	0.5	50,000,000	1:20,000	0.3	50%	91%
6	125	0.12	12,000,000	1:80,000	<0.1		
5	62	0.06	6,000,000	1:160,000	<0.1	6%	26%
3	16	0.015	1,000,000	1:640,000	<0.1		
2	4	0.0035	400,000	1:2,600,000	<0.1		
1.1	1	0.0008	100,000	1:10,000,000	<0.1	0.05%	

1. As reported in Ref [7](#); bold font indicates experimental data. Other data were calculated by extrapolation

2. As reported in Ref [6](#)

3. As reported in Ref [3](#)

4. As reported in Ref. [4](#)

specific cut-off of percent mutant DNA in the circulation. If this is the case, the data will be biased towards detection of tumors which are rather large (> 10 mm in diameter and have a mutant fraction of > 0.01%). In fact, >95% of their cases have mutant fraction > 0.1% (Figure 5 from Phallen *et al.*¹ which suggests tumor sizes of > 27 mm (Table 1). We anticipate that in a real case scenario, small and asymptomatic tumors will likely yield percent mutant DNA fraction of much less than 0.1%, thus making detection of these tumors highly unlikely, due to sampling error.

The sensitivity of the method proposed by Phallen *et al.* is further compromised by the number of genes tested for mutations (currently 58 genes). The predicted maximum sensitivity is about 80%, assuming best case scenario of 1 mutation detected to signify cancer. If we assume that the overall positivity of this test is 50%, then, the overall sensitivity will likely be less than 40%. This would make screening of asymptomatic individuals less efficient, since the majority of patients will receive incorrect screening results (false negatives) due to either lack of the targeted mutations or sampling error.

The specificity of this method is claimed to be very high due to detection of only driver mutations. However, as shown in Phallen *et al.*'s Supplementary Figure S3¹, the gene with the highest sensitivity for cancer detection of all of the 4 cancer types is p53. However, there are multiple reports demonstrating mutations of p53 in apparently healthy individuals, ranging from 0.2% to 11%^{8–11}. If we assume an overall specificity of 99% and an overall sensitivity of 40% with this test, and a prevalence of cancer around 1%, then in 1,000 screened individuals, there will be 4 true positive, 6 false negative, 10 false positive and 992 true negative results. The positive predictive value (PPV) will be only 29%. At 98% specificity, the PPV further drops to 20%. Not only would more than half the patients will receive incorrect (false negative) results but about 72–80% of tested positive patients would need to undergo additional diagnostic procedures to rule-in or rule-out cancer. Additional specificity concerns may include technical noise and somatic mutations stemming from other biological phenomena. Also, in this paper, no matched DNA was employed to

differentiate somatic from germline mutations, another important limitation¹².

Our analysis shows that measurement of ctDNA for early cancer diagnosis is problematic, not only due to the limited capability of deep whole genome sequencing to reveal mutations, but because the amount of circulating tumor DNA retrieved from a 10mL blood draw will be extremely small, or even nonexistent, thus making efficient diagnosis of cancer unlikely. Our analyses generally show that when the tumor diameter drops below 10 mm, not a single mutant DNA copy will be retrieved. We do not exclude the possibility of future additional technical improvements, so that mutant DNA could somehow be extracted from the whole circulation and subjected to such deep sequencing, to increase sensitivity, provided that the specificity of the test would be preserved to close to 100%. Otherwise, under a screening scenario, these methods are unlikely to work. Our analyses also show that a 1 mm tumor is likely associated with 1 copy of ctDNA in the entire circulation, further underlining the difficulty in detection. The possibility exists that early cancer detection may be possible in specific clinical contexts or specific tumor types using this method. Also, it may be possible to combine protein and molecular markers to increase sensitivity, as shown already for pancreatic cancer¹³.

We hope that our analyses will help to further understand the opportunities and limitations of ctDNA as a cancer biomarker, especially for early cancer detection.

Competing interests

No competing interests were disclosed.

Grant information

This work was funded by a grant to EPD from the Canadian Cancer Society (Grant #703873).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- Phallen J, Sausen M, Adleff V, *et al.*: **Direct detection of early-stage cancers using circulating tumor DNA.** *Sci Transl Med.* 2017; **9**(403): pii: eaan2415.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Garber K: **Ontario institute offers new model of cancer research.** *J Natl Cancer Inst.* 2008; **100**(14): 980–982.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Countercurrants Series, Narod SA: **Disappearing breast cancers.** *Curr Oncol.* 2012; **19**(2): 59–60.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Weedon-Fekjaer H, Lindqvist BH, Vatten LJ, *et al.*: **Breast cancer tumor growth estimated through mammography screening data.** *Breast Cancer Res.* 2008; **10**(3): R41.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Wan JCM, Massie C, Garcia-Corbacho J, *et al.*: **Liquid biopsies come of age: towards implementation of circulating tumour DNA.** *Nat Rev Cancer.* 2017; **17**(4): 223–238.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Del Monte U: **Does the cell number 10⁹ still really fit one gram of tumor tissue?** *Cell Cycle.* 2009; **8**(3): 505–506.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Abbosh C, Birkbak NJ, Wilson GA, *et al.*: **Phylogenetic ctDNA analysis depicts early-stage lung cancer evolution.** *Nature.* 2017; **545**(7655): 446–451.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Schwaederle M, Husain H, Fanta PT, *et al.*: **Detection rate of actionable mutations in diverse cancers using a biopsy-free (blood) circulating tumor cell DNA assay.** *Oncotarget.* 2016; **7**(9): 9707–9717.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Gormally E, Vineis P, Matullo G, *et al.*: **TP53 and KRAS2 mutations in plasma DNA of healthy subjects and subsequent cancer occurrence: a prospective study.** *Cancer Res.* 2006; **66**(13): 6871–6876.
[PubMed Abstract](#) | [Publisher Full Text](#)

10. Fernandez-Cuesta L, Perdomo S, Avogbe PH, *et al.*: **Identification of Circulating Tumor DNA for the Early Detection of Small-cell Lung Cancer.** *EBioMedicine*. 2016; **10**: 117–123.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
11. Newman AM, Lovejoy AF, Klass DM, *et al.*: **Integrated digital error suppression for improved detection of circulating tumor DNA.** *Nat Biotechnol*. 2016; **34**(5): 547–555.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
12. Jones S, Anagnostou V, Lytle K, *et al.*: **Personalized genomic analyses for cancer mutation discovery and interpretation.** *Sci Transl Med*. 2015; **7**(283): 283ra53.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
13. Cohen JD, Javed AA, Thoburn C, *et al.*: **Combined circulating tumor DNA and protein biomarker-based liquid biopsy for the earlier detection of pancreatic cancers.** *Proc Natl Acad Sci U S A*. 2017; **114**(38): 10202–10207.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 04 January 2018

<https://doi.org/10.5256/f1000research.14593.r29016>

© 2018 Alix-Panabières C. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Catherine Alix-Panabières

Laboratory of Rare Circulating Human Cells (LCCRH), Department of Pathology and Onco-Biology, University Medical Center of Montpellier, Montpellier, France

This commentary, very well written by the expert Eleftherios P Diamandis and his colleague Clare Fiala, is based on the last article of Phallen *et al.* who described a method for early cancer diagnosis by using circulating tumor DNA (ctDNA) sequenced without knowledge of tumor mutations.

In this commentary, the authors examined step by step the feasibility of this approach for detecting small, asymptomatic tumors, based on previously published empirical data. They nicely showed that measurement of ctDNA for early cancer diagnosis is problematic, not only due to the limited capability of deep whole genome sequencing to reveal mutations, but because the amount of circulating tumor DNA retrieved from a 10mL blood draw will be extremely small, or even nonexistent, thus making efficient diagnosis of cancer unlikely.

This commentary, very well documented, is important in the liquid biopsy field and needs to be indexed to understand the current opportunities and limitations of ctDNA as a biomarker for early detection of cancer.

Is the rationale for commenting on the previous publication clearly described?

Yes

Are any opinions stated well-argued, clear and cogent?

Yes

Are arguments sufficiently supported by evidence from the published literature or by new data and results?

Yes

Is the conclusion balanced and justified on the basis of the presented arguments?

Yes

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 14 December 2017

<https://doi.org/10.5256/f1000research.14593.r29018>

© 2017 Wald N et al. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Nicholas J. Wald

Wolfson Institute of Preventive Medicine, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London, UK

Robert W Old

Wolfson Institute of Preventive Medicine, Barts and the London School of Medicine and Dentistry, Queen Mary University of London, London, UK

This paper is a commentary on a study by Phallen *et al.* who developed and applied advanced DNA analysis techniques for circulating tumor DNA from patients with untreated breast, colorectal and ovarian cancer. They were able to detect cancer in about 60-70% of patients with Stage I or Stage II disease. Diamandis and Fiala's Commentary examines the feasibility of applying these techniques to small tumors that are not reliably detectable by other means. They extrapolate original data from Phallen *et al.*, making the assumption that the number of tumor-derived genomes in the circulation is proportional to tumor volume. Diamandis and Fiala reasonably conclude that tumors smaller than about 10mm in diameter result in less than one tumor-derived genome per 10ml blood draw and are therefore, in principle, undetectable in practice. This logic is probably conservatively based because small, low grade tumors are likely to produce proportionally less ctDNA than high grade tumors. This Commentary is a useful contribution to the literature.

Is the rationale for commenting on the previous publication clearly described?

Yes

Are any opinions stated well-argued, clear and cogent?

Yes

Are arguments sufficiently supported by evidence from the published literature or by new data and results?

Yes

Is the conclusion balanced and justified on the basis of the presented arguments?

Yes

Competing Interests: No competing interests were disclosed.

We confirm that we have read this submission and believe that we have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research