OPINION ARTICLE

Midlife interventions are critical in prevention, delay, or improvement of Alzheimer's disease and vascular cognitive impairment and dementia [version 1; peer review: 2 approved]

Sam Gandy^{1,2}, Tamas Bartfai³, Graham V. Lees⁴, Mary Sano²

¹Department of Neurology and NFL Neurological Center, Icahn School of Medicine at Mount Sinai, New York, NY, 10029, USA

²Department of Psychiatry and Alzheimer's Disease Research Center, Icahn School of Medicine at Mount Sinai, New York, NY, 10029, USA

³Department of Neurochemistry, Stockholm University, Stockholm, 114 18, Sweden

⁴Corpus Alienum Oy, Kirkkonummi, 02400, Finland

V1 First published: 03 Apr 2017, 6:413
<https://doi.org/10.12688/f1000research.11140.1>

Latest published: 03 Apr 2017, 6:413
<https://doi.org/10.12688/f1000research.11140.1>

Abstract

The basic strategy for focusing exclusively on genetically identified targets for intervening in late life dementias was formulated 30 years ago. Three decades and billions of dollars later, all efforts at disease-modifying interventions have failed. Over that same period, evidence has accrued pointing to dementias as late-life clinical phenotypes that begin as midlife pathologies. Effective prevention therefore may need to begin in midlife, in order to succeed. No current interventions are sufficiently safe to justify their use in midlife dementia prevention trials. Observational studies could be informative in testing the proposal that amyloid imaging and *APOEε4* genotype can predict those who are highly likely to develop Alzheimer's disease and in whom higher risk interventions might be justifiable. A naturally occurring, diet-responsive cognitive decline syndrome occurs in canines that closely resembles human Alzheimer's. Canine cognitive dysfunction could be useful in estimating how early intervention must begin in order to succeed. This model may also help identify and assess novel targets and strategies. New approaches to dementia prevention are urgently required, since none of the world's economies can sustain the costs of caring for this epidemic of brain failure that is devastating half of the over 85-year-olds globally.

Keywords

dementia, cognition, cognitive decline, cognitive impairment, amyloid, tau

Open Peer Review

Approval Status

	1	2
version 1
03 Apr 2017	view	view

1. **W.Sue T Griffin**, University of Arkansas for Medical Sciences, Little Rock, USA
Orwa Aboud, University of Arkansas for Medical Sciences, Little Rock, USA
2. **Domenico Pratico**, Temple University, Philadelphia, USA

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Sam Gandy (samuel.gandy@mssm.edu)

Competing interests: No competing interests were disclosed. The opinions expressed here are exclusively those of the authors.

Grant information: MS and SG would like to acknowledge the support of National Institute on Aging P50 Alzheimer's Disease Research Center (AG05138) to MS. SG would also like to acknowledge the support of the Louis B. Mayer Foundation, the Georgianne and Doctor Reza Khatib Foundation, VA MERIT (I01RX002333 and I01RX000684), and NIH Accelerating Medicines Partnership (U01AG046170) to Eric Schadt, Icahn School of Medicine.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2017 Gandy S *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. Data associated with the article are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

How to cite this article: Gandy S, Bartfai T, Lees GV and Sano M. **Midlife interventions are critical in prevention, delay, or improvement of Alzheimer's disease and vascular cognitive impairment and dementia [version 1; peer review: 2 approved]** F1000Research 2017, 6:413 <https://doi.org/10.12688/f1000research.11140.1>

First published: 03 Apr 2017, 6:413 <https://doi.org/10.12688/f1000research.11140.1>

We are not winning the fight against Alzheimer's disease

The first generation of the “amyloidocentric” approach to Alzheimer's has recently drawn to a close, and we are left with the same approved symptomatic treatments that we have had for the past 30 years (i.e., cholinesterase inhibitors). The Alzheimer's disease-modifying drug discovery field remains at a perfect 100% failure rate when it comes to new approvable disease-modifying interventions. The two most promising amyloid-reducing interventions of 2016, solanezumab¹ and verubecestat², recently failed to modify decline in mild Alzheimer's disease (AD) and were abandoned.

This is a very serious situation for society, as the disease burden of Alzheimer's continues to skyrocket globally, yet the private profit-based pharmaceutical companies cannot, and will not, continue working on disease modifying drugs for Alzheimer's unless convincing new scientific avenues are opened. What this means is that new drug targets must be identified, but for this to happen, we must be open-minded about the trap laid by the early success of identifying genes and molecular mechanisms of the familial Alzheimer's patients. These early targets presented challenges in druggability, but for many years, these targets seemed scientifically rock solid. Now, this certainty and billions of dollars in trial funding are gone, and many large, prestigious pharmaceutical companies have closed their Alzheimer's drug discovery programs.

It is time to admit that we are experiencing a rare phenomenon in drug development where the molecular mechanisms uncovered in familial cases of the disease have not helped us manage the sporadic form of the disease. In order to explain our repeated failures, we first blamed the quality of the drug candidates. More recently, we put forward a “kinetic argument”, causing trials to begin earlier and earlier in the course of the disease whilst making little or no effort to identify alternative or additional pathophysiologies beyond amyloidosis and tauopathy. The “kinetic argument” permitted us to remain focused on genetically-predicted drug targets as the most important drug targets, bringing to a halt virtually all other research. As those genetically derived targets continue to fail, experts are commonly overheard to say, “Of course that drug failed: the trial started too late in the progression of the disease; no one expected that to work.” This is revisionist history; it was only five years ago when scientists at several major pharma houses were convinced that the odds of success were high enough to invest \$50–100 million for phase 1 and 2 clinical trials, or up to \$1 billion each for phase 3 trials, even though each will take 4 years and at least 3 separate successful iterations are required. What was truly surprising to the pharmaceutical industry was that immunotherapies and enzyme inhibitors converging on the same target through entirely independent mechanisms yielded failure after failure, with no new insights. The situation was so surprising that most experts hid their disappointment. As recently as last year, clinicians were telling patients and families how these “anti-amyloid” antibodies and BACE inhibitors were “the most promising drugs ever to enter the pipeline”. To now turn around and say, “no one ever expected these drugs to work” is both disingenuous and hurtful to patients, who respond saying, “If no one expected that drug to work, then why did you recommend the trial to me?”

Each successively earlier trial failure condemns doctors and patients to another 5 to 7 years of purgatory while the next trial iteration moves five clicks earlier in enrollment age. Yet these trial designs move forward without sufficient ability to define the molecular pathology with precision at the level of the individual patient. This argues not only for diversifying the disease-modifying portfolio but also for redoubling efforts on symptomatic interventions that are also easier to get through the regulatory process.

Can we pinpoint how early is “early enough”?

Perhaps we should seek specific, empirical data about how early is “early enough” in humans. Patients with epilepsy and *APOEε4* alleles do not develop Alzheimer's yet they deposit plaques in their early 40s³. This rare but surprisingly early phenomenon argues strongly that we should initiate intervention to prevent amyloidosis or tauopathy not just a little earlier, as we are doing now, but far earlier; in other words, we should intervene in midlife, not later in life. This dovetails well with evidence that midlife risk factors lead to a late-life phenotypes. Midlife-onset hypertension is a risk factor for AD in late life; late life-onset hypertension appears to be *protective* of cognition⁴. What sorts of interventions might these be? Dietary interventions with small effect size but employed over decades might have important cumulative effects. Vaccines may provide lifelong or very long interventions when the proper antigens and adjuvants are identified and employed.

What might we do to refine our guess at what might be truly “early enough” for us to intervene? One might design an observational study of *APOEε4* carriers in their 40s^{3,5}. Annual amyloid and tau imaging of *APOEε4* carriers could be used to identify those with evidence, or at highest risk, of progression⁵. Once an *APOEε4* carrier becomes amyloid-imaging positive, one could imagine entering them either into an observational study or into an authentic trial employing reducers of amyloid, tau, or both. An advantage of this design is the possibility that for serially imaged subjects observed to change from negative to positive proteinopathy imaging, it will be possible to know that the proteinopathy has been present for 12 months or less. If we jump back in age as far as the mid 40s, and show that we can engage the proteinopathy targets at that early point, but yet *still* fail clinically, that will be a strong indication that “anti-proteinopathy-only” will never succeed.

Mixed pathology may be the most common underpinning for dementia

It is worth keeping in mind that even if we have impact on Alzheimer's pathology, the frequent concurrent presence of multiple pathologies will continue to confound. These days, there is more research on the relationship between vascular cognitive impairment and dementia (VCID) and Alzheimer's dementia, but not yet enough, even though defining this is probably profoundly important. Among African Americans with clinical Alzheimer's, 70% have mixed pathology at autopsy⁶. Despite much attention being given to insulin signaling and brain proteinopathy, virtually all the accumulated data indicate that dementia of type-2 diabetes is not Alzheimer's but is primarily vascular in origin⁷. Synucleinopathy, which is present in about 1/3 of Alzheimer's patients, induces a plethora of epigenetic changes in the brain transcriptome⁸. This means that, currently, perfect antemortem diagnosis is sometimes impossible. Certainly, the more mixed

dementia subjects there are in a trial intended to assess the efficacy of a drug that only treats Alzheimer's, the lower the sensitivity to see a signal will be. There is a good chance that each of the various underlying pathogenetic mechanisms will have their own optimum time window for intervention, and that this will have to be factored in as well. On the other hand, improved midlife cardiovascular health could benefit cognition and may delay *all* causes of dementia.

Immunology of cognitive decline in Alzheimer's disease

While genetics and, more recently, multi-scale network-level genomics⁹ have taught us, and continue to teach us, much about the molecular pathology of Alzheimer's, imaging and pathology have taught us just as much about the disappointingly poor clinicopathological and clinicoradiological correlation in this illness. Cognitive decline is poorly predicted by amyloid imaging in isolation, as was foretold by the Religious Orders study some 20 years ago¹⁰.

Are there truly new drug targets? What might we be missing in our formulation of the pathogenesis of sporadic Alzheimer's? Of the two dozen genes linked by genome-wide association studies (GWAS), two thirds are lipid- or immune-related¹¹, raising the possibility of a neuroinflammatory/neurodegenerative dementia-causing pathway that might be worthy of further research. DeStrooper has recently proposed an alternative formulation of AD as a clinical umbrella under which "feed-forward" pathogenesis scenarios lie (e.g., inflammation causes or exacerbates tauopathy; then, in turn, tauopathy aggravates inflammation)¹². This model fits the existing data at least as well as the classical linear amyloid hypothesis and explains some of the holes in the current predictive models. DeStrooper's formulation¹² includes scenarios where anti-proteinopathy drugs alone would probably be inadequate. Along the same lines, the *CRI* risk polymorphism is associated with increased risk for clinical Alzheimer's but in the setting of progression-related reduction in amyloidosis^{13,14}.

One other model might be worth investigating: Canine cognitive dysfunction (CCD)¹⁵ is the only naturally occurring mammalian dementia to mimic Alzheimer's disease. In CCD, diet and lifestyle have measurable impact on disease progression¹⁵. The CCD model could contribute to our understanding of how early intervention must begin in order to be effective. And with CCD, one could test drugs as prophylaxis *in vivo*. In the meantime, the importance and potential benefit of safe and easy diet and lifestyle changes for humans should be a topic of much stronger advocacy. In fact, strong evidence that better cardiovascular health reduces prevalence of both Alzheimer's and VCID is already beginning to emerge and

must be a clear part of patient education by clinicians^{16–23}. To its credit, the April 2017 issue of *Scientific American* heralds "Success in the Fight Against Alzheimer's", a review of dementia risk reduction benefit that can be realized through modification of diet and lifestyle beginning in midlife²⁴.

There may be more "unknown unknowns" yet to be revealed

Finally, it is important to emphasize that researchers tackling Alzheimer's and other dementias must remain clear-eyed, challenged, and worried that there are still many gaps in our understanding. A simple backward jump to earlier intervention may require 5 more iterations at the current pace, if indeed we should be intervening when subjects are in their 40s. If the disease can be primarily driven by lipid pathology or inflammatory pathology even with little or no protein aggregate pathology as could be inferred from the nature of the GWAS hits¹¹, and from the variability in amyloid-first phenotypic sequences vs. neurodegeneration-first phenotypic sequences²⁵, then efforts focused solely on purging clumped proteins from brains in which they may have lingered silently for decades may always come up short.

Acknowledgements

Dedicated to the memory of Kerstin Iverfeldt, PhD (1957–2017) who made important contributions to our understanding of the biology of APP and the amyloid β peptide.

Author contributions

All authors contributed to drafting and editing.

SG prepared the first draft and the final submitted version.

Competing interests

No competing interests were disclosed. The opinions expressed here are exclusively those of the authors.

Grant information

MS and SG would like to acknowledge the support of National Institute on Aging P50 Alzheimer's Disease Research Center (AG05138) to MS. SG would also like to acknowledge the support of the Louis B. Mayer Foundation, the Georgianne and Doctor Reza Khatib Foundation, VA MERIT (I01RX002333 and I01RX000684), and NIH Accelerating Medicines Partnership (U01AG046170) to Eric Schadt, Icahn School of Medicine.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

1. The Lancet Neurology: Solanezumab: too late in mild Alzheimer's disease? *Lancet Neurol.* 2017; **16**(2): 97.
[PubMed Abstract](#) | [Publisher Full Text](#)
2. Hawkes N: Merck ends trial of potential Alzheimer's drug verubecestat. *BMJ.* 2017; **356**: j845.
[PubMed Abstract](#) | [Publisher Full Text](#)

3. Gouras GK, Relkin NR, Sweeney D, *et al.*: **Increased apolipoprotein E epsilon 4 in epilepsy with senile plaques.** *Ann Neurol.* 1997; 41(3): 402–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
4. Corrada MM, Hayden KM, Paganini-Hill A, *et al.*: **Age of onset of hypertension and risk of dementia in the oldest-old: The 90+ Study.** *Alzheimers Dement.* 2017; 13(2): 103–110, pii: S1552-5260(16)32962-4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
5. Lim YY, Villemagne VL, Pietrzak RH, *et al.*: **APOE ε4 moderates amyloid-related memory decline in preclinical Alzheimer's disease.** *Neurobiol Aging.* 2015; 36(3): 1239–44.
[PubMed Abstract](#) | [Publisher Full Text](#)
6. Barnes LL, Leurgans S, Aggarwal NT, *et al.*: **Mixed pathology is more likely in black than white decedents with Alzheimer dementia.** *Neurology.* 2015; 85(6): 528–34.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
7. Stoeckel LE, Arvanitakis Z, Gandy S, *et al.*: **Complex mechanisms linking neurocognitive dysfunction to insulin resistance and other metabolic dysfunction [version 2; referees: 2 approved].** *F1000Res.* 2016; 5: 353.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
8. Desplats P, Spencer B, Coffee E, *et al.*: **Alpha-synuclein sequesters Dnmt1 from the nucleus: a novel mechanism for epigenetic alterations in Lewy body diseases.** *J Biol Chem.* 2011; 286(11): 9031–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
9. Wang M, Roussos P, McKenzie A, *et al.*: **Integrative network analysis of nineteen brain regions identifies molecular signatures and networks underlying selective regional vulnerability to Alzheimer's disease.** *Genome Med.* 2016; 8(1): 104.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
10. Snowdon DA: **Aging and Alzheimer's disease: lessons from the Nun Study.** *Gerontologist.* 1997; 37(2): 150–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
11. Lambert JC, Ibrahim-Verbaas CA, Harold D, *et al.*: **Meta-analysis of 74,046 individuals identifies 11 new susceptibility loci for Alzheimer's disease.** *Nat Genet.* 2013; 45(12): 1452–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
12. De Strooper B: **After Solanezumab: Where Should Alzheimer's Research Go?** *Alzforum.* 2017.
[Reference Source](#)
13. Thambisetty M, An Y, Nalls M, *et al.*: **Effect of complement CR1 on brain amyloid burden during aging and its modification by APOE genotype.** *Biol Psychiatry.* 2013; 73(5): 422–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
14. Gandy S, Haroutunian V, DeKosky ST, *et al.*: **CR1 and the “vanishing amyloid” hypothesis of Alzheimer's disease.** *Biol Psychiatry.* 2013; 73(5): 393–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
15. Pop V, Head E, Hill MA, *et al.*: **Synergistic effects of long-term antioxidant diet and behavioral enrichment on beta-amyloid load and non-amyloidogenic processing in aged canines.** *J Neurosci.* 2010; 30(29): 9831–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
16. Ngandu T, Lehtisalo J, Solomon A, *et al.*: **A 2 year multidomain intervention of diet, exercise, cognitive training, and vascular risk monitoring versus control to prevent cognitive decline in at-risk elderly people (FINGER): a randomised controlled trial.** *Lancet.* 2015; 385(9984): 2255–63.
[PubMed Abstract](#) | [Publisher Full Text](#)
17. Lobo A, Saz P, Marcos G, *et al.*: **Prevalence of dementia in a southern European population in two different time periods: the ZARADEMP Project.** *Acta Psychiatr Scand.* 2007; 116(4): 299–307.
[PubMed Abstract](#) | [Publisher Full Text](#)
18. Sekita A, Ninomiya T, Tanizaki Y, *et al.*: **Trends in prevalence of Alzheimer's disease and vascular dementia in a Japanese community: the Hisayama Study.** *Acta Psychiatr Scand.* 2010; 122(4): 319–25.
[PubMed Abstract](#) | [Publisher Full Text](#)
19. Rocca WA, Petersen RC, Knopman DS, *et al.*: **Trends in the incidence and prevalence of Alzheimer's disease, dementia, and cognitive impairment in the United States.** *Alzheimers Dement.* 2011; 7(1): 80–93.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
20. Schrijvers EM, Verhaaren BF, Koudstaal PJ, *et al.*: **Is dementia incidence declining?: Trends in dementia incidence since 1990 in the Rotterdam Study.** *Neurology.* 2012; 78(19): 1456–63.
[PubMed Abstract](#) | [Publisher Full Text](#)
21. Wu YT, Lee HY, Norton S, *et al.*: **Prevalence studies of dementia in mainland china, Hong Kong and taiwan: a systematic review and meta-analysis.** *PLoS One.* 2013; 8(6): e66252.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
22. Qiu C, von Strauss E, Bäckman L, *et al.*: **Twenty-year changes in dementia occurrence suggest decreasing incidence in central Stockholm, Sweden.** *Neurology.* 2013; 80(20): 1888–94.
[PubMed Abstract](#) | [Publisher Full Text](#)
23. Wimo A, Sjölund BM, Sködlunger A, *et al.*: **Cohort Effects in the Prevalence and Survival of People with Dementia in a Rural Area in Northern Sweden.** *J Alzheimers Dis.* 2016; 50(2): 387–96.
[PubMed Abstract](#) | [Publisher Full Text](#)
24. Kivipelto M, Häkansson K: **A Rare Success against Alzheimer's.** *Sci Am.* 2017; 316(4): 32–37.
[PubMed Abstract](#) | [Publisher Full Text](#)
25. Jack CR Jr, Wiste HJ, Weigand SD, *et al.*: **Different definitions of neurodegeneration produce similar amyloid/neurodegeneration biomarker group findings.** *Brain.* 2015; 138(Pt 12): 3747–59.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 19 April 2017

<https://doi.org/10.5256/f1000research.12018.r21956>

© 2017 Pratico D. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Domenico Pratico

Department of Pharmacology, Center for Translational Medicine, Lewis Katz School of Medicine, Temple University, Philadelphia, PA, USA

This is a very interesting and stimulating opinion article in which the authors try to make sense of the current state of the Alzheimer's disease (AD) pharmacotherapy.

They present in a very objective fashion the disappointing scenario we are facing in this area, and at the same time they provide very insightful food for thought on how the research should move forward.

I would like the authors to spend some additional words on the followings:

1. The multifactorial nature of sporadic AD, and because of that a multi-target and may be personalized approach versus a one-size-fits-all solution is more likely to work.
2. While the large GWAS studies have provided evidence for good genetic leads, there is strong evidence indicating that environmental factors (i.e., lifestyle, diet) can ultimately influence the clinical phenotype.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 18 April 2017

<https://doi.org/10.5256/f1000research.12018.r21939>

© 2017 Griffin W et al. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

W.Sue T Griffin

Department of Geriatrics, University of Arkansas for Medical Sciences, Little Rock, AR, USA

Orwa Aboud

Department of Geriatrics, University of Arkansas for Medical Sciences, Little Rock, AR, USA

Dr. Gandy and his colleagues have come to the heart of the matter – current anti-amyloid strategies have failed to change the course of Alzheimer’s disease (AD). This important article is written as an “Aha, we can now address the question of what other strategies might hold promise toward preventing or delaying onset of Alzheimer’s disease (AD).” They raise a number of very important questions, including how early should we treat; what is the role of multiple pathologies in Alzheimer pathogenesis; are the pathways that give rise to the neuronal dysfunction and loss noted in AD the same ones that obtain in the pathogenesis of other neurodegenerative diseases; are such pathways known and, if so, have the drivers of these pathways been identified? Most of these questions have been addressed in animal models and some have included evidence of the existence of such pathways in brain tissue from patients vs controls. A great deal of effort has been given to one of the areas of interest noted by the authors “Immunology of cognitive decline in AD.” The authors’ discussion of a recent review of the potential relationship between lifestyle changes such as diet and exercise and how this may have a beneficial effect as a middle-age combatant of cognitive decline is laudable. However, implementation of such lifestyle changing practices would require individuals to be either coerced or otherwise convinced to adhere to such practices. At present, even though there is convincing evidence of tangible benefits of exercise and diet¹, the data is not with us regarding adherence to such regimens. Therefore, we would like to suggest inflammation, both in the brain and in the periphery, as a prime target for early intervention in Alzheimer pathogenesis through the use of currently available drugs or drugs developed so as to have fewer or more tolerable side effects. Data from the Framingham study of 691 cognitively intact individuals provide evidence of spontaneous increases in production by blood monocytes of two pluripotent proinflammatory cytokines, interleukin-1 (IL-1) and TNF α ²; these increases were suggested to serve as markers that “strengthen” a link between inflammation and development of AD. In addition, an accumulation of epidemiological data has consistently sided with use of anti-inflammatory compounds as increasing the odds against being in the “Alzheimer group” and in favor of being in the non-demented, non-neuropathologically confirmed group. Data from the Rotterdam study of non-NSAID users vs NSAID users reported an adjusted relative risk ratio of 0.54 among NSAID users as support for the potential of NSAIDs to protect against development of AD³. In a larger study of the VA database (~50,000 clinically and neuropathologically confirmed AD patients vs ~200,000 non-AD patients), 5 or more years of ibuprofen use reduced the adjusted odds ratio to 0.56⁴. Further in the extended results of the ADAPT study, asymptomatic subjects who received naproxen had a reduced incidence of AD compared to those who received either celecoxib or placebo⁵.

In view of the evidence implicating inflammation as increasing risk for later development of AD, it seems prudent to first explore what we know at present regarding early events in Alzheimer pathogenesis as such events may presage development, or help us pinpoint “when is the best time to intervene” to forestall formation of the aggregate defining neuropathological hallmarks of AD. In the McGeers’ early studies of neuroinflammation in AD, immune markers were identified on plaque-associated microglia in AD brain⁶. The following three discoveries: one showing that activated glia in AD brain overexpress IL-1, a second showing that microglial activation with overexpression of interleukin-1 (IL-1) as well as astrocyte activation and overexpression of the neurite-growth stimulating cytokine S100B are prominent in fetuses, newborns, children, and adults⁷, and a third showing that IL-1 induces synthesis of APP in human cell cultures⁸ opened a

new field of investigation, viz., the potential of IL-1-directed pathways to promote neuropathogenesis. These two cytokines, IL-1 and S100B upregulate each other and both induce synthesis of β APP in vitro and in vivo. This β APP may then be cleaved for β -amyloid ($A\beta$) plaque formation or for sAPP α for further microglial activation and overexpression and release of IL-1 β ⁹⁻¹⁰. Moreover, IL-1 β , via its ability to induce synthesis as well as activation of MAPK-p38, increases production of hyperphosphorylated tau, for formation of the paired helical filaments in neurofibrillary tangles, as well as α -synuclein for formation of Lewy bodies¹¹. If such neuroinflammation, which has now been associated with immune challenges in the periphery is associated with risk for development of AD, it would seem prudent to develop an inflammatory index in order to decide when treatment should begin. In view of the appearance of neuroinflammation, noted as increased activation of glia and overexpression of IL-1 decades before frank pathology in Down's syndrome, perhaps erring on the side of caution, beginning treatment sooner rather than later may be preferable.

References

1. Imayama I, Ulrich CM, Alfano CM, Wang C, et al.: Effects of a caloric restriction weight loss diet and exercise on inflammatory biomarkers in overweight/obese postmenopausal women: a randomized controlled trial. *Cancer Res.* 2012; **72** (9): 2314-26 [PubMed Abstract](#) | [Publisher Full Text](#)
2. Tan ZS, Beiser AS, Vasan RS, Roubenoff R, et al.: Inflammatory markers and the risk of Alzheimer disease: the Framingham Study. *Neurology.* 2007; **68** (22): 1902-8 [PubMed Abstract](#) | [Publisher Full Text](#)
3. Andersen K, Launer LJ, Ott A, Hoes AW, et al.: Do nonsteroidal anti-inflammatory drugs decrease the risk for Alzheimer's disease? The Rotterdam Study. *Neurology.* 1995; **45** (8): 1441-5 [PubMed Abstract](#)
4. Vlad SC, Miller DR, Kowall NW, Felson DT: Protective effects of NSAIDs on the development of Alzheimer disease. *Neurology.* 2008; **70** (19): 1672-7 [PubMed Abstract](#) | [Publisher Full Text](#)
5. Breitner JC, Baker LD, Montine TJ, Meinert CL, et al.: Extended results of the Alzheimer's disease anti-inflammatory prevention trial. *Alzheimers Dement.* 2011; **7** (4): 402-11 [PubMed Abstract](#) | [Publisher Full Text](#)
6. McGeer PL, McGeer EG: History of innate immunity in neurodegenerative disorders. *Front Pharmacol.* 2011; **2**: 77 [PubMed Abstract](#) | [Publisher Full Text](#)
7. Griffin WS, Stanley LC, Ling C, White L, et al.: Brain interleukin 1 and S-100 immunoreactivity are elevated in Down syndrome and Alzheimer disease. *Proc Natl Acad Sci U S A.* 1989; **86** (19): 7611-5 [PubMed Abstract](#)
8. Goldgaber D, Goldfarb LG, Brown P, Asher DM, et al.: Mutations in familial Creutzfeldt-Jakob disease and Gerstmann-Sträussler-Scheinker's syndrome. *Exp Neurol.* 1989; **106** (2): 204-6 [PubMed Abstract](#)
9. Barger SW, Harmon AD: Microglial activation by Alzheimer amyloid precursor protein and modulation by apolipoprotein E. *Nature.* 1997; **388** (6645): 878-81 [PubMed Abstract](#) | [Publisher Full Text](#)
10. Li Y, Liu L, Barger SW, Griffin WS: Interleukin-1 mediates pathological effects of microglia on tau phosphorylation and on synaptophysin synthesis in cortical neurons through a p38-MAPK pathway. *J Neurosci.* 2003; **23** (5): 1605-11 [PubMed Abstract](#)
11. Griffin WS, Liu L, Li Y, Mrak RE, et al.: Interleukin-1 mediates Alzheimer and Lewy body pathologies. *J Neuroinflammation.* 2006; **3**: 5 [PubMed Abstract](#) | [Publisher Full Text](#)

Is the topic of the opinion article discussed accurately in the context of the current literature?

Yes

Are all factual statements correct and adequately supported by citations?

Yes

Are arguments sufficiently supported by evidence from the published literature?

Yes

Are the conclusions drawn balanced and justified on the basis of the presented arguments?

Partly

Competing Interests: No competing interests were disclosed.

We confirm that we have read this submission and believe that we have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reader Comment 15 May 2017

William Grant, Sunlight, Nutrition and Health Research Center, USA

The opinion article by Gandy et al. discusses some important issues regarding preventing, delaying, and /or preventing Alzheimer's disease (AD).¹ Near the end, they discuss the role of diet and lifestyle in affecting risk of AD, a topic I have studied for 20 years. In this commentary I add a few observations regarding that point.

From the study of AD prevalence rates with respect to national diets, it is found that diet appears to be the most important risk-modifying factor. For example, AD prevalence rates for elderly in Japan increased from 1% in 1985 to 7% in 2008, lagging the nutrition transition from the traditional Japanese diet to the Western diet by 15-25 years.² The major changes were a 5-fold increase in meat consumption and a 50% reduction in rice consumption. In a recent 10-country study, it was found that meat, eggs, fish, and cheese were highly correlated with AD prevalence.³

The Mediterranean diet, which is touted as reducing risk of AD by about 50%, was found to lie in the middle of the regression fit to AD prevalence vs. meat consumption. The mechanisms whereby eating animal products increases risk of AD include the effects of cholesterol and increasing the intake of heavy metals while reducing intake of beneficial trace minerals (calcium, magnesium, potassium), thereby leading to increased destruction of neurons by free radicals. Cooking meat at high, dry temperatures also increases production of advanced glycation end products (AGEs), which also increase risk of AD.⁴

The dietary links to AD are somewhat similar to those for coronary heart disease (CHD).⁵ Thus, the experience in reducing risk of CHD through diet and lifestyle changes should provide some guidance for preventing AD. A good example comes from Finland. A young physician, Pekka Puska, observed that the highest cardiovascular disease mortality rates in the world were in North Karelia and were associated with high intake of saturated fats and high serum cholesterol levels.⁶ He organized a community-based program that reduced intake of saturated fats from 20% to 12% in 2007. This project led to large reductions in CHD mortality rates, from 647/100,000/yr for men and 114/100,000/yr for women in 1969-72 to

from 289/100,000/yr for men and 36/100,000/yr for women in 1992.⁷

Higher sun exposure and vitamin D status are also associated with reduced risk of AD.⁸ Interestingly, a study in New York state found that those who had non-melanoma skin cancer had a greatly reduced incidence of AD.⁹

It is unlikely that public health policies in the U.S. and many other countries will lead to reduced consumption of animal products among those middle aged or older anytime soon. However, individuals can learn the advantages of whole-plant based diets and reduce their personal risk of AD, CHD, and, as an added bonus, many types of cancer.^{10,11}

References

1. Gandy S, Bartfai T, Lees GV, Sano M. [Midlife interventions are critical in prevention, delay, or improvement of Alzheimer's disease and vascular cognitive impairment and dementia.](#) F1000Res. 2017 Apr 3;6:413
2. Grant WB. Trends in diet and Alzheimer's disease during the nutrition transition in Japan and developing countries. J Alz Dis. 2014;38(3):611-20.
3. Grant WB. Using Multicountry Ecological and Observational Studies to Determine Dietary Risk Factors for Alzheimer's Disease. J Am Coll Nutr, 2016;35(5):476-489. <http://www.tandfonline.com/doi/full/10.1080/07315724.2016.1161566>
4. Perrone L, Grant WB. Observational and ecological studies of dietary advanced glycation end products in national diets and Alzheimer's disease incidence and prevalence. J Alz Dis. 2015;45: 965-79.
5. Rafique R, Amjad N. [Dietary predictors of early-onset ischaemic heart disease in a sample drawn from a Pakistani population.](#) Heart Asia. 2012;4(1):129-34.
6. Vartiainen E, Laatikainen T, Tapanainen H, Puska P. [Changes in Serum Cholesterol and Diet in North Karelia and All Finland.](#) Glob Heart. 2016;11(2):179-84.
7. Vartiainen E, Puska P, Pekkanen J, et al. [Changes in risk factors explain changes in mortality from ischaemic heart disease in Finland.](#) BMJ. 1994;309(6946):23-7.
8. Afzal S, Bojesen SE, Nordestgaard BG. [Reduced 25-hydroxyvitamin D and risk of Alzheimer's disease and vascular dementia.](#) Alzheimers Dement. 2014;10(3):296-302.
9. White RS, Lipton RB, Hall CB, Steinerman JR. [Nonmelanoma skin cancer is associated with reduced Alzheimer disease risk.](#) Neurology. 2013;80(21):1966-72.
10. Aune D, De Stefani E, Ronco A, Boffetta P, Deneo-Pellegrini H, Acosta G, Mendilaharsu M. [Meat consumption and cancer risk: a case-control study in Uruguay.](#) Asian Pac J Cancer Prev. 2009;10(3):429-36.
11. Grant WB. A multicountry ecological study of cancer incidence rates in 2008 with respect to various risk-modifying factors, Nutrients. 2013;6(1):163-189.

Competing Interests: I receive funding from Bio-Tech Pharmacal, Inc. for research on vitamin D.

Comments on this article

Version 1

Reader Comment 17 Aug 2017

Feng-Yen Li, Physicians Committee for Responsible Medicine, USA

The authors pinpointed a number of important caveats with the current therapeutic development pipeline for Alzheimer's disease and related dementia. They astutely recognized that "we must be open-minded about the trap laid by the early success of identifying genes and molecular mechanisms of the familial Alzheimer's patients." As indicated by the continuing failures of pharmacological developments, researchers are focusing on the wrong targets. Their advice to shift the research focus from anti-proteinopathy treatments to modifying midlife risk factors with lifestyle interventions is very sound, insightful, and well-supported: "In the meantime, the importance and potential benefit of safe and easy diet and lifestyle changes for humans should be a topic of much stronger advocacy." However, their suggestion to use canine models to further such studies may not be a wise recommendation as there will always be physiological differences between animals and humans that can limit translation. After all, despite the potential of the dietary and behavioral intervention to modulate amyloid-beta in the canine model, the cited study in the paper for this model stated that "amyloid-beta measures did not correlate with cognitive performance on any of the 6 tasks assessed, suggesting that modulation of amyloid-beta alone may be a relatively minor mechanism mediating cognitive benefits of the interventions." Plus, any findings in such animal models will need to be confirmed in humans. Given that dietary and lifestyle interventions can be directly tested in humans without safety concerns, it would best to complete such studies in humans rather than animal models to avoid the extra cost, time, and potential misleading conclusions.

The commentary posted by William Grant above pointed to strong epidemiological evidence for the association of AD prevalence and animal product consumption. He also makes the astute observation that the risk factors of AD are similar to those of CHD and provides lessons to be learned from studies demonstrating the feasibility of using dietary and lifestyle interventions to reduce CHD mortality. He further suggests a whole-plant based diet together with supplementation of important nutrients like vitamin D as an effective strategy to reduce the prevalence of many chronic diseases, including AD, CHD, and many cancers. Indeed, the effectiveness of such dietary interventions has already been shown to be highly effective even for reversing CHD, diabetes, and even cancers¹⁻⁵. A randomized controlled trial testing the efficacy of the whole food plant-based diet supplemented with key anti-inflammatory, neuroprotective nutrients (e.g. vitamin D, B12, algae-based omega-3, turmeric, etc) for the prevention or reversal AD and related dementia promises to help reverse the current epidemiological trends for this burdensome disease.

1. Esselstyn CB Jr, Gendy G, Doyle J, Golubic M, Roizen MF. A way to reverse CAD? J Fam Pract. 2014 Jul;63(7):356-364b.
2. Lee YM, Kim SA, Lee IK, Kim JG, Park KG, Jeong JY, Jeon JH, Shin JY, Lee DH. Effect of a Brown Rice Based Vegan Diet and Conventional Diabetic Diet on Glycemic Control of Patients with Type 2 Diabetes: A 12-Week Randomized Clinical Trial. PLoS One. 2016 Jun 2;11(6):e0155918.
3. Barnard ND, Cohen J, Jenkins DJ, Turner-McGrievy G, Gloede L, Green A, Ferdowsian H. A low-fat vegan diet and a conventional diabetes diet in the treatment of type 2 diabetes: a

- randomized, controlled, 74-wk clinical trial. Am J Clin Nutr. 2009 May;89(5):1588S-1596S.
4. Frattaroli J, Weidner G, Dnistrian AM, Kemp C, Daubenmier JJ, Marlin RO, Crutchfield L, Yglecias L, Carroll PR, Ornish D. Clinical events in prostate cancer lifestyle trial: results from two years of follow-up. Urology. 2008 Dec;72(6):1319-23
 5. Saxe GA, Major JM, Nguyen JY, Freeman KM, Downs TM, Salem CE. Potential attenuation of disease progression in recurrent prostate cancer with plant-based diet and stress reduction. Integr Cancer Ther. 2006 Sep;5(3):206-13.

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research