REVIEW

Selective vulnerabilities and biomarkers in neurocognitive aging [version 1; peer review: 4 approved]

Zachariah Reagh, Michael Yassa

Department of Neurobiology and Behavior, Center for the Neurobiology of Learning and Memory, Institute for Memory Impairments and Neurological Disorders, University of California, Irvine, CA, USA

V1 First published: 13 Apr 2017, 6(F1000 Faculty Rev):491
<https://doi.org/10.12688/f1000research.10652.1>

Latest published: 13 Apr 2017, 6(F1000 Faculty Rev):491
<https://doi.org/10.12688/f1000research.10652.1>

Abstract

As the world's population continues to age, an understanding of the aging brain becomes increasingly crucial. This review focuses on several recent ideas and findings in the study of neurocognitive aging, specifically focusing on episodic memory, and discusses how they can be considered and used to guide us moving forward. Topics include dysfunction in neural circuits, the roles of neurogenesis and inhibitory signaling, vulnerability in the entorhinal cortex, individual differences, and comorbidities. These avenues of study provide a brief overview of promising themes in the field and together provide a snapshot of what we believe will be important emerging topics in selective vulnerabilities in the aging brain.

Keywords

neurocognitive aging, episodic memory, entorhinal cortex

Open Peer Review

Approval Status

	1	2	3	4
version 1 13 Apr 2017				

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Peter R Rapp**, Laboratory of Behavioral Neuroscience, National Institute on Aging, Baltimore, Baltimore, USA
2. **Morgan D Barense**, Department of Psychology, University of Toronto, Toronto, Canada
3. **Heikki Tanila**, A.I. Virtanen Institute for Molecular Sciences, University of Eastern Finland, Kuopio, Kuopio, Finland
4. **Michela Gallagher**, Department of Psychological and Brain Sciences, Johns Hopkins University, Baltimore, USA

Any comments on the article can be found at the end of the article.

Corresponding authors: Zachariah Reagh (zreagh@uci.edu), Michael Yassa (myassa@uci.edu)

Competing interests: No competing interests were disclosed.

Grant information: Support for the authors comes from NIH R01 MH102392, R21 AG049220, and P50 AG16573 (to MAY) as well as NSF GRF DGE1232825, T32 AG96-33 and a fellowship from the Achievement Rewards for College Scientists Foundation and Roche Foundation (to ZMR).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2017 Reagh Z and Yassa M. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. Data associated with the article are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

How to cite this article: Reagh Z and Yassa M. **Selective vulnerabilities and biomarkers in neurocognitive aging [version 1; peer review: 4 approved]** F1000Research 2017, 6(F1000 Faculty Rev):491 <https://doi.org/10.12688/f1000research.10652.1>

First published: 13 Apr 2017, 6(F1000 Faculty Rev):491 <https://doi.org/10.12688/f1000research.10652.1>

Introduction

As of 2015, the global percentage of individuals 65 years of age or older was estimated to be 8.5%, which is projected to grow to 16.7% by 2050¹. Increasing age is a primary risk factor of many disease states, including dementias such as Alzheimer's disease (AD). Beyond disease risk, however, even 'healthy' aging significantly alters functional and structural properties of the brain. Furthermore, there is considerable individual variability in aged individuals in terms of both cognitive and neural measures^{2,3}. Thus, it is imperative that we understand age-related changes to the brain in both the presence and absence of frank pathology such as that of dementing illnesses. Such an understanding will provide valuable insight into 'healthy' aging as well as potential biomarkers for differentiating this trajectory from diseases.

In this review, we will discuss important prevailing themes that can serve to guide our thinking about neurocognitive aging. We will also highlight some recent findings that may offer promising directions in the search for biomarkers and potential targets of intervention:

1. Aging is a story of dysfunction rather than cell loss

This first point is perhaps the least contentious and has been important in guiding our thinking about how aging affects cognition. Though the brain undergoes volumetric changes with age, this is thought to reflect synaptic degradation rather than frank neuronal loss in the absence of disease. Moreover, loss of function in particular neuronal circuits may be far more striking and informative at early stages of the aging process. It is worth noting that this understanding forms an important background that frames the other findings we discuss below.

2. Neurogenesis is a complex biomarker

Until very recently, it has been fairly well accepted that neurogenesis significantly declines in the aged brain, which may account for declines in memory ability. However, recent findings in the human brain have complicated this view.

3. Letting go of your inhibition

Many areas of the brain—notably, regions involved in memory—are typically under tight inhibition. This inhibitory control is dysregulated in aging, which can have important consequences on circuit-level computations.

4. Entorhinal cortex: the gateway to neurodegeneration

A wealth of evidence points to entorhinal cortex (EC) as being among the earliest brain regions affected by aging, and recent studies suggest that certain subdivisions of the EC may be especially vulnerable. In addition to the EC, several new candidate regions of vulnerability, including thalamic nuclei and the locus coeruleus, have emerged in recent years.

5. Individual differences are informative

Individuals vary widely in terms of both cognition and neurobiology as they age. We often collapse across and ignore many such differences in any given experiment, but they likely provide important clues about a person's cognitive and clinical trajectories.

6. It is time to embrace and understand comorbidities

Brain aging is inexorably linked to other factors that influence the brain's health as well as the rest of the body. However, these comorbidities are often viewed as 'nuisance' variables that we attempt to exclude. These are important factors that should be considered for a full and valid understanding of the aging brain.

Aging is a story of dysfunction rather than cell loss

Neuronal loss in the medial temporal lobes (MTLs) is observed in histological studies of AD, which correlates with disease onset^{4,5}. Although hippocampal volume is often found to decrease in 'healthy' aging^{6,7}, there is little evidence to suggest that 'healthy' aging features comparable loss of cells⁸. What could account for this discrepancy? It is likely that such volumetric changes are due to more subtle losses of cellular complexity, including degeneration of synapses⁹ and axons¹⁰. This is consistent with studies using unbiased stereology in animal models of aged rodents, all of which show that there is little to no evidence of frank cell loss in the aging brain⁸. Additionally, in specific transgenic mouse models of AD, impairments in memory-guided tasks correlate more closely with synaptic dysfunction than neuronal loss¹¹. Though likely less drastic than cell death, these kinds of changes may contribute to subtle cognitive perturbations. For instance, the hallmark decline of episodic memory ability with aging may arise from highly specific synaptic loss observed in the perforant path between the EC and the hippocampus⁸. In humans, white matter loss specific to this projection correlates with poorer mnemonic discrimination (disambiguating overlapping events in memory), thought to rely on hippocampal pattern separation^{12,13}. In a typical mnemonic discrimination paradigm, participants encode a set of items and later are tasked with rejecting similar 'lure' items as being distinct from the studied set. Importantly, these discrimination paradigms have been shown to be highly sensitive to age-related decline in episodic memory ability.

Beyond morphological changes that do not correspond to neuronal loss per se, changes to the functional properties of brain systems have been reported that correlate with cognitive deficits. In the hippocampi of aged rats there are several functional changes that are associated with poorer memory performance. This includes alterations to excitatory synaptic transmission and synaptic plasticity across all hippocampal subfields¹⁴. Several studies have also noted that elevated levels of activity in the CA3 subfield correlate with impaired mnemonic discrimination^{15,16}, a finding mirrored in humans with high-resolution functional magnetic resonance imaging (fMRI)¹⁷. Behaviorally, mnemonic discrimination paradigms are sensitive to subtle age-related deficits across multiple domains (for example, objects, space, and time)^{18–20}. Other fMRI experiments examining activity during task-free rest report decreased correlations in activity among 'default mode' brain regions²¹, many of which are strongly implicated in memory. Importantly, many of these studies control for gray matter volume in their analyses such that global atrophy does not explain these disrupted correlational patterns among brain regions. Together, these findings suggest that noninvasive functional approaches such as behavioral studies and measures of brain activity at rest or driven by mechanistically targeted tasks are potentially very fruitful avenues of inquiry.

Neurogenesis is a complex biomarker

Though adult neurogenesis (the birth of new neurons in the mature brain) has been known for some time to occur in the dentate gyrus of the mammalian hippocampus²², the phenomenon has recently become a popular avenue of research²³. In rodents, nearly 10,000 new cells are generated per day²⁴ and only half of these survive²⁵. Among these surviving granule cells, most integrate into existing hippocampal circuits and may replace existing granule cells²⁶. Though the precise functions of adult neurogenesis are not known, disruption of neurogenesis is detrimental to certain mnemonic processes, primarily pattern separation²⁷. It must be noted, however, that many memory-guided behaviors are unaffected by this. Adult neurogenesis reportedly decreases with age in the rodent brain²⁸, leading some to propose it as a mechanism for curbing age-related memory loss and detecting pathological states along different aging trajectories²⁹.

A recent study using nuclear bomb test-derived ¹⁴C in genomic DNA found that although hippocampal neurogenesis does decline with age in humans, the effect is nowhere near as drastic as in rodents³⁰. Moreover, after age 20, the loss of neurogenesis dropped at a low and fairly monotonic rate even into advanced age. Although this does not refute a role for neurogenesis in explaining human age-related memory decline, the relationship is unclear. It must be noted that there are differences between human and rodent adult neurogenesis. For instance, compared with rodents³⁰, a greater number of newborn neurons in the human dentate gyrus integrate into existing circuits³¹. There is also evidence to suggest that the dynamics and behavioral relevance of neurogenesis differ between rats and mice³². Nonetheless, targeting neurogenesis as a mechanism for therapeutic intervention has some promise, but further research is needed to fully understand its role in the human brain across the life span.

Letting go of your inhibition

As previously noted, hyperactivity in hippocampal CA3 in aged animals is associated with diminished memory^{15,16}. This likely stems at least in part from reduced inhibitory signaling from somatostatin-positive GABAergic interneurons to CA3³³. Similarly, somatostatin-positive interneurons are selectively lost in the hilar region of the dentate gyrus in older animals with memory impairments³⁴. As hippocampal networks are typically under tight inhibition, such alterations can drastically affect mnemonic processes. For instance, elevated firing rates in the dentate gyrus may reduce the sparsity of firing in granule cells, thereby reducing its capacity for creating independent representations. The same change in the CA3 region may facilitate runaway excitation in recurrent collaterals and further interfere with the network's ability to represent new information. This is likely to lead to failures in pattern separation and instead biases the hippocampus toward erroneous generalization^{35,36} or the generation of 'false' memories.

In a recent series of studies in rats³⁷, humans³⁸, and transgenic mice³⁹ with AD pathology, a low dose of the anti-epileptic drug levetiracetam was found to reduce hippocampal hyperactivity in memory-impaired older individuals as well as rescue cognitive deficits to normal levels. Hippocampal hyperactivity is therefore a strong candidate biomarker in neurocognitive aging that may

differentiate 'healthy' from pathological aging. Moreover, although both the acute and long-term effects of the relevant drugs must be studied more extensively, targeting and reducing hyperactivity in specific brain networks provide a well-supported mechanistic avenue for treating cognitive decline.

It should be noted, however, that loss of inhibition is not universal across the aging brain but rather may be regionally specific. Some regions in fact exhibit *greater* inhibition in advanced age. For example, evidence in both primates⁴⁰ and rodents⁴¹ associates aging with increased inhibitory tone in the prefrontal cortex. Thus, inhibitory control is a broad and potentially very important issue to consider.

Entorhinal cortex: the gateway to neurodegeneration

Though the hippocampus is often studied in the context of dementia and is often predictive of AD⁴², histological⁴³ and neuroimaging⁴⁴ studies have shown that the EC is among the earliest brain regions affected by AD-related pathology, such as cortical thinning and the accumulation of hyperphosphorylated tau protein. Recent evidence suggests that there may be detectable changes in the functional properties of EC. An elegant study by Khan and colleagues⁴⁵ demonstrated that, across mice and humans, metabolism in the lateral portion of the EC (LEC) was affected in preclinical AD, consistent with other studies in rodent models of aging⁴⁶. Moreover, positron emission tomography (PET) imaging of hyperphosphorylated tau deposition in the EC is a promising biomarker⁴⁷.

Critically, although it has been known for some time that LEC and medial EC (MEC) are functionally dissociable in rodents^{48,49}, evidence in humans has only recently been demonstrated^{50–52} (though in humans the division may be more anterolateral versus posteromedial^{50,51}). Generally, cells in MEC are thought to represent spatial configurations via grid-like firing patterns⁴⁸, whereas those in LEC are thought to represent objects or items within those configurations⁴⁹. Task-driven dissociations between LEC and MEC⁵² provide a valuable opportunity to test key changes to the aging human brain. In particular, one might hypothesize that LEC shows selective functional impairments in aged adults, and the extent of this impairment might inform us as to one's neurocognitive trajectory. A recent study by Olsen and colleagues provides evidence for such a view via structural changes to EC subdivisions⁵³. Future studies can further address this question and can resolve ambiguities in behavioral outcomes such as whether functions of the LEC are broadly impacted by 'healthy' aging⁵⁴ or only in the case of a clinical trajectory⁵⁵.

In addition to the EC, several other regions have recently surfaced as candidate locales for neurodegenerative changes in AD. The first is the locus coeruleus, which is a major brainstem hub for producing norepinephrine and which recently was identified as a region that accumulates tau pathology early in AD⁵⁶ and may even transfer this pathology to the EC via a prion-like mechanism⁵⁷. Second, there is evidence that the thalamus⁵⁸, striatum⁵⁸, and basal forebrain cholinergic system⁵⁹ are also compromised as regions expressing early pathology and neurodegenerative changes in AD. A recent study reported evidence that basal forebrain degeneration may actually precede and dictate AD-related pathology in the EC⁶⁰. Moving forward, it will be important to elucidate which

regions exhibit neurodegenerative or volumetric change related to AD versus volumetric change related to synaptic and dendritic loss, which may be a part of ‘healthy’ aging.

Individual differences are informative

Our use of quotations around ‘healthy’ aging here is quite intentional. It is well noted that brain aging is not a uniform process but rather consists of many different cognitive and neurobiological phenotypes⁶¹. This raises an important question: what exactly constitutes a normal aging trajectory? Absent a disease state, cognitive abilities vary widely across older individuals^{2,62}, and there is evidence suggesting that early life education⁶³ and genetic factors⁶⁴ influence outcomes. Moreover, individuals with a common diagnosis may share a wide array of distinct symptoms or pathological phenotypes. In many experimental studies, it is common to attempt to reduce or even eliminate as many variables as possible for a ‘cleaner’ examination of a variable of interest. This is often a matter of cost or practicality in a given study, as more variables may necessitate larger sample sizes. Nonetheless, we suggest that attempting to understand these differences and relate them to experimental outcomes is a necessary step in understanding neurocognitive aging more broadly.

Several studies across rodents⁶⁵, primates^{66,67}, and humans^{18–20} have embraced individual differences in behavior, which sometimes can cleanly dissociate subgroups of aged individuals. In one novel and exciting approach, recent studies by Madan and Kensinger^{68,69} found that the structural complexity of certain brain regions (including MTL and striatum) is a more sensitive measure of age-related differences than volume and cortical thickness. This approach leverages individual variability rather than controlling for its effects as a set of confounding variables. If combined with extensive demographic information and sensitive behavioral data, a technique such as this might provide a powerful means of understanding factors that impact and arise from aging.

It is time to embrace and understand comorbidities

Related to individual variability, comorbidities provide a considerable challenge to characterizing a normal neurocognitive aging process. As many as one third of individuals in the US 65 years or older are diagnosed with at least two chronic diseases⁷⁰. Prominently, aging itself is the primary risk factor for AD and shares pathological features such as amyloid deposition⁷¹. One major factor is cerebrovascular infarction (stroke), which is fairly common to some extent in a large proportion of older individuals⁷². In addition to major stroke events, more chronic forms of cerebrovascular disease such as microbleeds, or ‘silent infarcts’, have been found to affect cognition absent any AD-related pathology⁷³. Relatedly, recent evidence has pointed to a strong role of inflammation in many age-related diseases, including arthritis, diabetes, dementia, and metabolic dysregulation⁷⁴. Aging upregulates a number of pro-inflammatory signaling molecules, which may dysregulate immune responses in the nervous system with damaging effects⁷⁵. Diabetes and metabolic syndrome—an increasingly widespread issue encompassing obesity, hypertension, insulin resistance, and related symptoms—have also been linked to age-related cognitive decline and AD⁷⁶. Finally, differences in neuropsychiatric health exist among aged adults. For example, many individuals go on to develop late-life depression⁷⁷, which

can affect not only cognition but core neurobiological processes as well.

As with individual differences, comorbidities are frequently treated as nuisance variables in a given study. Although there have been some efforts, a systematic understanding of comorbidities and how they can be leveraged to improve our grasp on the aging process has not been provided. One promising example comes from studies of the oldest old where the best indicator of the likelihood and severity of dementia was the number of concurrent pathologies⁷⁸. This is not a trivial undertaking, as the presence of any given condition—be it vascular disease or depression—likely influences the presentation of another condition such as dementia. Nonetheless, this is a major remaining issue that the field must soon tackle. One crucial thing to bear in mind is that nearly all aged human brains have some amount of vascular or parenchymal brain pathologies, some of which are often considered biomarkers for diseases such as AD. Thus, it is critical to come to understand which of these, where in the brain, and in what combinations constitute a serious threat to aging brain health.

Summary

With the world’s aging population on the rise, it is imperative that we have a better understanding of the ‘healthy’ aging brain and how it differs from various pathological states. This is no easy feat. We suggest that in order to make inroads here, we must first focus on brain networks and systems that are selectively more vulnerable than others. We focused on episodic memory systems as memory complaints are quite common in the elderly and memory loss is one of the earliest hallmarks of AD. We first remind the reader that, absent neurodegenerative disease, aging is associated with synaptic dysfunction and not cell loss per se. With this in mind, we discuss the putative roles of inhibition and neurogenesis. We discuss regional vulnerability to neurodegenerative change, focusing on the EC but also extending to other recently identified subcortical targets. Finally, we discuss an important role for research considering individual differences and disease comorbidities in understanding neurocognitive changes in the aging brain. While by no means exhaustive, these are some of the important themes we think should guide research in the area over the coming few years.

Competing interests

The authors declare that they have no competing interests.

Grant information

Support for the authors comes from NIH R01 MH102392, R21 AG049220, and P50 AG16573 (to MAY) as well as NSF GRF DGE1232825, T32 AG96-33 and a fellowship from the Achievement Rewards for College Scientists Foundation and Roche Foundation (to ZMR).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgments

We extend our thanks to our four reviewers for their helpful and insightful comments.

References

1. He W, Goodkind D, Kowal P: **An Aging World: 2015.** *International Population Reports.* 2016; P95/16-1.
[Reference Source](#)
2. MacDonald SW, Nyberg L, Backman L: **Intra-individual variability in behavior: links to brain structure, neurotransmission and neuronal activity.** *Trends Neurosci.* 2006; **29**(8): 474–80.
[PubMed Abstract](#) | [Publisher Full Text](#)
3. Raz N, Ghisletta P, Rodrigue KM, *et al.*: **Trajectories of brain aging in middle-aged and older adults: regional and individual differences.** *Neuroimage.* 2010; **51**(2): 501–11.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
4. Price JL, Ko AI, Wade MJ, *et al.*: **Neuron number in the entorhinal cortex and CA1 in preclinical Alzheimer disease.** *Arch Neurol.* 2001; **58**(9): 1395–402.
[PubMed Abstract](#) | [Publisher Full Text](#)
5. West MJ, Coleman PD, Flood DG, *et al.*: **Differences in the pattern of hippocampal neuronal loss in normal ageing and Alzheimer's disease.** *Lancet.* 1994; **344**(8925): 769–72.
[PubMed Abstract](#) | [Publisher Full Text](#)
6. **F** Raz N, Lindenberger U, Rodrigue KM, *et al.*: **Regional brain changes in aging healthy adults: general trends, individual differences and modifiers.** *Cereb Cortex.* 2005; **15**(11): 1676–89.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
7. Jack CR Jr, Petersen RC, Xu YC, *et al.*: **Medial temporal atrophy on MRI in normal aging and very mild Alzheimer's disease.** *Neurology.* 1997; **49**(3): 786–94.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
8. Burke SN, Barnes CA: **Neural plasticity in the ageing brain.** *Nat Rev Neurosci.* 2006; **7**(1): 30–40.
[PubMed Abstract](#) | [Publisher Full Text](#)
9. Hedden T, Gabrieli JD: **Insights into the ageing mind: a view from cognitive neuroscience.** *Nat Rev Neurosci.* 2004; **5**(2): 87–96.
[PubMed Abstract](#) | [Publisher Full Text](#)
10. Schliebs R, Arendt T: **The cholinergic system in aging and neuronal degeneration.** *Behav Brain Res.* 2011; **221**(2): 555–63.
[PubMed Abstract](#) | [Publisher Full Text](#)
11. Selkoe DJ: **Alzheimer's disease is a synaptic failure.** *Science.* 2002; **298**(5594): 789–91.
[PubMed Abstract](#) | [Publisher Full Text](#)
12. Yassa MA, Mattfeld AT, Stark SM, *et al.*: **Age-related memory deficits linked to circuit-specific disruptions in the hippocampus.** *Proc Natl Acad Sci U S A.* 2011; **108**(21): 8873–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
13. **F** Bennett IJ, Stark CE: **Mnemonic discrimination relates to perforant path integrity: An ultra-high resolution diffusion tensor imaging study.** *Neurobiol Learn Mem.* 2016; **129**: 107–12.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
14. Burke SN, Barnes CA: **Senescent synapses and hippocampal circuit dynamics.** *Trends Neurosci.* 2010; **33**(3): 153–61.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
15. **F** Wilson IA, Ikonen S, Gallagher M, *et al.*: **Age-associated alterations of hippocampal place cells are subregion specific.** *J Neurosci.* 2005; **25**(29): 6877–86.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
16. Wilson IA, Ikonen S, Gurevicius K, *et al.*: **Place cells of aged rats in two visually identical compartments.** *Neurobiol Aging.* 2005; **26**(7): 1099–106.
[PubMed Abstract](#) | [Publisher Full Text](#)
17. Yassa MA, Lacy JW, Stark SM, *et al.*: **Pattern separation deficits associated with increased hippocampal CA3 and dentate gyrus activity in nondemented older adults.** *Hippocampus.* 2011; **21**(9): 968–79.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
18. Stark SM, Yassa MA, Lacy JW, *et al.*: **A task to assess behavioral pattern separation (BPS) in humans: Data from healthy aging and mild cognitive impairment.** *Neuropsychologia.* 2013; **51**(12): 2442–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
19. Reagh ZM, Roberts JM, Ly M, *et al.*: **Spatial discrimination deficits as a function of mnemonic interference in aged adults with and without memory impairment.** *Hippocampus.* 2014; **24**(3): 303–14.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
20. Roberts JM, Ly M, Murray E, *et al.*: **Temporal discrimination deficits as a function of lag interference in older adults.** *Hippocampus.* 2014; **24**(10): 1189–96.
[PubMed Abstract](#) | [Publisher Full Text](#)
21. Damoiseaux JS, Beckmann CF, Arigita EJ, *et al.*: **Reduced resting-state brain activity in the "default network" in normal aging.** *Cereb Cortex.* 2008; **18**(8): 1856–64.
[PubMed Abstract](#) | [Publisher Full Text](#)
22. Bayer SA, Yackel JW, Puri PS: **Neurons in the rat dentate gyrus granular layer substantially increase during juvenile and adult life.** *Science.* 1982; **216**(4548): 890–2.
[PubMed Abstract](#) | [Publisher Full Text](#)
23. Gage FH: **Mammalian neural stem cells.** *Science.* 2000; **287**(5457): 1433–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
24. Cameron HA, McKay RD: **Adult neurogenesis produces a large pool of new granule cells in the dentate gyrus.** *J Comp Neurol.* 2001; **435**(4): 406–17.
[PubMed Abstract](#) | [Publisher Full Text](#)
25. **F** van Praag H, Schinder AF, Christie BR, *et al.*: **Functional neurogenesis in the adult hippocampus.** *Nature.* 2002; **415**(6875): 1030–4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
26. Dayer AG, Ford AA, Cleaver KM, *et al.*: **Short-term and long-term survival of new neurons in the rat dentate gyrus.** *J Comp Neurol.* 2003; **460**(4): 563–72.
[PubMed Abstract](#) | [Publisher Full Text](#)
27. **F** Clelland CD, Choi M, Romberg C, *et al.*: **A functional role for adult hippocampal neurogenesis in spatial pattern separation.** *Science.* 2009; **325**(5937): 210–3.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
28. Kuhn HG, Dickinson-Anson H, Gage FH: **Neurogenesis in the dentate gyrus of the adult rat: age-related decrease of neuronal progenitor proliferation.** *J Neurosci.* 1996; **16**(6): 2027–33.
[PubMed Abstract](#)
29. Hollands C, Bartolotti N, Lazarov O: **Alzheimer's Disease and Hippocampal Adult Neurogenesis: Exploring Shared Mechanisms.** *Front Neurosci.* 2016; **10**: 178.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
30. **F** Spalding KL, Bergmann O, Alkass K, *et al.*: **Dynamics of hippocampal neurogenesis in adult humans.** *Cell.* 2013; **153**(6): 1219–27.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
31. Ninkovic J, Mori T, Götz M: **Distinct modes of neuron addition in adult mouse neurogenesis.** *J Neurosci.* 2007; **27**(40): 10906–11.
[PubMed Abstract](#) | [Publisher Full Text](#)
32. **F** Snyder JS, Choe JS, Clifford MA, *et al.*: **Adult-born hippocampal neurons are more numerous, faster maturing, and more involved in behavior in rats than in mice.** *J Neurosci.* 2009; **29**(46): 14484–95.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
33. **F** Thomé A, Gray DT, Erickson CA, *et al.*: **Memory impairment in aged primates is associated with region-specific network dysfunction.** *Mol Psychiatry.* 2016; **21**(9): 1257–62.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
34. **F** Spiegel AM, Koh MT, Vogt NM, *et al.*: **Hilar interneuron vulnerability distinguishes aged rats with memory impairment.** *J Comp Neurol.* 2013; **521**(15): 3508–23.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
35. Wilson IA, Gallagher M, Eichenbaum H, *et al.*: **Neurocognitive aging: prior memories hinder new hippocampal encoding.** *Trends Neurosci.* 2006; **29**(12): 662–70.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
36. Yassa MA, Stark CE: **Pattern separation in the hippocampus.** *Trends Neurosci.* 2011; **34**(10): 515–25.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
37. Koh MT, Haberman RP, Foti S, *et al.*: **Treatment strategies targeting excess hippocampal activity benefit aged rats with cognitive impairment.** *Neuropsychopharmacology.* 2010; **35**(4): 1016–25.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
38. Bakker A, Krauss GL, Albert MS, *et al.*: **Reduction of hippocampal hyperactivity improves cognition in amnesic mild cognitive impairment.** *Neuron.* 2012; **74**(3): 467–74.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
39. **F** Sanchez PE, Zhu L, Verret L, *et al.*: **Levetiracetam suppresses neuronal network dysfunction and reverses synaptic and cognitive deficits in an Alzheimer's disease model.** *Proc Natl Acad Sci U S A.* 2012; **109**(42): E2895–903.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
40. Arnsten AF: **Catecholamine modulation of prefrontal cortical cognitive function.** *Trends Cogn Sci.* 1998; **2**(11): 436–47.
[PubMed Abstract](#) | [Publisher Full Text](#)
41. McQuail JA, Frazier CJ, Bizon JL: **Molecular aspects of age-related cognitive decline: the role of GABA signaling.** *Trends Mol Med.* 2015; **21**(7): 450–60.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
42. Wang L, Swank JS, Glick IE, *et al.*: **Changes in hippocampal volume and shape across time distinguish dementia of the Alzheimer type from healthy aging.** *Neuroimage.* 2003; **20**(2): 667–82.
[PubMed Abstract](#) | [Publisher Full Text](#)
43. Gómez-Isla T, Price JL, McKeel DW Jr, *et al.*: **Profound loss of layer II entorhinal cortex neurons occurs in very mild Alzheimer's disease.** *J Neurosci.* 1996; **16**(14): 4491–500.
[PubMed Abstract](#)
44. Whitwell JL, Przybelski SA, Weigand SD, *et al.*: **3D maps from multiple MRI illustrate changing atrophy patterns as subjects progress from mild cognitive impairment to Alzheimer's disease.** *Brain.* 2007; **130**(Pt 7): 1777–86.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

45. **F** Khan UA, Liu L, Provenzano FA, *et al.*: **Molecular drivers and cortical spread of lateral entorhinal cortex dysfunction in preclinical Alzheimer's disease.** *Nat Neurosci.* 2014; **17**(2): 304–11.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
46. Stranahan AM, Haberman RP, Gallagher M: **Cognitive decline is associated with reduced reelin expression in the entorhinal cortex of aged rats.** *Cereb Cortex.* 2011; **21**(2): 392–400.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
47. Cho H, Choi JY, Hwang MS, *et al.*: **Tau PET in Alzheimer disease and mild cognitive impairment.** *Neurology.* 2016; **87**(4): 375–83.
[PubMed Abstract](#) | [Publisher Full Text](#)
48. **F** Hargreaves EL, Rao G, Lee I, *et al.*: **Major dissociation between medial and lateral entorhinal input to dorsal hippocampus.** *Science.* 2005; **308**(5729): 1792–4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
49. Yoganasimha D, Rao G, Knierim JJ: **Lateral entorhinal neurons are not spatially selective in cue-rich environments.** *Hippocampus.* 2011; **21**(12): 1363–74.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
50. **F** Maass A, Berron D, Libby LA, *et al.*: **Functional subregions of the human entorhinal cortex.** *eLife.* 2015; **4**: e06426.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
51. **F** Navarro Schröder T, Haak KV, Zaragoza Jimenez NI, *et al.*: **Functional topography of the human entorhinal cortex.** *eLife.* 2015; **4**: e06738.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
52. Reagh ZM, Yassa MA: **Object and spatial mnemonic interference differentially engage lateral and medial entorhinal cortex in humans.** *Proc Natl Acad Sci U S A.* 2014; **111**(40): E4264–73.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
53. Olsen RK, Yeung LK, Noly-Gandon A, *et al.*: **Human anterolateral entorhinal cortex volumes are associated with preclinical cognitive decline.** *bioRxiv Preprint.* 2016.
[Publisher Full Text](#)
54. Reagh ZM, Ho HD, Leal SL, *et al.*: **Greater loss of object than spatial mnemonic discrimination in aged adults.** *Hippocampus.* 2016; **26**(4): 417–22.
[PubMed Abstract](#) | [Publisher Full Text](#)
55. **F** Fidalgo CO, Changoor AT, Page-Gould E, *et al.*: **Early cognitive decline in older adults better predicts object than scene recognition performance.** *Hippocampus.* 2016; **26**(12): 1579–92.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
56. Braak H, Thal DR, Ghebremedhin E, *et al.*: **Stages of the pathologic process in Alzheimer disease: age categories from 1 to 100 years.** *J Neuropathol Exp Neurol.* 2011; **70**(11): 960–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
57. Braak H, Del Tredici K: **Alzheimer's pathogenesis: is there neuron-to-neuron propagation?** *Acta Neuropathol.* 2011; **121**(5): 589–95.
[PubMed Abstract](#) | [Publisher Full Text](#)
58. Leh SE, Kälin AM, Schroeder C, *et al.*: **Volumetric and shape analysis of the thalamus and striatum in amnesic mild cognitive impairment.** *J Alzheimers Dis.* 2016; **49**(1): 237–49.
[PubMed Abstract](#) | [Publisher Full Text](#)
59. Kilimann I, Grothe M, Heinsen H, *et al.*: **Subregional basal forebrain atrophy in Alzheimer's disease: a multicenter study.** *J Alzheimers Dis.* 2014; **40**(3): 687–700.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
60. Schmitz TW, Nathan Spreng R, Alzheimer's Disease Neuroimaging Initiative: **Basal forebrain degeneration precedes and predicts the cortical spread of Alzheimer's pathology.** *Nat Commun.* 2016; **7**: 13249.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
61. Jagust W: **Vulnerable neural systems and the borderland of brain aging and neurodegeneration.** *Neuron.* 2013; **77**(2): 219–34.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
62. Wilson RS, Beckett LA, Barnes LL, *et al.*: **Individual differences in rates of change in cognitive abilities of older persons.** *Psychol Aging.* 2002; **17**(2): 179–93.
[PubMed Abstract](#) | [Publisher Full Text](#)
63. Bennett DA, Wilson RS, Schneider JA, *et al.*: **Education modifies the relation of AD pathology to level of cognitive function in older persons.** *Neurology.* 2003; **60**(12): 1909–15.
[PubMed Abstract](#) | [Publisher Full Text](#)
64. Deary IJ, Yang J, Davies G, *et al.*: **Genetic contributions to stability and change in intelligence from childhood to old age.** *Nature.* 2012; **482**(7384): 212–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
65. Gallagher M, Colantuoni C, Eichenbaum H, *et al.*: **Individual differences in neurocognitive aging of the medial temporal lobe.** *Age (Dordr).* 2006; **28**(3): 221–33.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
66. Rapp PR, Amaral DG: **Recognition memory deficits in a subpopulation of aged monkeys resemble the effects of medial temporal lobe damage.** *Neurobiol Aging.* 1991; **12**(5): 481–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
67. Hara Y, Rapp PR, Morrison JH: **Neuronal and morphological bases of cognitive decline in aged rhesus monkeys.** *Age (Dordr).* 2012; **34**(5): 1051–73.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
68. Madan CR, Kensinger EA: **Cortical complexity as a measure of age-related brain atrophy.** *Neuroimage.* 2016; **134**: 617–29.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
69. Madan CR, Kensinger EA: **Age-related differences in the structural complexity of subcortical and ventricular structures.** *Neurobiol Aging.* 2017; **50**: 87–95.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
70. Guralink JM, LaCroix A, Everett D, *et al.*: **Aging in the Eighties: The Prevalence of Comorbidity and its Association with Disability.** *Advance Data from Vital and Health Statistics.* 1989.
[Reference Source](#)
71. **F** Sperling RA, Johnson KA, Doraiswamy PM, *et al.*: **Amyloid deposition detected with florbetapir F 18 (¹⁸F-AV-45) is related to lower episodic memory performance in clinically normal older individuals.** *Neurobiol Aging.* 2013; **34**(3): 822–31.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
72. Hachinski VC, Lassen NA, Marshall J: **Multi-infarct dementia. A cause of mental deterioration in the elderly.** *Lancet.* 1974; **2**(7874): 207–10.
[PubMed Abstract](#) | [Publisher Full Text](#)
73. O'Brien JT, Erkinjuntti T, Reisberg B, *et al.*: **Vascular cognitive impairment.** *Lancet Neurol.* 2003; **2**(2): 89–98.
[PubMed Abstract](#) | [Publisher Full Text](#)
74. Yu BP, Chung HY: **The inflammatory process in aging.** *Rev Clin Gerontol.* 2006; **16**(3): 179–87.
[Publisher Full Text](#)
75. Chung HY, Sung B, Jung KJ, *et al.*: **The molecular inflammatory process in aging.** *Antioxid Redox Signal.* 2006; **8**(3–4): 572–81.
[PubMed Abstract](#) | [Publisher Full Text](#)
76. Razay G, Vreugdenhil A, Wilcock G: **The metabolic syndrome and Alzheimer disease.** *Arch Neurol.* 2007; **64**(1): 93–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
77. Alexopoulos GS, Buckwalter K, Olin J, *et al.*: **Comorbidity of late life depression: an opportunity for research on mechanisms and treatment.** *Biol Psychiatry.* 2002; **52**(6): 543–58.
[PubMed Abstract](#) | [Publisher Full Text](#)
78. **F** Kawas CH, Kim RC, Sonnen JA, *et al.*: **Multiple pathologies are common and related to dementia in the oldest-old: The 90+ Study.** *Neurology.* 2015; **85**(6): 535–42.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. Michela Gallagher

Department of Psychological and Brain Sciences, Johns Hopkins University, Baltimore, MD, USA

Competing Interests: Michela Gallagher is the Founder of AgeneBio, Inc.. She serves as a scientific consultant and has financial interests in that company. She is also an inventor on technology that is licensed from Johns Hopkins University to AgeneBio, Inc.. Johns Hopkins University owns AgeneBio, Inc. stock and is entitled to royalties on a drug described in this article. This arrangement has been reviewed and approved by the Johns Hopkins University in accordance with its conflict of interest policies.

2. Heikki Tanila

A.I. Virtanen Institute for Molecular Sciences, University of Eastern Finland, Kuopio, Kuopio, Finland

Competing Interests: No competing interests were disclosed.

3. Morgan D Barese

Department of Psychology, University of Toronto, Toronto, ON, Canada

Competing Interests: No competing interests were disclosed.

4. Peter R Rapp

Laboratory of Behavioral Neuroscience, National Institute on Aging, Baltimore, Baltimore, USA

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research