REVIEW

Immunosurveillance by human $\gamma\delta$ T lymphocytes: the emerging role of butyrophilins [version 1; peer review: 2 approved]

Dieter Kabelitz, Marcus Lettau, Ottmar Janssen

Institute of Immunology, University of Kiel and University Hospital Schleswig-Holstein Campus Kiel, Arnold-Heller-Str. 3, Building 17, D-24105 Kiel, Germany

V1 First published: 05 Jun 2017, 6(F1000 Faculty Rev):782
<https://doi.org/10.12688/f1000research.11057.1>
 Latest published: 05 Jun 2017, 6(F1000 Faculty Rev):782
<https://doi.org/10.12688/f1000research.11057.1>

Abstract

In contrast to conventional T lymphocytes, which carry an $\alpha\beta$ T-cell receptor and recognize antigens as peptides presented by major histocompatibility complex class I or class II molecules, human $\gamma\delta$ T cells recognize different metabolites such as non-peptidic pyrophosphate molecules that are secreted by microbes or overproduced by tumor cells. Hence, $\gamma\delta$ T cells play a role in immunosurveillance of infection and cellular transformation. Until recently, it has been unknown how the $\gamma\delta$ T-cell receptor senses such pyrophosphates in the absence of known antigen-presenting molecules. Recent studies from several groups have identified a unique role of butyrophilin (BTN) protein family members in this process, notably of BTN3A1. BTNs are a large family of transmembrane proteins with diverse functions in lipid secretion and innate and adaptive immunity. Here we discuss current models of how BTN molecules regulate $\gamma\delta$ T-cell activation. We also address the implications of these recent findings on the design of novel immunotherapeutic strategies based on the activation of $\gamma\delta$ T cells.

Keywords

lymphocytes, butyrophilins, Immunosurveillance, T-cell, immunotherapeutic

Open Peer Review

Approval Status

	1	2
version 1
05 Jun 2017		

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **John Trowsdale**, University of Cambridge, Cambridge, UK
2. **Richard D. Lopez**, Duke University, Durham, USA

Any comments on the article can be found at the end of the article.

Corresponding author: Dieter Kabelitz (Dietrich.Kabelitz@uksh.de)

Competing interests: The authors declare that they have no competing interests.

Grant information: Work from the authors' laboratory has been supported by the Deutsche Forschungsgemeinschaft (Ka 502/16-1, JA 610/7-1, and ExC 306 Inflammation-at-Interfaces).

Copyright: © 2017 Kabelitz D *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Kabelitz D, Lettau M and Janssen O. **Immunosurveillance by human $\gamma\delta$ T lymphocytes: the emerging role of butyrophilins [version 1; peer review: 2 approved]** F1000Research 2017, 6(F1000 Faculty Rev):782 <https://doi.org/10.12688/f1000research.11057.1>

First published: 05 Jun 2017, 6(F1000 Faculty Rev):782 <https://doi.org/10.12688/f1000research.11057.1>

Introduction

T lymphocytes are the specific effector cells of the adaptive immune system. T-cell differentiation takes place in the thymus, giving rise to large numbers of mature CD4 and CD8 T cells expressing a clonally variable $\alpha\beta$ T-cell receptor (TCR). Interaction with thymic stromal cells, transcription factors, and cytokines together drive the differentiation of early thymic progenitor cells into mature CD4 and CD8 $\alpha\beta$ T cells which recognize peptides presented in the context of major histocompatibility complex (MHC) class I (CD8 T cells) or class II (CD4 T cells)¹. The functional diversification of mature T cells into T helper type 1 (Th1), Th2, and Th17 cells and memory cell subsets is induced by the context-dependent interaction with neighboring cells (e.g. dendritic and epithelial cells) and transcriptional networks which are further modulated by metabolic and epigenetic processes²⁻⁵. In addition to these “conventional” T cells, T cells with a highly restricted canonical $\alpha\beta$ TCR repertoire exist which recognize non-peptide antigens in the absence of restriction by classical MHC molecules. Such “unconventional” human $\alpha\beta$ T cells include the invariant natural killer T (iNKT) cells expressing a V α 24-J α 18-encoded TCR and the mucosa-associated invariant T (MAIT) cells expressing a V α 7.2-J α 33-encoded TCR. iNKT cells recognize endogenous and exogenous (microbial) lipids presented by CD1 (specifically CD1d) molecules, whereas MAIT cells recognize small intermediates generated in the riboflavin (vitamin B2) metabolic pathway which are presented by the MHC-related 1 (MR1) molecules⁶. Since unconventional $\alpha\beta$ T cells are not dependent on the antigen processing machinery like conventional CD4 and CD8 T cells, they can rapidly perform effector functions upon ligand recognition. Both iNKT and MAIT cells are found in the blood and at increased numbers not only in mucosal tissue but also in the liver and are important players in local immunosurveillance and anti-bacterial immunity^{7,8}. More recently, innate lymphoid cells (ILCs) have been identified as innate homologs of differentiated effector T cells which do not express clonally rearranged TCR but share similar transcription factor and cytokine specifications⁹. Subsets of ILCs interact with innate and adaptive immune cells, epithelial cells, and microbiota and thereby contribute to tissue repair, metabolic homeostasis, and local inflammation¹⁰.

While it might appear that the above outlined arsenal of available immune cells should suffice to combat all dangerous (infectious and non-infectious) antigens, evolution has conserved yet another class of unconventional T cells, i.e. T lymphocytes carrying a CD3-associated $\gamma\delta$ TCR heterodimer rather than the $\alpha\beta$ TCR. It has been known for a long time that the major population of $\gamma\delta$ T cells found in the peripheral blood of adults specifically recognizes non-peptidic small microbial pyrophosphate molecules, again without requirement for a dedicated MHC class I, MHC class II, or CD1 presenting molecule¹¹⁻¹³. It thus remained a mystery for many years how such “phosphoantigens” (pAg) contained in crude bacterial lysates¹⁴, purified by preparative anion exchange chromatography¹⁵, or chemically synthesized¹⁶ can trigger such potent $\gamma\delta$ T-cell responses.

A landmark paper addressing the activation requirements of human $\gamma\delta$ T cells in response to pAg was published in 2012 by Harly and

co-workers¹⁷. These authors reported the unexpected finding that a member of the transmembrane butyrophilin (BTN) proteins was absolutely required for the activation of human $\gamma\delta$ T cells by microbial or endogenous pAg. On the grounds of these findings, several groups set out to study the precise role of BTN proteins at the molecular level. Surprisingly, these investigations resulted in quite controversial models, assigning an essential role to either the extracellular¹⁸ or the intracellular domain¹⁹ of a particular BTN3A isoform. In this review, we discuss the current knowledge on the interplay of human $\gamma\delta$ TCR with specific BTN proteins, both in terms of basic mechanisms of $\gamma\delta$ T-cell activation and with respect to improving future strategies of $\gamma\delta$ T-cell-based immunotherapies.

$\gamma\delta$ T cells: unconventional T lymphocytes linking innate and adaptive immunity

$\gamma\delta$ T cells account for approximately 2–5% of peripheral blood T cells in healthy adult donors but are present at much higher numbers in mucosal tissues, where they comprise 20–30% of intraepithelial lymphocytes in the small intestine²⁰. In contrast to $\alpha\beta$ T cells, there are only few variable (V) gene segments available in the germline genome which can be used during intrathymic TCR gene rearrangement to express functional TCR proteins. In humans, there are six expressed V γ genes (V γ 2, 3, 4, 5, 8, and 9) and a similarly limited number of V δ genes²¹. Nonetheless, $\gamma\delta$ TCR can display an enormous CDR3 loop diversity²². $\gamma\delta$ T cells expressing particular V γ V δ pairing are not randomly distributed but are preferentially located in certain compartments. Thus, the majority of $\gamma\delta$ T cells in peripheral blood express V δ 2 (paired almost exclusively with V γ 9), while intraepithelial $\gamma\delta$ T cells frequently express V δ 1 (or other non-V δ 2 segments) which can pair with different V γ elements¹³. Most $\gamma\delta$ T cells lack CD4 and CD8 surface expression, well in line with their MHC-independent ligand recognition (note, however, that a substantial proportion of $\gamma\delta$ T cells can express CD8 at low levels). Major efforts were made over the years to identify antigens and ligands that are specifically recognized by the $\gamma\delta$ TCR^{22,23}. A list of some currently identified ligands for human $\gamma\delta$ T cells is presented in Table 1. Obviously, the best-characterized ligands are prokaryotic and eukaryotic pAg, which are exclusively recognized by human V δ 2V γ 9 $\gamma\delta$ T cells²⁴. Such pyrophosphates

Table 1. Some examples of ligands that are specifically recognized by subsets of human $\gamma\delta$ T cells.

$\gamma\delta$ TCR	Ligand	References
V δ 2V γ 9	Prokaryotic pAg, HMBPP	15
	Eukaryotic pAg, IPP	26
	F1-ATPase + apolipoprotein A-I	28
	hMSH2	27
V δ 1	MICA	31
	CD1d-lipid	30
V δ 5	EPCR	29

Abbreviations: EPCR, endothelial protein C receptor; HMBPP, (E)-4-hydroxy-3-methyl-but-2-enyl pyrophosphate; hMSH2, human MutS homolog 2; IPP, isopentenyl pyrophosphate; MICA, major histocompatibility complex class I-related chain A; pAg, phosphoantigen

are intermediates of the eukaryotic mevalonate or the prokaryotic non-mevalonate (also termed Rohmer's) pathway of isoprenoid synthesis^{15,25,26}. Other ligands for V δ 2V γ 9 T cells include the ectopically expressed DNA mismatch repair protein hMSH2²⁷ and F1-ATPase together with apolipoprotein A-I²⁸. Some of the identified ligands for non-V δ 2 $\gamma\delta$ TCR include endothelial protein C receptor²⁹ and lipids bound to CD1d³⁰ but also the stress-inducible MHC class I-related chain A (MICA) molecules³¹. In all instances, recognition of respective ligands by the $\gamma\delta$ TCR is a rapid event and addresses all $\gamma\delta$ T cells carrying the appropriate TCR with little, if any, contribution of CDR3 variation. Since V δ 2V γ 9 cells comprise the vast majority (up to 95%) of peripheral blood $\gamma\delta$ T cells and all respond to pAg stimulation with no need for antigen processing, this implies that a large proportion (2–4%) of all peripheral blood T cells is rapidly activated (e.g. to produce cytokines including interferon- γ and tumor necrosis factor- α) upon encounter of such pAg¹³.

In addition to the TCR, $\gamma\delta$ T cells express other activating cell surface receptors, notably natural killer group 2 member D (NKG2D), which is a receptor for multiple stress-inducible MHC class I-related molecules including MICA/B and six members of the UL16 binding protein family (ULBP1-6)³². NKG2D is expressed on innate natural killer (NK) cells, some CD8 and CD4 T cells, and essentially all $\gamma\delta$ T cells. While normal cells usually do not express NKG2D ligands, cell surface expression is induced by cell stress, DNA damage, and cellular transformation. Upon ligand binding, NKG2D transmits cellular activation via the PI3-kinase pathway, resulting in cytokine production and

triggering of cytotoxic activity³³. NKG2D ligands can be released from the surface of tumor cells via protease-mediated shedding or via exosome secretion, and soluble NKG2D ligands may block NKG2D receptor activation and thereby serve as a tumor immune escape mechanism³⁴.

Furthermore, $\gamma\delta$ T cells can also express some receptors specifically associated with the innate immune system, notably Toll-like receptors (TLRs), and corresponding TLR ligands can co-stimulate $\gamma\delta$ T-cell activation³⁵. Conceivably, such effects might be primarily mediated via monocytic and/or dendritic cells when heterogeneous cell populations are investigated³⁶, but it has also been demonstrated that purified $\gamma\delta$ T cells express certain TLRs and directly respond to TLR ligand co-stimulation^{37,38}. To summarize, $\gamma\delta$ T cells express receptors of both the innate (e.g. NKG2D and TLR) and the adaptive (TCR) immune system, and the outcome of functional responses is regulated through integration of various signaling pathways (Figure 1). There is another feature of human V δ 2V γ 9 T cells which further places them as a link between the innate and adaptive immune systems: as initially reported by Bernhard Moser's group, activated $\gamma\delta$ T cells can serve as antigen-presenting cells to specifically stimulate peptide-specific $\alpha\beta$ T cells³⁹. Importantly, $\gamma\delta$ T cells can even take up antigen particles, process such antigens intracellularly, and load corresponding peptides onto MHC class I molecules for cross-presentation to antigen-specific CD8 $\alpha\beta$ T cells, a process normally restricted to "professional" antigen-presenting cells such as dendritic cells⁴⁰. The antigen-presenting capacity of $\gamma\delta$ T cells may help to initiate a subsequent tumor antigen-specific CD8 T-cell response once $\gamma\delta$

Figure 1. Three classes of receptors regulating human $\gamma\delta$ T-cell activation. (1) The activating natural killer (NK) receptor NK group 2 member D (NKG2D) recognizes stress-inducible ligands including major histocompatibility complex class I-related chain A (MICA)/MICB and UL16 binding protein family (ULBP) 1–6 and triggers cytotoxic activity and cytokine production via the phosphoinositide 3 (PI3)-kinase pathway. (2) The CD3-associated T-cell receptor (TCR) recognizes ligands including “phosphoantigens” (pAg) in a butyrophilin (BTN) 3A-dependent way and human MutS homolog 2 (hMSH2) (V δ 2V γ 9 TCR), or lipids bound to CD1d and endothelial protein C receptor (EPCR) (non-V δ 2 TCR). (3) Pattern recognition receptors including Toll-like receptors (TLRs) sense conserved microbial ligands such as acetylated lipids (TLR2/6 heterodimer) or poly I:C (intracellular TLR3) and co-stimulate $\gamma\delta$ T-cell activation via the nuclear factor kappa-light-chain-enhancer of activated B cells (NF- κ B) pathway. APC, antigen-presenting cell.

T cells have killed opsonized tumor cells and taken up apoptotic tumor cell fragments⁴¹.

$\gamma\delta$ T cells: important players in anti-tumor immunity

Many solid tumors and leukemia/lymphoma cells are quite susceptible to $\gamma\delta$ T-cell-mediated lysis. In contrast to $\alpha\beta$ T cells, $\gamma\delta$ T cells recognize tumor cells not on the basis of tumor-specific antigenic peptides presented in the context of MHC class I or class II molecules but rather sense cell-surface-expressed stress molecules and/or metabolites of the dysregulated mevalonate pathway. Moreover, $\gamma\delta$ T cells can make use of both the TCR and the NKG2D receptor to recognize and kill tumor cells^{42,43}. Interestingly, the sensitivity of tumor cells to lysis by V δ 2V γ 9 $\gamma\delta$ T cells can be pharmacologically manipulated. Nitrogen-containing bisphosphonates (N-BPs) such as zoledronic acid are in clinical use to treat diseases associated with bone resorption. In addition to their anti-resorptive bone activity, N-BPs also interfere with the mevalonate metabolic pathway where $\gamma\delta$ T-cell-stimulating pyrophosphates are generated²⁵. N-BPs block an enzyme downstream of the synthesis of isopentenyl pyrophosphate (IPP), leading to increased accumulation of IPP and thereby to $\gamma\delta$ T-cell activation⁴⁴. Therefore, pretreatment of tumor cells with N-BP increases their susceptibility to $\gamma\delta$ T-cell-mediated lysis⁴⁵. Application of N-BPs to patients also induces *in vivo* activation of $\gamma\delta$ T cells⁴⁶, and in fact some clinical responses have been noted in small-scale studies in cancer patients given intravenous N-BPs together with low-dose interleukin-2⁴⁷. Moreover, $\gamma\delta$ T cells have also been adoptively transferred to cancer patients, with no obvious major adverse effects but some clinical responses in a few patients⁴⁷. The efficacy of tumor cell killing by $\gamma\delta$ T cells can be further increased by specifically targeting $\gamma\delta$ T cells to tumor cells via antibody-mediated cellular cytotoxicity (ADCC)⁴⁸ or bispecific antibody constructs^{49,50}. While V δ 2V γ 9 $\gamma\delta$ T cells can be easily activated and expanded to large cell numbers by activation with pAg or N-BPs, it should be

kept in mind that non-V δ 2 subsets of $\gamma\delta$ T cells might also have potent anti-tumor activity, and protocols for selective expansion of those $\gamma\delta$ T cells are in development⁵¹. Attempts to explore the anti-tumor capacity of $\gamma\delta$ T cells in a clinical setting were boosted by the recent demonstration in a large patient cohort that the proportion of $\gamma\delta$ T cells among tumor-infiltrating immune cells was the best positive predictive parameter across a multitude of human tumor entities⁵². On the other hand, however, it must be considered that $\gamma\delta$ T cells might also negatively regulate anti-tumor immune responses. For instance, it has been demonstrated that $\gamma\delta$ T cells infiltrating into human breast cancer have a regulatory activity and inhibit $\alpha\beta$ T-cell responses⁵³. Moreover, other potentially tumor-promoting activities of $\gamma\delta$ T cells have been reported in colorectal and pancreatic cancer^{54,55}. Overall, however, it appears that $\gamma\delta$ T cells are interesting and promising candidates for cellular immunotherapy supplementing other strategies such as NK cells and chimeric antigen receptor (CAR) T cells^{56,57}.

Butyrophilins: a large family of proteins with immunomodulatory functions

BTNs were originally described as plasma-membrane-associated glycoproteins in the lactating mammary glands of many species which constitute a major component of the milk fat globule membrane⁵⁸. The type 1 transmembrane BTN proteins belong to the immunoglobulin (Ig) superfamily and typically consist of extracellular Ig-like domains (IgV and IgC), a transmembrane domain, and, in some but not all cases, an intracellular B30.2 signaling domain^{59–61}. BTN and BTN-like (BTNL) proteins are variably related to the B7 family of costimulatory molecules (e.g., CD80 and CD86) which supports the role of (at least) some BTN members in the immune system⁶². The genes are clustered in two regions on human chromosome 6: *BTN* telomeric to HLA class I genes and *BTNL* near the HLA-DR genes. An additional *BTNL* gene cluster is located on human chromosome 5q35⁶¹. The protein domain structure of some functionally important BTN and BTNL

Figure 2. Domain structure of butyrophilin (BTN) proteins. Typically, BTN and BTN-like (BTNL) proteins consist of extracellular immunoglobulin V (IgV)- and IgC-like domains, a transmembrane domain, and a cytosolic B30.2 (or PRYSPRY) domain. Three isoforms of BTN3A differ in their cytosolic structure (BTN3A1: B30.2 domain; BTN3A2: no B30.2 domain; BTN3A3: B30.2 domain plus additional stretch of amino acids). BTNL2 has two tandem repeats of IgV and IgC domains and lacks the cytosolic B30.2 domain.

members is shown in [Figure 2](#). The cytosolic B30.2 domain (also termed PRYSPRY) and the homologous SPRY domain are present in many cellular proteins, including tripartite motif molecules (TRIM), where they potentially interact with diverse intracellular molecules including NOD2, retroviral capsids, or Fc parts of IgG^{63,64}. Given that BTN molecules have multiple roles in innate and adaptive immunity, it comes as no surprise that *BTN* gene polymorphisms may influence disease susceptibility. As an example, *BTN3A2* has been shown to be associated with susceptibility to type I diabetes⁶⁵, and more examples are discussed in [60](#). Interestingly, genetic variants in *BTN* genes can also alter susceptibility to infection, as has been demonstrated for a selection of hepatitis C virus genotypes and subsequent disease progression⁶⁶.

BTN proteins are widely expressed in immune cells and epithelial cells and can exert a multitude of immunoregulatory activities^{60,61}. While a few binding partners have been identified (such as DC-SIGN [CD209] on dendritic cells and monocytes for BTN2A1⁶⁷), specific receptors are largely unknown. It also appears that the overall effect of specific BTN molecules depends on experimental conditions and respective reagents (recombinant proteins, soluble or immobilized antibodies, cell-surface-expressed molecules, etc.). The extracellular domain of the human BTNL protein BTNL8 co-stimulates proliferation and cytokine production of anti-CD3 antibody-stimulated CD4 and CD8 $\alpha\beta$ T cells, and a putative BTNL8 receptor on the cell surface of resting T cells was detected by flow cytometry with a BTNL8-Fc fusion protein⁶⁸. Olive's group has generated a number of monoclonal antibodies (mAb) directed against the extracellular domain of BTN3A molecules (also termed CD277), which do not differentiate between the three isoforms BTN3A1, BTN3A2, and BTN3A3⁶². They showed that anti-CD277 mAb clone 20.1 co-stimulated cytokine production and early signaling cascades in purified human CD4 and CD8 T cells when immobilized together with anti-CD3 mAb in cell culture plates or on microbeads, pointing to an $\alpha\beta$ T cell co-stimulating activity of BTN3A⁶⁹. In contrast, another anti-CD277 mAb specific for a different epitope in the extracellular region of BTN3A (clone 232-5) rather inhibited T-cell activation when added to anti-CD3 mAb activated CD4 or CD8 T cells⁷⁰. While these studies indicated that CD277 expressed on T cells can transmit positive or negative co-stimulatory signals, other studies showed that CD277 overexpressed on antigen-presenting cells profoundly inhibited T-cell proliferation and cytokine production⁷¹. Using a Myc-tagged extracellular CD277 construct, Cubillos-Ruiz also obtained evidence for the expression of a CD277-binding protein on activated but not resting human T cells⁷¹. Together with their observation of strong CD277 expression on most of the analyzed ovarian cancer tissues, these authors argued that CD277 is a negative regulator of human T-cell activation with relevance for the immunosuppressive tumor microenvironment⁷¹. This view, however, does not integrate the fact that T cells themselves (like most, if not all, other immune cells) strongly express CD277^{62,69,72}.

Other BTN/BTNL members with reported modulatory activity on $\alpha\beta$ T-cell activation include BTN2A2 and BTNL2. The extracellular part of BTN2A2 (i.e. a BTN2A2-Fc fusion protein) inhibited early signaling events in anti-CD3 plus anti-CD28 mAb

stimulated murine T cells and prevented cell cycle entry. Interestingly, BTN2A2-Fc also induced *de novo* expression of FoxP3 in anti-CD3/anti-CD28 mAb activated naive CD4 T cells, suggesting that BTN2A2 may attenuate T-cell activation through multiple pathways, including the induction of FoxP3-expressing regulatory T cells (Tregs)⁷³. Recently, *BTN2A2*^{-/-} mice were reported to exhibit enhanced T-cell responses as shown by greater severity of T-cell-dependent models of autoimmunity (experimental autoimmune encephalomyelitis [EAE]) but also enhanced response to tumor vaccination⁷⁴. Quite similar to BTN2A2, BTNL2 also inhibits murine T-cell activation and co-stimulates FoxP3 induction and thus Treg induction when applied as immobilized BTNL2-Fc fusion protein⁷⁵. Interestingly, *BTNL2* was found to be upregulated during acute-phase malaria infection, pointing to a possible feedback loop between inhibitory BTN/BTNL molecules and T-cell activation in inflammation and infection⁷⁶.

In addition to immune cells, BTN and BTNL proteins are expressed in the intestine and regulate tissue integrity, local immune responses, and inflammation. Recent gene expression data point to a correlation of upregulated intestinal *BTN/BTNL* gene expression with inflammatory bowel diseases, in line with an important role of BTN/BTNL proteins in shaping local T-cell responses⁷⁷. In murine intestinal epithelial cells, BTNL1 and BTNL6 form heteromeric complexes which enhance the proliferative activity of intraepithelial lymphocytes, specifically of a subset of $\gamma\delta$ T cells expressing the V γ 7V δ 4 TCR^{78,79}. Interestingly, it was recently shown that human gut epithelial cells express BTNL3 and BTNL8, which together also regulate tissue-specific $\gamma\delta$ T cells, in this case intestinal $\gamma\delta$ T cells expressing a V γ 4 TCR⁸⁰. Taken together, it is obvious that BTN and BTNL proteins regulate multiple T-cell responses in a negative or positive manner. One of the unsolved questions here is how such signals are transmitted to T cells (i.e. the nature of putative receptors), an issue which needs further investigation.

The puzzling role of BTN3A1 in $\gamma\delta$ T-cell activation

As already mentioned, an indispensable role of CD277/BTN3A in the activation of human V δ 2V γ 9 $\gamma\delta$ T cells by microbial or tumor-derived pAg was reported by Harly and colleagues¹⁷. These authors used CD277 knockdown and domain-shuffling strategies to demonstrate the importance of the BTN3A1 isoform (carrying the cytosolic B30.2 domain) in this process. They went on to show that the anti-CD277 mAb 20.1 (used in immobilized form in previous studies to demonstrate co-stimulatory activity on CD4 and CD8 $\alpha\beta$ T cells⁶⁹) could selectively activate V δ 2V γ 9 T cells when added in soluble form together with interleukin-2 to peripheral blood mononuclear cells, and furthermore sensitized a broad range of tumor cells to killing by $\gamma\delta$ T cells¹⁷. They also identified another anti-CD277 antibody termed 103.2, which binds to a different epitope on the extracellular part of BTN3A and specifically inhibited V δ 2V γ 9 T-cell activation by pAg, N-BP, or agonistic mAb 20.1^{17,81}. The essential role of BTN3A1 for pAg stimulation of human $\gamma\delta$ T cells was confirmed by several other reports^{18,19,82–86}. While the original study by Harly *et al.* described the importance of BTN3A/CD277 for human $\gamma\delta$ T-cell activation by pyrophosphate antigens, the molecular mechanism was not yet precisely identified. Subsequently, two largely conflicting models

Figure 3. Alternative models of the role of butyrophilin (BTN) 3A molecules in phosphoantigen (pAg)-mediated $\gamma\delta$ T-cell activation. **A:** pAg directly bind to the extracellular immunoglobulin V (IgV) domain of BTN3A1 and are then “presented” to the $\gamma\delta$ TCR; as-yet-unidentified transporter molecules would shuffle pAg (generated within tumor cells due to the dysregulated mevalonate pathway) from the cytosol to the extracellular compartment for presentation by BTN3A1^{18,87}. **B:** pAg bind to the cytosolic B30.2 domain, leading to the recruitment of linker proteins including perioplakin and the activation of the RhoB GTPase, which together induce a spatial redistribution of BTN3A1 molecules recognized by the $\gamma\delta$ TCR^{19,85,91}. APC, antigen-presenting cell; TCR, T-cell receptor.

were proposed, i.e. a “presenting” versus a “pyrophosphate sensing” function of CD277 (Figure 3). Vavassori and co-workers reported that pAg IPP and HMBPP could directly bind with low affinity to the recombinant extracellular IgV domain of BTN3A1; furthermore, they also observed weak binding of recombinant soluble V δ 2V γ 9 TCR to immobilized BTN3A1 molecules, which was further facilitated by IPP¹⁸. These observations were in line with BTN3A1 serving as an antigen-presenting molecule for pAg to be specifically recognized by the human V δ 2V γ 9 TCR (Figure 3A^{18,87}). However, direct binding of pAg to the extracellular domain of BTN3A1 could not be verified by other groups^{19,82,84}. Instead, Sandstrom and co-workers demonstrated that the cytosolic B30.2 domain could directly bind several $\gamma\delta$ T-cell-stimulating pAg through a positively charged surface pocket¹⁹, an observation which was confirmed by others^{84–86,88}. Currently, most available data thus support the pyrophosphate-sensing function of the cytosolic B30.2 domain^{89,90}. How, then, can binding of pyrophosphates to the cytosolic domain of a trans-membrane protein (BTN3A1) translate into TCR-dependent selective activation of a specific (V δ 2V γ 9) $\gamma\delta$ T-cell subset? Recent progress in the field has helped to elucidate some of the molecular mechanisms. Overall, it appears that an “inside-out” signaling mechanism induced by intracellular pAg sensing conveys a spatial redistribution or conformational change of the extracellular CD277 domain, which is then somehow recognized by the V δ 2V γ 9 TCR^{89,91}. Recently, crucial steps along this pathway have been identified. The cytoskeletal adaptor protein perioplakin has been shown to interact with a membrane-proximal intracellular part of BTN3A1⁸⁵. Perioplakin is a member of a family of cytoskeletal linker proteins that interact with various membrane-associated proteins and are involved in cytoskeletal (re)organization⁹². It is

thus conceivable that upon pAg binding the interaction of cytosolic parts with perioplakin and possibly additional adaptor proteins contributes to the spatial rearrangement of BTN3A1⁸⁵. Another step in this process is the small GTPase RhoB, which was recently identified in a genome-wide screen as an important component in BTN3A1-dependent tumor cell recognition by V δ 2V γ 9 T cells⁹¹. RhoB interacts with and regulates the membrane mobility of BTN3A1, and intracellular redistribution of RhoB in different tumor cells correlated with their recognition by $\gamma\delta$ T cells. These results point to a correlation in tumor cells between the dysregulated mevalonate pathway (which also controls small GTPases), RhoB activity, accumulation of pyrophosphates, and sensitivity to $\gamma\delta$ T-cell killing⁹¹.

Taken together, it is now clear that BTN3A/CD277 is required for pAg-mediated activation of V δ 2V γ 9 $\gamma\delta$ T cells and that perioplakin and RhoB have important roles in spatial rearrangement of BTN3A1 following intracellular pAg sensing (Figure 3B). However, some pieces of the puzzle are still unsolved. While a membrane reorganization of BTN3A1 (which can apparently also be induced by agonistic anti-CD277 antibodies⁸⁹) is a crucial step, it is not yet known what precisely then the $\gamma\delta$ TCR recognizes. Moreover, and in contrast to initial studies¹⁷, it appears that it is not just the BTN3A1 isoform and its cytosolic B30.2 domain which are involved in pAg-mediated $\gamma\delta$ T cell activation—BTN3A2 and BTN3A3 isoforms have been implicated⁸⁵. Furthermore, using Chinese hamster ovary cells retrovirally transduced with the human *BTN3A1* gene and additionally harboring, or not, human chromosome 6, Riaño and co-workers obtained evidence for a role for additional genes in the BTN3A1-dependent activation of $\gamma\delta$ T cells by pAg⁸³. Therefore, we can expect to witness the

discovery of additional new players before we fully understand how cell-surface-rearranged BTN3A molecules and pAg exclusively activate human V δ 2V γ 9 T cells. Along this line, it will be important to study in more detail the role of various accessory cells in this process. BTN3A is widely expressed on leukocytes, yet only monocytes serve as accessory cells for three mechanistically different stimuli for V δ 2V γ 9 T cells, i.e. N-BP, pAg, and agonistic anti-CD277 mAb 20.1⁷². In co-cultures with purified $\gamma\delta$ T cells, purified CD277-positive CD4 T cells can “present” pAg HMBPP to $\gamma\delta$ T cells⁷². Given that HMBPP is most likely not directly presented by extracellular BTN3A1 domains^{89,90}, how then do pAg enter the cell to initiate $\gamma\delta$ T-cell activation following binding to cytosolic B30.2? A putative transporter molecule has been postulated⁸⁷ (Figure 3A) and recently an energy-dependent uptake of HMBPP was demonstrated⁸⁶, but the precise molecular mechanisms remain to be clarified.

Concluding remarks

BTN and BTNL have emerged as potent immunomodulatory proteins. The T-cell-inhibitory activity of some BTN/BTNL members suggests that they might be novel targets for checkpoint inhibitors, in addition to established checkpoint proteins such as CTLA-4, PD1, and PD-L1⁹³. Some BTN/BTNL proteins have a unique role in recruiting and activating particular subsets of unconventional $\gamma\delta$ T cells. The recently discovered role of BTNL3 and BTNL8 for shaping V γ 4 $\gamma\delta$ T cells in the human gut might be

relevant for the loss of mucosal barrier function in inflammatory bowel diseases. The perhaps most fascinating example is the *ménage à trois* of BTN3A, pyrophosphate molecules, and the human V δ 2V γ 9 TCR. The availability of anti-BTN3A/CD277 antibodies which selectively activate (e.g. mAb 20.1) or inhibit (e.g. mAb 103.2) V δ 2V γ 9 $\gamma\delta$ T cells opens new avenues for $\gamma\delta$ T-cell-directed immunotherapies. In a pre-clinical xenotransplantation model of acute myeloid leukemia, therapeutic application of mAb 20.1 enhanced the therapeutic efficacy of adoptively transferred V δ 2V γ 9 T cells⁹⁴. Therefore, humanized agonistic anti-BTN3A/CD277 antibodies might be a novel and highly specific approach to activate tumor-reactive $\gamma\delta$ T cells *in vivo*. Vice versa, humanized inhibitory anti-BTN3A/CD277 antibodies might represent potent reagents for selective silencing of V δ 2V γ 9 T cells in clinical conditions where they might contribute to the disease process, e.g. in autoimmune diseases⁹⁵.

Competing interests

The authors declare that they have no competing interests.

Grant information

Work from the authors' laboratory has been supported by the Deutsche Forschungsgemeinschaft (Ka 502/16-1, JA 610/7-1, and ExC 306 Inflammation-at-Interfaces).

References

- Blom B, Spits H: **Development of human lymphoid cells.** *Annu Rev Immunol.* 2006; **24**: 287–320.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Dimeloe S, Burgener AV, Grähler J, *et al.*: **T-cell metabolism governing activation, proliferation and differentiation; a modular view.** *Immunology.* 2017; **150**(1): 35–44.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
- De Obaldia ME, Bhandoola A: **Transcriptional regulation of innate and adaptive lymphocyte lineages.** *Annu Rev Immunol.* 2015; **33**: 607–42.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
- Wang C, Collins M, Kuchroo VK: **Effector T cell differentiation: are master regulators of effector T cells still the masters?** *Curr Opin Immunol.* 2015; **37**: 6–10.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Durek P, Nordström K, Gasparoni G, *et al.*: **Epigenomic Profiling of Human CD4⁺ T Cells Supports a Linear Differentiation Model and Highlights Molecular Regulators of Memory Development.** *Immunity.* 2016; **45**(5): 1148–61.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
- Mori L, Lepore M, De Libero G: **The Immunology of CD1- and MR1-Restricted T Cells.** *Annu Rev Immunol.* 2016; **34**: 479–510.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Gao Y, Williams AP: **Role of Innate T Cells in Anti-Bacterial Immunity.** *Front Immunol.* 2015; **6**: 302.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
- Godfrey DI, Uldrich AP, McCluskey J, *et al.*: **The burgeoning family of unconventional T cells.** *Nat Immunol.* 2015; **16**(11): 1114–23.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
- Kang J, Malhotra N: **Transcription factor networks directing the development, function, and evolution of innate lymphoid effectors.** *Annu Rev Immunol.* 2015; **33**: 505–38.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Tait Wojno ED, Artis D: **Emerging concepts and future challenges in innate lymphoid cell biology.** *J Exp Med.* 2016; **213**(11): 2229–48.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
- Constant P, Davodeau F, Peyrat MA, *et al.*: **Stimulation of human gamma delta T cells by nonpeptidic mycobacterial ligands.** *Science.* 1994; **264**(5156): 267–70.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Morita CT, Beckman EM, Bukowski JF, *et al.*: **Direct presentation of nonpeptide prenyl pyrophosphate antigens to human gamma delta T cells.** *Immunity.* 1995; **3**(4): 495–507.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Bonneville M, O'Brien RL, Born WK: **Gammadelta T cell effector functions: a blend of innate programming and acquired plasticity.** *Nat Rev Immunol.* 2010; **10**(7): 467–78.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kabelitz D, Bender A, Schondelmaier S, *et al.*: **A large fraction of human peripheral blood gamma/delta + T cells is activated by Mycobacterium tuberculosis but not by its 65-kD heat shock protein.** *J Exp Med.* 1990; **171**(3): 667–79.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Hintz M, Reichenberg A, Altincicek B, *et al.*: **Identification of (E)-4-hydroxy-3-methyl-but-2-enyl pyrophosphate as a major activator for human gammadelta T cells in Escherichia coli.** *FEBS Lett.* 2001; **509**(2): 317–22.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
- Espinosa E, Belmant C, Pont F, *et al.*: **Chemical synthesis and biological activity of bromohydrin pyrophosphate, a potent stimulator of human gamma delta T cells.** *J Biol Chem.* 2001; **276**(21): 18337–44.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Harly C, Guillaume Y, Nedellec S, *et al.*: **Key implication of CD277/butyrophilin-3 (BTN3A) in cellular stress sensing by a major human $\gamma\delta$ T-cell subset.** *Blood.* 2012; **120**(11): 2269–79.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
- Vavassori S, Kumar A, Wan GS, *et al.*: **Butyrophilin 3A1 binds phosphorylated antigens and stimulates human $\gamma\delta$ T cells.** *Nat Immunol.* 2013; **14**(9): 908–16.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
- Sandstrom A, Peigné CM, Léger A, *et al.*: **The intracellular B30.2 domain of butyrophilin 3A1 binds phosphoantigens to mediate activation of human V γ 9V δ 2 T cells.** *Immunity.* 2014; **40**(4): 490–500.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**

20. Kalyan S, Kabelitz D: **Defining the nature of human $\gamma\delta$ T cells: a biographical sketch of the highly empathetic.** *Cell Mol Immunol.* 2013; 10(1): 21–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
21. Hinz T, Wesch D, Halary F, *et al.*: **Identification of the complete expressed human TCR V gamma repertoire by flow cytometry.** *Int Immunol.* 1997; 9(8): 1065–72.
[PubMed Abstract](#) | [Publisher Full Text](#)
22. Chien YH, Meyer C, Bonneville M: **$\gamma\delta$ T cells: first line of defense and beyond.** *Annu Rev Immunol.* 2014; 32: 121–55.
[PubMed Abstract](#) | [Publisher Full Text](#)
23. Vantourout P, Hayday A: **Six-of-the-best: unique contributions of $\gamma\delta$ T cells to immunology.** *Nat Rev Immunol.* 2013; 13(2): 88–100.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
24. Bukowski JF, Morita CT, Tanaka Y, *et al.*: **V gamma 2V delta 2 TCR-dependent recognition of non-peptide antigens and Daudi cells analyzed by TCR gene transfer.** *J Immunol.* 1995; 154(3): 998–1006.
[PubMed Abstract](#)
25. Kabelitz D: **Small molecules for the activation of human gammadelta T cell responses against infection.** *Recent Pat Antiinfect Drug Discov.* 2008; 3(1): 1–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
26. Tanaka Y, Morita CT, Tanaka Y, *et al.*: **Natural and synthetic non-peptide antigens recognized by human gamma delta T cells.** *Nature.* 1995; 375(6527): 155–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
27. Dai Y, Chen H, Mo C, *et al.*: **Ectopically expressed human tumor biomarker MutS homologue 2 is a novel endogenous ligand that is recognized by human $\gamma\delta$ T cells to induce innate anti-tumor/virus immunity.** *J Biol Chem.* 2012; 287(20): 16812–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
28. **F** Scotet E, Martinez LO, Grant E, *et al.*: **Tumor recognition following Vgamma9Vdelta2 T cell receptor interactions with a surface F1-ATPase-related structure and apolipoprotein A-I.** *Immunity.* 2005; 22(1): 71–80.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
29. **F** Willcox CR, Pitard V, Netzer S, *et al.*: **Cytomegalovirus and tumor stress surveillance by binding of a human $\gamma\delta$ T cell antigen receptor to endothelial protein C receptor.** *Nat Immunol.* 2012; 13(9): 872–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
30. Uldrich AP, Le Nours J, Pellicci DG, *et al.*: **CD1d-lipid antigen recognition by the $\gamma\delta$ TCR.** *Nat Immunol.* 2013; 14(11): 1137–45.
[PubMed Abstract](#) | [Publisher Full Text](#)
31. Xu B, Pizarro JC, Holmes MA, *et al.*: **Crystal structure of a gammadelta T-cell receptor specific for the human MHC class I homolog MICA.** *Proc Natl Acad Sci U S A.* 2011; 108(6): 2414–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
32. Lanier LL: **NKG2D Receptor and Its Ligands in Host Defense.** *Cancer Immunol Res.* 2015; 3(2): 575–82.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
33. El-Gazzar A, Groh V, Spies T: **Immunobiology and conflicting roles of the human NKG2D lymphocyte receptor and its ligands in cancer.** *J Immunol.* 2013; 191(4): 1509–15.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
34. Chitadze G, Bhat J, Lettau M, *et al.*: **Generation of soluble NKG2D ligands: proteolytic cleavage, exosome secretion and functional implications.** *Scand J Immunol.* 2013; 78(2): 120–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
35. Wesch D, Peters C, Oberg H, *et al.*: **Modulation of $\gamma\delta$ T cell responses by TLR ligands.** *Cell Mol Life Sci.* 2011; 68(14): 2357–70.
[PubMed Abstract](#) | [Publisher Full Text](#)
36. Kunzmann V, Kretschmar E, Herrmann T, *et al.*: **Polyinosinic-polycytidylic acid-mediated stimulation of human gammadelta T cells via CD11c dendritic cell-derived type I interferons.** *Immunology.* 2004; 112(3): 369–77.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
37. Wesch D, Beetz S, Oberg H, *et al.*: **Direct costimulatory effect of TLR3 ligand poly(I:C) on human gamma delta T lymphocytes.** *J Immunol.* 2006; 176(3): 1348–54.
[PubMed Abstract](#) | [Publisher Full Text](#)
38. Pietschmann K, Beetz S, Welte S, *et al.*: **Toll-like receptor expression and function in subsets of human gammadelta T lymphocytes.** *Scand J Immunol.* 2009; 70(3): 245–55.
[PubMed Abstract](#) | [Publisher Full Text](#)
39. **F** Brandes M, Willmann K, Moser B: **Professional antigen-presentation function by human gammadelta T Cells.** *Science.* 2005; 309(5732): 264–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
40. Brandes M, Willmann K, Bioley G, *et al.*: **Cross-presenting human gammadelta T cells induce robust CD8 α alphabeta T cell responses.** *Proc Natl Acad Sci U S A.* 2009; 106(7): 2307–12.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
41. Himoudi N, Morgenstern DA, Yan M, *et al.*: **Human $\gamma\delta$ T lymphocytes are licensed for professional antigen presentation by interaction with opsonized target cells.** *J Immunol.* 2012; 188(4): 1708–16.
[PubMed Abstract](#) | [Publisher Full Text](#)
42. Rincon-Orozco B, Kunzmann V, Wrobel P, *et al.*: **Activation of V gamma 9V delta 2 T cells by NKG2D.** *J Immunol.* 2005; 175(4): 2144–51.
[PubMed Abstract](#) | [Publisher Full Text](#)
43. Wrobel P, Shojaei H, Schitteck B, *et al.*: **Lysis of a broad range of epithelial tumour cells by human gamma delta T cells: involvement of NKG2D ligands and T-cell receptor- versus NKG2D-dependent recognition.** *Scand J Immunol.* 2007; 66(2–3): 320–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
44. Roelofs AJ, Jauhainen M, Monkkonen H, *et al.*: **Peripheral blood monocytes are responsible for gammadelta T cell activation induced by zoledronic acid through accumulation of IPP/DMAPP.** *Br J Haematol.* 2009; 144(2): 245–50.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
45. Braza MS, Klein B: **Anti-tumour immunotherapy with V γ 9V δ 2 T lymphocytes: from the bench to the bedside.** *Br J Haematol.* 2013; 160(2): 123–32.
[PubMed Abstract](#) | [Publisher Full Text](#)
46. **F** Dieli F, Vermijlen D, Fulfaro F, *et al.*: **Targeting human {gamma}delta T cells with zoledronate and interleukin-2 for immunotherapy of hormone-refractory prostate cancer.** *Cancer Res.* 2007; 67(15): 7450–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
47. **F** Buccheri S, Guggino G, Caccamo N, *et al.*: **Efficacy and safety of $\gamma\delta$ T cell-based tumor immunotherapy: a meta-analysis.** *J Biol Regul Homeost Agents.* 2014; 28(1): 81–90.
[PubMed Abstract](#) | [F1000 Recommendation](#)
48. Capietto A, Martinet L, Fournie J: **Stimulated $\gamma\delta$ T cells increase the *in vivo* efficacy of trastuzumab in HER-2 $^{+}$ breast cancer.** *J Immunol.* 2011; 187(2): 1031–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
49. Oberg H, Peipp M, Kellner C, *et al.*: **Novel bispecific antibodies increase $\gamma\delta$ T-cell cytotoxicity against pancreatic cancer cells.** *Cancer Res.* 2014; 74(5): 1349–60.
[PubMed Abstract](#) | [Publisher Full Text](#)
50. Oberg H, Kellner C, Gonnermann D, *et al.*: **$\gamma\delta$ T cell activation by bispecific antibodies.** *Cell Immunol.* 2015; 296(1): 41–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
51. Almeida AR, Correia DV, Fernandes-Platzgummer A, *et al.*: **Delta One T Cells for Immunotherapy of Chronic Lymphocytic Leukemia: Clinical-Grade Expansion/ Differentiation and Preclinical Proof of Concept.** *Clin Cancer Res.* 2016; 22(23): 5795–804.
[PubMed Abstract](#) | [Publisher Full Text](#)
52. **F** Gentles AJ, Newman AM, Liu CL, *et al.*: **The prognostic landscape of genes and infiltrating immune cells across human cancers.** *Nat Med.* 2015; 21(8): 938–45.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
53. Ye J, Ma C, Wang F, *et al.*: **Specific recruitment of $\gamma\delta$ regulatory T cells in human breast cancer.** *Cancer Res.* 2013; 73(20): 6137–48.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
54. **F** Wu P, Wu D, Ni C, *et al.*: **$\gamma\delta$ T17 cells promote the accumulation and expansion of myeloid-derived suppressor cells in human colorectal cancer.** *Immunity.* 2014; 40(5): 785–800.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
55. **F** Daley D, Zambirinis CP, Seifert L, *et al.*: **$\gamma\delta$ T Cells Support Pancreatic Oncogenesis by Restraining $\alpha\beta$ T Cell Activation.** *Cell.* 2016; 166(6): 1485–1499. e15.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
56. Poggi A, Zocchi MR: **$\gamma\delta$ T Lymphocytes as a First Line of Immune Defense: Old and New Ways of Antigen Recognition and Implications for Cancer Immunotherapy.** *Front Immunol.* 2014; 5: 575.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
57. **F** Zou C, Zhao P, Xiao Z, *et al.*: **$\gamma\delta$ T cells in cancer immunotherapy.** *Oncotarget.* 2017; 8(5): 8900–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
58. Heid HW, Winter S, Bruder G, *et al.*: **Butyrophilin, an apical plasma membrane-associated glycoprotein characteristic of lactating mammary glands of diverse species.** *Biochim Biophys Acta.* 1983; 728(2): 228–38.
[PubMed Abstract](#) | [Publisher Full Text](#)
59. Afrache H, Gouret P, Ainouche S, *et al.*: **The butyrophilin (BTN) gene family: from milk fat to the regulation of the immune response.** *Immunogenetics.* 2012; 64(11): 781–94.
[PubMed Abstract](#) | [Publisher Full Text](#)
60. Arnett HA, Viney JL: **Immune modulation by butyrophilins.** *Nat Rev Immunol.* 2014; 14(8): 559–69.
[PubMed Abstract](#) | [Publisher Full Text](#)
61. Rhodes DA, Reith W, Trowsdale J: **Regulation of Immunity by Butyrophilins.** *Annu Rev Immunol.* 2016; 34: 151–72.
[PubMed Abstract](#) | [Publisher Full Text](#)
62. Compte E, Pontarotti P, Collette Y, *et al.*: **Frontline: Characterization of BT3 molecules belonging to the B7 family expressed on immune cells.** *Eur J Immunol.* 2004; 34(8): 2089–99.
[PubMed Abstract](#) | [Publisher Full Text](#)
63. Rhodes DA, de Bono B, Trowsdale J: **Relationship between SPRY and B30.2 protein domains. Evolution of a component of immune defence?** *Immunology.*

- 2005; 116(4): 411–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
64. D'Cruz AA, Babon JJ, Norton RS, *et al.*: **Structure and function of the SPRY/B30.2 domain proteins involved in innate immunity.** *Protein Sci.* 2013; 22(1): 1–10.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 65. Viken MK, Blomhoff A, Olsson M, *et al.*: **Reproducible association with type 1 diabetes in the extended class I region of the major histocompatibility complex.** *Genes Immun.* 2009; 10(4): 323–33.
[PubMed Abstract](#) | [Publisher Full Text](#)
 66. Ampuero J, del Campo JA, Rojas L, *et al.*: **Fine-mapping butyrophilin family genes revealed several polymorphisms influencing viral genotype selection in hepatitis C infection.** *Genes Immun.* 2015; 16(5): 297–300.
[PubMed Abstract](#) | [Publisher Full Text](#)
 67. Malcherek G, Mayr L, Roda-Navarro P, *et al.*: **The B7 homolog butyrophilin BTN2A1 is a novel ligand for DC-SIGN.** *J Immunol.* 2007; 179(6): 3804–11.
[PubMed Abstract](#) | [Publisher Full Text](#)
 68. Chapoval AI, Smithson G, Brunick L, *et al.*: **BTNL8, a butyrophilin-like molecule that costimulates the primary immune response.** *Mol Immunol.* 2013; 56(4): 819–28.
[PubMed Abstract](#) | [Publisher Full Text](#)
 69. Messal N, Mamessier E, Sylvain A, *et al.*: **Differential role for CD277 as a co-regulator of the immune signal in T and NK cells.** *Eur J Immunol.* 2011; 41(12): 3443–54.
[PubMed Abstract](#) | [Publisher Full Text](#)
 70. Yamashiro H, Yoshizaki S, Tadaki T, *et al.*: **Stimulation of human butyrophilin 3 molecules results in negative regulation of cellular immunity.** *J Leukoc Biol.* 2010; 88(4): 757–67.
[PubMed Abstract](#) | [Publisher Full Text](#)
 71. Cubillos-Ruiz JR, Martinez D, Scarlett UK, *et al.*: **CD277 is a negative co-stimulatory molecule universally expressed by ovarian cancer microenvironmental cells.** *Oncotarget.* 2010; 1(5): 329–38.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 72. Nerdal PT, Peters C, Oberg HH, *et al.*: **Butyrophilin 3A/CD277-Dependent Activation of Human $\gamma\delta$ T Cells: Accessory Cell Capacity of Distinct Leukocyte Populations.** *J Immunol.* 2016; 197(8): 3059–68.
[PubMed Abstract](#) | [Publisher Full Text](#)
 73. Ammann JU, Cooke A, Trowsdale J: **Butyrophilin Btn2a2 inhibits TCR activation and phosphatidylinositol 3-kinase/Akt pathway signaling and induces Foxp3 expression in T lymphocytes.** *J Immunol.* 2013; 190(10): 5030–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 74. **F** Sarter K, Leimgruber E, Gobet F, *et al.*: **Btn2a2, a T cell immunomodulatory molecule coregulated with MHC class II genes.** *J Exp Med.* 2016; 213(2): 177–87.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 75. Swanson RM, Gavin MA, Escobar SS, *et al.*: **Butyrophilin-like 2 modulates B7 costimulation to induce Foxp3 expression and regulatory T cell development in mature T cells.** *J Immunol.* 2013; 190(5): 2027–35.
[PubMed Abstract](#) | [Publisher Full Text](#)
 76. Subramaniam KS, Spaulding E, Ivan E, *et al.*: **The T-Cell Inhibitory Molecule Butyrophilin-Like 2 Is Up-regulated in Mild *Plasmodium falciparum* Infection and Is Protective During Experimental Cerebral Malaria.** *J Infect Dis.* 2015; 212(8): 1322–31.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 77. **F** Lebrero-Fernández C, Wenzel UA, Akeus P, *et al.*: **Altered expression of Butyrophilin (BTN) and BTN-like (BTNL) genes in intestinal inflammation and colon cancer.** *Immun Inflamm Dis.* 2016; 4(2): 191–200.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 78. **F** Lebrero-Fernández C, Bergström JH, Pelaseyed T, *et al.*: **Murine Butyrophilin-Like 1 and Btl6 Form Heteromeric Complexes in Small Intestinal Epithelial Cells and Promote Proliferation of Local T Lymphocytes.** *Front Immunol.* 2016; 7: 1.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 79. Lebrero-Fernández C, Bas-Forsberg A: **The ontogeny of Butyrophilin-like (Btl) 1 and Btl6 in murine small intestine.** *Sci Rep.* 2016; 6: 31524.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 80. **F** Di Marco Barros R, Roberts NA, Dart RJ, *et al.*: **Epithelia Use Butyrophilin-like Molecules to Shape Organ-Specific $\gamma\delta$ T Cell Compartments.** *Cell.* 2016; 167(1): 203–218.e17.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 81. Palakodeti A, Sandstrom A, Sundaresan L, *et al.*: **The molecular basis for modulation of human V γ 9V δ 2 T cell responses by CD277/butyrophilin-3 (BTN3A)-specific antibodies.** *J Biol Chem.* 2012; 287(39): 32780–90.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 82. Wang H, Henry O, Distefano MD, *et al.*: **Butyrophilin 3A1 plays an essential role in prenyl pyrophosphate stimulation of human V γ 2V δ 2 T cells.** *J Immunol.* 2013; 191(3): 1029–42.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 83. **F** Riaño F, Karunakaran MM, Starick L, *et al.*: **V γ 9V δ 2 TCR-activation by phosphorylated antigens requires butyrophilin 3 A1 (BTN3A1) and additional genes on human chromosome 6.** *Eur J Immunol.* 2014; 44(9): 2571–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 84. **F** Wang H, Morita CT: **Sensor Function for Butyrophilin 3A1 in Prenyl Pyrophosphate Stimulation of Human V γ 2V δ 2 T Cells.** *J Immunol.* 2015; 195(10): 4583–94.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 85. **F** Rhodes DA, Chen H, Price AJ, *et al.*: **Activation of human $\gamma\delta$ T cells by cytosolic interactions of BTN3A1 with soluble phosphoantigens and the cytoskeletal adaptor periplakin.** *J Immunol.* 2015; 194(5): 2390–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 86. **F** Kilcollins AM, Li J, Hsiao CC, *et al.*: **HMBPP Analog Prodrugs Bypass Energy-Dependent Uptake To Promote Efficient BTN3A1-Mediated Malignant Cell Lysis by V γ 9V δ 2 T Lymphocyte Effectors.** *J Immunol.* 2016; 197(2): 419–28.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 87. De Libero G, Lau SY, Mori L: **Phosphoantigen Presentation to TCR $\gamma\delta$ Cells, a Conundrum Getting Less Gray Zones.** *Front Immunol.* 2014; 5: 679.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 88. Hsiao CC, Lin X, Barney RJ, *et al.*: **Synthesis of a phosphoantigen prodrug that potentially activates V γ 9V δ 2 T-lymphocytes.** *Chem Biol.* 2014; 21(8): 945–54.
[PubMed Abstract](#) | [Publisher Full Text](#)
 89. Gu S, Nawrocka W, Adams EJ: **Sensing of Pyrophosphate Metabolites by V γ 9V δ 2 T Cells.** *Front Immunol.* 2015; 5: 688.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 90. Harly C, Peigne C, Scotet E: **Molecules and Mechanisms Implicated in the Peculiar Antigenic Activation Process of Human V γ 9V δ 2 T Cells.** *Front Immunol.* 2014; 5: 657.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 91. **F** Sebestyen Z, Scheper W, Vyborova A, *et al.*: **RhoB Mediates Phosphoantigen Recognition by V γ 9V δ 2 T Cell Receptor.** *Cell Rep.* 2016; 15(9): 1973–85.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 92. Boczonadi V, Maatta A: **Functional Analysis of Periplakin and Envoplakin, Cytoskeletal Linkers, and Cornified Envelope Precursor Proteins.** *Meth Enzymol.* 2016; 569: 309–29.
[PubMed Abstract](#) | [Publisher Full Text](#)
 93. **F** Menon S, Shin S, Dy G: **Advances in Cancer Immunotherapy in Solid Tumors.** *Cancers (Basel).* 2016; 8(12): pii: E106.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 94. **F** Benyamini A, Le Roy A, Mamessier E, *et al.*: **BTN3A molecules considerably improve V γ 9V δ 2 cells-based immunotherapy in acute myeloid leukemia.** *Oncotarget.* 2016; 5(10): e1146843.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 95. Paul S, Shilpi, Lal G: **Role of gamma-delta ($\gamma\delta$) T cells in autoimmunity.** *J Leukoc Biol.* 2015; 97(2): 259–71.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. **Richard D. Lopez**

Division of Cellular Therapy/Bone Marrow Transplantation, Duke University, Durham, NC, USA

Competing Interests: No competing interests were disclosed.

2. **John Trowsdale**

Pathology Department, Tennis Court Road, University of Cambridge, Cambridge, UK

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research