REVIEW

Human papillomavirus vaccination: the population impact

[version 1; peer review: 3 approved]

Lai-yang Lee^{1,2}, Suzanne M. Garland¹⁻⁴

¹Department of Microbiology and Infectious Diseases, Royal Women's Hospital, Parkville, Victoria, Australia

²Department of Microbiology, Royal Children's Hospital, Parkville, Victoria, Australia

³Department of Infection and Immunity, Murdoch Childrens Research Institute, Royal Children's Hospital, Victoria, Australia

⁴Department of Obstetrics Gynaecology, University of Melbourne, Parkville, Victoria, Australia

V1 First published: 12 Jun 2017, 6(F1000 Faculty Rev):866
<https://doi.org/10.12688/f1000research.10691.1>

Latest published: 12 Jun 2017, 6(F1000 Faculty Rev):866
<https://doi.org/10.12688/f1000research.10691.1>

Abstract

We currently have the knowledge and experience to prevent much of human papillomavirus (HPV)-related disease burden globally. In many countries where prophylactic HPV vaccination programs have been adopted as highly effective public health programs with good vaccine coverage, we are already seeing, in real-world settings, reduction of vaccine-related HPV-type infections, genital warts and cervical pre-cancers with potential reductions in vulvar, vaginal and anal pre-cancers. Moreover, we are seeing a change in cervical screening paradigms, as HPV-based screening programs now have strong evidence to support their use as more sensitive ways to detect underlying cervical abnormalities, as compared with conventional cervical cytology. This article describes the impact of prophylactic vaccination on these outcomes and in settings where these vaccines have been implemented in national immunisation programs. Given the successes seen to date and the availability of essential tools, there has been a global push to ensure that every woman has access to effective cervical screening and every girl has the opportunity for primary prevention through vaccination. A gender-neutral approach by offering vaccination to young boys has also been adopted by some countries and is worthy of consideration given that HPV-related cancers also affect males. Furthermore, vaccination of young boys has the advantage of reducing the risk of HPV transmission to sexual partners, lowering the infectious pool of HPV in the general population and ultimately HPV-related diseases for both genders. Therefore, it is appropriate that all countries consider and promote national guidelines and programs to prevent HPV-related diseases.

Keywords

HPV vaccination , prophylactics , cervical cancer , prevention

Open Peer Review

Approval Status

	1	2	3
version 1 12 Jun 2017			

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Eileen Dunne**, Centers for Disease Control and Prevention, Atlanta, USA
Thailand MOPH CDC Collaboration, Bangkok, Thailand
2. **Linda Niccolai**, Yale School of Public Health, New Haven, USA
3. **Mark Jit**, London School of Hygiene and Tropical Medicine, London, UK
Public Health England, London, UK

Any comments on the article can be found at the end of the article.

Corresponding author: Suzanne M. Garland (Suzanne.garland@thewomens.org.au)

Competing interests: SG has received speaking fees from Merck Sharp & Dohme (MSD) and Sanofi Pasteur MSD for work performed in her personal time. Merck paid for her travel and accommodation to present at HPV Advisory Board meetings. Lai-yang Lee declares that she has no competing interests.

Grant information: SG has received grants to her institution from the Commonwealth Department of Health for HPV genoprevalence surveillance post-vaccination, Merck and GlaxoSmithKline to perform phase 3 clinical vaccine trials, Merck to evaluate HPV in RRP post-vaccination program, Seqirus Australia for funding of the Australian Cervical Cancer Typing Study, and the Victoria Cancer Agency for a study on monitoring and effectiveness of the Australian cervical cancer program, a translation to reduction in vaccine-related HPV infection and pre-cancerous cervical lesions study (the Vaccine Against Cervical Cancer Impact and Effectiveness study) plus a study on associations and typing of early-onset cancers in young women.

Copyright: © 2017 Lee Ly and Garland SM. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Lee Ly and Garland SM. **Human papillomavirus vaccination: the population impact [version 1; peer review: 3 approved]** F1000Research 2017, 6(F1000 Faculty Rev):866 <https://doi.org/10.12688/f1000research.10691.1>

First published: 12 Jun 2017, 6(F1000 Faculty Rev):866 <https://doi.org/10.12688/f1000research.10691.1>

Introduction

Human papillomavirus infection and disease association

Human papillomavirus (HPV) is the commonest sexually transmitted infection, and the resultant diseases have significant morbidity and mortality¹. Although most HPV infections are transient, persistent infections are a prerequisite for pre-cancerous lesions and ultimately cancer.²⁻⁵ It is highly likely that latent infection (as infection dormant in the basal cells not readily detectable by diagnostic assays, but reversible)⁶ or subclinical infections (or both) exist and this is particularly relevant when an individual is immunocompromised since it increases the risk of developing symptomatic disease. Oncogenic HPVs were shown by Harald zur Hausen to be the causative agent of cervical cancer in the early 1980s by virtue of molecular epidemiology; for this work, zur Hausen shared the Nobel Prize in Physiology or Medicine in 2008⁷. Of the many genotypes specifically infecting the anogenital area, HPV16 and HPV18 are the commonest high-risk or oncogenic genotypes in cervical cancer and are responsible for approximately 50% of high-grade cervical dysplasias and 70% of cases of cervical cancer, the fourth most common cancer in females globally^{8,9}. Oncogenic HPVs cause almost 100% of cervical cancers, 90% of anal, 70% of vaginal, 40% of vulvar, 50% of penile and 13% to 72% of oropharyngeal cancers, and HPV16 predominates in all of these non-cervical HPV-related cancers. HPV6 and HPV11, which are classified as low-risk genotypes, cause 90% of genital warts as well as the rare but debilitating recurrent respiratory papillomatosis (RRP)^{10,11}.

Prophylactic human papillomavirus vaccines

Given the heavy disease burden of cervical cancer, prophylactic HPV vaccines were developed to target the commonest high- and low-risk HPV genotypes. Two such vaccines were first licensed for clinical use in 2006 following phase 3 clinical trials, which showed efficacy, safety and immunogenicity against vaccine-related HPV types¹²⁻¹⁵. Currently available vaccines now include bivalent (2vHPV and targets HPV 16/18)^{12,13}, quadrivalent (4vHPV and targets HPV 16/18/6/11)^{14,15} and nonavalent (targets HPV 16/18/6/11 as well as the five next most common oncogenic types found in cervical cancers, 31/33/45/52/58) vaccines¹⁶. Whilst many countries have licensed these vaccines, worldwide introduction into national immunisation programs differs by country, and some vaccines are available only on a user-pays basis and therefore entail an out-of-pocket expense. This has contributed to the resultant generally poor and inequitable uptake. For the greatest impact of these vaccines, the aim is to vaccinate adolescent females prior to sexual debut and to gain high coverage of the target population⁹. Some countries have implemented catch-up programs for older females or routine vaccination of adolescent males (or both) to increase overall population coverage and enhance herd protection¹⁷.

Vaccine effectiveness and impact

Clinical trials indicated efficacy of the vaccines against the HPV types included in the respective vaccines as well as some modest cross-protection against some non-vaccine but phylogenetically related strains (HPV 31, 33 for HPV16-related types and HPV45 for 18-related types)^{12,18}. The true benefit of vaccination can be seen only from real-life impact and effectiveness once vaccination has been included in public health programs. To appreciate this requires good surveillance and observation of changes in

genoprevalence within vaccine-eligible age cohorts as well as of disease outcomes. Some of the challenges here have included the fact that HPV cannot be cultured by traditional means, so DNA typing in the general community before and after vaccination is required to determine molecular genoprevalence. Although a number of assays to determine the presence of HPV are currently available¹⁹, their appropriateness is based on whether they are meant to be used for clinical needs (to detect underlying disease in which case assays are designed with clinical cut-offs and are less sensitive) or for pure epidemiological purposes (these use analytical viral endpoints and are highly sensitive)^{20,21}. Furthermore, as HPV infection is not notifiable, nor are many of the associated clinical manifestations in most countries, individual surveillance systems have been required to measure the impact and effectiveness of the vaccines. In addition, as the times from infection to the various disease manifestations differ and may be weeks to months (genital warts) to years (pre-cancers) or decades (cancer), the impact of vaccination on each disease varies. Consequently, in the 10 years since the introduction of both 4vHPV and 2vHPV vaccines, measures of vaccine impact have largely been determined by individual observational studies in countries with higher vaccine coverage, although a number of large, countrywide surveillance activities are now occurring. Thus, a direct comparison between the multiple studies published to date is complicated as the population studied, the duration of study and the endpoints observed or measured (or both) differ. Monitoring impact is difficult, particularly as some endpoints such as pre-cancerous lesions require a robust pre-cancer screening program to be in place or are not notifiable or have a prolonged interval between infection and development of diseases, particularly for cancers which usually occur after decades of persistent infection with a high-risk HPV type.

In this article, we define vaccine impact as the effect of public health programs of HPV vaccines at a population level and the measurement thereof of reduction in disease burden (before and after initiation of a vaccination program) and largely focus on this. In contrast, vaccine effectiveness is examined at an individual level²². It is largely observational and determines the effect of vaccination observed in populations after a program commences and compares outcomes in those vaccinated with those unvaccinated.

Vaccine impact will be determined by the type of vaccine program (an ongoing routine vaccination program to a specific age group versus inclusion of a catch-up program which may be limited in time and/or target a narrow or wide number of age cohorts), vaccine coverage, which population is targeted (for example, a program focused entirely on females, a gender-neutral approach, or targeted vaccination such as the HIV-positive population), the duration of vaccine protection, the duration of follow-up after vaccination, and type of surveillance, if any, of various outcome measures.

Vaccine impact on human papillomavirus infection

Australia was one of the first countries to implement a fully government-funded, population-based HPV vaccination program. It commenced in 2007 as an ongoing school-based program with a three-dose course of the 4vHPV vaccine, targeting females in the first year of high school at age 12 to 13 years, with a catch-up for those ages 12 to 26 years until December 2009²³. In 2013, the Australian

government extended the program to include 12- to 13-year-old males, including a 2-year catch-up vaccination program for those ages 14 to 15 years^{24,25}. Overall, the program has been well received and high vaccine coverage rates have been achieved; the greatest rates have been from those within the school-based cohorts: over 70% have received all three doses for young girls and just under 70% have received all three doses for boys²⁴⁻²⁶. Consequently, Australian researchers were among the first to report reductions in the prevalence of vaccine-type HPV infections, by 86% in 18- to 24-year-olds who had received three vaccine doses and 76% for those who had received one or two doses^{22,27,28}. In general, Tabrizi *et al.* found that the greatest decline in vaccine-type HPV prevalence corresponded to the age group which had the highest vaccine coverage, even in the unvaccinated but eligible-age population, highly suggestive of herd protection²⁸. Furthermore, Australia is already seeing herd protection in young males as a result of the female program²⁹.

Other countries that have achieved high coverage with 4vHPV or 2vHPV vaccines also report rapid and large declines in vaccine-related HPV infections. For example, in Scotland, which also has a school-based program, which targets 12- to 13-year-old girls as an ongoing program and had a limited 3-year catch-up from 2008 to 2011 for 13- to 17-year-olds, coverage rates for the 2vHPV vaccine have been even greater, at up to 90%, and reported declines in vaccine-related infections were from 29.8% to 13.6%³⁰. In contrast, in the USA, coverage of the 4vHPV vaccine has been much lower, at 40% for females and 22% for males; yet despite this, national impact studies from the USA 4 years after the introduction of HPV vaccines noted a 56% decrease in HPV vaccine types (from

cervical-vaginal samples from 14- to 19-year-olds)^{31,32}. Although the early vaccination initiation uptake rate was low (at 17%) for females 19 to 26 years old in 2009, by 2012 the uptake in the USA had doubled to 34%, resulting in a further 45% decrease in the prevalence of vaccine-type HPV³³, or an overall 64% decrease in vaccine-type HPV prevalence after 6 years³². A less marked, but still significant, reduction of 34% prevalence over the same time frame was noted in the 20- to 24-year-old group³². In sexually active US women ages 14 to 24 in the post-vaccination era, those who had received at least one HPV vaccine dose had a 2.1% reduction of vaccine-type HPV prevalence compared with 16.9% in the same age group of unvaccinated women³².

Genital warts

Genital warts are a common condition, which affects up to 10% of the female population under 45 years old and which usually develops 2 to 3 months after infection with low-risk HPV genotypes, primarily HPV 6 and 11^{10,34}. Australia was the first to report a reduction in genital warts, and the reduction was larger and faster than had been expected by researchers^{35,36}, and the 4vHPV vaccine provided up to 92% reduction in HPV-associated genital warts³⁷. Interestingly, observations from early use of 2vHPV vaccines noted an unexpected decrease in genital warts. It is speculated that this response is due to vaccination resulting in a cell-mediated immune response which confers a moderate amount of protection against some low-risk HPV types^{38,39}.

Since the commencement of 4vHPV vaccination, there has been a substantial reduction in reports of genital warts in countries with 4vHPV vaccination programs (Table 1)^{35-37,40-43}. A Belgian study

Table 1. Summary of human papillomavirus vaccination programs, outcomes and cervical screening programs.

	Australia	USA	United Kingdom	Denmark	Scotland
Delivery route	School	Clinic	Mostly school	Clinic	School
HPV vaccine program commencement	2007	2006	2008	2009	2008
Vaccine type	4vHPV Ongoing	4vHPV 2006-2016 9vHPV December 2016	2vHPV 2008-2012 4vHPV 2012	4vHPV	2vHPV 2008-2012 4vHPV 2012
Three-dose schedule	Yes				
Two-dose schedule ^a	No	Yes October 2016	Yes September 2014	Yes 2014	Yes 2014
Females	Yes	Yes	Yes	Yes	Yes
Target age, years	12-13	11-12	12-13	12	12-13
Males (routine)	Yes from 2013	Yes from 2011	No	No	No
Target age males, years	12-13	11-12	-	-	-

	Australia	USA	United Kingdom	Denmark	Scotland
Catch-up program	12-26 ♀ 2007-2009 14-15 years old ♂ 2013-2015	13-26 ♀ Ongoing	Up to 18 ♀ 2009-2011	13 -15 ♀ 2009 2nd catch-up 20-27 ♀ 2012	13-17 ♀ 2008-2011
Estimated coverage	2015	2015	2013/2014	2015	
Three doses	77.4% ♀ 66.4% ♂	41.9% ♀ 28.1% ♂	86.7%	82%	12-13 years old 90% Catch-up cohort 66%
At least one dose	85.6% ♀ 77% ♂	62.8% ♀ 49.8% ♂	91.1%	90%	>90%
Vaccine-type HPV infection reduction	18-26 years old 3x dose 86% 2x dose 76%	2010 14-19 ♀ 56% 2012 14-19 years old ♀ 64% 20-24 ♀ 34%	16-18 years old HPV 16/18 prevalence reduce from 19.1% to 6.5%	Not available	HPV 16/18 prevalence reduce from 29.8% to 13.6%
Genital warts (types 6/11) Reduction	Up to 92%	<21 years old 34.8% >21 years old 10%	2vHPV 20.8% 4vHPV- 38.9% ♀ 30.2% ♂	2013 <18 ♀ up to 55.1% <18 years old ♂ up to 36.6%	Since 4vHPV vaccine use, decline in wart prescription (personal communication, Kevin Pollock, 5 January 2017)
CIN/ Adenocarcinoma Reduction	Low-grade 34% High-grade 47% <20 years old 54% 20-24 years old 37%	HPV16/18 CIN2+ ♀ vaccinated >24 months before PAP versus unvaccinated ♀, adjusted prevalence ratio of 0.67	Not available	Atypia <18 years old 33.4% (annual percentage) 18-20 years old 12.6% CIN2+ 18-20 years old 14%	CIN1 29% CIN2 50% CIN3 55%
Screening program: Type and interval age commencement	Cervical cytology 18-69 every 2 years	Cervical cytology 21-65 every 3 years	Cervical cytology 25-49 every 3 years 50-64 every 5 years January 2016: UK National Screening Committee recommend HPV primary screening (to commence in the near future) ⁹⁰	Cervical cytology 23-49 every 3 years 50-65 every 5 years	Cervical cytology 25-49 every 3 years 50-64 every 5 years
Comments	1 May 2017 to be changed to HPV DNA every 5 years for 25-74 years old		Since 2014, two-dose schedule <15 years old >15 years old on initiation, three- dose schedule	Since 2014, two-dose schedule <15 years old >15 years old on initiation, three-dose schedule	Screening prior to 2013, 20-60 years old every 3 years
Additional references	25,91	92-96	97-100	101,102	103,104

^aLess than 15 years at the time of first dose: two-dose regimen of a prime and a boost separated by a minimum of 6 months⁷³.

2vHPV, bivalent human papillomavirus vaccine; 4vHPV, quadrivalent human papillomavirus vaccine; 9vHPV, nonavalent human papillomavirus vaccine; CIN, cervical intraepithelial neoplasia; HPV, human papillomavirus; PAP, Papanicolaou test.

noted a decrease of 8.1% in genital warts in the general population, and the highest decreases were in 16- to 22-year-old males and females between the pre- and post-vaccine era⁴⁴. Australia, with its high vaccine coverage rates, reported the greatest worldwide decline, of up to 92% in the under-21 age group³⁷. Several studies have highlighted that the greatest efficacy of vaccination against genital warts was when the first dose of vaccine was given at a younger age⁴⁴ with less of an effect at an older age⁴⁰.

Cervical cytology and histological abnormalities

Genital warts have a short incubation period and hence this was the first disease reduction noted from vaccination. The second disease outcome to see reductions was cervical dysplasia, which has an intermediate incubation period (peak age of 26 to 30 years)⁴⁵. Again, this was first reported in an ecological study in Victoria, Australia⁴⁶ and was made possible because Australia had a HPV vaccine register, comprehensive cervical cytology reporting as well as the ability (following legislation) for the registers to be linked. Australia's national HPV vaccination programs have led to a 34% decline in low-grade and a 47% decline in high-grade cervical intraepithelial neoplasia (CIN) and adenocarcinoma *in situ*, and the largest reductions were in the vaccinated younger age group^{22,47}. Women under the age of 20 had a prevalence decline from 10.9/1000 screened women to 5.0/1000 over a period of 10 years, and prevalence in 20- to 24-year-olds decreased from 21.5/1000 screened to 13.5/1000 in a similar time period. In the over-30-year-old age group, the prevalence of high-grade CIN has continued to slowly rise⁴⁸. Similar declines have been observed in Scotland and Denmark⁴⁹⁻⁵² (Table 1). There are randomised controlled phase 2/3 trials indicating that, despite diverse populations and geographical locations, HPV vaccines covering HPV16/18 provided some cross-protection against non-vaccine strain HPV and could protect women against cervical and non-cervical HPV-related infections⁵³.

Cervical cancer

At present, it is too early to show clinical disease or population-based disease reduction for cervical cancer, as this is a disease peaking in the mid-30s and mid-40s in Australia⁵⁴, although the peak has shifted to the late 20s to early 30s in some countries⁵⁵. One would expect to see reductions within the next few years in the under-30-year-olds, particularly in those countries with high coverage of catch-up programs extending to 26 years of age, such as Australia and Denmark.

Based on the PATRICIA trial (Papilloma Trial against Cancer in young Adults) of the 2vHPV vaccine, it has been estimated that with 50% vaccine coverage, there could be a worldwide reduction of cervical cancer incidence of 246,086 cases annually but that with 90% coverage, there could be up to 442,955 cases averted⁵⁶. With the greater coverage provided by the nonavalent vaccine, it is proposed that with high coverage, 90% to 93% of all cervical cancers would be prevented^{57,58}.

The incidence of non-cervical HPV-related cancers is generally low and peaks at a later age compared with cervical cancer⁵⁹. Thus, more time and large population-based effectiveness studies with various cancers as endpoints will be required before impact can be determined. There are early data sets currently reviewing the

implication of vaccine-type HPV reduction and the effect on persistent anal infections and intraepithelial neoplasia⁶⁰.

Impediments and challenges for the future

Vaccine uptake and coverage. Although the effect of population-wide vaccination has been observed for 10 years, the reported findings may not accurately reflect future impact and this is for several reasons. Some countries have been slow to adopt HPV vaccination or vaccine coverage has been low (or both), even in high-income countries. Vaccine uptake rates in 13- to 17-year-olds in the USA in 2013 were 57% (one dose) and 38% (three doses)³². In Australia, between 2007 and 2015, vaccine uptake rates were 85.6% (one dose) and 77.4% (three doses) for females turning 15 years of age⁶¹. Successful strategies to increase coverage have included publically supported health promotion and missed-doses catch-up vaccine campaigns⁶², client reminders, recall programs, provider assessment and feedback interventions. Low- and middle-income countries often have a high HPV disease burden: however, only 15% of these countries have adopted a vaccination program¹⁸. Focusing on these countries and giving reassurance regarding vaccine safety will substantially impact HPV vaccination rates and outcomes⁶³.

Safety. Concern regarding vaccine safety is an issue that often arises. The US Centers for Disease Control and Prevention (CDC), through the Vaccine Adverse Event Reporting System, classified only 7% of adverse events as serious from over 90 million doses of HPV vaccines. The CDC found no causal association between HPV vaccines and ovarian failure, Guillain-Barré syndrome or postural orthostatic tachycardia syndrome. Overall, HPV vaccines have a good safety profile, but ongoing monitoring will be required as 9vHPV vaccines have been introduced⁶⁴⁻⁶⁷. In Japan, poor or inappropriate information (or both), including reports on complex regional pain syndrome, has resulted in the government withdrawing support for HPV vaccination programs⁶⁸⁻⁷⁰. Despite comprehensive independent evaluations of safety^{71,72}, including endorsement of safety by bodies such as the World Health Organization (WHO)⁷³, the International Papillomavirus Society⁶⁶ and the CDC⁷⁴, the Japanese government has continued to refuse to re-implement population-wide vaccination⁶⁹.

Ideal age for vaccination. In many respects, if the HPV vaccines could be administered in childhood, they could use infrastructure already in place for the Expanded Program on Immunization vaccines, as in general these see high uptake even in those countries with low resources. However, the immunity developed by HPV vaccination needs to be robust for several decades to ensure that infection does not occur when young people become sexually active and therefore to prevent diseases in the longer term. Although some studies have shown detectable neutralising antibodies 10 years after vaccination, there are not yet any data about antibody persistence beyond that²². It is currently believed that the antibody responses made by the vaccines are strong and long-lasting and that only small titres are in fact required for protection⁷⁵.

Over the last 10 years, there have been changes in sexual behaviour, such as lowered median age of sexual debut, increased exposure from the increase in the number of sexual partners, and choice of orientation of sexual partner; however, these trends vary

from country to country. The ability of vaccine programs in different countries to be flexible will allow these changes to be accounted for in the future. As shown in [Table 1](#), different delivery programs/routes for the vaccines result in different outcomes in different countries. A single worldwide approach is not appropriate, and programs need to be tailored to each country's population.

Dosing. New information regarding number of doses of vaccine necessary for long-lasting immunity is becoming available. The current WHO recommendation is that with respect to HPV-vaccine-related-type antibody responses, two doses of HPV vaccine for females 9 to 14 years old is non-inferior to three doses in adult women as long as the two doses are at least 6 months apart⁷³. This decision was based largely on neutralising antibody outcome and more recent findings of reduced HPV-vaccine-type-related disease outcomes, but again good surveillance systems should be able to monitor impact on disease endpoints with longer time intervals from the point of vaccination. Furthermore, this may help to address concerns regarding reduced efficacy⁷⁶ to ensure that a third dose could be administered if surveillance shows breakthrough disease. For females older than 15 years, those who are immunocompromised or HIV-positive, a three-dose regimen is still recommended^{73,77}.

Gender-neutral approaches. Although females have been the primary target of HPV vaccine, extending the program towards males may have some benefit. A 2011 Australian study predicted that, based on the female-only vaccination program, there could be up to a 68% decrease in male HPV infections by 2050 and a 14% decrease in head, neck and anogenital cancers. The effect on oropharyngeal cancers is unclear⁷⁸. This prediction is based on heterosexual relationships offering herd protection. A high-risk HPV group, men who have sex with men, are less likely to benefit from a female-only vaccination program⁷⁹. A gender-neutral approach may not only reduce HPV-related diseases in males but also reduce the infection and transmission to females, ultimately reducing the pool of infectious virus in a community. Male vaccination would confer an even greater benefit in settings where female vaccination rates are low. Hence, vaccination of males has recently been incorporated into several national programs. Measuring vaccine effectiveness in the context of gender-neutral programs will bring further challenges²⁴.

Multivalent human papillomavirus vaccines. Some countries have implemented or have plans to commence the use of 9vHPV vaccines to offer protection against the additional five next most common oncogenic HPV types after HPV 16/18 in an effort to further improve health outcomes. This move is supported by studies which indicate potential cost savings and health benefits that could result from universal nonavalent vaccination programs^{16,80,81}.

Alternative vaccine policies: "HPV FASTER" potential to impact disease earlier

Consideration for vaccinating women who are already sexually active but may not have been infected with all HPV types covered in the vaccines could well reduce the burden of disease more rapidly. This concept is based on data from phase 3 trials indicating

that 2vHPV and 4vHPV vaccines are efficacious, immunogenic and safe in women ages 26 to 45 years old with a 90% vaccine efficacy in protection against cervical pre-cancer in HPV-DNA-negative women (regardless of serostatus) and 50% vaccine efficacy in women who had previously been exposed to HPV⁸²⁻⁸⁴. Moreover, in trials in younger women, it was shown that vaccines were not efficacious when a woman was HPV-positive for a particular type⁸⁵, although it was likely for women who were HPV-negative but antibody-positive from natural infection that they would gain efficacy against later disease^{86,87}. The importance of a coordinated screening and vaccination program cannot be overlooked and this has prompted the HPV-FASTER concept. This protocol intends to offer women up to the age of 45 years HPV vaccination and incorporates HPV testing after the age of 30 for screening, with ongoing follow-up and early management of high-risk HPV-positive women. Further studies will be required to investigate the acceptability and feasibility of such an approach, most appropriate cut-off age of offering vaccination, ideal screening commencement, cessation and interval times to refine the protocol⁸⁸.

Surveillance. A necessary condition for accurately monitoring the impact and effectiveness of vaccine programs is the existence of diligent and ongoing surveillance⁸⁹. For HPV, this requires surveillance of HPV vaccination coverage and safety, HPV infection genoprevalence, cervical dysplasia incidence, cervical (as well as other anogenital and oropharyngeal cancer, ideally with genotyping of cancers) cancer registries, as well as surveillance for non-cancer outcomes such as warts and RRP. Ideally, this would include the ability for the various data sources to be linked. Encouragement in participation in screening programs is essential, particularly as new cervical screening programs that incorporate DNA testing as the primary screen usually have longer intervals between screens compared with those with cervical cytology as primary. Also, the vaccinated population may not recognise that, despite vaccination, there is a need for ongoing monitoring. With so many changes, education of the community, clinicians and women in particular is critical.

Conclusions

Overall, the implementation of HPV vaccine public health programs has resulted in major decreases in vaccine-type HPV infection prevalence as well as associated disease incidence in countries that have introduced them. Administration of vaccines is ideally done prior to sexual debut for both males and females for the greatest impact. Herd and cross-protection contribute further to the impact of the vaccines. More time for observation is required to determine the effect on cancer rates, as carcinomas develop decades after infection acquisition. Although we have the tools to markedly impact HPV-related disease, we also face the challenge of ensuring that all young people, regardless of their geography, social status, or immunocompromised status, are eligible for and can access these vaccines. To observe the vaccine impact and effectiveness of national HPV vaccination programs, support is required from accurate vaccine uptake surveillance, preferably through vaccine registers in addition to genoprevalence and HPV-related disease endpoint surveillance, including cancer registries and pre-cancer screening programs.

Competing interests

SG has received speaking fees from Merck Sharp & Dohme (MSD) and Sanofi Pasteur MSD for work performed in her personal time. Merck paid for her travel and accommodation to present at HPV Advisory Board meetings. Lai-yang Lee declares that she has no competing interests.

Grant information

SG has received grants to her institution from the Commonwealth Department of Health for HPV genoprevalence surveillance post-vaccination, Merck and GlaxoSmithKline to perform phase 3 clinical vaccine trials, Merck to evaluate HPV in RRP post-vaccination

program, Seqirus Australia for funding of the Australian Cervical Cancer Typing Study, and the Victoria Cancer Agency for a study on monitoring and effectiveness of the Australian cervical cancer program, a translation to reduction in vaccine-related HPV infection and pre-cancerous cervical lesions study (the Vaccine Against Cervical Cancer Impact and Effectiveness study) plus a study on associations and typing of early-onset cancers in young women.

Acknowledgements

We would like to thank David Mesher, Kevin Pollock, Charles Lacey and Susanne Krüger Kjaer for their prompt replies, insightful comments and information about their respective countries' vaccine programs.

References

- Forman D, de Martel C, Lacey CJ, *et al.*: **Global burden of human papillomavirus and related diseases.** *Vaccine*. 2012; **30**(Suppl 5): F12–23.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Walboomers JM, Jacobs MV, Manos MM, *et al.*: **Human papillomavirus is a necessary cause of invasive cervical cancer worldwide.** *J Pathol*. 1999; **189**(1): 12–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Giuliano AR, Harris R, Sedjo RL, *et al.*: **Incidence, prevalence, and clearance of type-specific human papillomavirus infections: The Young Women's Health Study.** *J Infect Dis*. 2002; **186**(4): 462–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Monsonog J, Bosch FX, Coursaget P, *et al.*: **Cervical cancer control, priorities and new directions.** *Int J Cancer*. 2004; **108**(3): 329–33.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Wright TC Jr: **Natural history of HPV infections.** *J Fam Pract*. Quadrant Healthcom, Inc; 2009; **58**(9 Suppl HPV): S3–7.
[PubMed Abstract](#)
- Gravitt PE: **Evidence and impact of human papillomavirus latency.** *Open Virol J*. 2012; **6**(Suppl 2): 198–203.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Hampton T: **Nobel Prize honors HIV, HPV discoveries.** *JAMA*. 2008; **300**(18): 2109.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Clifford GM, Smith JS, Plummer M, *et al.*: **Human papillomavirus types in invasive cervical cancer worldwide: a meta-analysis.** *Br J Cancer*. 2003; **88**(1): 63–73.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- World Health Organisation: **Media Centre: Human Papillomavirus (HPV) and cervical cancer.** 2016.
[Reference Source](#)
- Garland SM, Steben M, Sings HL, *et al.*: **Natural history of genital warts: analysis of the placebo arm of 2 randomized phase III trials of a quadrivalent human papillomavirus (types 6, 11, 16, and 18) vaccine.** *J Infect Dis*. 2009; **199**(6): 805–14.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Somers GR, Tabrizi SN, Borg AJ, *et al.*: **Juvenile laryngeal papillomatosis in a pediatric population: a clinicopathologic study.** *Pediatr Pathol Lab Med*. 1997; **17**(1): 53–64.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Wheeler CM, Castellsagué X, Garland SM, *et al.*: **Cross-protective efficacy of HPV-16/18 AS04-adjuvanted vaccine against cervical infection and precancer caused by non-vaccine oncogenic HPV types: 4-year end-of-study analysis of the randomised, double-blind PATRICIA trial.** *Lancet Oncol*. 2012; **13**(1): 100–10.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lehtinen M, Paavonen J, Wheeler CM, *et al.*: **Overall efficacy of HPV-16/18 AS04-adjuvanted vaccine against grade 3 or greater cervical intraepithelial neoplasia: 4-year end-of-study analysis of the randomised, double-blind PATRICIA trial.** *Lancet Oncol*. 2012; **13**(1): 89–99.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Garland SM, Hernandez-Avila M, Wheeler CM, *et al.*: **Quadrivalent vaccine against human papillomavirus to prevent anogenital diseases.** *N Engl J Med*. 2007; **356**(19): 1928–43.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- FUTURE II Study Group: **Quadrivalent vaccine against human papillomavirus to prevent high-grade cervical lesions.** *N Engl J Med*. 2007; **356**(19): 1915–27.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Joura EA, Giuliano AR, Iversen OE, *et al.*: **A 9-valent HPV vaccine against infection and intraepithelial neoplasia in women.** *N Engl J Med*. 2015; **372**(8): 711–23.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Brisson M, Bénard É, Drolet M, *et al.*: **Population-level impact, herd immunity, and elimination after human papillomavirus vaccination: A systematic review and meta-analysis of predictions from transmission-dynamic models.** *Lancet Public Health*. 2016; **1**(1): e8–e17.
[Publisher Full Text](#) | [F1000 Recommendation](#)
- Drolet M, Bénard E, Boily MC, *et al.*: **Population-level impact and herd effects following human papillomavirus vaccination programmes: a systematic review and meta-analysis.** *Lancet Infect Dis*. 2015; **15**(5): 565–80.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Poljak M, Kocjan BJ, Oštrbenk A, *et al.*: **Commercially available molecular tests for human papillomaviruses (HPV): 2015 update.** *J Clin Virol*. 2016; **76**(Suppl 1): S3–S13.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Snijders PJ, van den Brule AJ, Meijer CJ: **The clinical relevance of human papillomavirus testing: relationship between analytical and clinical sensitivity.** *J Pathol*. 2003; **201**(1): 1–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Pagliusi SR, Garland SM: **International standard reagents for HPV detection.** *Dis Markers*. 2007; **23**(4): 283–96.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Garland SM, Kjaer SK, Muñoz N, *et al.*: **Impact and Effectiveness of the Quadrivalent Human Papillomavirus Vaccine: A Systematic Review of 10 Years of Real-world Experience.** *Clin Infect Dis*. 2016; **63**(4): 519–27.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Garland SM, Skinner SR, Brotherton JM: **Adolescent and young adult HPV vaccination in Australia: achievements and challenges.** *Prev Med*. 2011; **53**(Suppl 1): S29–35.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Garland SM, Molesworth EG, Machalek DA, *et al.*: **How to best measure the effectiveness of male human papillomavirus vaccine programmes?** *Clin Microbiol Infect*. 2015; **21**(9): 834–41.
[PubMed Abstract](#) | [Publisher Full Text](#)
- National HPV Vaccination Program Register: **HPV vaccination coverage by dose number (Australia) for males by age group in mid 2014.** [cited 2017 Jan 8].
[Reference Source](#)
- Brotherton JM, Liu B, Donovan B, *et al.*: **Human papillomavirus (HPV) vaccination coverage in young Australian women is higher than previously estimated: independent estimates from a nationally representative mobile**

- phone survey. *Vaccine*. 2014; **32**(5): 592–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
27. **F** Tabrizi SN, Brotherton JM, Kaldor JM, *et al.*: **Fall in human papillomavirus prevalence following a national vaccination program.** *J Infect Dis*. 2012; **206**(11): 1645–51.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 28. Tabrizi SN, Brotherton JM, Kaldor JM, *et al.*: **Assessment of herd immunity and cross-protection after a human papillomavirus vaccination programme in Australia: a repeat cross-sectional study.** *Lancet Infect Dis*. 2014; **14**(10): 958–66.
[PubMed Abstract](#) | [Publisher Full Text](#)
 29. Machalek DA, Chow EP, Garland SM, *et al.*: **Human Papillomavirus Prevalence in Unvaccinated Heterosexual Men After a National Female Vaccination Program.** *J Infect Dis*. 2017; **215**(2): 202–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
 30. **F** Kavanagh K, Pollock KG, Potts A, *et al.*: **Introduction and sustained high coverage of the HPV bivalent vaccine leads to a reduction in prevalence of HPV 16/18 and closely related HPV types.** *Br J Cancer*. 2014; **110**(11): 2804–11.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 31. **F** Markowitz LE, Hariri S, Lin C, *et al.*: **Reduction in human papillomavirus (HPV) prevalence among young women following HPV vaccine introduction in the United States, National Health and Nutrition Examination Surveys, 2003–2010.** *J Infect Dis*. 2013; **208**(3): 385–93.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 32. **F** Markowitz LE, Liu G, Hariri S, *et al.*: **Prevalence of HPV After Introduction of the Vaccination Program in the United States.** *Pediatrics*. 2016; **137**(3): e20151968.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 33. **F** Berenson AB, Laz TH, Rahman M: **Reduction in Vaccine-Type Human Papillomavirus Prevalence Among Women in the United States, 2009–2012.** *J Infect Dis*. 2016; **214**(12): 1961–4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 34. Kjaer SK, Tran TN, Sparen P, *et al.*: **The burden of genital warts: a study of nearly 70,000 women from the general female population in the 4 Nordic countries.** *J Infect Dis*. 2007; **196**(10): 1447–54.
[PubMed Abstract](#) | [Publisher Full Text](#)
 35. **F** Donovan B, Franklin N, Guy R, *et al.*: **Quadrivalent human papillomavirus vaccination and trends in genital warts in Australia: analysis of national sentinel surveillance data.** *Lancet Infect Dis*. 2011; **11**(1): 39–44.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 36. **F** Read TR, Hocking JS, Chen MY, *et al.*: **The near disappearance of genital warts in young women 4 years after commencing a national human papillomavirus (HPV) vaccination programme.** *Sex Transm Infect*. 2011; **87**(7): 544–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 37. **F** Ali H, Donovan B, Wand H, *et al.*: **Genital warts in young Australians five years into national human papillomavirus vaccination programme: national surveillance data.** *BMJ*. 2013; **346**: f2032.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 38. Howell-Jones R, Soldan K, Wetten S, *et al.*: **Declining genital Warts in young women in england associated with HPV 16/18 vaccination: an ecological study.** *J Infect Dis*. 2013; **208**(9): 1397–403.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 39. Szarewski A, Skinner SR, Garland SM, *et al.*: **Efficacy of the HPV-16/18 AS04-adjuvanted vaccine against low-risk HPV types (PATRICIA randomized trial): an unexpected observation.** *J Infect Dis*. 2013; **208**(9): 1391–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 40. **F** Dochez C, Bogers JJ, Verhelst R, *et al.*: **HPV vaccines to prevent cervical cancer and genital warts: an update.** *Vaccine*. 2014; **32**(14): 1595–601.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 41. Baandrup L, Blomberg M, Dehlendorf C, *et al.*: **Significant decrease in the incidence of genital warts in young Danish women after implementation of a national human papillomavirus vaccination program.** *Sex Transm Dis*. 2013; **40**(2): 130–5.
[PubMed Abstract](#)
 42. **F** Bollerup S, Balduz-Felskov B, Blomberg M, *et al.*: **Significant Reduction in the Incidence of Genital Warts in Young Men 5 Years into the Danish Human Papillomavirus Vaccination Program for Girls and Women.** *Sex Transm Dis*. 2016; **43**(4): 238–42.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 43. Blomberg M, Dehlendorf C, Munk C, *et al.*: **Strongly decreased risk of genital warts after vaccination against human papillomavirus: nationwide follow-up of vaccinated and unvaccinated girls in Denmark.** *Clin Infect Dis*. 2013; **57**(7): 929–34.
[PubMed Abstract](#) | [Publisher Full Text](#)
 44. **F** Dominiak-Felden G, Gobbo C, Simondon F: **Evaluating the Early Benefit of Quadrivalent HPV Vaccine on Genital Warts in Belgium: A Cohort Study.** *PLoS One*. 2015; **10**(7): e0132404.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 45. Australian Government. Australian Institute of Health and Welfare: **Cervical screening in Australia 2013–2014.** [cited 2017 Jan 8].
[Reference Source](#)
 46. Brotherton JM, Fridman M, May CL, *et al.*: **Early effect of the HPV vaccination programme on cervical abnormalities in Victoria, Australia: an ecological study.** *Lancet*. 2011; **377**(9783): 2085–92.
[PubMed Abstract](#) | [Publisher Full Text](#)
 47. Gertig DM, Brotherton JM, Budd AC, *et al.*: **Impact of a population-based HPV vaccination program on cervical abnormalities: a data linkage study.** *BMC Med*. 2013; **11**: 227.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 48. Brotherton JM, Saville AM, May CL, *et al.*: **Human papillomavirus vaccination is changing the epidemiology of high-grade cervical lesions in Australia.** *Cancer Causes Control*. 2015; **26**(6): 953–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
 49. **F** Pollock KG, Kavanagh K, Potts A, *et al.*: **Reduction of low- and high-grade cervical abnormalities associated with high uptake of the HPV bivalent vaccine in Scotland.** *Br J Cancer*. 2014; **111**(9): 1824–30.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 50. **F** Balduz-Felskov B, Dehlendorf C, Munk C, *et al.*: **Early impact of human papillomavirus vaccination on cervical neoplasia—nationwide follow-up of young Danish women.** *J Natl Cancer Inst*. 2014; **106**(3): djt460.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 51. **F** Balduz-Felskov B, Dehlendorf C, Junge J, *et al.*: **Incidence of cervical lesions in Danish women before and after implementation of a national HPV vaccination program.** *Cancer Causes Control*. 2014; **25**(7): 915–22.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 52. **F** Balduz-Felskov B, Munk C, Nielsen TS, *et al.*: **Trends in the incidence of cervical cancer and severe precancerous lesions in Denmark, 1997–2012.** *Cancer Causes Control*. 2015; **26**(8): 1105–16.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 53. **F** Skinner SR, Apter D, De Carvalho N, *et al.*: **Human papillomavirus (HPV)-16/18 AS04-adjuvanted vaccine for the prevention of cervical cancer and HPV-related diseases.** *Expert Rev Vaccines*. 2016; **15**(3): 367–87.
[PubMed Abstract](#) | [F1000 Recommendation](#)
 54. Australian Government Cancer Australia: **Cervical cancer in Australia.** [cited 2017 Jan 8].
[Reference Source](#)
 55. Cancer Research UK: **Cervical cancer incidence statistics.** [cited 2017 Mar 21].
[Reference Source](#)
 56. **F** van Kriekinge G, Castellsague X, Cibula D, *et al.*: **Estimation of the potential overall impact of human papillomavirus vaccination on cervical cancer cases and deaths.** *Vaccine*. 2014; **32**(6): 733–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 57. **F** Riethmuller D, Jacquard A, Lacau St Guily J, *et al.*: **Potential impact of a nonavalent HPV vaccine on the occurrence of HPV-related diseases in France.** *BMC Public Health*. 2015; **15**: 453.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 58. **F** Luckett R, Feldman S: **Impact of 2-, 4- and 9-valent HPV vaccines on morbidity and mortality from cervical cancer.** *Hum Vaccin Immunother*. 2016; **12**(6): 1332–42.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 59. **F** Buchanan TR, Graybill WS, Pierce JY: **Morbidity and mortality of vulvar and vaginal cancers: Impact of 2-, 4-, and 9-valent HPV vaccines.** *Hum Vaccin Immunother*. 2016; **12**(6): 1352–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 60. **F** Stier EA, Chigurupati NL, Fung L: **Prophylactic HPV vaccination and anal cancer.** *Hum Vaccin Immunother*. 2016; **12**(6): 1348–51.
[PubMed Abstract](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 61. National HPV Vaccination Program Register: **HPV Vaccination Coverage.** [cited 2017 Jan 13].
[Reference Source](#)
 62. Garland SM: **The Australian experience with the human papillomavirus vaccine.** *Clin Ther*. 2014; **36**(1): 17–23.
[PubMed Abstract](#) | [Publisher Full Text](#)
 63. Oliver K, Frawley A, Garland E: **HPV vaccination: Population approaches for improving rates.** *Hum Vaccin Immunother*. 2016; **12**(6): 1589–93.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 64. **F** Gee J, Weinbaum C, Sukumaran L, *et al.*: **Quadrivalent HPV vaccine safety review and safety monitoring plans for nine-valent HPV vaccine in the United States.** *Hum Vaccin Immunother*. 2016; **12**(6): 1406–17.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 65. **F** Naud PS, Roteli-Martins CM, Carvalho NS, *et al.*: **Sustained efficacy, immunogenicity, and safety of the HPV-16/18 AS04-adjuvanted vaccine: final analysis of a long-term follow-up study up to 9.4 years post-vaccination.** *Hum Vaccin Immunother*. 2014; **10**(8): 2147–62.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 66. Garland SM, Stanley M, Brotherton J, *et al.*: **IPVS Policy statement on safety of HPV vaccines.** *Papillomavirus Res*. 2016; **2**: 9–10.
[Publisher Full Text](#)
 67. Centers for Disease Control and Prevention: **Frequently asked Questions about HPV Vaccine Safety.** [cited 2017 Mar 13].
[Reference Source](#)
 68. Larson HJ, Wilson R, Hanley S, *et al.*: **Tracking the global spread of vaccine**

- sentiments: the global response to Japan's suspension of its HPV vaccine recommendation. *Hum Vaccin Immunother*. 2014; 10(9): 2543–50.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
69. Tanaka Y, Ueda Y, Egawa-Takata T, *et al.*: Outcomes for girls without HPV vaccination in Japan. *Lancet Oncol*. 2016; 17(7): 868–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
 70. Hanley SJ, Yoshioka E, Ito Y, *et al.*: HPV vaccination crisis in Japan. *Lancet*. 2015; 385(9987): 2571.
[PubMed Abstract](#) | [Publisher Full Text](#)
 71. Vichnin M, Bonanni P, Klein NP, *et al.*: An Overview of Quadrivalent Human Papillomavirus Vaccine Safety: 2006 to 2015. *Pediatr Infect Dis J*. 2015; 34(9): 983–91.
[PubMed Abstract](#) | [Publisher Full Text](#)
 72. Safety of HPV Vaccines: Global Vaccine Safety. [cited 2017 Jan 8].
[Reference Source](#)
 73. World Health Organisation: Immunisation, Vaccines and Biologicals: Human Papillomavirus (HPV). 2016, [cited 2017 Jan 3].
[Reference Source](#)
 74. Centers for Disease Control and Prevention: Human Papillomavirus (HPV) Vaccine Safety. [cited 2017 Jan 14].
[Reference Source](#)
 75. **F** Stanley M: Immune responses to human papillomavirus. *Vaccine*. 2006; 24(Suppl 1): S16–22.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 76. Basu P, Bhatla N, Ngoma T, *et al.*: Less than 3 doses of the HPV vaccine-Review of efficacy against virological and disease end points. *Hum Vaccin Immunother*. 2016; 12(6): 1394–402.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 77. Australian Technical Advisory Group on Immunisation (ATAGI): The Australian Immunisation Handbook 10th ed (2016 update). Canberra: Australian Government Department of Health. 2016.
[Reference Source](#)
 78. Smith MA, Lew JB, Walker RJ, *et al.*: The predicted impact of HPV vaccination on male infections and male HPV-related cancers in Australia. *Vaccine*. 2011; 29(48): 9112–22.
[PubMed Abstract](#) | [Publisher Full Text](#)
 79. **F** Smith MA, Canfell K: Incremental benefits of male HPV vaccination: accounting for inequality in population uptake. *PLoS One*. 2014; 9(8): e101048.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 80. **F** Langeron N, Petry KU, Jacob J, *et al.*: An estimate of the public health impact and cost-effectiveness of universal vaccination with a 9-valent HPV vaccine in Germany. *Expert Rev Pharmacoecon Outcomes Res*. 2017; 17(1): 85–98.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 81. **F** Drolet M, Laprise JF, Boily MC, *et al.*: Potential cost-effectiveness of the nonavalent human papillomavirus (HPV) vaccine. *Int J Cancer*. 2014; 134(9): 2264–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 82. Skinner SR, Szarewski A, Romanowski B, *et al.*: Efficacy, safety, and immunogenicity of the human papillomavirus 16/18 AS04-adjuvanted vaccine in women older than 25 years: 4-year interim follow-up of the phase 3, double-blind, randomised controlled VIVIANE study. *Lancet*. 2014; 384(9961): 2213–27.
[PubMed Abstract](#) | [Publisher Full Text](#)
 83. Wheeler CM, Skinner SR, Del Rosario-Raymundo MR, *et al.*: Efficacy, safety, and immunogenicity of the human papillomavirus 16/18 AS04-adjuvanted vaccine in women older than 25 years: 7-year follow-up of the phase 3, double-blind, randomised controlled VIVIANE study. *Lancet Infect Dis*. 2016; 16(10): 1154–68.
[PubMed Abstract](#) | [Publisher Full Text](#)
 84. **F** Munoz N, Manalastas R Jr, Pitisuttithum P, *et al.*: Safety, immunogenicity, and efficacy of quadrivalent human papillomavirus (types 6, 11, 16, 18) recombinant vaccine in women aged 24–5 years: a randomised, double-blind trial. *Lancet*. 2009; 373(9679): 1949–57.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 85. Haupt RM, Wheeler CM, Brown DR, *et al.*: Impact of an HPV6/11/16/18 L1 virus-like particle vaccine on progression to cervical intraepithelial neoplasia in seropositive women with HPV16/18 infection. *Int J Cancer*. 2011; 129(11): 2632–42.
[PubMed Abstract](#) | [Publisher Full Text](#)
 86. Olsson SE, Kjaer SK, Sigurdsson K, *et al.*: Evaluation of quadrivalent HPV 6/11/16/18 vaccine efficacy against cervical and anogenital disease in subjects with serological evidence of prior vaccine type HPV infection. *Hum Vaccin*. 2009; 5(10): 696–704.
[PubMed Abstract](#) | [Publisher Full Text](#)
 87. Szarewski A, Poppe WA, Skinner SR, *et al.*: Efficacy of the human papillomavirus (HPV)-16/18 AS04-adjuvanted vaccine in women aged 15–25 years with and without serological evidence of previous exposure to HPV-16/18. *Int J Cancer*. 2012; 131(1): 106–16.
[PubMed Abstract](#) | [Publisher Full Text](#)
 88. Bosch FX, Robles C, Díaz M, *et al.*: HPV-FASTER: broadening the scope for prevention of HPV-related cancer. *Nat Rev Clin Oncol*. 2016; 13(2): 119–32.
[PubMed Abstract](#) | [Publisher Full Text](#)
 89. Brotherton JM, Kaldor JM, Garland SM: Monitoring the control of human papillomavirus (HPV) infection and related diseases in Australia: towards a national HPV surveillance strategy. *Sex Health*. 2010; 7(3): 310–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
 90. Stubbs R: Public Health England Screening-HPV primary screening in the cervical screening programme. [cited 2017 Mar 13].
[Reference Source](#)
 91. ICO Information Centre on HPV and Cancer. hpvcentre.net. [cited 2017 Jan 13].
[Reference Source](#)
 92. Centers for Disease Control and Prevention: Human Papillomavirus (HPV). [cited 2017 Jan 8].
[Reference Source](#)
 93. Centers for Disease Control and Prevention: Human Papillomavirus(HPV)ACIP Vaccine Recommendations. [cited 2017 Jan 13].
[Reference Source](#)
 94. National, Regional State, and Selected Local Area Vaccination Coverage Among Adolescents Aged 13–17years-United States 2015 MMWR Aug 26 2016. [cited 2017 Jan 13].
[Reference Source](#)
 95. NIH National Cancer Institute: Human Papillomavirus (HPV) Vaccine. 2016, [cited 2017 Jan 13].
[Reference Source](#)
 96. Powell SE, Hariri S, Steinau M, *et al.*: Impact of human papillomavirus (HPV) vaccination on HPV 16/18-related prevalence in precancerous cervical lesions. *Vaccine*. 2012; 31(1): 109–13.
[PubMed Abstract](#) | [Publisher Full Text](#)
 97. NHS: HPV Vaccine. 2014, [cited 2017 Jan 13].
[Reference Source](#)
 98. NHS: Cervical screening. 2015, [cited 2017 Jan 13].
[Reference Source](#)
 99. Public Health England: Human Papillomavirus (HPV) Vaccine Coverage in England, 2008/09 to 2013/14, A review of the full six years of the three-dose schedule. 2015, [cited 2017 Jan 13].
[Reference Source](#)
 100. Public Health England: HPR volume 10 issue 22:news (8July). [cited 2017 Jan 13].
[Reference Source](#)
 101. ICO Information Centre on HPV and Cancer: Human Papillomavirus and Related Diseases Report: Denmark, December 15 2016. [cited 2017 Jan 8].
[Reference Source](#)
 102. Blomberg M, Dehlendorff C, Sand C, *et al.*: Dose-Related Differences in Effectiveness of Human Papillomavirus Vaccination Against Genital Warts: A Nationwide Study of 550,000 Young Girls. *Clin Infect Dis*. 2015; 61(5): 676–82.
[PubMed Abstract](#) | [Publisher Full Text](#)
 103. Immunisation Scotland: HPV Vaccine. 2016, [cited 2017 Jan 13].
[Reference Source](#)
 104. Information Services Division Publication Report: HPV Immunisation Statistics Scotland, School year 2015/2016. 2016, [cited 2017 Jan 13].
[Reference Source](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. Mark Jit

¹ Department of Infectious Disease Epidemiology, London School of Hygiene and Tropical Medicine, London, UK

² National Infections Service - Colindale, Public Health England, London, UK

Competing Interests: No competing interests were disclosed.

2. Linda Niccolai

Yale School of Public Health, New Haven, CT, USA

Competing Interests: No competing interests were disclosed.

3. Eileen Dunne

¹ Division of HIV/AIDS Prevention, Centers for Disease Control and Prevention, Atlanta, Georgia, USA

² HIV/STD Research Program, Thailand MOPH CDC Collaboration, Bangkok, Thailand

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research