BRIEF REPORT

Real time portable genome sequencing for global food security [version 1; peer review: 1 approved, 1 approved with reservations]

Laura Boykin ¹, Ammar Ghalab¹, Bruno Rossitto De Marchi ^{1,2}, Anders Savill¹, James M. Wainaina¹, Tonny Kinene¹, Stephen Lamb¹, Myriam Rodrigues¹, Monica Kehoe³, Joseph Ndunguru⁴, Fred Tairo⁴, Peter Sseruwagi⁴, Charles Kayuki⁴, Deogratius Mark⁴, Joel Erasto⁴, Hilda Bachwenkizi⁴, Titus Alicai⁵, Geoffrey Okao-Okuja⁵, Phillip Abridrabo ⁵, Emmanuel Ogwok ⁵, John Francis Osingada⁵, Jimmy Akono⁵, Elijah Ateka⁶, Brenda Muga⁶, Samuel Kiarie⁶

¹School of Molecular Sciences, The University of Western Australia, Crawley, Western Australia, 6009, Australia

²School of Agriculture, Dept. of Plant Protection, São Paulo State University (UNESP), Botucatu (SP), CEP 18610-307, Brazil

³Crop Protection Branch, Department of Agriculture and Food Western Australia, South Perth, Western Australia, 6151, Australia

⁴Mikocheni Agricultural Research Institute (MARI), Dar es Salaam, Tanzania

⁵National Crops Resources Research Institute (NaCRRI), Kampala, Uganda

⁶Jomo Kenyatta University of Agriculture and Technology (JKUAT), Nairobi, Kenya

V1 First published: 18 Jul 2018, 7:1101
<https://doi.org/10.12688/f1000research.15507.1>
 Latest published: 18 Jul 2018, 7:1101
<https://doi.org/10.12688/f1000research.15507.1>

Abstract

Crop losses due to viral diseases and pests are major constraints on food security and income for millions of households in sub-Saharan Africa (SSA). Such losses can be reduced if plant diseases and pests are correctly diagnosed and identified early. Currently, accurate diagnosis for definitive identification of plant viruses and their vectors in SSA mostly relies on standard PCR and next generation sequencing technologies (NGS). However, it can take up to 6 months before results generated using these approaches are available. The long time taken to detect or identify viruses impedes quick, within-season decision-making necessary for early action, crop protection advice and disease control measures by farmers. This ultimately compounds the magnitude of crop losses and food shortages suffered by farmers. The MinION portable pocket DNA sequencer was used, to our knowledge globally for the first time, to sequence whole plant virus genomes. We used this technology to identify the begomoviruses causing the devastating cassava mosaic virus, which is ravaging smallholder farmers' crops in sub-Saharan Africa.

Open Peer Review

Approval Status

	1	2
version 1		
18 Jul 2018	view	view

1. **Ruth E Timme** , U.S. Food and Drug Administration, College Park, USA
2. **Alfonso Benítez-Páez** , Institute of Agrochemistry and Food Technology-Spanish National Research Council (IATA-CSIC), Valencia, Spain

Any reports and responses or comments on the article can be found at the end of the article.

Keywords

cassava, uganda, kenya, tanzania, nanopore, minion, SDG2

This article is included in the **Nanopore Analysis** gateway.

This article is included in the **Agriculture, Food and Nutrition** gateway.

Corresponding author: Laura Boykin (laura.boykin@uwa.edu.au)

Author roles: **Boykin L:** Conceptualization, Data Curation, Formal Analysis, Funding Acquisition, Investigation, Methodology, Project Administration, Resources, Software, Supervision, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Ghalab A:** Data Curation, Formal Analysis, Investigation, Methodology, Writing – Original Draft Preparation, Writing – Review & Editing; **Rossitto De Marchi B:** Data Curation, Formal Analysis, Investigation, Methodology, Software, Validation, Writing – Original Draft Preparation, Writing – Review & Editing; **Savill A:** Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Software, Supervision, Validation, Visualization, Writing – Review & Editing; **M. Wainaina J:** Data Curation, Investigation, Methodology, Writing – Original Draft Preparation, Writing – Review & Editing; **Kinene T:** Data Curation, Formal Analysis, Investigation, Methodology, Software, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Lamb S:** Data Curation, Formal Analysis, Investigation, Methodology, Software, Validation, Writing – Review & Editing; **Rodrigues M:** Data Curation, Formal Analysis, Investigation, Methodology, Writing – Original Draft Preparation; **Kehoe M:** Conceptualization, Formal Analysis, Investigation, Methodology, Project Administration, Resources, Software, Supervision, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Ndunguru J:** Conceptualization, Formal Analysis, Funding Acquisition, Investigation, Methodology, Project Administration, Resources, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing; **Tairo F:** Conceptualization, Project Administration, Resources, Software, Supervision, Validation, Writing – Original Draft Preparation, Writing – Review & Editing; **Sseruwagi P:** Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Project Administration, Resources, Software, Supervision, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Kayuki C:** Data Curation, Formal Analysis, Investigation, Methodology, Project Administration, Resources, Software, Supervision, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Mark D:** Data Curation, Formal Analysis, Investigation, Methodology, Software, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Erasto J:** Data Curation, Formal Analysis, Investigation, Methodology, Software, Validation, Visualization, Writing – Original Draft Preparation; **Bachwenkizi H:** Formal Analysis, Investigation, Methodology, Software, Validation, Visualization, Writing – Original Draft Preparation; **Alicai T:** Conceptualization, Data Curation, Formal Analysis, Funding Acquisition, Investigation, Methodology, Project Administration, Resources, Software, Supervision, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Okao-Okuja G:** Data Curation, Formal Analysis, Investigation, Methodology, Project Administration, Software, Supervision, Validation, Visualization, Writing – Original Draft Preparation; **Abidrabo P:** Data Curation, Formal Analysis, Investigation, Methodology, Software, Validation, Visualization, Writing – Original Draft Preparation; **Ogwok E:** Data Curation, Formal Analysis, Investigation, Methodology, Software, Validation, Visualization, Writing – Review & Editing; **Francis Osingada J:** Data Curation, Formal Analysis, Investigation, Methodology, Software, Validation, Visualization, Writing – Original Draft Preparation; **Akono J:** Formal Analysis, Investigation, Methodology, Software, Validation, Visualization, Writing – Original Draft Preparation; **Ateka E:** Conceptualization, Data Curation, Formal Analysis, Funding Acquisition, Investigation, Methodology, Project Administration, Resources, Software, Supervision, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Muga B:** Data Curation, Formal Analysis, Investigation, Methodology, Software, Supervision, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Kiarie S:** Formal Analysis, Investigation, Methodology, Validation, Visualization, Writing – Original Draft Preparation

Competing interests: No competing interests were disclosed.

Grant information: This work was part-funded by the Crawford Fund, Australia (grant number WA-803-2017).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2018 Boykin L *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Boykin L, Ghalab A, Rossitto De Marchi B *et al.* **Real time portable genome sequencing for global food security [version 1; peer review: 1 approved, 1 approved with reservations]** F1000Research 2018, 7:1101 <https://doi.org/10.12688/f1000research.15507.1>

First published: 18 Jul 2018, 7:1101 <https://doi.org/10.12688/f1000research.15507.1>

Introduction

The United Nations has listed Zero Hunger as one of the 17 global sustainable development goals to end extreme poverty by 2030. Plant viruses are a major constraint to crop production globally, causing an estimated \$30 billion in damage¹ and leaving millions of people food-insecure². In Africa, agriculture employs up to 50% of the workforce, yet only contributes 15% to the GDP on average³, suggesting that there is low productivity and limited value addition. This can be addressed through continued innovation in the fields of science and technology, as suggested in the Science Agenda for Agriculture in Africa (S3A)⁴. Sustainable management of plant viruses and their associated vectors must include efficient diagnostics for surveillance, detection and identification to inform disease management, including the development and strategic deployment of virus resistant varieties. To date, researchers have been utilizing conventional methods such as; PCR, qPCR, high-throughput sequencing (RNA-Seq, DNA-Seq) and Sanger sequencing for pathogen identification. However, these methods are both costly and time consuming, delaying timely control actions. The emergence of new tools for real-time diagnostics, such as the Oxford Nanopore MinION, have recently proven useful for the early detection of Ebola⁵ and Zika^{6,7}, even in poorly resourced laboratories. For the first time globally, the MinION portable pocket DNA sequencer was used to sequence whole plant virus genomes. We used this technology to identify the begomoviruses causing the devastating cassava mosaic virus, which is ravaging smallholder farmers' crops in sub-Saharan Africa. Cassava, a carbohydrate crop from which tapioca originates, is a major source of calories for over 800 million people worldwide. With this technology, farmers struggling with diseased crops can take immediate, restorative action to improve their livelihoods based on information about the health of their plants, generated using a portable, real-time DNA sequencing device.

Portable DNA sequencing technology has great potential to reduce the risk of community crop failure and help improve livelihoods of millions of people, especially in low resourced communities. Plant diseases are a major cause of low crop productivity and viruses such as tobacco mosaic virus, tomato mosaic, tomato spotted wilt, potato leaf roll, Potato virus X and Y in potato, papaya mosaic, citrus tristeza, chilli leaf curl, and banana bunchy top have been implicated. In particular, cassava viruses are among the world's greatest risk to food insecurity. Losses caused by cassava mosaic disease (CMD) and cassava brown streak disease are estimated at US \$2-3 billion annually⁸.

Results and discussion

We visited smallholder farmers in Tanzania, Uganda and Kenya (Table 1) who are suffering yield shortages due to cassava virus infections. We utilized the MinION to test infected material and farmers were informed within 48 hours of the specific strain of the virus that was infecting their cassava, and a resistant cassava variety was deployed. The advantages of adopting this technology far outweigh the challenges (Table 2). Cassava mosaic begomoviruses were in high enough concentration that reads of whole genomes were obtained without an enrichment step (Table 1). As expected, the viral reads increased with

the severity of the symptoms observed (Table 1). We detected a dual infection for a leaf sample with the severity score of 5 in Uganda. In addition, one asymptomatic plant in Tanzania had one viral read detected. The shortest time to obtain a viral read was 15 s (severity score 5) and the longest was 4 h 11 m 15 s (severity score 1).

MinION sequencing is superior to traditional methods of PCR identification, given its generation of whole genome sequences which enable the identification of the plant virus strain even if it becomes mutated or divergent, as it is not biased using primers that rely on known virus sequences. With regards to cassava, there are three major advantages of this technology. Firstly, improved diagnostics are required and real-time whole-genome sequencing will help develop diagnostic primers that are up-to-date. Secondly, this technology will assist with the development of resistant cassava varieties and will allow breeders to immediately test the varieties they are developing against different viral strains. Lastly, it ensures the delivery of the correct healthy uninfected planting material to farmers. In addition, we could detect virus in a plant before it showed symptoms (Table 1). Utilizing traditional PCR methods, three samples collected from farmer 1's field in Tanzania tested positive for EACMV and none were positive for ACMV. The asymptomatic sample from Mikocheni Agricultural Research Institute (MARI) tested negative for both ACMV and EACMV. There were eight fresh cassava leaf samples from Uganda found to be dually infected with ACMV and EACMV-UG using conventional PCR primers for ACMV and EACMV-UG. The primers used in this PCR yield products of 1000 bp and 1500 bp for ACMV and EACMV-UG, respectively. In total, 12 Kenyan samples were tested and all but two (barcode 2 and 3) were found to be positive using conventional PCR (Table 1). Further studies are needed to verify our results regarding the sensitivity of the protocol for early detection of CMD in cassava, but these results are very promising for ensuring farmers receive clean planting material through the early detection of viral infection.

Nanopore sequencing technology has wide applications globally, but in East Africa these include: (a) crop improvement by screening for virus resistant germplasm and genetic diversity during breeding; (b) indexing of cassava planting materials for virus presence or absence to ensure that only clean materials in multiplication fields are distributed to farmers; (c) detection and identification of alternative plant species for cassava-infecting begomoviruses, so that farmers are advised to remove and/or grow crops away from such plants as a management strategy; and (d) virus and biodiversity studies.

Methods

Sample collection and DNA extraction

In Tanzania, three cassava mosaic disease (CMD) symptomatic cassava leaf samples (Figure 1, Table 1) were collected from the smallholder cassava farmer 1's field in Bagamoyo. Disease severity was assessed as described by Legg *et al.*², where 1 is healthy and 5 shows severe symptoms of the disease, including leaf distortion and stunting of the plant. One more

Table 1. Nanopore results of cassava mosaic begomovirus-infected cassava leaves from Tanzania, Uganda and Kenya. Disease severity described by Legg *et al.*², where 1 is healthy and 5 shows severe symptoms of the disease including leaf distortion and stunting of the plant (Figure 1).

Farmer	CMD Severity	Lab ID	Barcode	Location	Concentration, ng/ul	Run Time	Time to first geminiviral read	# Nanopore reads	# Viral Reads	PCR Result	Length of 1 st virus read, bp	Closest Species Match
Tanzania												
1	3	3	1	Kiromo-Kitonga Bagamoyo	3,378.40	24 hrs	47s	63,653	950	+	969	EACMV-KE
1	2	5	2	Kiromo-Kitonga Bagamoyo	3,385.20	24 hrs	32m57s	8,844	43	+	971	EACMV-KE
1	4	6	3	Kiromo-Kitonga Bagamoyo	4,845.20	24 hrs	28s	32,540	338	+	775	EACMV-KE
1	1	8	4	MARI	1,752.70	24 hrs	4h11m15s	20,005	1	-	2213	EACMV-KE
Uganda												
2	4	1	1	Wakiso	603.2	24 hrs	15s	9,859	244	+	957	EACMV-UG
2	3	2	2	Wakiso	711.3	24 hrs	43s	8,482	98	+	780	EACMV-UG
2	5	3	3	Wakiso	603.6	24 hrs	17m14s	5,562	298	+	801	EACMV-UG AND ACMV-UG
2	2	4	4	Wakiso	598.2	24 hrs	11s	17,146	371	+	789	ACMV-UG
2	3	5	5	Wakiso	610	24 hrs	33s	7,999	250	+	2264	ACMV-UG
2	1	6A	6	Wakiso	448	24 hrs	2h14m50s	9,972	18	+	920	ACMV-KE
2	3	6B	7	Wakiso	490.9	24 hrs	2m52s	9,662	119	+	1418	ACMV-KE
2	1	7	8	NaCRRI	710.6	24 hrs	-	7,881	none	-	-	none
Kenya												
3	2	13	12	Eldoret	44.3	17hr 17min	3h4m9s	30	22	+	754	ACMV-KE
4	1	1	1	Ngoliba	2433.4	17hr 17min	20m17s	3,298	1,760	+	1145	ACMV-NG AND EACMV-UG
4	1	2	2	Ngoliba	2624.2	17hr 17min	37m23s	3,232	1,838	-	1778	ACMV-UG AND EACMV-UG
5	1	3	3	Ngoliba	1812.5	17hr 17min	10m3s	3,787	2,108	-	2723	EACMV-UG
Rainbow Hotel	1	5	4	Ruiru	3066.3	17hr 17min	57m9s	2,536	1,423	+	905	EACMV-UG
JKUAT	3	NF19C2	5	Siaya	767.59	17hr 17min	50s	2,672	1,493	+	1985	ACMV-KE
JKUAT	4	CF13B2	6	Kilifi	1143.7	17hr 17min	28s	3,870	2,221	+	847	EACMV-KE AND ACMV-KE
JKUAT	3	EF23C2	7	Kitui	1213.8	17hr 17min	18m6s	2,964	1,629	+	766	EACMV-KE
JKUAT	3	EF20C1	8	Kitui	1023	17hr 17min	54m2s	5,120	2,906	+	1887	EACMV-UG
JKUAT	3	WF2S1	9	Vihiga	1111.9	17hr 17min	23m7s	5,095	3,021	+	2233	ACMV-NG AND EACMV-UG
JKUAT	3	WF20C2	10	Busia	929.36	17hr 17min	36m1s	4,652	2,838	+	1966	ACMV-UG AND EACMV-UG
JKUAT	3	CF36C2	11	Taita taveta	1032.3	17hr 17min	55m	4,366	2,541	+	2089	EACMV-KE

JKUAT, Jomo Kenyatta University of Agriculture and Technology.

Table 2. Advantages and challenges of using the MinION portable DNA sequencer in Tanzania, Uganda and Kenya.

Advantages	Challenges
1. Rapid – obtain results timely to support quick decision making on disease management	1. Power
2. Expenses on sending samples for sequencing abroad removed	2. Internet
3. High resolution of results and reliability	3. Cost
4. Detection of mixed infections	4. Computer access
5. Detection of virus in latently infected plants	5. Shipping
6. Discovery of unknown viruses	
7. Minimising sample degradation and loss during shipping	
8. Virus-indexing for safe movement of germplasm	
9. Better plant health regulation, inspection/phytosanitary – improved international trade

Figure 1. Cassava plant infected with cassava mosaic begomoviruses.

asymptomatic leaf sample was collected at MARI, Dar es Salaam. In total, seven CMD symptomatic plants were collected from farmer 2's farm in Wakiso district in Uganda. Both Tanzanian and Ugandan samples were collected in September 2017. A total of 12 samples from Kenya were collected in February 2018 from various sources (Table 1). High-quality DNA was isolated using the cetyl trimethylammonium bromide method⁹. Each DNA sample (Table 1) was quantified and the purity checked using a NanoDrop 2000c UV–vis Spectrophotometer (Thermo Fisher Scientific, Wilmington, DE, USA) was used to check the purity and quantity of DNA for each sample, and results were recorded in Table 1.

Nanopore library preparation and sequencing

In Tanzania and Uganda, the Rapid Barcoding kit SQK-RBK001 and MinION 9.4.1 flow cells (Oxford Nanopore) were used to process genomic DNA extracted using a standard CTAB method⁹. We utilized the Rapid Barcoding kit SQK-RBK004 with 9.4.1 flow cells in Kenya. DNA was diluted to 700 ng as specified in the library protocol. The SQK-RBK001 (Sept 2017) and/or

the SQK-RBK004 (Feb 2018) protocols were performed as described by the manufacturer. In Tanzania and Uganda, the MinION was run for 24 h instead of the recommended 48 h, and in Kenya we had a total run time of approximately 17 h due to power interruptions.

Nanopore bioinformatics

In Tanzania and Uganda, Albacore 2.0.2 was used for base calling. In Kenya, Albacore 2.1.10 was used and the scripts were modified to reflect the newest rapid barcoding kit RBK004. Fastq files were imported into Geneious¹⁰ and a local blast database of all known cassava mosaic begomovirus whole genomes were downloaded from GenBank (program set to megablast, maximum hits=1, scoring (match mismatch 1-2, max e-value 1e-1, word size 28, gap cost=linear) and a local blast was performed on each of the reads generated using the Nanopore device. There are other bioinformatics pipelines that are used for Nanopore data, but the focus of our study was to use a simple, efficient analyses pipeline, which was a local blast database and a well-curated reference dataset¹¹. To ensure we did not miss other begomoviruses or pathogens, we confirmed these in-field results by performing a post diagnostic blast of reads on the Nimbus Cloud at the Pawsey Supercomputing Center (Kensington, Australia) with blast 2.2.31 against the full NCBI nucleotide database to confirm results. For reference, the specific database used was { \$ blastcmd -db nt/nt -info } Database: Nucleotide collection (nt) 46,853,753 sequences; 170,830,796,758 total bases Date: Feb 20, 2018 5:04 PM. The data were processed into a blast archive using a blast script with the following parameters (Script attached) { \$blastn -query "\$file" -db /mnt/nucdb/nt/nt -outfmt 11 -culling_limit 10 -out "out.\$file.asn" -num_threads 17 } then converted into XML (for loading into Geneious) and HTML for viewing where possible.

Verification of DNA sequencing results

Traditional PCR was used to verify our DNA sequencing results. In Tanzania and Kenya, two primer pairs: EAB 555F/EAB 555R¹² and JSP001/JSP002¹³, which amplify 556 bp and 774 bp, respectively, were used to detect East African CMVs

(EACMV) and African CMVs (ACMV), respectively. Specifically, high quality DNA were isolated using CTAB method as described by Lodhi *et al.*⁹. Electrophoresis and spectrophotometric measurements were used to check the quality and quantity of DNA. PCR reactions were carried out in a 25 µl volume, with 12.5 µl (1X final concentration) of One Taq PCR 2X Master Mix (New England Biolabs), One Taq DNA Polymerase in an optimized buffer with 1.5 mM Mg²⁺ and 0.2 mM each dNTPs in 1X final concentration. A total of 0.2 µM (0.5 µl of each) of primer sets were used in the reaction and nuclease-free water were added to 25µl final reaction volume. The template used was less than 500 ng per reaction. PCR was carried out in 2720 Thermal Cycler (Applied Biosystems) with the following programme: 1 cycle of 3 min at 94°C, then 30 cycles at 94°C for 45 s, 56°C for 45 s, 72°C for 1 min, and a final cycle at 72°C for 7 min. Electrophoresis of the PCR product was run on 1% agarose gel stained in ethidium bromide in 1X TAE buffer in a submarine gel electrophoresis unit and visualized using a BioDoc-It 210 Imaging System m-20V Transilluminator (Thomas Scientific).

In Uganda, the presence of ACMV and EACMV in each sample was detected using a pair of specific primers for ACMV, ACMV-AL1/F and ACMV-ARO/R, and primers specific for EACMV-UG2, UV-AL1/F and ACMV-CP/R3¹⁴. Specifically, the presence of ACMV and EACMV in each sample was detected using a pair of specific primers for ACMV, ACMV-AL1/F (5'-GCC GGA ATC CCT AAC ATT ATC -3') and ACMV-ARO/R (5'-GCT CGT ATG TAT CCT CTA AGG CCT G -3') and specific for EACMV-UG2, UV-AL1/F (5'-TGT CTT CTG GGA CTT GTG TG -3') and ACMV-CP/R3 (5'- TGC CTC CTG ATG ATT ATA TGT C -3') described by Zhou *et al.*¹⁴. The primers amplify about 1000 bp for ACMV and 1500 bp for EACMV-UG2 of the Coat Protein and AV2 gene sequences of the ACMV and EACMV genomes, respectively. The PCR reaction was set up using GoTaq® Green Master Mix (Promega, Madison USA). Each of the 25 µl PCR mix contained 12.0 µl of 2X GoTaq® Green Master Mix [containing GoTaq® DNA polymerase, 2X Green GoTaq® Reaction Buffer (pH 8.5), 400 µM dATP, 400 µM dGTP, 400 µM dCTP, 400 µM dTTP and 3 mM MgCl₂], 1.0 µl of forward

primer, 1.0 µl of reverse primer, 10.0 µl of nuclease-free water (Amresco, Ohio, USA) and 1.0 µl of DNA. PCR was performed using a Biometra professional thermocycler (Biometra, Gottingen, Germany) programmed as follows: 94 °C for 2 min for initial denaturation followed by 30 cycles of 94 °C for 1 min, 60 °C for 1.5 min, 72°C for 2 min and 72°C for 10 min for denaturation, annealing, extension and final extension, respectively. PCR amplicons were separated by electrophoresis in a 1× Tris-acetate-EDTA (TAE) buffer in a 1.2% agarose gel, stained with ethidium bromide (0.1 mg/ml) and visualized using a U:Genius3 (Syngene, Cambridge, UK) gel documentation system.

The image in **Figure 1** was captured in Kiromo-Kitonga Bagamoyo, Tanzania by J.N. using a Samsung s8 smartphone.

An earlier version of this article can be found on bioRxiv (DOI: <https://doi.org/10.1101/314526>).

Data availability

Raw data for this study are available on figshare: DOI: <https://doi.org/10.6084/m9.figshare.6667409>¹¹.

Competing interests

No competing interests were disclosed.

Grant information

This work was part-funded by the Crawford Fund, Australia (grant number WA-803-2017).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgements

Funding for the Kenyan trip was provided by the Crawford Fund. We also thank the participants from the University of Eldoret who assisted in the preparation of libraries for the Kenyan samples.

References

- Subramanya Sastry K, Zitter TA: **Management of Virus and Viroid Diseases of Crops in the Tropics**. In *Plant Virus and Viroid Diseases in the Tropics Vol. 2 Epidemiology and Management*. 2014; 149–480.
[Publisher Full Text](#)
- Legg JP, Thresh JM: **Cassava mosaic virus disease in East Africa: a dynamic disease in a changing environment**. *Virus Res*. 2000; **71**(1–2): 135–149.
[PubMed Abstract](#) | [Publisher Full Text](#)
- AFDB: **African Development Bank**. Annual Report. 2016.
[Reference Source](#)
- FARA: **Science agenda for agriculture in Africa (S3A): "Connecting Science" to transform agriculture in Africa**. Forum for Agricultural Research in Africa (FARA), Accra, Ghana. 2014.
[Reference Source](#)
- Quick J, Loman NJ, Duraffour S, *et al.*: **Real-time, portable genome sequencing for Ebola surveillance**. *Nature*. 2016; **530**(7589): 228–232.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Faria NR, Sabino EC, Nunes MR, *et al.*: **Mobile real-time surveillance of Zika virus in Brazil**. *Genome Med*. 2016; **8**(1): 97.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Quick J, Grubaugh ND, Pullan ST, *et al.*: **Multiplex PCR method for MinION and Illumina sequencing of Zika and other virus genomes directly from clinical samples**. *Nat Protoc*. 2017; **12**(6): 1261–1276.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Scholthof KB, Adkins S, Czosnek H, *et al.*: **Top 10 plant viruses in molecular plant pathology**. *Mol Plant Pathol*. 2011; **12**(9): 938–954.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lodhi MA, Ye GN, Weeden NF, *et al.*: **A simple and efficient method for DNA extraction from grapevine cultivars and Vitis species**. *Plant Mol Biol Rep*. 1994;

- 12(1): 6–13.
[Publisher Full Text](#)
10. Kearsse M, Moir R, Wilson A, *et al.*: **Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data.** *Bioinformatics*. 2012; **28**(12): 1647–1649.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
11. Boykin L: **Nanopore sequencing of cassava from Tanzania, Uganda and Kenya.** *figshare*. Fileset. 2018.
[Data Source](#)
12. Fondong VN, Pita JS, Rey ME, *et al.*: **Evidence of synergism between African cassava mosaic virus and a new double-recombinant geminivirus infecting cassava in Cameroon.** *J Gen Virol*. 2000; **81**(Pt 1): 287–297.
[PubMed Abstract](#) | [Publisher Full Text](#)
13. Pita JS, Fondong VN, Sangaré A, *et al.*: **Genomic and biological diversity of the African cassava geminiviruses.** *Euphytica*. 2001; **120**: 115–125.
[Publisher Full Text](#)
14. Zhou X, Liu Y, Calvert L, *et al.*: **Evidence that DNA-A of a geminivirus associated with severe cassava mosaic disease in Uganda has arisen by interspecific recombination.** *J Gen Virol*. 1997; **78**(Pt 8): 2101–2111.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 24 August 2018

<https://doi.org/10.5256/f1000research.16909.r36979>

© 2018 Benítez-Páez A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Alfonso Benítez-Páez

Microbial Ecology, Nutrition & Health Research Unit, Institute of Agrochemistry and Food Technology-Spanish National Research Council (IATA-CSIC), Valencia, Spain

Summary of the article

Boykin et al present a pilot study aiming the application of real-time DNA sequencing for the detection of ACMV and EACMV in cassava plants in multiple crops of African East countries. This study represents the successful approaching of the most valuable feature of the MinION nanopore sequencing platform, its portability. At the same time, the authors made the maximum use of the singularities of the system by the translational application of their results, thus preventing the spread of plant viruses and to improve the crop efficiency by timely advising of farmers. This last exercise really highlights the value of such technology, particularly in the epidemiological surveillance and control of pathogens.

Notwithstanding, I have some minor concerns that if addressed they would constitute an added value to the approach described.

- 1) I strongly recommend that authors store and make publicly available the genetic information retrieved from different sequencing runs to a specialized repository such as ENA or GenBank.
- 2) It would be very informative for future studies in this field that Table 1 contains additional information the average or median values of the sequence identity derived from the comparison between nanopore reads and reference sequences. In a similar way, they should declare the level of relationship between ACMV and EACMV, in terms of genome-wide nucleotide identity, in order to disclose any potential misidentification given the high error rate of nanopore-derived DNA reads.
- 3) The authors must be aware that there is a strong effect derived from sequencing kits used, different from Uganda/Tanzania and Kenya. They should make some correlations between the CMD severity scoring and DNA reads retrieved independently accordingly to the kits used.

4) In the same line of thoughts than above, it would be very elegant that authors will estimate the maximum time for expecting a viral DNA read just for setting a threshold and optimize the sequencing time. I have noticed that for CMD severity = 1, it took maximum 4h to retrieve viral DNA reads, using the sequencing kit SQK-RBK001. Another different history was the utilization of SQK-RBK004 in Kenya, where apparently there is not a correlation between CMD severity and viral DNA reads retrieved. In the last cases, the apparently not symptomatic plants were detected as positive in less than one hour. The setting of a time threshold for a proper detection (getting enough number of reads to estimate reliable identification) would be useful to speed up the farmers' advising and consequently the reduction of risks for the spread.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: My expertise areas are the microbial genomics, nanopore DNA sequencing technology, molecular biology, and the massive DNA data analysis.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Reader Comment 13 Nov 2018

Edwin karoney, Masinde Muliro University of Science and Technology, Kenya

Great work! Outstanding technology!

Competing Interests: No competing interests were disclosed.

Reviewer Report 26 July 2018

<https://doi.org/10.5256/f1000research.16909.r36183>

© 2018 Timme R. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Ruth E Timme

Center for Food Safety and Applied Nutrition, U.S. Food and Drug Administration, College Park, MD, USA

Boykin and co-authors present results from a pilot study exploring the use of MinION technology to detect a plant viral pathogen in real-time. Their data shows a huge advantage of using this hand-held sequence technology over the standard PCR methods. The authors propose implementing a MinION quality-control step on the plants before they are distributed, so that CMD-infected plants can be removed from the supply chain before they reach local farmers. This report is short, but its impact appears to be far-reaching. Most of my comments are minor editorial suggestions, but overall the writing and readability is excellent.

Minor revisions:

Public availability of the DNA sequence data. While the authors technically made the sequence files public by posting to FigShare, the standard repository for DNA sequence data is the INSDC (NCBI/EBI/DBJ). I highly urge the authors to create a BioProject at NCBI or EBI that houses the raw and assembled sequences (fasta files) for this effort so that other researchers in this area can easily build off this important work.

Editorial comments:

Results and Discussion

"We utilized the MinION to test infected material and farmers were informed within 48 hours of the specific strain of the virus that was infecting their cassava, and a resistant cassava variety was deployed." Consider converting to two sentences. One about using the MinION and the second to cover the response. Were resistant cassava plants really deployed within 48 hrs? wow.

"MinION sequencing is superior to traditional methods of PCR identification, given its generation of whole genome sequences which enable the identification of the plant virus strain even if it becomes mutated or divergent, as it is not biased using primers that rely on known virus sequences." Consider a minor re-write: "In general MinION sequencing is superior to traditional PCR methods of identification because the virus can be detected even when the PCR primers don't work, and 2) entire viral genome sequence is generated enabling the identification of the specific viral strain, along with other molecular information, which allows for a much higher resolution of surveillance.

"In addition, we could detect virus in a plant before it showed symptoms (Table 1)." Change to present tense to match the rest of the paragraph?

"Utilizing traditional PCR methods, three samples collected from farmer 1's field in Tanzania tested positive for EACMV and none were positive for ACMV." Define EACMV and ACMV before abbreviation.

Methods:

"In Tanzania, three cassava mosaic disease (CMD) symptomatic cassava leaf samples (Figure 1, Table 1) were collected from the smallholder cassava farmer 1's field in Bagamoyo." CMD already defined in Intro.

"In Tanzania and Kenya, two primer pairs: EAB 555F/EAB 555F12 and JSP001/JSP00213, which amplify 556 bp and 774 bp, respectively, were used to detect East African CMVs (EACMV) and African CMVs (ACMV), respectively." Use the abbreviations here after adding the full names to the Results.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: I'm a Research Microbiologist for the US Food and Drug Administration. My research focuses on the validation of phylogenetic tools for molecular surveillance. I also manage a genome surveillance network for foodborne pathogens called the GenomeTrakr, that collects

and publishes WGS data from food and environmental isolates.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research