REVIEW

Recent advances in understanding of the pathogenesis of ANCA-associated vasculitis [version 1; peer review: 2 approved]

Maria Prendecki , Charles D. Pusey

Department of Medicine, Imperial College London, Hammersmith Campus, Du Cane Road, London, W12 0NN, UK

V1 **First published:** 19 Jul 2018, 7(F1000 Faculty Rev):1113
<https://doi.org/10.12688/f1000research.14626.1>
Latest published: 19 Jul 2018, 7(F1000 Faculty Rev):1113
<https://doi.org/10.12688/f1000research.14626.1>

Abstract

Anti-neutrophil cytoplasm antibody (ANCA)-associated vasculitides (AAV) are rare systemic autoimmune diseases characterised by inflammation of small blood vessels. Recent developments have been made in our understanding of the pathogenesis of these diseases, including the pathogenic role of ANCA, neutrophils and monocytes as mediators of injury, dysregulation of the complement system, and the role of T and B cells. Current treatment strategies for AAV are based on broad immunosuppression, which may have significant side effects. Advances in understanding of the pathogenesis of disease have led to the identification of new therapeutic targets which may lead to treatment protocols with less-toxic side effects. The aim of this review is to summarise current information and recent advances in understanding of the pathogenesis of AAV.

Keywords

ANCA, Vasculitis

Open Peer Review

Approval Status

	1	2
version 1
19 Jul 2018		

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Minghui Zhao**, Peking University First Hospital, Beijing, China
2. **Augusto Vaglio**, University Hospital of Parma, Parma, Italy

Any comments on the article can be found at the end of the article.

Corresponding author: Maria Prendecki (m.prendecki@imperial.ac.uk)

Author roles: Prendecki M: Writing – Original Draft Preparation; Pusey CD: Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: We acknowledge support from the NIHR Imperial Biomedical Research Centre.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2018 Prendecki M and Pusey CD. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Prendecki M and Pusey CD. **Recent advances in understanding of the pathogenesis of ANCA-associated vasculitis [version 1; peer review: 2 approved]** F1000Research 2018, 7(F1000 Faculty Rev):1113 <https://doi.org/10.12688/f1000research.14626.1>

First published: 19 Jul 2018, 7(F1000 Faculty Rev):1113 <https://doi.org/10.12688/f1000research.14626.1>

Introduction and background

The anti-neutrophil cytoplasm antibody (ANCA)-associated vasculitides (AAV) are a group of systemic autoimmune diseases characterised by inflammation of small blood vessels with multi-organ involvement, including the kidney, lung, nerves, gut, and ear, nose, and throat (ENT). Until 1979, it was assumed that rapidly progressive glomerulonephritis (RPGN) was caused by circulating immune complexes or anti-glomerular basement membrane (anti-GBM) antibodies. However, Stilmant *et al.* observed that many cases had no evidence of glomerular deposition of complement or immunoglobulin and were pauci-immune¹. Subsequently, antibody binding to neutrophil cytoplasm was shown by using serum from patients with crescentic glomerulonephritis for indirect immunofluorescence². The two main target antigens of ANCA were then identified as proteinase-3 (PR3) and myeloperoxidase (MPO), which are present in the granules of neutrophils and lysosomes of monocytes³⁻⁵. There are differing clinical syndromes associated with ANCA: granulomatosis with polyangiitis (GPA), microscopic polyangiitis (MPA), eosinophilic GPA (EGPA), and renal limited vasculitis. Around 10% of patients are ANCA negative⁶. The aim of this review is to provide an overview of the current information and recent advances in understanding of the pathogenesis of AAV focussing on MPA and GPA rather than EGPA.

AAV is uncommon; its incidence in Europe is reported to be 13 to 20 cases per million⁷. There is a slight male preponderance, and incidence increases with age, although peak incidence has been reported variously as 55 to 64, 65 to 74, and more than 75 years⁷⁻⁹. AAV is rarer in non-Caucasian or non-Asian populations, and there are differences in the incidence of different clinical phenotypes between populations. When a Japanese and UK population were directly compared, the overall incidence of AAV was similar but GPA was much less common in Japan¹⁰. There is certainly a genetic basis for AAV, and this may explain some of the population differences. Two large genome-wide association studies showed an association between AAV and genetic factors, and there was a stronger genetic association with ANCA specificity than clinical syndrome, suggesting that MPO-ANCA and PR3-ANCA may be defining differing diseases. In anti-PR3 AAV, there were associations with *HLA-DP*, *PRTN3* (the gene encoding proteinase-3), and *SERPINA1* (the gene encoding α 1-antitrypsin, a circulating inhibitor of PR3); anti-MPO AAV was associated mainly with *HLA-DQ* polymorphisms^{11,12}. There are reports of an association between *HLA-DRB1*15* and PR3-ANCA in African-Americans, and *HLA DPB1*0401* has also been associated with PR3-ANCA disease^{13,14}. Several studies have shown an association with a single-nucleotide polymorphism (SNP) in *PTPN22* (the gene encoding a lymphoid-specific phosphatase which is involved in T-cell activation) and GPA, although whether this SNP is also associated with MPA is less clear^{15,16}.

There are several reported environmental associations with AAV. Infection may precede disease relapse and nasal carriage of staphylococci correlates with disease relapse in patients with anti-PR3 AAV and ENT disease¹⁷. A mechanism of molecular mimicry whereby an immune response against microbial antigens

cross-reacts with self tissue has been proposed¹⁸. An atypical ANCA, anti-human lysosome-associated membrane protein-2 (anti-LAMP-2) antibody, was first identified in patients with pauci-immune glomerulonephritis (GN) in 1995 and has 100% sequence homology with FimH, a bacterial adhesion protein on Gram-negative bacteria. Rats immunised with FimH develop GN and antibodies which react to human and rat LAMP-2¹⁸. However, the clinical association has been reproduced in some but not other laboratories^{19,20}. An alternative proposal involves complementary peptides of the auto-antigen. Patients with anti-PR3 AAV have been shown to have circulating antibodies to both PR3 and anti-sense complementary PR3 peptides (cPr3), suggesting that an initial immune response may be against the anti-sense peptide leading to the development of anti-idiotypic antibodies which recognise PR3²¹.

Other environmental risk factors identified include silica, heavy metal exposure and drugs which can induce ANCA, including propylthiouracil, hydralazine, and levamisole-contaminated cocaine²²⁻²⁴.

Pathogenicity of ANCA

ANCA have been shown to be pathogenic in several clinical and pre-clinical studies. There is a reported case of maternal-foetal transfer of anti-MPO ANCA resulting in neonatal renal disease and pulmonary haemorrhage shortly after birth²⁵. Levels of ANCA have been shown to correlate with disease activity in some but not all case series with better correlation in patients with renal disease²⁶. Removal of antibodies with plasma exchange has been shown to improve prognosis in severe AAV²⁷, and depletion of B cells with rituximab has been shown to be effective in induction and maintenance of remission²⁸⁻³⁰.

Some of the best evidence for the pathogenesis of ANCA comes from a passive transfer model of anti-MPO AAV. MPO-deficient mice are immunised with mouse MPO and develop high-titre anti-MPO antibodies. Transfer of these antibodies into wild-type or Rag2 mice (which lack lymphocytes) results in the mice developing severe vasculitis with crescentic GN and pulmonary haemorrhage, demonstrating that MPO-ANCA alone are sufficient to induce disease³¹. Neutrophils were shown to have an essential role in disease pathogenesis in this model; depletion of neutrophils prior to transfer of antibodies prevented the development of disease³². ANCA have also been shown to be pathogenic in an autoimmune rat model of AAV, experimental autoimmune vasculitis in the susceptible WKY rat strain. Rats are immunised with human MPO in complete Freund's adjuvant and also receive two doses of intraperitoneal pertussis toxin as an immune adjuvant. Animals develop polyclonal anti-MPO antibodies with pauci-immune vasculitis and pulmonary haemorrhage. Intra-vital imaging in this model showed increased leucocyte adhesion and transmigration at the endothelium in response to CXCL1, and this could also be observed in healthy animals following infusion of anti-MPO IgG isolated from rats with disease, supporting a role for the pathogenicity of ANCA³³. The pathogenic role of PR3-ANCA is less well defined and this is owing, at least in part, to the difficulty in developing animal models of anti-PR3 AAV. An attempt to create a passive transfer model analogous to the one using anti-MPO

antibodies resulted in no features of vasculitis and only a mild inflammatory response to tumour necrosis factor (TNF) in the skin³⁴. This is potentially due to a lack of PR3 expression on the surface of unstimulated mouse neutrophils and a lesser degree of sequence homology between mouse and human PR3 than there is for MPO³⁵.

Despite evidence for the pathogenicity of ANCA, the relationship between ANCA and active vasculitis is complex and ANCA are not always pathogenic. ANCA can persist in remission, can recur without evidence of clinical relapse, and have been identified in healthy individuals. Natural anti-MPO antibodies are of lower avidity and titre than are antibodies from patients with AAV³⁶. The IgG subclass of ANCA may also be important. *In vitro*, IgG3-ANCA has been shown to be more effective than other IgG subclasses at activating neutrophils, although in other clinical studies the IgG subclass of ANCA did not correlate with disease severity^{37,38}. Epitope mapping to identify the pathogenic epitopes of both PR3 and MPO have been carried out. One study using epitope excision and mass spectrometry identified a linear epitope on MPO at residue 447–459 that was limited to patients with disease; interestingly, when the three-dimensional structure of MPO was visualised, this epitope was close to epitopes seen in individuals with natural antibodies, leading the authors to suggest that pathogenic ANCA arise by a process of epitope spreading. In this study, IgG purified from patients with ANCA-negative vasculitis was able to bind to an MPO epitope, and it was suggested that competition for binding in immunoassays by a fragment of caeruloplasmin may be why ANCA cannot be detected in these patients³⁹.

ANCA-induced activation of neutrophils and monocytes

The ability of ANCA to bind to and activate neutrophils causing degranulation and production of reactive oxygen species (ROS) was first shown nearly 30 years ago⁴⁰. Since then, several *in vitro* studies have shown that neutrophils which have been primed with TNF α , lipopolysaccharide (LPS), or complement (C5a) undergo activation and degranulation and mediate endothelial cell damage when stimulated with MPO or PR3-ANCA^{41,42}. ANCA binding to neutrophils has also been shown to activate intracellular signalling pathways leading to altered adhesion molecule expression and conformational changes which promote neutrophil adhesion and transmigration at the vascular endothelium⁴³. Both the ANCA antigen-binding site and binding to Fc γ receptors on the surface of primed neutrophils and monocytes have been identified as mechanisms by which ANCA activates these cells.

ANCA have also been shown to be mediators of NETosis, a form of neutrophil cell death with release of neutrophil extracellular traps (NETs). NETs have a DNA backbone with a variety of pro-inflammatory proteins, including histones, high-mobility group box 1 (HMGB-1), neutrophil elastase, calprotectin, MPO, and PR3⁴⁴. NETs have been shown to be present at sites of tissue damage in AAV, and patients also have increased levels of NETs in the circulation⁴⁵. NETs may play a pathogenic role in AAV; they can cause activation of dendritic cells and autoreactive B cells, endothelial damage, and complement

activation^{46,47}. NETs may also play a role in the loss of tolerance to ANCA antigens; one study has shown that dendritic cells activated by NETs induce loss of tolerance to both MPO and PR3⁴⁸. Neutrophils from patients with AAV undergo more spontaneous NETosis than those from healthy controls, but ANCA can also induce this process. The exact mechanism by which ANCA induce NETosis is unclear but is thought to require binding of both Fc γ receptors and the ANCA target antigen on the cell surface⁴⁹.

Although many studies have focussed on neutrophils and their interactions with ANCA in the pathogenesis of disease, monocytes may also play a role in mediating AAV. Monocytes express ANCA antigens, and stimulation of monocytes *in vitro* with ANCA leads to cytokine production and generation of ROS^{50,51}. Monocytes from patients with AAV have been shown to express higher levels of CD14, the LPS receptor, than monocytes from patients in remission or healthy controls, suggesting an increased cell activation state in patients with AAV⁵². Circulating monocytes from patients with active AAV have also been shown to express higher levels of cell surface markers which are essential for interaction between leucocytes and the endothelium⁵³. Recent studies have shown that monocytes and macrophages are the predominant cells in glomeruli in renal biopsies from patients with AAV^{54,55}. In one study using the passive transfer model of mouse anti-MPO AAV, depleting monocytes decreased glomerular crescent formation but had no effect on urinary abnormalities⁵⁶.

Complement and AAV

There is increasing evidence for a role for complement in the pathogenesis of AAV. In the antibody transfer model of mouse anti-MPO AAV, mice deficient in C5 or those depleted of complement by pre-treatment with cobra venom did not develop disease. C4-deficient mice were not protected, suggesting a role for the alternative rather than the classic pathway⁵⁷. Mice deficient in C5aR are protected from disease, and mice with knock-in of the human C5a receptor treated with an antagonist of human C5aR (CCX168; avacopan) showed decreased disease severity^{58,59}. There is also evidence for a role for complement from *in vitro* studies; C5a can prime neutrophils to respond to stimulation by ANCA, and this may be due to its actions at the C5aR⁴². The interaction of C5a with its other receptor, C5L2, is more complex, and some studies report that it has a pro-inflammatory role *in vitro* but knockout of C5L2 resulted in more-severe disease in mouse anti-MPO AAV^{59,60}. It has also been shown that both ANCA-stimulated neutrophils and NETs can activate the alternative pathway of the complement system, leading to a positive feedback loop^{47,57}. There is evidence of complement deposition, such as C3d and factor B, at sites of tissue inflammation in patients with AAV and kidney deposition of Bb (a marker of activation of the alternative pathway) correlated with pathological severity of disease⁶¹. Plasma levels of C3a, C5a, soluble C5b-9, and Bb were higher in patients with active AAV than in those in remission^{62,63}. In one study, patients with lower circulating C3 levels were shown to have poorer outcomes in terms of both patient and renal survival⁶⁴. Blockade of C5 cleavage with eculizumab has been reported as treatment for AAV in one case report. It was used as

add-on therapy to cyclophosphamide with good renal recovery, although unfortunately the patient developed non-Hodgkin lymphoma, thought to be unrelated to the eculizumab, and died from sepsis following chemotherapy⁶⁵. A recently published phase II trial has shown that avacopan was effective in replacing high-dose glucocorticoids for induction of remission when added to cyclophosphamide or rituximab⁶⁶. A phase III trial of this treatment approach is currently recruiting (ADVOCATE, ClinicalTrials.gov Identifier: NCT02994927).

B cells and AAV

B cells have a central role in AAV in that they produce ANCA, and levels of activated B cells have been shown to correlate with disease activity⁶⁷. Depletion of B cells with rituximab has been shown to be effective in inducing and maintaining disease remission^{28,29}. The return of B cells after rituximab may predict relapse of AAV, and it has been shown that following induction of remission with rituximab and cyclophosphamide, the return of B cells has a high negative predictive value for relapse but a poor positive predictive value^{68,69}. It may be that the phenotype of the repopulating B cells is important in predicting relapse, and one study suggested that those who repopulate with a low percentage of CD5⁺ B cells have a shorter time to relapse⁷⁰. Several studies have shown differences in B-cell subsets between patients with AAV and healthy controls. One study reported a memory B-cell subset with higher CD19 expression in patients with AAV, suggesting that these may represent autoreactive B cells⁷¹. Regulatory B (B_{reg}) cells skew T-cell differentiation towards regulatory T (T_{reg}) cells and away from T helper 1 (T_{H1}) and T_{H17} phenotypes and decrease B cells which are producing ANCA⁷². Several studies have shown decreased B_{reg} cells in patients with AAV as defined by cell surface markers such as CD5, CD24, and CD38^{73,74}. *In vitro*, neutrophils stimulated with ANCA release B-cell survival factors such as B lymphocyte stimulator (BLyS) and a proliferation-inducing ligand (APRIL). In one study, incubating B cells with supernatant from ANCA-stimulated neutrophils or with recombinant BLyS resulted in increased B-cell survival⁷⁵. Several studies have reported higher levels of BLyS in patients with AAV, and some have shown that levels correlated with disease activity and ANCA titre and decreased following treatment^{76,77}. Following rituximab treatment, serum BLyS levels have been shown to increase both in patients with AAV and in patients with other auto-immune diseases^{75,78}. One study has shown that a SNP in BLyS predicted which patients were more likely to relapse following rituximab and had earlier return of B cells after treatment. The authors suggest that this SNP may result in higher baseline BLyS or increase in BLyS after B-cell depletion⁷⁹. This may suggest a potential role for targeting BLyS as maintenance treatment of AAV following induction treatment with rituximab. A phase III trial which added anti-BLyS treatment with belimumab to azathioprine and steroids for maintenance of remission did not show any reduction in risk of relapse; however, in the subgroup of patients who received rituximab as an induction agent, belimumab did reduce relapse rate, although this was not significant⁸⁰.

T-cell immunity and AAV

T cells are present in glomeruli and the tubulointerstitium in renal biopsy tissue from patients with AAV, suggesting that T-cell responses are pathogenic. Studies have shown that patients

with AAV have defective T_{reg} cell suppressive function⁸¹; one study has also shown increased frequency of a CD4⁺ T-cell subset that is resistant to the suppressor effects of T_{reg} cells⁸². In a small group of patients, anti-thymocyte globulin was used as a successful treatment for refractory GPA⁸³. Additionally, differential T_H cell polarisation has been described in AAV, such that patients with active and systemic disease are more likely to have a T_{H2} response⁸⁴.

In the mouse passive transfer model of anti-MPO AAV, CD4⁺ T cells have been used to transfer disease, demonstrating a role for T cells in pathogenesis. Mice pre-immunised with CD4⁺ T cells from MPO-immunised, B-cell-deficient, MPO-deficient mice developed greater severity of GN after induction of disease with MPO-ANCA compared with mice immunised with OVA-sensitised CD4⁺ cells⁸⁵. In a model of anti-MPO AAV in which mice are immunised with MPO followed by a subnephritogenic dose of anti-GBM globulin, depletion of CD4⁺ cells decreased disease severity with no effect on ANCA titres⁸⁶. This model of disease has been used to identify pathogenic epitopes for both CD4⁺ and CD8⁺ T cells, and these epitopes have been used to induce disease^{87,88}.

The T_{H17} axis may also be involved in the development of ANCA; serum interleukin-23 (IL-23) and IL-17 are raised in the serum of patients with acute AAV, and in one study IL-23 levels correlated with disease activity⁸⁹. IL-23 induces T-cell differentiation into the T_{H17} subset and enhances the production of IL-17 from these cells. Stimulation of neutrophils by ANCA has been shown to induce the production of IL-17⁹⁰, and in one study IL-17-deficient mice were protected from MPO-ANCA-induced disease⁹¹.

Granuloma formation

Granulomatous disease is frequently seen in isolated and systemic GPA. Early granuloma formation is typified by activated neutrophils forming micro-abscesses and only scattered multinucleated giant cells. Later granulomas consist of a central necrotic area with multinucleated giant cells at the margin and surrounding dendritic cells, T lymphocytes, B lymphocytes, and plasma cells forming a follicular structure of ectopic lymphoid tissue^{92,93}. The mechanisms initiating granuloma formation have not been fully identified, but there is some evidence that granulomatous inflammation is being driven by T cells producing T_{H1} cytokines⁹⁴. It has also been shown that APRIL and BLyS are present in granulomas along with activated B cells, leading some authors to suggest that close association of B cells with PR3-positive cells within granulomas could lead to initiation or maintenance of anti-PR3 responses⁹⁵. In an *in vivo* model of xenografted nasal mucosa from patients with GPA to mice, tissue damage was shown to be mediated by fibroblasts⁹⁶.

Conclusions

The pathogenesis of AAV is complex and remains incompletely understood. Recent advances have been made in our understanding of the mechanisms of both the development of auto-immunity and inflammation leading to tissue damage. Our understanding of the generation of the auto-immune response is incomplete but may well involve molecular mimicry and dysregulation of both B and T cells. There is substantial evidence

for the pathogenicity of ANCA, and neutrophils are both the target of ANCA and mediators of endothelial injury. NETs in particular have been shown to mediate tissue damage but also could be involved in the loss of tolerance to ANCA. Advances in understanding the role of the alternative pathway of the complement system in AAV have led to clinical trials of novel therapeutic agents. Further understanding of the mechanisms of disease may lead to the use of other novel therapeutics such as molecules to block NETosis, BlyS inhibitors, or monoclonal antibodies against IL-17 or IL-23.

Competing interests

The authors declare that they have no competing interests.

Grant information

We acknowledge support from the NIHR Imperial Biomedical Research Centre.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- Stilmant MM, Bolton WK, Sturgill BC, *et al.*: **Crescentic glomerulonephritis without immune deposits: clinicopathologic features.** *Kidney Int.* 1979; **15**(2): 184–95.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Davies DJ, Moran JE, Niall JF, *et al.*: **Segmental necrotising glomerulonephritis with antineutrophil antibody: possible arbovirus aetiology?** *Br Med J (Clin Res Ed).* 1982; **285**(6342): 606.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Falk RJ, Jennette JC: **Anti-neutrophil cytoplasmic autoantibodies with specificity for myeloperoxidase in patients with systemic vasculitis and idiopathic necrotizing and crescentic glomerulonephritis.** *N Engl J Med.* 1988; **318**(25): 1651–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Niles JL, McCluskey RT, Ahmad MF, *et al.*: **Wegener's granulomatosis autoantigen is a novel neutrophil serine proteinase.** *Blood.* 1989; **74**(6): 1888–93.
[PubMed Abstract](#)
- van der Woude FJ, Rasmussen N, Lobatto S, *et al.*: **Autoantibodies against neutrophils and monocytes: tool for diagnosis and marker of disease activity in Wegener's granulomatosis.** *Lancet.* 1985; **1**(8426): 425–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Eisenberger U, Fakhouri F, Vanhille P, *et al.*: **ANCA-negative pauci-immune renal vasculitis: histology and outcome.** *Nephrol Dial Transplant.* 2005; **20**(7): 1392–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Watts RA, Lane SE, Benthall G, *et al.*: **Epidemiology of systemic vasculitis: a ten-year study in the United Kingdom.** *Arthritis Rheum.* 2000; **43**(2): 414–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Gonzalez-Gay MA, Garcia-Porrúa C, Guerrero J, *et al.*: **The epidemiology of the primary systemic vasculitides in northwest Spain: implications of the Chapel Hill Consensus Conference definitions.** *Arthritis Rheum.* 2003; **49**(3): 388–93.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Mohammad AJ, Jacobsson LT, Westman KW, *et al.*: **Incidence and survival rates in Wegener's granulomatosis, microscopic polyangiitis, Churg-Strauss syndrome and polyarteritis nodosa.** *Rheumatology (Oxford).* 2009; **48**(12): 1560–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Fujimoto S, Watts RA, Kobayashi S, *et al.*: **Comparison of the epidemiology of anti-neutrophil cytoplasmic antibody-associated vasculitis between Japan and the U.K.** *Rheumatology (Oxford).* 2011; **50**(10): 1916–20.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lyons PA, Rayner TF, Trivedi S, *et al.*: **Genetically distinct subsets within ANCA-associated vasculitis.** *N Engl J Med.* 2012; **367**(3): 214–23.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Merkel PA, Xie G, Monach PA, *et al.*: **Identification of Functional and Expression Polymorphisms Associated With Risk for Antineutrophil Cytoplasmic Autoantibody-Associated Vasculitis.** *Arthritis Rheumatol.* 2017; **69**(5): 1054–66.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Cao Y, Schmitt JL, Yang J, *et al.*: **DRB1*15 allele is a risk factor for PR3-ANCA disease in African Americans.** *J Am Soc Nephrol.* 2011; **22**(6): 1161–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Hilhorst M, Arndt F, Joseph Kemna M, *et al.*: **HLA-DPB1 as a Risk Factor for Relapse in Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: A Cohort Study.** *Arthritis Rheumatol.* 2016; **68**(7): 1721–30.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Carr EJ, Niederer HA, Williams J, *et al.*: **Confirmation of the genetic association of CTLA4 and PTPN22 with ANCA-associated vasculitis.** *BMC Med Genet.* 2009; **10**: 121.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Marotana D, Maritati F, Malerba G, *et al.*: **PTPN22 R620W polymorphism in the ANCA-associated vasculitides.** *Rheumatology (Oxford).* 2012; **51**(5): 805–12.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Popa ER, Stegeman CA, Abdulahad WH, *et al.*: **Staphylococcal toxic-shock-syndrome-toxin-1 as a risk factor for disease relapse in Wegener's granulomatosis.** *Rheumatology (Oxford).* 2007; **46**(6): 1029–33.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kain R, Exner M, Brandes R, *et al.*: **Molecular mimicry in pauci-immune focal necrotizing glomerulonephritis.** *Nat Med.* 2008; **14**(10): 1088–96.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Roth AJ, Brown MC, Smith RN, *et al.*: **Anti-LAMP-2 antibodies are not prevalent in patients with antineutrophil cytoplasmic autoantibody glomerulonephritis.** *J Am Soc Nephrol.* 2012; **23**(3): 545–55.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Peschel A, Basu N, Benharkou A, *et al.*: **Autoantibodies to hLAMP-2 in ANCA-negative pauci-immune focal necrotizing GN.** *J Am Soc Nephrol.* 2014; **25**(3): 455–63.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Pendergraft WF 3rd, Preston GA, Shah RR, *et al.*: **Autoimmunity is triggered by cPR-3(105-201), a protein complementary to human autoantigen proteinase-3.** *Nat Med.* 2004; **10**(1): 72–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Pendergraft WF 3rd, Herlitz LC, Thornley-Brown D, *et al.*: **Nephrotoxic effects of common and emerging drugs of abuse.** *Clin J Am Soc Nephrol.* 2014; **9**(11): 1996–2005.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- de Lind van Wijngaarden RA, van Rijn L, Hagen EC, *et al.*: **Hypotheses on the etiology of antineutrophil cytoplasmic autoantibody associated vasculitis: the cause is hidden, but the result is known.** *Clin J Am Soc Nephrol.* 2008; **3**(1): 237–52.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Yu F, Chen M, Gao Y, *et al.*: **Clinical and pathological features of renal involvement in propylthiouracil-associated ANCA-positive vasculitis.** *Am J Kidney Dis.* 2007; **49**(5): 607–14.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Bansal PJ, Tobin MC: **Neonatal microscopic polyangiitis secondary to transfer of maternal myeloperoxidase-antineutrophil cytoplasmic antibody resulting in neonatal pulmonary hemorrhage and renal involvement.** *Ann Allergy Asthma Immunol.* 2004; **93**(4): 398–401.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Fussner LA, Hummel AM, Schroeder DR, *et al.*: **Factors Determining the Clinical Utility of Serial Measurements of Antineutrophil Cytoplasmic Antibodies Targeting Proteinase 3.** *Arthritis Rheumatol.* 2016; **68**(7): 1700–10.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Jayne DR, Gaskin G, Rasmussen N, *et al.*: **Randomized trial of plasma exchange or high-dosage methylprednisolone as adjunctive therapy for severe renal vasculitis.** *J Am Soc Nephrol.* 2007; **18**(7): 2180–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Jones RB, Tervaert JW, Hauser T, *et al.*: **Rituximab versus cyclophosphamide in ANCA-associated renal vasculitis.** *N Engl J Med.* 2010; **363**(3): 211–20.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Stone JH, Merkel PA, Spiera R, *et al.*: **Rituximab versus cyclophosphamide**

- for ANCA-associated vasculitis. *N Engl J Med*. 2010; 363(3): 221–32.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
30. **F** Guillevin L, Pagnoux C, Karras A, *et al.*: Rituximab versus azathioprine for maintenance in ANCA-associated vasculitis. *N Engl J Med*. 2014; 371(19): 1771–80.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 31. Xiao H, Heeringa P, Hu P, *et al.*: Antineutrophil cytoplasmic autoantibodies specific for myeloperoxidase cause glomerulonephritis and vasculitis in mice. *J Clin Invest*. 2002; 110(7): 955–63.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 32. Xiao H, Heeringa P, Liu Z, *et al.*: The role of neutrophils in the induction of glomerulonephritis by anti-myeloperoxidase antibodies. *Am J Pathol*. 2005; 167(1): 39–45.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 33. Little MA, Smyth CL, Yadav R, *et al.*: Antineutrophil cytoplasm antibodies directed against myeloperoxidase augment leukocyte-microvascular interactions *in vivo*. *Blood*. 2005; 106(6): 2050–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
 34. Pfister H, Ollert M, Fröhlich LF, *et al.*: Antineutrophil cytoplasmic autoantibodies against the murine homolog of proteinase 3 (Wegener autoantigen) are pathogenic *in vivo*. *Blood*. 2004; 104(5): 1411–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
 35. Little MA: L7. Animal models of PR3-ANCA vasculitis: approaches and controversies. *Presse Med*. 2013; 42(4 Pt 2): 512–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
 36. Xu PC, Cui Z, Chen M, *et al.*: Comparison of characteristics of natural autoantibodies against myeloperoxidase and anti-myeloperoxidase autoantibodies from patients with microscopic polyangiitis. *Rheumatology (Oxford)*. 2011; 50(7): 1236–43.
[PubMed Abstract](#) | [Publisher Full Text](#)
 37. Nowack R, Grab I, Flores-Suarez LF, *et al.*: ANCA titres, even of IgG subclasses, and soluble CD14 fail to predict relapses in patients with ANCA-associated vasculitis. *Nephrol Dial Transplant*. 2001; 16(8): 1631–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
 38. Mulder AH, Heeringa P, Brouwer E, *et al.*: Activation of granulocytes by anti-neutrophil cytoplasmic antibodies (ANCA): a Fc gamma RII-dependent process. *Clin Exp Immunol*. 1994; 98(2): 270–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 39. **F** Roth AJ, Ooi JD, Hess JJ, *et al.*: Epitope specificity determines pathogenicity and detectability in ANCA-associated vasculitis. *J Clin Invest*. 2013; 123(4): 1773–83.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 40. Falk RJ, Terrell RS, Charles LA, *et al.*: Anti-neutrophil cytoplasmic autoantibodies induce neutrophils to degranulate and produce oxygen radicals *in vitro*. *Proc Natl Acad Sci U S A*. 1990; 87(11): 4115–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 41. Savage CO, Gaskin G, Pusey CD, *et al.*: Anti-neutrophil cytoplasm antibodies can recognize vascular endothelial cell-bound anti-neutrophil cytoplasm antibody-associated autoantigens. *Exp Nephrol*. 1993; 1(3): 190–5.
[PubMed Abstract](#)
 42. Schreiber A, Xiao H, Jennette JC, *et al.*: C5a receptor mediates neutrophil activation and ANCA-induced glomerulonephritis. *J Am Soc Nephrol*. 2009; 20(2): 289–98.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 43. Radford DJ, Luu NT, Hewins P, *et al.*: Antineutrophil cytoplasmic antibodies stabilize adhesion and promote migration of flowing neutrophils on endothelial cells. *Arthritis Rheum*. 2001; 44(12): 2851–61.
[PubMed Abstract](#) | [Publisher Full Text](#)
 44. **F** Fuchs TA, Abed U, Goosmann C, *et al.*: Novel cell death program leads to neutrophil extracellular traps. *J Cell Biol*. 2007; 176(2): 231–41.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 45. Kessenbrock K, Krumbholz M, Schönermarck U, *et al.*: Netting neutrophils in autoimmune small-vessel vasculitis. *Nat Med*. 2009; 15(6): 623–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 46. Villanueva E, Yalavarthi S, Berthier CC, *et al.*: Netting neutrophils induce endothelial damage, infiltrate tissues, and expose immunostimulatory molecules in systemic lupus erythematosus. *J Immunol*. 2011; 187(1): 538–52.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 47. **F** Wang H, Wang C, Zhao MH, *et al.*: Neutrophil extracellular traps can activate alternative complement pathways. *Clin Exp Immunol*. 2015; 181(3): 518–27.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 48. Sangaliti S, Tripodo C, Chiodoni C, *et al.*: Neutrophil extracellular traps mediate transfer of cytoplasmic neutrophil antigens to myeloid dendritic cells toward ANCA induction and associated autoimmunity. *Blood*. 2012; 120(15): 3007–18.
[PubMed Abstract](#) | [Publisher Full Text](#)
 49. Kettritz R, Jennette JC, Falk RJ: Crosslinking of ANCA-antigens stimulates superoxide release by human neutrophils. *J Am Soc Nephrol*. 1997; 8(3): 386–94.
[PubMed Abstract](#)
 50. Weidner S, Neupert W, Goppelt-Strube M, *et al.*: Antineutrophil cytoplasmic antibodies induce human monocytes to produce oxygen radicals *in vitro*. *Arthritis Rheum*. 2001; 44(7): 1698–706.
[PubMed Abstract](#) | [Publisher Full Text](#)
 51. Jennette JC, Falk RJ: ANCAs are also antimonocyte cytoplasmic autoantibodies. *Clin J Am Soc Nephrol*. 2015; 10(1): 4–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 52. Tarzi RM, Liu J, Schneider S, *et al.*: CD14 expression is increased on monocytes in patients with anti-neutrophil cytoplasm antibody (ANCA)-associated vasculitis and correlates with the expression of ANCA autoantigens. *Clin Exp Immunol*. 2015; 181(1): 65–75.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 53. Haller H, Eichhorn J, Pieper K, *et al.*: Circulating leukocyte integrin expression in Wegener's granulomatosis. *J Am Soc Nephrol*. 1996; 7(1): 40–8.
[PubMed Abstract](#)
 54. Weidner S, Carl M, Riess R, *et al.*: Histologic analysis of renal leukocyte infiltration in antineutrophil cytoplasmic antibody-associated vasculitis: importance of monocyte and neutrophil infiltration in tissue damage. *Arthritis Rheum*. 2004; 50(11): 3651–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
 55. **F** Zhao L, David MZ, Hyjek E, *et al.*: M2 macrophage infiltrates in the early stages of ANCA-associated pauci-immune necrotizing GN. *Clin J Am Soc Nephrol*. 2015; 10(1): 54–62.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 56. **F** Rousselle A, Kettritz R, Schreiber A: Monocytes Promote Crescent Formation in Anti-Myeloperoxidase Antibody-Induced Glomerulonephritis. *Am J Pathol*. 2017; 187(9): 1908–15.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 57. Xiao H, Schreiber A, Heeringa P, *et al.*: Alternative complement pathway in the pathogenesis of disease mediated by anti-neutrophil cytoplasmic autoantibodies. *Am J Pathol*. 2007; 170(1): 52–64.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 58. **F** Huigen D, van Esch A, Xiao H, *et al.*: Inhibition of complement factor C5 protects against anti-myeloperoxidase antibody-mediated glomerulonephritis in mice. *Kidney Int*. 2007; 71(7): 646–54.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 59. **F** Xiao H, Dairaghi DJ, Powers JP, *et al.*: C5a receptor (CD88) blockade protects against MPO-ANCA GN. *J Am Soc Nephrol*. 2014; 25(2): 225–31.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 60. Hao J, Wang C, Yuan J, *et al.*: A pro-inflammatory role of C5L2 in C5a-primed neutrophils for ANCA-induced activation. *PLoS One*. 2013; 8(6): e66305.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 61. Xing GQ, Chen M, Liu G, *et al.*: Complement activation is involved in renal damage in human antineutrophil cytoplasmic autoantibody associated pauci-immune vasculitis. *J Clin Immunol*. 2009; 29(3): 282–91.
[PubMed Abstract](#) | [Publisher Full Text](#)
 62. Gou SJ, Yuan J, Wang C, *et al.*: Alternative complement pathway activation products in urine and kidneys of patients with ANCA-associated GN. *Clin J Am Soc Nephrol*. 2013; 8(11): 1884–91.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 63. **F** Manenti L, Vaglio A, Gnappi E, *et al.*: Association of Serum C3 Concentration and Histologic Signs of Thrombotic Microangiopathy with Outcomes among Patients with ANCA-Associated Renal Vasculitis. *Clin J Am Soc Nephrol*. 2015; 10(12): 2143–51.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 64. Augusto JF, Langs V, Demiselle J, *et al.*: Low Serum Complement C3 Levels at Diagnosis of Renal ANCA-Associated Vasculitis Is Associated with Poor Prognosis. *PLoS One*. 2016; 11(7): e0158871.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 65. Manenti L, Urban ML, Maritati F, *et al.*: Complement blockade in ANCA-associated vasculitis: an index case, current concepts and future perspectives. *Intern Emerg Med*. 2017; 12(6): 727–31.
[PubMed Abstract](#) | [Publisher Full Text](#)
 66. **F** Jayne DRW, Bruchfeld AN, Harper L, *et al.*: Randomized Trial of C5a Receptor Inhibitor Avacopan in ANCA-Associated Vasculitis. *J Am Soc Nephrol*. 2017; 28(9): 2756–67.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 67. Popa ER, Stegeman CA, Bos NA, *et al.*: Differential B- and T-cell activation in Wegener's granulomatosis. *J Allergy Clin Immunol*. 1999; 103(5 Pt 1): 885–94.
[PubMed Abstract](#) | [Publisher Full Text](#)
 68. **F** Alberici F, Smith RM, Jones RB, *et al.*: Long-term follow-up of patients who received repeat-dose rituximab as maintenance therapy for ANCA-associated vasculitis. *Rheumatology (Oxford)*. 2015; 54(7): 1153–60.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 69. McAdoo SP, Medjeral-Thomas N, Gopaluni S, *et al.*: Long-term follow-up of a combined rituximab and cyclophosphamide regimen in renal anti-neutrophil cytoplasm antibody-associated vasculitis. *Nephrol Dial Transplant*. 2018.
[PubMed Abstract](#) | [Publisher Full Text](#)
 70. **F** Bunch DO, Mendoza CE, Aybar LT, *et al.*: Gleaning relapse risk from B cell phenotype: decreased CD5⁺ B cells portend a shorter time to relapse after B cell depletion in patients with ANCA-associated vasculitis. *Ann Rheum Dis*. 2015; 74(9): 1784–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 71. Culton DA, Nicholas MW, Bunch DO, *et al.*: Similar CD19 dysregulation in two autoantibody-associated autoimmune diseases suggests a shared mechanism

- of B-cell tolerance loss. *J Clin Immunol*. 2007; **27**(1): 53–68.
[PubMed Abstract](#) | [Publisher Full Text](#)
72. **F** Jennette JC, Falk RJ: **B cell-mediated pathogenesis of ANCA-mediated vasculitis**. *Semin Immunopathol*. 2014; **36**(3): 327–38.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
 73. Todd SK, Pepper RJ, Draibe J, *et al.*: **Regulatory B cells are numerically but not functionally deficient in anti-neutrophil cytoplasm antibody-associated vasculitis**. *Rheumatology (Oxford)*. 2014; **53**(9): 1693–703.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 74. Wilde B, Thewissen M, Damoiseaux J, *et al.*: **Regulatory B cells in ANCA-associated vasculitis**. *Ann Rheum Dis*. 2013; **72**(8): 1416–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
 75. **F** Holden NJ, Williams JM, Morgan MD, *et al.*: **ANCA-stimulated neutrophils release BlyS and promote B cell survival: a clinically relevant cellular process**. *Ann Rheum Dis*. 2011; **70**(12): 2229–33.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
 76. Sanders JS, Huitma MG, Kallenberg CG, *et al.*: **Plasma levels of soluble interleukin 2 receptor, soluble CD30, interleukin 10 and B cell activator of the tumour necrosis factor family during follow-up in vasculitis associated with proteinase 3-antineutrophil cytoplasmic antibodies: associations with disease activity and relapse**. *Ann Rheum Dis*. 2006; **65**(11): 1484–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 77. Krumbholz M, Specks U, Wick M, *et al.*: **BAFF is elevated in serum of patients with Wegener's granulomatosis**. *J Autoimmun*. 2005; **25**(4): 298–302.
[PubMed Abstract](#) | [Publisher Full Text](#)
 78. Pollard RP, Abdulahad WH, Vissink A, *et al.*: **Serum levels of BAFF, but not APRIL, are increased after rituximab treatment in patients with primary Sjogren's syndrome: data from a placebo-controlled clinical trial**. *Ann Rheum Dis*. 2013; **72**(1): 146–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
 79. **F** Alberici F, Smith RM, Fonseca M, *et al.*: **Association of a *TNFSF13B* (BAFF) regulatory region single nucleotide polymorphism with response to rituximab in antineutrophil cytoplasmic antibody-associated vasculitis**. *J Allergy Clin Immunol*. 2017; **139**(5): 1684–1687.e10.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
 80. Jayne D, Blockmans D, Luqmani R, *et al.*: **Efficacy and Safety of Belimumab in Combination with Azathioprine for Remission Maintenance in Granulomatosis with Polyangiitis and Microscopic Polyangiitis: A Multicenter Randomized, Placebo-Controlled Study (abstract)**. *Arthritis Rheumatol*. 2017; **69**(suppl 10).
[Reference Source](#)
 81. Morgan MD, Day CJ, Piper KP, *et al.*: **Patients with Wegener's granulomatosis demonstrate a relative deficiency and functional impairment of T-regulatory cells**. *Immunology*. 2010; **130**(1): 64–73.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 82. Free ME, Bunch DO, McGregor JA, *et al.*: **Patients with antineutrophil cytoplasmic antibody-associated vasculitis have defective Treg cell function exacerbated by the presence of a suppression-resistant effector cell population**. *Arthritis Rheum*. 2013; **65**(7): 1922–33.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 83. Schmitt WH, Hagen EC, Neumann I, *et al.*: **Treatment of refractory Wegener's granulomatosis with antithymocyte globulin (ATG): an open study in 15 patients**. *Kidney Int*. 2004; **65**(4): 1440–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
 84. Schönermarck U, Csernok E, Trabandt A, *et al.*: **Circulating cytokines and soluble CD23, CD26 and CD30 in ANCA-associated vasculitides**. *Clin Exp Rheumatol*. 2000; **18**(4): 457–63.
[PubMed Abstract](#)
 85. Gan PY, Holdsworth SR, Kitching AR, *et al.*: **Myeloperoxidase (MPO)-specific CD4⁺ T cells contribute to MPO-anti-neutrophil cytoplasmic antibody (ANCA) associated glomerulonephritis**. *Cell Immunol*. 2013; **282**(1): 21–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
 86. Ruth A, Kitching AR, Kwan RY, *et al.*: **Anti-neutrophil cytoplasmic antibodies and effector CD4⁺ cells play nonredundant roles in anti-myeloperoxidase crescentic glomerulonephritis**. *J Am Soc Nephrol*. 2006; **17**(7): 1940–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
 87. **F** Chang J, Eggenhuizen P, O'Sullivan KM, *et al.*: **CD8⁺ T Cells Effect Glomerular Injury in Experimental Anti-Myeloperoxidase GN**. *J Am Soc Nephrol*. 2017; **28**(1): 47–55.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
 88. Ooi JD, Chang J, Hickey MJ, *et al.*: **The immunodominant myeloperoxidase T-cell epitope induces local cell-mediated injury in antimyeloperoxidase glomerulonephritis**. *Proc Natl Acad Sci U S A*. 2012; **109**(39): E2615–24.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 89. Nogueira E, Hamour S, Sawant D, *et al.*: **Serum IL-17 and IL-23 levels and autoantigen-specific Th17 cells are elevated in patients with ANCA-associated vasculitis**. *Nephrol Dial Transplant*. 2010; **25**(7): 2209–17.
[PubMed Abstract](#) | [Publisher Full Text](#)
 90. Hoshino A, Nagao T, Nagi-Miura N, *et al.*: **MPO-ANCA induces IL-17 production by activated neutrophils *in vitro* via classical complement pathway-dependent manner**. *J Autoimmun*. 2008; **31**(1): 79–89.
[PubMed Abstract](#) | [Publisher Full Text](#)
 91. Gan PY, Steinmetz OM, Tan DS, *et al.*: **Th17 cells promote autoimmune anti-myeloperoxidase glomerulonephritis**. *J Am Soc Nephrol*. 2010; **21**(6): 925–31.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 92. Mueller A, Holl-Ulrich K, Gross WL: **Granuloma in ANCA-associated vasculitides: another reason to distinguish between syndromes?** *Curr Rheumatol Rep*. 2013; **15**(11): 376.
[PubMed Abstract](#) | [Publisher Full Text](#)
 93. Schönermarck U, Csernok E, Gross WL: **Pathogenesis of anti-neutrophil cytoplasmic antibody-associated vasculitis: challenges and solutions 2014**. *Nephrol Dial Transplant*. 2015; **30** Suppl 1: i46–52.
[PubMed Abstract](#) | [Publisher Full Text](#)
 94. Csernok E, Trabandt A, Müller A, *et al.*: **Cytokine profiles in Wegener's granulomatosis: predominance of type 1 (Th1) in the granulomatous inflammation**. *Arthritis Rheum*. 1999; **42**(4): 742–50.
[PubMed Abstract](#) | [Publisher Full Text](#)
 95. Zhao Y, Odell E, Choong LM, *et al.*: **Granulomatosis with polyangiitis involves sustained mucosal inflammation that is rich in B-cell survival factors and autoantigen**. *Rheumatology (Oxford)*. 2012; **51**(9): 1580–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
 96. Kesel N, Köhler D, Herich L, *et al.*: **Cartilage destruction in granulomatosis with polyangiitis (Wegener's granulomatosis) is mediated by human fibroblasts after transplantation into immunodeficient mice**. *Am J Pathol*. 2012; **180**(5): 2144–55.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. **Augusto Vaglio**

Nephrology Unit, University Hospital of Parma, Parma, Italy

Competing Interests: No competing interests were disclosed.

2. **Minghui Zhao**

Institute of Nephrology, Peking University First Hospital, Beijing, China

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research