REVIEW

Recent advances in understanding bile duct remodeling and fibrosis [version 1; peer review: 2 approved]

Marinda Scrushy¹, April O'Brien^{1,2}, Shannon Glaser¹⁻³

¹Texas A&M University College of Medicine, Temple, TX, USA

²Central Texas Veterans Research Center, Temple, TX, USA

³Digestive Diseases Research Center, Baylor Scott & White Health, Temple, TX, USA

V1 First published: 31 Jul 2018, 7(F1000 Faculty Rev):1165
<https://doi.org/10.12688/f1000research.14578.1>

Latest published: 31 Jul 2018, 7(F1000 Faculty Rev):1165
<https://doi.org/10.12688/f1000research.14578.1>

Abstract

Cholestatic liver disease encompasses a detrimental group of diseases that are non-discriminatory in nature. These diseases occur over every age range from infancy (biliary atresia) to geriatrics (hepatitis). They also cover both genders in the form of primary sclerosing cholangitis in men and primary biliary cholangitis in women. Oftentimes, owing to the disease progression and extensive scarring, the treatment of last resort becomes a liver transplant. In this review, we will briefly discuss and explore new avenues of understanding in the progression of cholestatic liver disease and possible therapeutic targets for intervention. The greater our understanding into the idiopathic nature of cholestatic liver disease, the better our chances of discovering treatment options to halt or reverse the progression, reducing or eliminating the need for expensive and risky transplants.

Keywords

cholestasis, cholangiocytes, ductular reaction, bile duct

Open Peer Review

Approval Status

	1	2
version 1 31 Jul 2018

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Tohru Itoh**, Institute for Quantitative Biosciences, The University of Tokyo, Tokyo, Japan
2. **Leo van Grunsven**, Vrije Universiteit Brussel, Brussels, Belgium

Any comments on the article can be found at the end of the article.

Corresponding author: Shannon Glaser (sglaser@medicine.tamhsc.edu)

Author roles: **Scrushy M:** Writing – Original Draft Preparation, Writing – Review & Editing; **O'Brien A:** Writing – Original Draft Preparation, Writing – Review & Editing; **Glaser S:** Conceptualization, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This material is the result of work supported by resources at the Central Texas Veterans Health Care System. Portions of this work was supported by a VA Merit Award (5I01BX002192) and National Institutes of Health R01 Grant (DK110035) to Dr. Glaser.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2018 Scrushy M *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Scrushy M, O'Brien A and Glaser S. **Recent advances in understanding bile duct remodeling and fibrosis [version 1; peer review: 2 approved]** F1000Research 2018, 7(F1000 Faculty Rev):1165 <https://doi.org/10.12688/f1000research.14578.1>

First published: 31 Jul 2018, 7(F1000 Faculty Rev):1165 <https://doi.org/10.12688/f1000research.14578.1>

Introduction

The hepatobiliary system, a crucial part of the digestive system, is made up of the liver, gall bladder, and bile ducts, a complex network of tubular intra- and extra-hepatic ducts that converge to drain bile into the duodenum to aid in fat digestion and the transportation of wastes¹. The bile ducts are lined by a subset of liver epithelial cells termed cholangiocytes. Cholangiocytes are responsible for the modification of bile fluidity and alkalinity after secretion of the major bile osmolytes and various constituents by the hepatocyte¹⁻³. These ducts, or channels, are part of a larger three-dimensional structure known as the biliary tree, which branches into the surrounding liver parenchyma². There is a wide array of disease states specifically targeting the cholangiocytes, termed cholangiopathies, many of which are induced by cholestasis. Cholestasis is defined as a bile flow decrease caused by either defective hepatocyte secretions or the physical obstruction of bile through the intra- and extra-hepatic bile ducts⁴.

Chronic cholestasis causes destruction of the bile ducts, strictures, and fibrosis. Chronic fibrosis and inflammation can progress to permanent scarring leading to hepatic failure, necessitating the need for a liver transplant. Mouse models of cholestasis include the mechanical model of bile duct ligation (BDL), as well as the 3,5-diethoxycarbonyl-1,4-dihydrocollidine (DDC) diet and the *Mdr2* or *abcb4* knockout mouse. In the BDL mouse model, a surgical incision is made in the abdomen, and after careful manipulation of the liver, the bile duct is exposed and isolated from the adjoining hepatic artery and portal vein, allowing it to be tied off by surgical sutures. The incision is then closed and cholestasis is allowed to develop⁵. As a consequence of BDL, adaptive remodeling of bile ducts as well as an increase in ductular tissue density has been observed^{2,6}. The DDC model was initially used to reproduce steatohepatitis but also has been shown to reproduce conditions observed in chronic cholestatic diseases such as ductular proliferation as well as ductal and liver fibrosis⁷. Alternatively, the *Mdr2* *abc4*^{-/-} knockout mice mimic cholestasis by inducing a deficit in canalicular phospholipid flippase leading to the absence of phospholipids in bile. This causes bile leakage that has been shown to lead to bile duct and portal fibrosis⁸.

This response is coined the “ductular reaction” and takes place after a cholestatic-inducing injury causes the normally quiescent cholangiocytes to take on a reactive phenotype. This involves an epithelial-to-mesenchymal transition (EMT), cholangiocyte proliferation, and a change to a neuroendocrine phenotype where cholangiocytes begin to secrete and respond to a number of peptides and hormones. This reactive phenotype brings about changes in the extra-cellular matrix (ECM) that leads to the collapse of the bile duct architecture with eventual fibrosis and strictures⁹. It is hypothesized that the proliferation and other morphologic changes observed after BDL are the liver’s attempt to increase both the surface area and function of cholangiocytes⁶. The aim of this review is to summarize current research on the cholangiocyte’s response to cholestatic injury, focusing on the role of proliferation as well as autocrine and paracrine signaling molecules and what role these play in the overall change in bile duct morphology.

Discussion of recent literature

The remodeling of the bile ducts in response to injury results in an increase in elongation, diameter, and branching of the ducts as well as increased intra-ductal mass^{2,6}. As cholestasis is defined as the buildup of bile in the ductular system, it can be hypothesized that this remodeling serves to find alternative routes to release the bile. This would function not only to reduce its toxic effects on resident cholangiocytes but also to continue its contribution to overall organismal homeostasis through the breakdown of fats for digestion^{4,6,10}. Using three-dimensional confocal imaging and reconstruction techniques, Vartak *et al.* described the bile duct changes they saw in BDL mice, demonstrating that the biliary tree transforms by increasing proliferation, branching, and corrugation. These changes led to an overall increase in surface area⁶. This increase in surface area is significant as cholangiocytes function via secretion and re-absorption of electrolytes to form and modify bile, a mechanism that requires significant surface area^{1,3}.

One of the key mechanisms involved in bile duct remodeling is cholangiocyte proliferation. As cholangiocytes are not normally mitotically active, this switch is seen as a response to insult¹¹. Numerous studies have outlined that various physiologic chemicals, including gastrointestinal hormones, angiogenic factors, sex hormones, neuropeptides and neurotransmitters, cytokines, and growth factors, play a role in initiating or attenuating biliary proliferation^{12,13}. One of the main mechanisms by which proliferation is increased or attenuated in response to these chemicals involves the modulation of cAMP levels, as large cholangiocytes proliferate through a cAMP-dependent mechanism¹².

Another possible key role of proliferation is that encroaching ductules deliver hepatic progenitors to the liver parenchyma in order to facilitate hepatocyte regeneration². These liver progenitor cells express both hepatic and biliary markers and have been shown to be involved in hepatic repair after chronic liver injury. Many growth factors and cytokines, including fibroblast growth factor 7 (FGF7), tumor necrosis factor-like weak induce of apoptosis (TWEAK), hepatocyte growth factor, and the Wnt/B-catenin pathway, have been shown to regulate liver progenitor cell function¹⁴⁻¹⁷.

FGF7 is a fibroblast growth factor that has been shown to attenuate liver markers of cholestasis when overexpressed in the DDC mouse model. Takase *et al.* also demonstrated that it was capable of reversing chronic liver injury induced in the DDC model. They increased FGF7 expression after three weeks of liver injury and again found decreased cholestatic markers¹⁶. Both FGF7 and TWEAK induce liver progenitor cell proliferation, and FGF7 plays a role in adaptive remodeling by increasing bile duct density^{2,15}. The effects of hepatic growth factor are mediated by the tyrosine kinase receptor c-Met. Studies in a c-Met knockout mouse showed that there was an increase in bile acid production leading to increased cholestasis and liver injury. Furthermore, mice lacking c-Met had a decreased proliferative capacity of ductal epithelial cells and thus liver regeneration¹⁷. Wnt signaling functions through a B-catenin-dependent pathway and non-B-catenin-dependent pathway. Okabe *et al.* demonstrated

that, using the DDC mouse model, Wnt signaling via the B-catenin signaling pathway not only increased biliary proliferation but may play a potential role in hepatocyte-to-biliary trans-differentiation¹⁴.

Another increasing interest in the story of ductular remodeling is the hepatocyte-to-cholangiocyte trans-differentiation, or metaplasia. Yanger *et al.* demonstrated through the Notch signaling pathway a transition of hepatocyte to biliary epithelial cells that share morphological, molecular, and structural similarities with their native counterparts¹⁸. Additional studies have suggested a role for this metaplasia as a means of escaping injury and expanding the bile ducts while retaining the ability to revert back to their native phenotype. Owing in part to the large percentage of the cellular population comprised by hepatocytes, it has been suggested that this plasticity provides a protective element in times of stress where hepatocytes are the primary targets (that is, hepatitis virus). Furthermore, it has been previously demonstrated that human hepatocytes *in vivo* appear phenotypically plastic following liver injury¹⁹.

Recent research has focused on the role that microRNAs (miRNAs) may play in inducing the proliferative phenotype of the injured cholangiocyte. miRNAs are non-coding RNAs that function to post-transcriptionally silence gene expression by binding to the untranslated region of the mRNA, causing destabilization and the prevention of protein expression²⁰. Research has started to focus on miRNA-21 because of its upregulation in multiple cancer cell types²¹. miRNA-21 has been found to be upregulated in the livers and bile ducts of BDL mice as well as human primary sclerosing cholangitis (PSC) liver samples. miRNA-21 also promotes proliferation in BDL mice and decreases apoptosis. Additionally, *in vitro* studies have demonstrated that it promotes fibrosis through the activation of hepatic stellate cells via a Smad-7 inhibitory mechanism¹¹.

Changes in ECM proteins, such as integrins, also play a significant role in proliferation. Integrins are a specific type of trans-membrane cellular matrix protein that functions to anchor intra-cellular components to the ECM, attaching to specific cytoskeletal molecules such as laminin and fibronectin^{20,22}. The structural nature of integrins, when bound, allows for the initiation of signaling cascades that regulate cellular functions such as proliferation and morphological change²⁰. One specific integrin, $\alpha\text{v}\beta\text{6}$, has been found to be upregulated in injured epithelial cells and plays a role in initiating fibrosis by activating transforming growth factor beta 1 (TGF- β1)²². Connective tissue growth factor (CTGF) is another important ECM component of which cholangiocytes are among the primary sources. CTGF and integrin $\alpha\text{v}\beta\text{6}$ work together to modulate cellular matrix interactions and activate key players in the ductular reaction and fibrosis²³.

The initiation of ductular reactions is characterized by cholangiocyte changes at the molecular level as well. A characteristic of reactive cholangiocytes is a change to a neuroendocrine phenotype as evidenced by their expression of neuroendocrine

markers such as chromogranin A and N-CAM²⁴. This phenotypic switch is characterized by increases in both secretion and responsiveness to certain neurohormones and neuropeptides. This induces a critical autocrine and paracrine signaling network that contributes to overall bile duct remodeling by tipping the balance between proliferation and bile duct loss as well as increasing fibrosis²⁵. Recent research has shown that neuropeptide Y (NPY) plays an important role in this process. Cholangiocytes express NPY receptors as well as secrete and respond to NPY during cholestasis with increasing proliferation²⁶. Other neuroendocrine players that upregulate cholangiocyte proliferation include substance P, galanin, gonadotropin-releasing hormone (GnRH), and insulin-like growth factor 1 (IGF1)^{12,27,28}. Alternatively, melatonin and serotonin have also been shown to attenuate the ductular response^{24,29}.

Galanin has recently been shown to be upregulated in the biliary system of mice that have undergone BDL. Furthermore, when galanin was administered to BDL rats, cholangiocyte proliferation was increased because of binding of the galanin receptor 1 with subsequent increases in cAMP²⁸. New research has also outlined a role for IGF1 in the reactive nature of cholangiocytes. Cholangiocytes have previously been shown to express the IGF1 receptor. A recent study demonstrated that in response to increased IGF1 in a rat model of cholestatic liver disease, the cholangiocytes increased both their proliferation and ECM deposition. These upregulated processes led to an increase in fibrosis, a key step in bile duct modulation in cholangiopathies. This ductular reaction may play a role in the activation of surrounding fibroblasts, which also respond with an increase in ECM³⁰.

Another key proliferative player is GnRH, which increases cholangiocyte proliferation³¹. Additionally, melatonin has been shown to attenuate cholangiocyte proliferation and recent research has investigated how these two neurohormones interact to alter the cholangiocyte response^{25,29,32}. Cholangiocytes express GnRH receptors 1 and 2, respectively, and through the action of GnRH, cholangiocyte proliferation has been shown to increase via a cAMP-dependent mechanism. This mechanism takes place primarily through the action of the GnRH receptor 1³¹. Melatonin has a regulatory function on GnRH secretion, attenuating its proliferative effects by decreasing its expression in cholangiocytes²⁵.

An important consequence of chronic cholestasis is bile duct fibrosis. Key mediators of the ductular reaction, such as neuropeptides and neurohormones, that upregulate cholangiocyte proliferation also play a role in upregulating bile duct fibrosis³³. Cholangiocytes play a direct key role by activating and secreting pro-fibrotic cytokines. Furthermore, cholangiocytes themselves can produce ECM components, such as collagen I and III³⁰. The EMT is a key event, triggered by injury, where cholangiocytes revert to a mesenchymal phenotype, allowing them to play a key role in fibrosis³⁰. EMT involves functional and morphological changes such as loss of cell-to-cell adhesion properties and apical-basal polarity³⁴. This transition is

considered one of the first steps in developing the “reactive” cholangiocyte phenotype. Although cholangiocytes have been shown to express some markers of EMT, this remains an area of controversy as recent lineage-tracing studies have demonstrated no concrete evidence of this change³⁵.

Conclusions

In summary, research has made great advances in the field of cholestatic liver injury. These advances are a result of both hard work and dedication from the liver research community as a whole with new advances in the understanding of the roles played by the cholangiocyte during injury and repair through subsequent signaling molecules and their mechanisms. These mechanisms, such as EMT, the role of various hormones in the pro-fibrogenic phenotype, and the role of cAMP in proliferation,

provide valuable insight and potential therapeutic targets in the fight against liver disease.

Competing interests

The authors declare that they have no competing interests.

Grant information

This material is the result of work supported by resources at the Central Texas Veterans Health Care System. Portions of this work was supported by a VA Merit Award (5I01BX002192) and National Institutes of Health R01 Grant (DK110035) to Dr. Glaser.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- Yoo KS, Lim WT, Choi HS: **Biology of Cholangiocytes: From Bench to Bedside.** *Gut Liver.* 2016; **10**(5): 687–98.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Kaneko K, Kamimoto K, Miyajima A, et al.: **Adaptive remodeling of the biliary architecture underlies liver homeostasis.** *Hepatology.* 2015; **61**(6): 2056–66.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Strazzabosco M: **Transport systems in cholangiocytes: their role in bile formation and cholestasis.** *Yale J Biol Med.* 1997; **70**(4): 427–34.
[PubMed Abstract](#) | [Free Full Text](#)
- Kumar V, AA, Aster J: **Liver and Gallbladder.** In *Robbins and Cotran Pathologic Basis of Disease* E. Saunders, Editor. 2015; 821–881.
- Tag CG, Sauer-Lehnen S, Weiskirchen S, et al.: **Bile duct ligation in mice: induction of inflammatory liver injury and fibrosis by obstructive cholestasis.** *J Vis Exp.* 2015; (96): e52438.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Vartak N, Damle-Vartak A, Richter B, et al.: **Cholestasis-induced adaptive remodeling of interlobular bile ducts.** *Hepatology.* 2016; **63**(3): 951–64.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Fickert P, Stöger U, Fuchsichler A, et al.: **A new xenobiotic-induced mouse model of sclerosing cholangitis and biliary fibrosis.** *Am J Pathol.* 2007; **171**(2): 525–36.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Popov Y, Patsenker E, Fickert P, et al.: **Mdr2 (Abcb4)-/- mice spontaneously develop severe biliary fibrosis via massive dysregulation of pro- and antifibrogenic genes.** *J Hepatol.* 2005; **43**(6): 1045–54.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Jirouskova M, Nepomucka K, Oyman-Eyrlimez G, et al.: **Plectin controls biliary tree architecture and stability in cholestasis.** *J Hepatol.* 2018; **68**(5): 1006–17.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Xia X, Francis H, Glaser S, et al.: **Bile acid interactions with cholangiocytes.** *World J Gastroenterol.* 2006; **12**(22): 3553–63.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kennedy LL, Meng F, Venter JK, et al.: **Knockout of microRNA-21 reduces biliary hyperplasia and liver fibrosis in cholestatic bile duct ligated mice.** *Lab Invest.* 2016; **96**(12): 1256–67.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Hall C, Sato K, Wu N, et al.: **Regulators of Cholangiocyte Proliferation.** *Gene Expr.* 2017; **17**(2): 155–71.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Jensen K, Marzioni M, Munshi K, et al.: **Autocrine regulation of biliary pathology by activated cholangiocytes.** *Am J Physiol Gastrointest Liver Physiol.* 2012; **302**(5): G473–83.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Okabe H, Yang J, Sylakowski K, et al.: **Wnt signaling regulates hepatobiliary repair following cholestatic liver injury in mice.** *Hepatology.* 2016; **64**(5): 1652–66.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Jakubowski A, Ambrose C, Parr M, et al.: **TWEAK induces liver progenitor cell proliferation.** *J Clin Invest.* 2005; **115**(9): 2330–40.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Takase HM, Itoh T, Ino S, et al.: **FGF7 is a functional niche signal required for stimulation of adult liver progenitor cells that support liver regeneration.** *Genes Dev.* 2013; **27**(2): 169–81.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ishikawa T, Factor VM, Marquardt JU, et al.: **Hepatocyte growth factor/c-met signaling is required for stem-cell-mediated liver regeneration in mice.** *Hepatology.* 2012; **55**(4): 1215–26.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Yanger K, Zong Y, Maggs LR, et al.: **Robust cellular reprogramming occurs spontaneously during liver regeneration.** *Genes Dev.* 2013; **27**(7): 719–24.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Tarlow BD, Pelz C, Naugler WE, et al.: **Bipotent adult liver progenitors are derived from chronically injured mature hepatocytes.** *Cell Stem Cell.* 2014; **15**(5): 605–18.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Mitchell RN: **The Cell as a Unit of Health and Disease.** In *Robbins and Cotran Pathologic Basis of Disease.* Elsevier. 2015; 1–29.
- Chen J, Wang X: **MicroRNA-21 in breast cancer: diagnostic and prognostic potential.** *Clin Transl Oncol.* 2014; **16**(3): 225–33.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Peng ZW, Ikenaga N, Liu SB, et al.: **Integrin α v β 6 critically regulates hepatic progenitor cell function and promotes ductular reaction, fibrosis, and tumorigenesis.** *Hepatology.* 2016; **63**(1): 217–32.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Pi L, Robinson PM, Jorgensen M, et al.: **Connective tissue growth factor and integrin α v β 6: a new pair of regulators critical for ductular reaction and biliary fibrosis in mice.** *Hepatology.* 2015; **61**(2): 678–91.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Omenetti A, Yang L, Gainetdinov RR, et al.: **Paracrine modulation of cholangiocyte serotonin synthesis orchestrates biliary remodeling in adults.** *Am J Physiol Gastrointest Liver Physiol.* 2011; **300**(2): G303–15.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- McMillin M, DeMorrow S, Glaser S, et al.: **Melatonin inhibits hypothalamic gonadotropin-releasing hormone release and reduces biliary hyperplasia and fibrosis in cholestatic rats.** *Am J Physiol Gastrointest Liver Physiol.* 2017; **313**(5): G410–G418.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- DeMorrow S, Meng F, Venter J, et al.: **Neuropeptide Y inhibits biliary hyperplasia of cholestatic rats by paracrine and autocrine mechanisms.** *Am J Physiol Gastrointest Liver Physiol.* 2013; **305**(3): G250–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Meng F, DeMorrow S, Venter J, et al.: **Overexpression of membrane**

- metalloendopeptidase inhibits substance P stimulation of cholangiocarcinoma growth. *Am J Physiol Gastrointest Liver Physiol.* 2014; **306**(9): G759–68.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
28. **F** McMillin M, Frampton G, Grant S, *et al.*: **The Neuropeptide Galanin Is Up-Regulated during Cholestasis and Contributes to Cholangiocyte Proliferation.** *Am J Pathol.* 2017; **187**(4): 819–30.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 29. Renzi A, Glaser S, DeMorrow S, *et al.*: **Melatonin inhibits cholangiocyte hyperplasia in cholestatic rats by interaction with MT1 but not MT2 melatonin receptors.** *Am J Physiol Gastrointest Liver Physiol.* 2011; **301**(4): G634–43.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 30. Sokolović A, Rodríguez-Ortígoza CM, Bloemendaal LT, *et al.*: **Insulin-like growth factor 1 enhances bile-duct proliferation and fibrosis in Abcb4(-/-) mice.** *Biochim Biophys Acta.* 2013; **1832**(6): 697–704.
[PubMed Abstract](#) | [Publisher Full Text](#)
 31. Ray D, Han Y, Franchitto A, *et al.*: **Gonadotropin-releasing hormone stimulates biliary proliferation by paracrine/autocrine mechanisms.** *Am J Pathol.* 2015; **185**(4): 1061–72.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 32. Han Y, Onori P, Meng F, *et al.*: **Prolonged exposure of cholestatic rats to complete dark inhibits biliary hyperplasia and liver fibrosis.** *Am J Physiol Gastrointest Liver Physiol.* 2014; **307**(9): G894–904.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 33. Gouw AS, Clouston AD, Theise ND: **Ductular reactions in human liver: diversity at the interface.** *Hepatology.* 2011; **54**(5): 1853–63.
[PubMed Abstract](#) | [Publisher Full Text](#)
 34. **F** Fabris L, Brivio S, Cadamuro M, *et al.*: **Revisiting Epithelial-to-Mesenchymal Transition in Liver Fibrosis: Clues for a Better Understanding of the “Reactive” Biliary Epithelial Phenotype.** *Stem Cells Int.* 2016; **2016**: 2953727.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 35. **F** Chu AS, Diaz R, Hui J, *et al.*: **Lineage tracing demonstrates no evidence of cholangiocyte epithelial-to-mesenchymal transition in murine models of hepatic fibrosis.** *Hepatology.* 2011; **53**(5): 1685–95.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. **Leo van Grunsven**

Liver Cell Biology Laboratory, Vrije Universiteit Brussel, Brussels, Belgium

Competing Interests: No competing interests were disclosed.

2. **Tohru Itoh**

Laboratory of Stem Cell Therapy, Institute for Quantitative Biosciences, The University of Tokyo, Tokyo, Japan

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research