Check for updates

RESEARCH ARTICLE

REVISED *Aloe barbadensis* Miller leaf exudate is a potential treatment for bovine mastitis [version 2; peer review: 1 approved, 1 not approved]

Samira de Aquino Leite Fiordalisi¹, Luciana Aparecida Honorato², Shirley Kuhnen³

¹Programa de Pós-Graduação em Agroecossistemas, Universidade Federal de Santa Catarina, Florianópolis, Santa Catarina, 88034-000, Brazil

²Centro de Ciências Biológicas, Universidade Federal de Santa Catarina, Florianópolis, Santa Catarina, 88034-000, Brazil

³Laboratório de Morfofisiologia e Bioquímica Animal - LABIMA, Departamento de Zootecnia e Desenvolvimento Rural, Universidade Federal de Santa Catarina, Florianópolis, Santa Catarina, 88034-000, Brazil

V2 First published: 14 Aug 2018, 7:1285
<https://doi.org/10.12688/f1000research.15671.1>
 Latest published: 26 Oct 2018, 7:1285
<https://doi.org/10.12688/f1000research.15671.2>

Abstract

Background: *Aloe barbadensis* Miller, also known as *Aloe vera* is a well-known phytotherapeutic, and parts of its leaves are used for a wide range of medicinal purposes. This study seeks to assess the *in vitro* antimicrobial and cytotoxic effects of leaf exudate (LE) from *A. barbadensis* leaves against *Staphylococcus aureus* and MAC-T bovine mammary epithelial cells.

Methods: Seasonal LE samples were collected, and the effect on total phenolic and aloin contents was determined. Antimicrobial activity of LE was evaluated using the broth microdilution method, and toxicity to MAC-T cells was determined by MTT assay.

Results: Samples collected during different seasons of the year showed a seasonal effect on the chemical profile of LE ($P < 0.05$). However, despite these chemical variations, we found no differences in antimicrobial activity against *S. aureus*. For all studied samples, the minimum inhibitory concentration (MIC) was 1,000 µg/mL. Furthermore, we found an elevated cytotoxic effect of LE on MAC-T cells with a significant reduction in cellular viability at 7.8 µg/mL ($P < 0.05$) and an IC₅₀ of 91.89 µg/mL.

Conclusions: Despite the antimicrobial effects of LE, the high toxicity for MAC-T cells suggests that it is unsuitable for intramammary use, but does have potential as a topical antimicrobial.

Keywords

phytotherapy, *Staphylococcus aureus*, MAC-T cells

Open Peer Review

Approval Status

	1	2
version 2		
(revision)
26 Oct 2018	view	view

version 1		
14 Aug 2018
	view	view

1. **Gaspar Diaz-Muñoz** , Federal University of Minas Gerais, Belo Horizonte, Brazil

2. **Liliana Cardemil** , Universidad de Chile, Santiago, Chile

Carlos Salinas, Universidad de Chile, Santiago, Chile

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Shirley Kuhnen (shirley.kuhnen@ufsc.br)

Author roles: **Fiordalisi SdAL:** Conceptualization, Data Curation, Investigation, Methodology, Writing – Original Draft Preparation;

Honorato LA: Conceptualization, Investigation, Methodology, Writing – Original Draft Preparation; **Kuhnen S:** Conceptualization, Funding Acquisition, Investigation, Methodology, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by the CNPq (National Council on Scientific and Technological Development; Brasilia, Brazil) (403415/2013-6, Edital 39/2013).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2018 Fiordalisi SdAL *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. Data associated with the article are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

How to cite this article: Fiordalisi SdAL, Honorato LA and Kuhnen S. *Aloe barbadensis* Miller leaf exudate is a potential treatment for bovine mastitis [version 2; peer review: 1 approved, 1 not approved] F1000Research 2018, 7:1285

<https://doi.org/10.12688/f1000research.15671.2>

First published: 14 Aug 2018, 7:1285 <https://doi.org/10.12688/f1000research.15671.1>

REVISED Amendments from Version 1

We provide a new [Figure 1](#) which shows the chemical structure of the aloin. We have corrected typos throughout the paper.

See referee reports

Figure 1. Chemical structure of the Aloin A and Aloin B.

Introduction

Bovine mastitis, which is characterized by inflammation of the mammary gland, is the most frequent infection found in dairy herds worldwide¹. The treatment recommended for mastitis is the administration of intramammary antimicrobials. However, control of infections caused by *Staphylococcus aureus*, the principal etiological agent of bovine mastitis², is very difficult. In addition to inactivating several antimicrobials, this microorganism can also survive in the intracellular environment after phagocytosis. As a consequence, the cure rate of mastitis caused by *S. aureus* is low, with a high incidence of recurrence³. As such, interest in the search for methods of control and prevention has increased, including the identification of new antimicrobials^{4–7}.

In vivo methods are still commonly used to study bovine mastitis, but *in vitro* testing has been recommended⁸. Based on *in vitro* models, studies have produced a wide range of results, from identifying the prevalence of etiological agents of mastitis to evaluating the direct effects of products on the susceptibility of studied microorganisms^{9,10}. Among these, *in vitro* tests on antimicrobials are some of the most widely used⁸. *In vitro* studies with bovine mammary gland explants or mammary epithelial cells (MEC) are commonly used to assess the different functions of mammary glands, such as the response to initial infection^{11,12}. Recently, primary cultures of mammary explants and MECs were also suggested as adequate models in the search for new therapeutic agents^{5,13}. In the case of mastitis, such *in vitro* methods can help evaluate the toxicity of antimicrobials, enabling the determination of safe doses and minimizing the potential risks during *in vivo* validation.

Aloe vera (*Aloe barbadensis* Miller) is a plant widely used and recognized for its antimicrobial, anti-inflammatory, wound-healing, antitumor, and antioxidant pharmacological properties¹⁴. Yet, until now, few studies have reported on its potential as a treatment for bovine mastitis. Most research on the pharmaceutical potential of *Aloe vera* has studied the mucilaginous gel, commonly known as aloe vera gel, that is rich in complex carbohydrates, particularly, acemannans^{15,16}. However, along with the gel, a yellow exudate with a strong odor and bitter taste, known as leaf exudate (LE), can also be extracted from the leaves¹⁷. Its release occurs as soon as the leaves are cut and it can be found within the phloem vessels^{18,19}. Despite being composed of large amounts of 1,8-dihydroxyanthraquinone derivatives and their glycosides, the industry that uses aloe vera gel as a raw material considers LE a residue. Among the anthraquinones found in LE, the major compounds are a mixture of two readily oxidizable diastereomers, aloin A and aloin B ([Figure 1](#)),

which are sometimes undesirable because of their toxic and cathartic potential. However, these compounds may be of therapeutic interest in the control of antimicrobial and tumor cell proliferation^{20,21}.

Thus, the current study seeks to investigate the potential of LE from *Aloe vera* leaves in the control of bovine mastitis through *in vitro* models that evaluate the antimicrobial effects against *S. aureus* and cytotoxicity to MAC-T cells.

Methods

Sampling and extraction of leaf exudate

A total of 30 plant samples were collected from 3-year-old *Aloe vera* (*Aloe barbadensis*) at random from a commercial grower (Naturama Sucos Integraais do Brasil Ltda®; Paulo Lopes, SC, Brazil) in March, June, September and December of 2015, and one leaf was taken from the mid-position of each plant. Thus, 30 leaves in total were collected for each month, representing each season. Leaves were cut at the base and maintained vertically for 3 h in a beaker to collect the LE at room temperature. Subsequently, the LE was lyophilized and stored at -20 °C. LEs of six plants were combined for a total of five repetitions for each season of the year.

Chemical profile of leaf exudate

Total phenolics. The total phenolic content was determined using the colorimetric method of Folin-Ciocalteu¹⁴ and an external standard curve of gallic acid (10–100 µg/mL) ($y = 0.0197x$ / $r^2 = 0.987$). The results were expressed in µg of gallic acid equivalents (GAE)/mg of extract (µg of GAE/mg). All tests were performed in triplicate.

Aloin. The aloin content in LE was obtained on an UHPLC Thermo Scientific UltiMate 3000 RS Dual System (Thermo Fisher Scientific, San Jose, CA), using a Thermo Scientific C18 reverse-phase column (4.6 × 250 mm; 5 µm; 120Å (Acclaim™120, Thermo Scientific®) at 40°C, operating at 240, 260, 280, and 320 nm. The mobile phase was eluted at 1 mL/min flow rate, using a methanol/water (70/30) mixture²². The identification of aloin was based on a comparison of the chromatographic profile and retention time with the commercial standard (Sigma-Aldrich, St. Louis, MO, USA/ B6906). After the addition of the standard, samples were co-chromatographed to confirm identification of the compound. Aloin content was determined through an external standard curve of barbaloin ($y = 302.73x$ / $r^2 = 0.9822$) and the result expressed in µg of aloin per mg of the sample (µg/mg).

Antimicrobial activity

Antimicrobial activity was evaluated using a broth microdilution method according to the Clinical and Laboratory Standards Institute Manual²³. We tested six different concentrations of LE (4000 to 125 µg/mL) against the standard strain of *S. aureus* ATCC 25923 (Collection of Reference Microorganisms on Health Surveillance, Fundação Oswaldo Cruz, Fiocruz, Brazil) and seven strains of mastitic milk isolates. Milk samples from cows were submitted to the California Mastitis Test (CMT). CMT-positive milk samples were plated on blood agar supplemented with 5% sterile ovine blood and incubated for 24–48 h at 37 °C. Gram-positive, catalase-positive, and rabbit plasma coagulase-positive samples were biochemically confirmed as *Staphylococcus aureus*²⁴. Each strain was considered one repetition of the experiment with five replicates/repetition. As such, we conducted eight repetitions for each LE sample.

The minimum inhibitory concentration (MIC) was determined through visual analysis of turbidity after 24 h of incubation on plates at 37 °C in addition to spectrophotometric reading at 600 nm to determine the percentage of inhibition of bacterial growth, using a previously described method²³.

Because LE is an exudate with a yellow color that easily oxidizes to a dark coloration¹⁴, we confirmed the MIC through a colorimetric method. We added 50 µL of resazurin dye (100 µg/mL, Sigma-Aldrich, St. Louis, MO, USA) to each well after reading the plates by spectrophotometer (600 nm). The plates were left to incubate at 37 °C for an additional 30 min²⁵.

Cytotoxicity of leaf exudate to MAC-T cells

Mammary epithelial cells of the MAC-T (Mammary Alveolar Cells-T) lineage were maintained in culture, as indicated by the supplier (Banco de Células do Rio de Janeiro, Brazil). Briefly, MAC-T cells were cultivated in Dulbecco's Modified Eagle's Medium (DMEM) and supplemented with 100 U/mL of penicillin, 100 µg/mL of streptomycin, 20% (V/V) heat-inactivated fetal bovine serum (FBS, Gibco, CA, USA), 4 mmol L⁻¹-glutamine (Synth), 4.5 g/L glucose (Sigma-Aldrich, St. Louis, MO, USA), 1 mmol of sodium pyruvate (Sigma-Aldrich, St. Louis, MO, USA), 1.5 g/L sodium bicarbonate (VetecTM Sigma-Aldrich, St. Louis, MO, USA), 5 µg/mL insulin (Sigma-Aldrich, St. Louis, MO, USA), and 1 µg/mL hydrocortisone (Sigma-Aldrich, St. Louis, MO, USA) at 37 °C and 5% CO₂ in a humidified incubator. We changed the medium every 48 h. After reaching confluence, the cells were treated with 0.25% trypsin with 1 mmol EDTA (Gibco, CA, USA) to prepare the cellular suspension (10⁵ cells/mL). The suspension was transferred a 96-well microplate (100 µL/well), followed by incubation (24 h) in culture conditions for adherence. Subsequently, varying concentrations (2000, 1000, 500, 250, 125, 62.5, 31.3, 15.6, 7.8 and 3.9 µg/mL) of LE from the summer samples were added for 24 h, and cytotoxicity was determined based on the MTT method (Sigma-Aldrich, St. Louis, MO, USA)²⁶. The formed formazan was dissolved with dimethyl sulfoxide (DMSO) to give a purple color with characteristic absorption at 540 nm. Intensity of purple color is directly proportional to the cell number,

thus indicating cell viability. The experiments were performed in triplicate.

Statistical analysis

Data were expressed as the mean ± standard deviation (SD), of at least three independent experiments. We analyzed the data using analysis of variance with a Tukey adjustment (GraphPad Prism 5.0). We considered the effects statistically significant for P<0.05. The inhibitory concentrations capable of reducing cellular viability by 50% (IC₅₀) were calculated using a nonlinear regression of data obtained from the cellular viability tests with GraphPad Prism 5.0 software. The average accumulated precipitation (mm) in the study region was calculated using available data²⁷.

Results and discussion

Leaf exudate chemical profile

The total phenolic and aloin content of LE from the aloe vera leaves varied based on the season during which the samples were collected. The highest levels were found in the samples taken during the summer and the lowest levels from the spring samples (P<0.05). The accumulation of total phenolics in the summer LE seems to be associated with the climatic conditions during the collection period (Figure 2a, b). Precipitation indices were lower during the summer months (January to March)²⁷ (Figure 2a), possibly causing hydric stress in the plants. In the spring, the lower total phenolic content of LE coincides with a period of greatest precipitation that year. Aloe is a plant comprised of 96% water; thus, its chemical composition is heavily influenced by precipitation levels²⁸, as well as other factors, such as the period of flowering²⁹.

The husk of *Aloe* leaves has greater levels of total phenolics compared to the leaf interior and internal parenchyma. In the literature, these values range from 12.06 to 20.86 µg GAE/mg leaf, depending on the species¹⁴. Among the various phenolic compounds in the leaves of *Aloe*, anthraquinones are noteworthy, particularly aloin. Anthraquinones are free in phloem vessels directly below the leaf epidermis, and aloin, in particular, is distributed throughout the plant as part of its defense mechanism³⁰. In the present study, we found the highest levels of aloin in the summer samples and the lowest in the spring samples (Figure 2 b, c). These results are correlated with the total phenolic content found in the studied samples (Figure 2). Previous studies have also suggested the effect of seasonality on aloin content, and its synthesis is strongly influenced by precipitation levels. Dry periods have been correlated with greater content of aloin in the analysis of aloe vera leaves^{15,31,32}. However, other factors can influence aloin content of aloe vera leaves, including cultivation conditions, age, and plant health³³. For example, higher levels of barbaloin, isobarbaloin, and aloin in *Aloe* sp. plants were found during periods of the year with higher temperatures³².

Antimicrobial activity

Despite significant differences in the levels of total phenolics and aloin in the LE samples (Figure 2b, c), these levels

Figure 2. Seasonal differences in rainfall, phenolic content and aloin content. (A) Average accumulated precipitation (mm) in the study region (Paulo Lopes, SC, Brazil) during 2015; Source: INMET²⁶. (B) Total phenolic content (µg GAE/mg) of Aloe vera leaf LE from different seasons of the year (average of five repetitions $\pm$ SD) ($P < 0.05$). (C) Average content (µg/mg) of aloin (average of three independent injections $\pm$ SD) in samples of LE collected from Aloe vera leaves during different seasons of the year. Data points with the same letter above them are not significantly different from each other ($P < 0.05$ indicates a significant difference).

did not influence antimicrobial activity against *S. aureus*. For all LE samples, the MIC was 1,000 µg/mL, as confirmed by resazurin oxidation. The concentrations below of 500 µg/mL were incapable of reducing bacterial growth to values greater than 70%. The effect of other concentrations between 500 and 1,000 µg/mL was not included in the study (Figure 3).

The effectiveness of LE from aloe vera leaves as an antimicrobial agent has been demonstrated for a wide variety of Gram-positive and Gram-negative bacteria, including *S. aureus* and others^{34,35}. In the literature, the MIC of aloe vera extracts against *S. aureus* varies. Previous studies have shown lower (195 µg/mL), similar (1560 µg/mL), and higher (5,000 µg/mL) MIC values compared to those in the present study^{36–38}. This variation may be related to diverse factors, such as the *Aloe* species studied,

the part of the aloe leaf used in the tests, and the type of extraction and resuspension vehicle used. In the current study, the LE samples were collected directly from the cut leaf without any type of posterior extraction of the compounds of interest. Some solvents are capable of extracting certain compounds that may possess greater antimicrobial activity than others³⁹; however, resuspension in water may be the easiest way to use aloe vera leaf subproducts, making it accessible, even to the producer.

An interesting aspect to consider in the present study is that the concentration of total phenolics and aloin in the LE samples does not seem to affect antimicrobial activity. By contrast³⁶, the antimicrobial and anti-inflammatory activity of aloe vera LE has been associated with the concentration of phenolic and aloin

Figure 3. Percentage of inhibition (mean $\pm$ SD) of the standard strain of *S. aureus* ATCC (25923) and seven strains of mastitic milk isolates by different concentrations of LE samples taken during different seasons of 2015.

compounds, suggesting that older leaves have higher levels of these compounds, and as such, have greater biological activity and defense against microorganisms and herbivores.

For Fabry *et al.*⁴⁰, the potentially useful activity defined for crude plant extracts with organic solvents is considered good when MIC values are $<8000 \mu\text{g/mL}$, while Gibbons⁴¹ suggests that phytochemical isolates must have MIC values $<1,000 \mu\text{g/mL}$. As such, the antimicrobial action of the LE samples in the present study can be considered good, even though neither extraction nor isolation of the principal components took place.

Cytotoxicity of leaf exudate

The LE showed high toxicity to MAC-T lineage cells, causing significant reduction in cellular viability at concentrations greater than $7.8 \mu\text{g/mL}$ (Figure 4). At higher concentrations, such as $500 \mu\text{g/mL}$, the reduction in the percentage of viable cells was greater than 80%. The IC_{50} was $91.89 \mu\text{g/mL}$. It is worth noting that the MIC of *S. aureus* growth was $1,000 \mu\text{g/mL}$ (Figure 3), a concentration that had a strong effect on the viability of mammary epithelial cells (Figure 4). This result is significant because it suggests that caution must be exercised when considering the intramammary use of LE in order to avoid inflammation, owing to the death of epithelial cells. However, future studies are needed with aloin purified to determine the inhibition of bacterial growth and the degree of toxicity on mammary cells.

In an *in vivo* situation, the administration of a toxic product to bovine mammary glands can result in the development of inflammation⁴², which is more severe than that caused by the infection of pathogens⁴². In these cases, the attempt to combat inflammation leads to the formation of connective tissue at the affected site, which can diminish the alveolar area responsible for the synthesis of milk and, consequently, reduce milk production. In more severe cases, the loss of complete mammary glandular function, or even death of the animal, can occur^{43,44}.

The MAC-T cellular lineage⁴⁵ is an established model that has been frequently used in the investigation of mammary glandular functions and mediators of inflammatory processes⁴⁶. Nonetheless, studies reporting on the effects of *Aloe* sp. extract, or fractions on this type of cell, are scarce. The toxic effects of aloe vera LE on other types of cells are discussed in the literature and have been associated with the presence of aloin and aloe emodin⁴⁷. These anthraquinones induce the apoptosis of cells caused by a reduction in the proportion of cells in the mitotic phase⁴⁸. Another hypothesis is that disruptions to the cell cycle and cellular differentiation, stimulation of the immune system, and antioxidant activity also have an anti-proliferative effect⁴⁹.

While the results found for MAC-T cells show that aloe vera LE has a high toxic potential for bovine mammary glands, the topical use of this product on the external area of the udder, for example pre- and post-dipping, or on instruments used during

Figure 4. Percent of cellular viability (mean ± SD) of the MAC-T lineage after exposure to different concentrations of LE samples in the summer of 2015. Data shown are an average of three independent experiments. Data points with the same letter above them are not significantly different from each other ($P < 0.05$ indicates a significant difference).

the management of milking, can be suggested. In this case, its potential as a disinfectant should be investigated, and cytotoxicity tests on epithelial cells will be performed.

Furthermore, it is important to highlight that the compounds present in the *Aloe vera* LE liquid oxidize easily in the presence of light, oxygen, and at room temperature⁵⁰. As such, very high concentrations are required in order to achieve antimicrobial efficacy against *S. aureus*, concentrations that would be toxic to mammary epithelial cells. Thus, we suggest the standardization of a methodology that can preserve and conserve these oxidative compounds, such as nanoencapsulation, which can maintain the desired antimicrobial activity and diminish the toxic effects.

Dataset 1. Raw data concerning the phytochemical characteristics of leaf exudate and its antimicrobial/cytotoxicity activity

<http://dx.doi.org/10.5256/f1000research.15671.d213901>

Conclusion

Although seasonality interferes with the chemical composition of *aloe vera* LE, the seasonal samples we evaluated did not

differ in relation to antimicrobial activity with a MIC of 1,000 µg/mL found for all samples. At this concentration, *aloe vera* LE shows strong toxic effects on bovine mammary epithelial cells of the MAC-T lineage. Despite the demonstrated antimicrobial activity of *aloe vera* LE, we suggest caution in recommending its intramammary use to treat bovine mastitis; instead, the topical use of this product on an external area, such as the udder, should be investigated.

Data availability

Dataset 1. Raw data concerning the phytochemical characteristics of leaf exudate and its antimicrobial/cytotoxicity activity. DOI: <https://doi.org/10.5256/f1000research.15671.d213901>⁵¹.

Grant information

This work was supported by the CNPq (National Council on Scientific and Technological Development; Brasilia, Brazil) (403415/2013-6, Edital 39/2013).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- Bradley A: **Bovine mastitis: an evolving disease.** *Vet J.* 2002; **164**(2): 116–128. [PubMed Abstract](#) | [Publisher Full Text](#)
- Watts JL: **Etiological agents of bovine mastitis.** *Vet Microbiol.* 1988; **16**(1): 41–66. [PubMed Abstract](#) | [Publisher Full Text](#)
- Anderson KL, Azizoglu RO: **Detection and Causes of Bovine Mastitis with Emphasis on *Staphylococcus aureus*.** *Encyclopedia of Agriculture and Food Systems.* Elsevier Ltd.; 2014; **2**: 435–440. [Publisher Full Text](#)
- Mushtaq S, Shah AM, Shah A, et al.: **Bovine mastitis: An appraisal of its**

- alternative herbal cure. *Microb Pathog*. 2018; 114: 357–361.
[PubMed Abstract](#) | [Publisher Full Text](#)
5. Fiordalisi SAL, Honorato LA, Loiko MR, *et al.*: The effects of Brazilian propolis on etiological agents of mastitis and the viability of bovine mammary gland explants. *J Dairy Sci*. 2016; 99(3): 2308–2318.
[PubMed Abstract](#) | [Publisher Full Text](#)
 6. Barlow J: Mastitis therapy and antimicrobial susceptibility: a multispecies review with a focus on antibiotic treatment of mastitis in dairy cattle. *J Mammary Gland Biol Neoplasia*. 2011; 16(4): 383–407.
[PubMed Abstract](#) | [Publisher Full Text](#)
 7. Collado R, Prenafeta A, González-González L, *et al.*: Probing vaccine antigens against bovine mastitis caused by *Streptococcus uberis*. *Vaccine*. 2016; 34(33): 3848–3854.
[PubMed Abstract](#) | [Publisher Full Text](#)
 8. Basso KM, Bracarense APFRL: Explantes teciduais: Um modelo redescoberto na experimentação animal. *Semin Agrar*. 2013; 34(6Supl2): 3951–3958.
[Publisher Full Text](#)
 9. De Medeiros ES, Veiga M, Wilton J, *et al.*: Avaliação *in vitro* da eficácia de desinfetantes comerciais utilizados no pré e pós-dipping frente amostras de *Staphylococcus* spp. isoladas de mastite bovina. *Pesq Vet Bras*. 2009; 29(1): 71–75.
[Publisher Full Text](#)
 10. Diaz MAN, Rossi CC, Mendonça VR, *et al.*: Screening of medicinal plants for antibacterial activities on *Staphylococcus aureus* strains isolated from bovine mastitis. *Rev Bras Farmacogn*. 2010; 20(5): 724–728.
[Publisher Full Text](#)
 11. Rabot A, Wellnitz O, Meyer HH, *et al.*: Use and relevance of a bovine mammary gland explant model to study infection responses in bovine mammary tissue. *J Dairy Res*. 2007; 74(1): 93–99.
[PubMed Abstract](#) | [Publisher Full Text](#)
 12. Boudjellab N, Chan-Tang HS, Zhao X: Bovine interleukin-1 expression by cultured mammary epithelial cells (MAC-T) and its involvement in the release of MAC-T derived interleukin-8. *Comp Biochem Physiol A Mol Integr Physiol*. 2000; 127(2): 191–199.
[PubMed Abstract](#) | [Publisher Full Text](#)
 13. Wang K, Jin XL, Liu J, *et al.*: Potential of dietary propolis in protecting bovine mammary epithelial cells against mastitis pathogens using *in vitro* models. *J Nutr Intermed Metab*. 2016; 4: 39.
[Publisher Full Text](#)
 14. Lucini L, Pellizzoni M, Pellegrino R, *et al.*: Phytochemical constituents and *in vitro* radical scavenging activity of different *Aloe* species. *Food Chem*. 2015; 170: 501–507.
[PubMed Abstract](#) | [Publisher Full Text](#)
 15. Rodríguez-González VM, Femenia A, González-Laredo RF, *et al.*: Effects of pasteurization on bioactive polysaccharide acemannan and cell wall polymers from *Aloe barbadensis* Mill. *Carbohydr Polym*. 2011; 86(4): 1675–1683.
[Publisher Full Text](#)
 16. Ray A, Gupta SD, Ghosh S: Evaluation of anti-oxidative activity and UV absorption potential of the extracts of *Aloe vera* L. gel from different growth periods of plants. *Ind Crops Prod*. 2013; 49: 712–719.
[Publisher Full Text](#)
 17. Sánchez-Machado DI, López-Cervantes J, Sendón R, *et al.*: *Aloe vera*: Ancient knowledge with new frontiers. *Trends Food Sci Technol*. 2017; 61: 94–102.
[Publisher Full Text](#)
 18. Fanali S, Aturki Z, D'Orazio G, *et al.*: Analysis of Aloe-based phytotherapeutic products by using nano-LC-MS. *J Sep Sci*. 2010; 33(17–18): 2663–2670.
[PubMed Abstract](#) | [Publisher Full Text](#)
 19. Chinchilla N, Carrera C, Durán AG, *et al.*: *Aloe barbadensis*: How a miraculous plant becomes reality. *Phytochem Rev*. 2013; 12(4): 581–602.
[Publisher Full Text](#)
 20. Hamman JH: Composition and applications of *Aloe vera* leaf gel. *Molecules*. 2008; 13(8): 1599–1616.
[PubMed Abstract](#) | [Publisher Full Text](#)
 21. Duval J, Pecher V, Poujol M, *et al.*: Research advances for the extraction, analysis and uses of anthraquinones: A review. *Ind Crops Prod*. 2016; 94: 812–833.
[Publisher Full Text](#)
 22. Mandrioli R, Mercolini L, Ferranti A, *et al.*: Determination of aloe emodin in *Aloe vera* extracts and commercial formulations by HPLC with tandem UV absorption and fluorescence detection. *Food Chem*. 2011; 126(1): 387–393.
[Publisher Full Text](#)
 23. CLSI: Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria That Grow Aerobically. Approved Standard-Tenth Edition. CLSI document M07-A10. 2015; 35(2): 1–87.
[Reference Source](#)
 24. CLSI: Abbreviated Identification of Bacteria and Yeast; Approved Guideline. CLSI document M35A. 2002; 22(18): 1–19.
[Reference Source](#)
 25. Bauer J, Siala W, Tulkens PM, *et al.*: A combined pharmacodynamic quantitative and qualitative model reveals the potent activity of daptomycin and delafloxacin against *Staphylococcus aureus* biofilms. *Antimicrob Agents Chemother*. 2013; 57(6): 2726–2737.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 26. Mosmann T: Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol Methods*. 1983; 65(1–2): 55–63.
[PubMed Abstract](#) | [Publisher Full Text](#)
 27. INMET-Instituto Nacional de Meteorologia. Disponível em: Acesso em: janeiro de 2017.
[Reference Source](#)
 28. Patel K, Patel DK: Medicinal importance, pharmacological activities, and analytical aspects of hispidulin: A concise report. *J Tradit Complement Med*. 2016; 7(3): 360–366.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 29. Koshioka (Née Nishimura) M, Koshioka M, Takino Y, *et al.*: Studies on the evaluation of *Aloe arborescens* Mill. Var. *Natalensis* berger and aloe extract (JP IX). *Pharm Biol*. 1982; 20(2): 53–59.
[Publisher Full Text](#)
 30. Wintola OA, Afolayan AJ: The foliar anatomy and micromorphology of *Aloe ferox* Mill. (Asphodelaceae). *Afr J Tradit Complement Altern Med*. 2014; 11(2): 350–357.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 31. Gutterman Y, Chauser-Volfson E: Peripheral defence strategy: Variation of barbaloin content in the succulent leaf parts of *Aloe arborescens* Miller (Liliaceae). *Bot J Linn Soc*. 2000; 132(4): 385–395.
[Publisher Full Text](#)
 32. Beppu H, Kawai K, Shimpo K, *et al.*: Studies on the components of *Aloe arborescens* from Japan - Monthly variation and differences due to part and position of the leaf. *Biochem Syst Ecol*. 2004; 32(9): 783–795.
[Publisher Full Text](#)
 33. Azaroual L, Liaizid A, Barbero GF, *et al.*: Improved Chromatographic Methods for Determination of Bioactive Compounds from *Aloe vera* Leaves. *ISRN Chromatogr*. 2012; 609095, 7.
[Publisher Full Text](#)
 34. Habeeb F, Shakir E, Bradbury F, *et al.*: Screening methods used to determine the anti-microbial properties of *Aloe vera* inner gel. *Methods*. 2007; 42(4): 315–320.
[PubMed Abstract](#) | [Publisher Full Text](#)
 35. Pellizzoni M, Ruzickova G, Kalhotka L, *et al.*: Antimicrobial activity of different *Aloe barbadensis* Mill. and *Aloe arborescens* Mill. leaf fractions. *J Med Plants Res*. 2012; 6(10): 1975–1982.
[Reference Source](#)
 36. Ndhlala AR, Amoo SO, Stafford GI, *et al.*: Antimicrobial, anti-inflammatory and mutagenic investigation of the South African tree aloe (*Aloe barberae*). *J Ethnopharmacol*. 2009; 124(3): 404–408.
[PubMed Abstract](#) | [Publisher Full Text](#)
 37. Cardoso FL, Murakami C, Mayworm MAS, *et al.*: Análise sazonal do potencial antimicrobiano e teores de flavonoides e quinonas de extratos foliares de *Aloe arborescens* Mill., Xanthorrhoeaceae. *Brazilian J Pharmacogn*. 2010; 20(1): 35–40.
[Publisher Full Text](#)
 38. Soyelu OT, Masika PJ: Traditional remedies used for the treatment of cattle wounds and myiasis in Amatola Basin, Eastern Cape Province, South Africa. *Onderstepoort J Vet Res*. 2009; 76(4): 393–397.
[PubMed Abstract](#) | [Publisher Full Text](#)
 39. Amoo SO, Aremu AO, Van Staden J: Unraveling the medicinal potential of South African *Aloe* species. *J Ethnopharmacol*. 2014; 153(1): 19–41.
[PubMed Abstract](#) | [Publisher Full Text](#)
 40. Fabry W, Okemo PO, Ansorg R: Antibacterial activity of East African medicinal plants. *J Ethnopharmacol*. 1998; 60(1): 79–84.
[PubMed Abstract](#) | [Publisher Full Text](#)
 41. Gibbons S: Anti-staphylococcal plant natural products. *Nat Prod Rep*. 2004; 21(2): 263–277.
[PubMed Abstract](#) | [Publisher Full Text](#)
 42. Troncarelli MZ, Langoni H, Brandão HM, *et al.*: Safety of a nanopropolis formulation intended for intramammary treatment of bovine mastitis in organic dairy herds. *Rev Bras Hig e Sanidade Anim*. 2014; 8(5): 517–45.
[Reference Source](#)
 43. Cunningham BK: Fisiologia Veterinária. Elsevier Saunders. 2009; 718.
[Reference Source](#)
 44. Dyce KM, Sack WO, Wensing CJG: Tratado de Anatomia Veterinária. Tratado de Anatomia Veterinária. 2004; 519–522.
 45. Huynh HT, Robitaille G, Turner JD: Establishment of bovine mammary epithelial cells (MAC-T): an *in vitro* model for bovine lactation. *Exp Cell Res*. 1991; 197(2): 191–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
 46. Lahouassa H, Moussay E, Rainard P, *et al.*: Lipopolysaccharides, cytokines, and nitric oxide affect secretion of prostaglandins and leukotrienes by bovine mammary gland during experimentally induced mastitis *in vivo* and *in vitro*. *Vet J*. 2014; 77(1): 90–99.
 47. Coran SA, Bartolucci G, Bambagiotti-Alberti M: Selective determination of aloin in different matrices by HPTLC densitometry in fluorescence mode. *J Pharm Biomed Anal*. 2011; 54(2): 422–425.
[PubMed Abstract](#) | [Publisher Full Text](#)
 48. Esmat AY, Tomasetto C, Rio MC: Cytotoxicity of a natural anthraquinone (Aloin) against human breast cancer cell lines with and without ErbB-2:

- topoisomerase IIalpha coamplification. *Cancer Biol Ther.* 2006; 5(1): 97–103.
[PubMed Abstract](#) | [Publisher Full Text](#)
49. Tomasin R, Gomes-Marcondes MC: **Oral administration of *Aloe vera* and honey reduces Walker tumour growth by decreasing cell proliferation and increasing apoptosis in tumour tissue.** *Phyther Res.* 2011; 25(4): 619–623.
[PubMed Abstract](#) | [Publisher Full Text](#)
50. Freitas VS, Rodrigues RAF, Gaspi FOG: **Propriedades farmacológicas da *Aloe vera* (L.) Burm. f.** *Rev Bras Plantas Med.* 2014; 16(2): 299–307.
[Publisher Full Text](#)
51. Fiordalisi SdAL, Honorato LA, Kuhn S: **Dataset 1 in: *Aloe barbadensis* Miller leaf exudate is a potential treatment for bovine mastitis.** *F1000Research.* 2018.
<http://www.doi.org/10.5256/f1000research.15671.d213901>

Open Peer Review

Current Peer Review Status:

Version 2

Reviewer Report 30 October 2018

<https://doi.org/10.5256/f1000research.18337.r39900>

© 2018 Cardemil L. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Liliana Cardemil

Department of Biology, Faculty of Science, Universidad de Chile, Santiago, Chile

I am not satisfied with the responses of the authors. Such an important compound has to be tested in these experiments. For scientists even if the research is for applied uses, it is fundamental to have the results of the control to get the necessary pieces of evidence that the other components of the exudate are producing the effects on cell viability and on the inhibition of bacterial growth. Therefore, I have not approved the manuscript. They should run again the experiments with such important compound present in the leaf exudate.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to state that I do not consider it to be of an acceptable scientific standard, for reasons outlined above.

Reviewer Report 26 October 2018

<https://doi.org/10.5256/f1000research.18337.r39901>

© 2018 Diaz-Muñoz G. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Gaspar Diaz-Muñoz

Department of Chemistry, Federal University of Minas Gerais, Belo Horizonte, Brazil

I read the revised manuscript entitled: "***Aloe barbadensis* Miller leaf exudate is a potential treatment for bovine mastitis**". The manuscript is very interesting. In general, I found it clearly written and errors were found as listed in the attached file that already were corrected. Overall, I

believe that this is otherwise a very good manuscript and can be published in F1000Research.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 14 September 2018

<https://doi.org/10.5256/f1000research.17100.r37690>

© 2018 Cardemil L et al. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Liliana Cardemil

Department of Biology, Faculty of Science, Universidad de Chile, Santiago, Chile

Carlos Salinas

Department of Biology, Faculty of Science, Universidad de Chile, Santiago, Chile

The work is interesting since the infection of bovine mammary epithelial cells by *Staphylococcus aureus* it's a problem in cattle farming.

However, the study they performed is incomplete. I do not understand, why the investigators did not try the Aloin that they purified to determine the inhibition of bacterial growth and the degree of toxicity on mammary cells with this compound as they did with the total leaf exudate? The leaf exudate has too many compounds. Not only phenols, the exudate may have amino acids, anthraquinones, chlorophyll, and other pigments, etc. At least testing the Aloin then you could know if this compound has an antibacterial effect by itself and/or, causes the toxicity of mammary epithelial cells. If the results are positive for Aloin as antimicrobial without causing toxicity they will know that Aloin can be used for the mastitis treatment. If not, they will know that the leaf exudate has to be free of Aloin to treat the mastitis.

The literature of the manuscript is missing important new and relevant literature on the uses of *Aloe barbadensis* Miller as a prebiotic and inhibitor of colon cancer development.

In my opinion the need to perform the analyses considering the aloin. In my opinion, the manuscript should be approved after the authors complete these analyses.

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Partly

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Partly

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Prebiotic molecules from Aloe vera

We confirm that we have read this submission and believe that we have an appropriate level of expertise to state that we do not consider it to be of an acceptable scientific standard, for reasons outlined above.

Reader Comment 22 Oct 2018

Shirley Kuhnen,

- We chose not to include a control with Aloin in the study. We acknowledge that there are many works involving the purification of compounds from natural products in the field of Pharmacology. However, our objective was to test a complex matrix for its use (in natura) by organic or agroecological small producers of milk. There are several studies in the literature with complex matrices derived from medicinal plants that do not include a control as mentioned by the reviewer. The use of a complex matrix has some advantages, among them the lesser emergence of resistance among microorganisms. We also point out that there are no methodological errors in the study such as the lack of the controls required by the CSLI Guidelines and Standards. Therefore, the lack of a control with Aloin does not invalidate the results obtained in the present study and the work should not be rejected for this reason.
- The literature suggested doesn't fit in our paper because it's related to polysaccharides' biological effects.
- The assay was performed according to CSLI Guidelines and Standards. This does not require control with isolated compounds. However, to cater to the suggestion, we added in the text that this may be a line of future research.

Competing Interests: I have no competing interests.

Reviewer Report 14 September 2018

<https://doi.org/10.5256/f1000research.17100.r37692>

© 2018 Diaz-Muñoz G. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Gaspar Diaz-Muñoz

Department of Chemistry, Federal University of Minas Gerais, Belo Horizonte, Brazil

I read with interest the manuscript entitled: "***Aloe barbadensis* Miller leaf exudate is a potential treatment for bovine mastitis**". The manuscript is very interesting. In general, I found clearly written and only a few typos and errors were found as listed in the [attached file](#) that already were corrected.

I did not understand why in the discussion of antimicrobial activity, the authors cited only the results of concentrations between 500 and 1000 ug/mL. In the methodology cited, the use of six concentrations between 500 and 4000 µg/mL.

I think the authors should consider the possibility of Aloe exudate being toxic to epithelial cells as it showed toxicity to the MAC T cells. I suggest making a cytotoxicity test for this cell before suggesting this exudate as a disinfectant.

Overall, I believe that this otherwise very good manuscript could benefit of a minor revision before indexing in F1000Research. I have attached a commented copy of the manuscript for the authors to consider in the revision.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Not applicable

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reader Comment 22 Oct 2018

Shirley Kuhnien,

1- Six concentrations of leaf exudate were tested. However, under 500 ug/mL, the reduction of microbial growth was lower than 70%, varying between seasons. The statement was modified.

2- Thank you very much. You are correct. This information was included in the text. The statements were modified.

3- Thank you very much. The suggestions were included in the new version.

Competing Interests: I have no competing interests .

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research