REVIEW

Recent advances in the understanding and treatment of pemphigus and pemphigoid [version 1; peer review: 2 approved]

Jun Yamagami

Department of Dermatology, Keio University School of Medicine, 35 Shinanomachi, Shinjuku-ku, Tokyo, 160-8582, Japan

V1 First published: 30 Aug 2018, 7(F1000 Faculty Rev):1360
<https://doi.org/10.12688/f1000research.14474.1>
 Latest published: 30 Aug 2018, 7(F1000 Faculty Rev):1360
<https://doi.org/10.12688/f1000research.14474.1>

Abstract

Pemphigus and pemphigoid are characterized as autoimmune blistering diseases in which immunoglobulin G autoantibodies cause blisters and erosions of the skin or mucosa or both. Recently, understanding of the pathophysiology of pemphigus and pemphigoid has been furthered by genetic analyses, characterization of autoantibodies and autoreactive B cells, and elucidation of cell-cell adhesion between keratinocytes. For the management of pemphigus and pemphigoid, the administration of systemic corticosteroids still represents the standard treatment strategy; however, evidence of the efficacy of therapies not involving corticosteroids, such as those employing anti-CD20 antibodies, is increasing. The goal should be to develop antigen-specific immune suppression-based treatments.

Keywords

pemphigus, pemphigoid

Open Peer Review

Approval Status

	1	2
version 1
30 Aug 2018		

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Ralf Ludwig**, University of Lübeck, Lübeck, Germany
2. **Soner Uzun**, Akdeniz University, Antalya, Turkey

Any comments on the article can be found at the end of the article.

Corresponding author: Jun Yamagami (yamagamijun@keio.jp)

Author roles: Yamagami J: Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: Jun Yamagami is supported by Grants-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology of Japan and the Keio Gijuku Academic Development Fund.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2018 Yamagami J. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Yamagami J. **Recent advances in the understanding and treatment of pemphigus and pemphigoid** [version 1; peer review: 2 approved] F1000Research 2018, 7(F1000 Faculty Rev):1360 <https://doi.org/10.12688/f1000research.14474.1>

First published: 30 Aug 2018, 7(F1000 Faculty Rev):1360 <https://doi.org/10.12688/f1000research.14474.1>

Introduction

Autoimmune blistering diseases are induced by autoantibodies against components of the skin. There are two main types of autoimmune blistering disease: pemphigus (blister in the epidermis) and pemphigoid (subepidermal blistering). In pemphigus, autoantibodies target desmoglein 1 (Dsg 1) and Dsg 3, which play a major role in desmosomes essential for cell–cell adhesion between keratinocytes and cause blister formation with acantholysis¹. In pemphigoid, autoantibodies target molecules involved in connecting basal epithelial cells to the basement membrane in hemidesmosomes, such as type XVII collagen (COL17, BP180), dystonin-e (BP230), type VII collagen (COL7), p200, and laminin 332^{2,3}. This article reviews recent advances in our understanding of the pathophysiology and treatment of pemphigus and pemphigoid.

Pemphigus

Pathophysiology

Pemphigus is considered one of the best-characterized human autoimmune diseases. Recent studies have shed light on the genetic susceptibility to pemphigus, characterized by the presence of Dsg-reactive autoantibodies, and revealed the mechanisms underlying blister formation⁴.

Genetic susceptibility. Several human leukocyte antigen (HLA) alleles, such as *HLA-DRB1*0402* and *HLA-DQB1*0503*, have been suggested as risk factors for pemphigus vulgaris (PV)^{5–7}. Although recent microarray analyses have indicated that HLA status may be a key driver of autoantibody expression and could be related to disease activity via the oxidative stress that occurs during PV, the correlation between the HLA genetic profile and overall clinical profile remains unclear^{8,9}. A non-HLA marker encoding ST18, a molecule that regulates apoptosis and inflammation, has been investigated as a new candidate gene associated with PV. *ST18*-associated variants may predispose to PV in a population-specific manner, and an association between *ST18* and PV was found in Jewish and Egyptian, but not German, populations¹⁰. In one report, PV serum-induced secretion of key inflammatory molecules by keratinocytes was related to a PV-associated functional risk variant residing within the *ST18* promoter region¹¹.

Characterization of autoantibodies. Investigations of anti-Dsg autoantibodies and autoreactive B cells have helped to uncover the mechanisms underlying the development of pemphigus and the production of autoantibodies^{12–15}. Recent studies suggested that B cells with *V_HI-46* heavy chain gene usage might be prone to Dsg3 autoreactivity. Dsg3-specific B cells that use *V_HI-46* demonstrated relatively few somatic mutations and required few, or none, of these mutations to bind Dsg3¹⁶. Furthermore, in patients with PV, cross-reactivity of B cells with *V_HI-46* to both Dsg3 and VP6, a rotavirus capsid protein, suggests that the anti-Dsg B-cell repertoire can cross-react with foreign antigens¹⁷. Previously, anti-Dsg1 immunoglobulin G (IgG) monoclonal antibodies that can cross-react with LJM11, a sand fly antigen, were reported in the endemic form of pemphigus foliaceus¹⁸. These findings indicate that the immune response to foreign antigens may lead to autoimmunity, in turn resulting in the development of pemphigus.

Long-term analysis of autoreactive B-cell repertoires in patients with pemphigus revealed the presence of identical anti-Dsg antibody clones before and after treatment, suggesting that targeting specific sets of autoreactive B cells may be a feasible therapeutic strategy¹⁹. However, proteomic analysis of pemphigus autoantibodies gave us a different perspective by indicating that changes in the proportions of autoantibodies occur over time²⁰. Further research is necessary to fully delineate the mechanisms of autoantibody production in pemphigus. In addition to Dsg, other autoimmune targets have been found, and their pathogenic roles might be elucidated in future investigation²¹.

Blister formation by autoantibodies. A direct pathogenic role of autoantibodies in pemphigus has been clearly established. There are two major routes to acantholytic blister formation: autoantibody-mediated steric hindrance of desmosomal adhesion^{12,15,22} and activation of particular cell signaling pathways such as the p38 mitogen-activated protein kinase (MAPK) pathway^{23–26}. Studies have shown that pathogenic anti-Dsg monoclonal antibodies can bind directly to residues that mediate adhesion and that polyclonal antibodies contribute to acantholysis in a non-redundant way; thus, all antibodies have the potential to cause blistering via synergistic mechanisms^{27,28}. Advances in understanding of the crystal structure of desmocollins and Dsgs²⁹ and in techniques for high-resolution imaging of the skin will further clarify the pathophysiological regulation of keratinocyte cell–cell adhesion in pemphigus^{26,30}.

Treatment

The goal in pemphigus treatment is to maintain complete remission, defined as the absence of new or established lesions³¹. Ideally, all systemic therapy should be stopped; however, remission achieved by minimal therapy, prednisone (≤10 mg/day), or minimal adjuvant therapy (or a combination of these) may be a more realistic goal in the management of pemphigus^{32,33}.

Because of the rarity of pemphigus, published guidelines for its management rely mostly on expert consensus, except for a few evidence-based controlled studies. Treatments recommended by European and Japanese guidelines include corticosteroids and immunosuppressive reagents, such as azathioprine and cyclophosphamide, plasmapheresis, intravenous immunoglobulin (IVIG), and rituximab; most therapies aim to improve symptoms by reducing serum autoantibodies, either directly or through generalized immune suppression^{33,34}. Recent studies have highlighted the advantages of rituximab, a monoclonal anti-CD20 antibody that targets CD20⁺ B cells, in the treatment of pemphigus^{35–38}. The results of multicenter prospective randomized trials of rituximab, as a first-line treatment for moderate to severe cases of pemphigus, were recently published in France³⁹. At 2 years after treatment, 41 (89%) of 46 patients assigned to rituximab plus short-term prednisone were in complete remission and off therapy versus 15 (34%) of 44 patients assigned to the prednisone-alone treatment. Significantly more serious adverse events (that is, of grade 3–4) were reported in the prednisone-alone group than in the rituximab-plus-prednisone

group. These results suggest that first-line use of rituximab for patients with pemphigus is more effective, and causes fewer adverse events, than corticosteroids alone. As such, the treatment strategy for pemphigus may change worldwide.

The ideal therapy for pemphigus would eliminate only those autoantibodies produced by Dsg-specific B cells. A recent study suggested the feasibility of such targeted therapy through the use of chimeric antigen receptor T (CART) cells⁴⁰. In PV, autoantigen-based chimeric immunoreceptors can direct T cells to kill autoreactive B cells according to the specificity of the B-cell receptor (BCR). Human T cells engineered to express chimeric autoantibody receptor (CAAR), consisting of Dsg3 fused to CD137-CD3 ζ signaling domains (Dsg3 CAAR-T cells), exhibit specific cytotoxicity against cells expressing anti-Dsg3 BCRs *in vitro* and *in vivo*.

Pemphigoid

Pathophysiology

There are three major types of pemphigoid: bullous pemphigoid (BP), which is the most common autoimmune bullous disorder and is characterized by widespread erythema and blisters on the skin; mucous membrane pemphigoid, which affects mainly the oral mucosa; and epidermolysis bullosa acquisita, which is a bullous disease associated with autoimmunity to COL7, the major component of the anchoring fibrils of the dermal–epidermal junction.

Disease development. Although the genetic background and mechanism underlying the development of pemphigoid remain to be precisely determined, several studies have suggested a genetic predisposition in its etiology^{41–43}. A German study showed that a polymorphism in the mitochondrially encoded ATP synthase 8 gene (*MT-ATP8*) may have an association with BP pathogenesis, which suggests a contribution of mitochondrial abnormalities to BP immunopathology⁴⁴. Furthermore, a recent study showed that gene polymorphism in *CYP2D6*, which is among the cytochrome P450 isoenzymes, may influence the risk of drug-induced BP⁴⁵. Dipeptidyl peptidase-IV (DPP-4) inhibitors have been increasingly implicated as a cause of drug-induced BP. Recently, a strong association between the use of DPP-4 inhibitors and the risk of BP was reported⁴⁶. Although the mechanism is still unclear, several reports revealed that patients with BP associated with the use of DPP-4 inhibitors tended to exhibit a non-inflammatory clinical phenotype and possessed IgG autoantibodies against full-length COL17, albeit outside of the non-collagenous 16A domain (NC16A) (the major IgG epitope in typical BP)^{47,48}.

Blister formation by autoantibodies. Following binding of IgG autoantibodies to their target antigens in the basement membrane zone (BMZ), complements and other factors induce neutrophil and eosinophil chemotaxis and mast cell degranulation as well as release their proteolytic enzymes, such as metallopeptidase-9 and neutrophil elastase^{49–51}. Although these complement-dependent mechanisms are believed to be essential for the

inflammatory cascade that causes subepidermal blistering, evidence supporting blister induction via direct interaction between autoantibodies and autoantigens (complement-independent mechanism) has recently been reported^{52–57}.

IgG4 is assumed to lack the ability to fix complement, and BP cases in which IgG4 is dominant and C3 deposition at the BMZ is lacking have been reported⁵⁸. Additionally, passive transfer of IgG4 and IgG1 autoantibodies in patients with BP has been shown to induce blister formation in neonatal C3-deficient COL17-humanized mice without complement activation⁵⁹. These reports suggest that IgG4 autoantibodies may contribute to complement-independent blister formation in some cases of BP.

Recent studies also revealed that skin fragility in BP is induced by macropinocytosis of immune complexes after the binding of autoantibodies to COL17 on the cell surface⁶⁰. Studies showing that antibodies targeting areas outside of NC16A had less capacity to deplete COL17 on keratinocytes than antibodies targeting NC16A suggest that the pathogenicity of autoantibodies in BP could be epitope dependent^{61,62}. Recent studies have offered some biomarkers for more efficient diagnosis and prediction of prognosis in pemphigoid^{63,64}.

Treatment

Systemic glucocorticoids remain the standard treatment for pemphigoid. Superpotent topical corticosteroids have shown efficacy in BP but may not be practical for daily use as whole-body ointments^{65,66}. British guidelines have mentioned anti-inflammatory antibiotics (that is, tetracyclines) for treating BP, but no clear evidence supporting this recommendation was provided⁶⁷. Recently, a randomized controlled trial comparing doxycycline with prednisolone as an initial treatment for BP was published⁶⁸. Participants were randomly assigned to a doxycycline (200 mg/day) or a prednisolone (0.5 mg/kg per day) group, and 83 (74%) of 112 patients had three or fewer blisters at 6 weeks compared with 92 (91%) of 101 patients on prednisolone. The rate of severe, life-threatening, or fatal events at 52 weeks were 18% (22 of 121) for those on doxycycline and 36% (41 of 113) for those on prednisolone. This study provided important evidence that doxycycline is not inferior to oral prednisolone for short-term blister control in BP and also is significantly safer in the long term. Although disease severity should be carefully evaluated on a case-by-case basis and the mechanism of action of doxycycline is still unclear, this agent likely will be widely used as a first-line treatment for BP.

Most therapies that have shown efficacy against pemphigus have also been tried for pemphigoid. A randomized double-blind trial of IVIG showed that the disease activity score on day 15 was significantly lower in severe BP versus the baseline value, suggesting that IVIG could be a good option for patients resistant to corticosteroids^{69,70}. In a retrospective study, first-line combination therapy with rituximab and corticosteroids was recommended for cases of moderate to severe BP; the regimen achieved a higher rate of complete remission compared with treatment by rituximab^{71,72}.

Accumulating evidence suggests a pathogenic role for IgE in BP. IgE deposition in the BMZ in patients with BP has frequently been observed^{73,74}, and a correlation between serum levels of IgE autoantibodies against COL17 and BP disease activity has been reported⁷⁵. High-affinity IgE receptor (FcεRI)-bound IgE on mast cells and eosinophils may be involved in the degranulation of such cells following their capture of shed extracellular fragments of COL17, in turn resulting in the development of urticarial erythema in BP⁷⁶. Reports of BP cases successfully treated with anti-IgE monoclonal antibodies also support the potential of IgE as a therapeutic target in BP^{77,78}.

Closing remarks

The precise mechanism that triggers autoantibody production in patients with pemphigus and pemphigoid remains unknown. Elucidating the pathogenesis of these diseases via repertoire analysis and study of the mechanism of activation

of autoreactive B cells is a promising approach. Furthermore, clarification of the mechanism underlying blister formation may yield a novel treatment target. Therapeutic approaches that do not involve corticosteroids, such as those based on anti-CD20 antibodies, are gradually being established. A goal of treatments for pemphigus and pemphigoid should be targeted depletion only of antigen-specific T/B cells or autoantibodies.

Grant information

Jun Yamagami is supported by Grants-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology of Japan and the Keio Gijuku Academic Development Fund.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- Stanley JR, Amagai M: **Pemphigus, bullous impetigo, and the staphylococcal scalded-skin syndrome.** *N Engl J Med.* 2006; 355(17): 1800–10.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kasperkiewicz M, Zillikens D, Schmidt E: **Pemphigoid diseases: pathogenesis, diagnosis, and treatment.** *Autoimmunity.* 2012; 45(1): 55–70.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Goletz S, Zillikens D, Schmidt E: **Structural proteins of the dermal-epidermal junction targeted by autoantibodies in pemphigoid diseases.** *Exp Dermatol.* 2017; 26(12): 1154–62.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Kasperkiewicz M, Ellebrecht CT, Takahashi H, et al.: **Pemphigus.** *Nat Rev Dis Primers.* 2017; 3: 17026.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Yan L, Wang JM, Zeng K: **Association between HLA-DRB1 polymorphisms and pemphigus vulgaris: a meta-analysis.** *Br J Dermatol.* 2012; 167(4): 768–77.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Erning R, Hennerici T, Bäcklund J, et al.: **Pathogenic IgG antibodies against desmoglein 3 in pemphigus vulgaris are regulated by HLA-DRB1*04:02-restricted T cells.** *J Immunol.* 2014; 193(9): 4391–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Li S, Zhang Q, Wang P, et al.: **Association between HLA-DQB1 polymorphisms and pemphigus vulgaris: A meta-analysis.** *Immunol Invest.* 2018; 47(1): 101–12.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Sinha AA: **The genetics of pemphigus.** *Dermatol Clin.* 2011; 29(3): 381–91, vii.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Shah AA, Dey-Rao R, Seiffert-Sinha K, et al.: **Increased oxidative stress in pemphigus vulgaris is related to disease activity and HLA-association.** *Autoimmunity.* 2016; 49(4): 248–57.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Sarig O, Bercovici S, Zoller L, et al.: **Population-specific association between a polymorphic variant in ST18, encoding a pro-apoptotic molecule, and pemphigus vulgaris.** *J Invest Dermatol.* 2012; 132(7): 1798–805.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Vodo D, Sarig O, Geller S, et al.: **Identification of a Functional Risk Variant for Pemphigus Vulgaris in the ST18 Gene.** *PLoS Genet.* 2016; 12(5): e1006008.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Tsunoda K, Ota T, Aoki M, et al.: **Induction of pemphigus phenotype by a mouse monoclonal antibody against the amino-terminal adhesive interface of desmoglein 3.** *J Immunol.* 2003; 170(4): 2170–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Payne AS, Ishii K, Kacir S, et al.: **Genetic and functional characterization of human pemphigus vulgaris monoclonal autoantibodies isolated by phage display.** *J Clin Invest.* 2005; 115(4): 888–99.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Yamagami J, Payne AS, Kacir S, et al.: **Homologous regions of autoantibody heavy chain complementarity-determining region 3 (H-CDR3) in patients with pemphigus cause pathogenicity.** *J Clin Invest.* 2010; 120(11): 4111–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Di Zenzo G, Di Lullo G, Corti D, et al.: **Pemphigus autoantibodies generated through somatic mutations target the desmoglein-3 cis-interface.** *J Clin Invest.* 2012; 122(10): 3781–90.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Cho MJ, Lo AS, Mao X, et al.: **Shared VH1-46 gene usage by pemphigus vulgaris autoantibodies indicates common humoral immune responses among patients.** *Nat Commun.* 2014; 5: 4167.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Cho MJ, Ellebrecht CT, Hammers CM, et al.: **Determinants of VH1-46 Cross-Reactivity to Pemphigus Vulgaris Autoantigen Desmoglein 3 and Rotavirus Antigen VP6.** *J Immunol.* 2016; 197(4): 1065–73.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Qian Y, Jeong JS, Maldonado M, et al.: **Cutting Edge: Brazilian pemphigus foliaceus anti-desmoglein 1 autoantibodies cross-react with sand fly salivary LJM11 antigen.** *J Immunol.* 2012; 189(4): 1535–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Hammers CM, Chen J, Lin C, et al.: **Persistence of anti-desmoglein 3 IgG⁺ B-cell clones in pemphigus patients over years.** *J Invest Dermatol.* 2015; 135(3): 742–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Chen J, Zheng Q, Hammers CM, et al.: **Proteomic Analysis of Pemphigus Autoantibodies Indicates a Larger, More Diverse, and More Dynamic Repertoire than Determined by B Cell Genetics.** *Cell Rep.* 2017; 18(1): 237–47.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Seiffert-Sinha K, Khan S, Attwood K, et al.: **Anti-Thyroid Peroxidase Reactivity Is Heightened in Pemphigus Vulgaris and Is Driven by Human Leukocyte Antigen Status and the Absence of Desmoglein Reactivity.** *Front Immunol.* 2018; 9: 625.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Heupel WM, Zillikens D, Drenckhahn D, et al.: **Pemphigus vulgaris IgG directly inhibit desmoglein 3-mediated transinteraction.** *J Immunol.* 2008; 181(3): 1825–34.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Mao X, Sano Y, Park JM, et al.: **p38 MAPK activation is downstream of the loss of intercellular adhesion in pemphigus vulgaris.** *J Biol Chem.* 2011; 286(2): 1283–91.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)

24. Bektas M, Jolly PS, Berkowitz P, *et al.*: **A pathophysiologic role for epidermal growth factor receptor in pemphigus acantholysis.** *J Biol Chem.* 2013; **288**(13): 9447–56.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
25. Schiögl E, Radeva MY, Vielmuth F, *et al.*: **Keratin Retraction and Desmoglein3 Internalization Independently Contribute to Autoantibody-Induced Cell Dissociation in Pemphigus Vulgaris.** *Front Immunol.* 2018; **9**: 858.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
26. Lotti R, Shu E, Petrachi T, *et al.*: **Soluble Fas Ligand Is Essential for Blister Formation in Pemphigus.** *Front Immunol.* 2018; **9**: 370.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
27. Kawasaki H, Tsunoda K, Hata T, *et al.*: **Synergistic pathogenic effects of combined mouse monoclonal anti-desmoglein 3 IgG antibodies on pemphigus vulgaris blister formation.** *J Invest Dermatol.* 2006; **126**(12): 2621–30.
[PubMed Abstract](#) | [Publisher Full Text](#)
28. **F** Saito M, Stahley SN, Caughman CY, *et al.*: **Signaling dependent and independent mechanisms in pemphigus vulgaris blister formation.** *PLoS One.* 2012; **7**(12): e50696.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
29. **F** Harrison OJ, Brasch J, Lasso G, *et al.*: **Structural basis of adhesive binding by desmocollins and desmogleins.** *Proc Natl Acad Sci U S A.* 2016; **113**(26): 7160–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
30. **F** Stahley SN, Warren MF, Feldman RJ, *et al.*: **Super-Resolution Microscopy Reveals Altered Desmosomal Protein Organization in Tissue from Patients with Pemphigus Vulgaris.** *J Invest Dermatol.* 2016; **136**(1): 59–66.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
31. Murrell DF, Peña S, Joly P, *et al.*: **Diagnosis and Management of Pemphigus: recommendations by an International Panel of Experts.** *J Am Acad Dermatol.* 2018; pii: S0190-9622(18)30207-X.
[PubMed Abstract](#) | [Publisher Full Text](#)
32. Murrell DF, Dick S, Ahmed AR, *et al.*: **Consensus statement on definitions of disease, end points, and therapeutic response for pemphigus.** *J Am Acad Dermatol.* 2008; **58**(6): 1043–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
33. Committee for Guidelines for the Management of Pemphigus Disease, Amagai M, Tanikawa A, *et al.*: **Japanese guidelines for the management of pemphigus.** *J Dermatol.* 2014; **41**(6): 471–86.
[PubMed Abstract](#) | [Publisher Full Text](#)
34. Herli M, Jedlickova H, Karpati S, *et al.*: **Pemphigus. S2 Guideline for diagnosis and treatment—guided by the European Dermatology Forum (EDF) in cooperation with the European Academy of Dermatology and Venereology (EADV).** *J Eur Acad Dermatol Venereol.* 2015; **29**(3): 405–14.
[PubMed Abstract](#) | [Publisher Full Text](#)
35. **F** Joly P, Mouquet H, Roujeau JC, *et al.*: **A single cycle of rituximab for the treatment of severe pemphigus.** *N Engl J Med.* 2007; **357**(6): 545–52.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
36. **F** Lunardon L, Tsai KJ, Probert KJ, *et al.*: **Adjuvant rituximab therapy of pemphigus: a single-center experience with 31 patients.** *Arch Dermatol.* 2012; **148**(9): 1031–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
37. **F** Wang HH, Liu CW, Li YC, *et al.*: **Efficacy of rituximab for pemphigus: a systematic review and meta-analysis of different regimens.** *Acta Derm Venereol.* 2015; **95**(8): 928–32.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
38. **F** Musette P, Bouaziz JD: **B Cell Modulation Strategies in Autoimmune Diseases: New Concepts.** *Front Immunol.* 2018; **9**: 622.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
39. **F** Joly P, Maho-Vaillant M, Prost-Squarcioni C, *et al.*: **First-line rituximab combined with short-term prednisone versus prednisone alone for the treatment of pemphigus (Ritux 3): A prospective, multicentre, parallel-group, open-label randomised trial.** *Lancet.* 2017; **389**(10083): 2031–40.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
40. **F** Ellebrecht CT, Bhoj VG, Nace A, *et al.*: **Reengineering chimeric antigen receptor T cells for targeted therapy of autoimmune disease.** *Science.* 2016; **353**(6295): 179–84.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
41. Delgado JC, Turbay D, Yunis EJ, *et al.*: **A common major histocompatibility complex class II allele HLA-DQB1* 0301 is present in clinical variants of pemphigoid.** *Proc Natl Acad Sci U S A.* 1996; **93**(16): 8569–71.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
42. Setterfield J, Theron J, Vaughan RW, *et al.*: **Mucous membrane pemphigoid: HLA-DQB1*0301 is associated with all clinical sites of involvement and may be linked to ant basement membrane IgG production.** *Br J Dermatol.* 2001; **145**(3): 406–14.
[PubMed Abstract](#) | [Publisher Full Text](#)
43. **F** Sadik CD, Bischof J, van Beek N, *et al.*: **Genomewide association study identifies GALT as susceptibility gene for mucous membrane pemphigoid.** *Exp Dermatol.* 2017; **26**(12): 1214–20.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
44. Hirose M, Schilf P, Benoit S, *et al.*: **Polymorphisms in the mitochondrially encoded ATP synthase 8 gene are associated with susceptibility to bullous pemphigoid in the German population.** *Exp Dermatol.* 2015; **24**(9): 715–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
45. Rychlik-Sych M, Barańska M, Wojtczak A, *et al.*: **The impact of the CYP2D6 gene polymorphism on the risk of pemphigoid.** *Int J Dermatol.* 2015; **54**(12): 1396–401.
[PubMed Abstract](#) | [Publisher Full Text](#)
46. **F** Béné J, Moulis G, Bennani I, *et al.*: **Bullous pemphigoid and dipeptidyl peptidase IV inhibitors: a case-noncase study in the French Pharmacovigilance Database.** *Br J Dermatol.* 2016; **175**(2): 296–301.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
47. **F** Izumi K, Nishie W, Mai Y, *et al.*: **Autoantibody Profile Differentiates between Inflammatory and Noninflammatory Bullous Pemphigoid.** *J Invest Dermatol.* 2016; **136**(11): 2201–10.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
48. Horikawa H, Kurihara Y, Funakoshi T, *et al.*: **Unique clinical and serological features of bullous pemphigoid associated with dipeptidyl peptidase-4 inhibitors.** *Br J Dermatol.* 2018; **178**(6): 1462–3.
[PubMed Abstract](#) | [Publisher Full Text](#)
49. Lo Schiavo A, Ruocco E, Brancaccio G, *et al.*: **Bullous pemphigoid: etiology, pathogenesis, and inducing factors: facts and controversies.** *Clin Dermatol.* 2013; **31**(4): 391–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
50. Le Jan S, Plée J, Vallerand D, *et al.*: **Innate immune cell-produced IL-17 sustains inflammation in bullous pemphigoid.** *J Invest Dermatol.* 2014; **134**(12): 2908–17.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
51. Plée J, Le Jan S, Giustiniani J, *et al.*: **Integrating longitudinal serum IL-17 and IL-23 follow-up, along with autoantibodies variation, contributes to predict bullous pemphigoid outcome.** *Sci Rep.* 2015; **5**: 18001.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
52. **F** Iwata H, Kamio N, Aoyama Y, *et al.*: **IgG from patients with bullous pemphigoid depletes cultured keratinocytes of the 180-kDa bullous pemphigoid antigen (type XVII collagen) and weakens cell attachment.** *J Invest Dermatol.* 2009; **129**(4): 919–26.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
53. Natsuga K, Nishie W, Shinkuma S, *et al.*: **Antibodies to pathogenic epitopes on type XVII collagen cause skin fragility in a complement-dependent and -independent manner.** *J Immunol.* 2012; **188**(11): 5792–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
54. **F** Iwata H, Ujiiie H: **Complement-independent blistering mechanisms in bullous pemphigoid.** *Exp Dermatol.* 2017; **26**(12): 1235–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
55. Ludwig RJ, Vanhoorelbeke K, Leyboldt F, *et al.*: **Mechanisms of Autoantibody-Induced Pathology.** *Front Immunol.* 2017; **8**: 603.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
56. **F** Mihai S, Hirose M, Wang Y, *et al.*: **Specific Inhibition of Complement Activation Significantly Ameliorates Autoimmune Blistering Disease in Mice.** *Front Immunol.* 2018; **9**: 535.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
57. **F** Karsten CM, Beckmann T, Holtsche MM, *et al.*: **Tissue Destruction in Bullous Pemphigoid Can Be Complement Independent and May Be Mitigated by C5aR2.** *Front Immunol.* 2018; **9**: 488.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
58. **F** Dainichi T, Nishie W, Yamagami Y, *et al.*: **Bullous pemphigoid suggestive of complement-independent blister formation with anti-BP180 IgG4 autoantibodies.** *Br J Dermatol.* 2016; **175**(1): 187–90.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
59. Ujiiie H, Sasaoka T, Izumi K, *et al.*: **Bullous pemphigoid autoantibodies directly induce blister formation without complement activation.** *J Immunol.* 2014; **193**(8): 4415–28.
[PubMed Abstract](#) | [Publisher Full Text](#)
60. **F** Iwata H, Kamaguchi M, Ujiiie H, *et al.*: **Macropinocytosis of type XVII collagen induced by bullous pemphigoid IgG is regulated via protein kinase C.** *Lab Invest.* 2016; **96**(12): 1301–10.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
61. **F** Wada M, Nishie W, Ujiiie H, *et al.*: **Epitope-Dependent Pathogenicity of Antibodies Targeting a Major Bullous Pemphigoid Autoantigen Collagen XVII/BP180.** *J Invest Dermatol.* 2016; **136**(5): 938–46.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
62. **F** Imafuku K, Iwata H, Kamaguchi M, *et al.*: **Autoantibodies of non-inflammatory bullous pemphigoid hardly deplete type XVII collagen of keratinocytes.** *Exp Dermatol.* 2017; **26**(12): 1171–4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
63. **F** Giusti D, Le Jan S, Gatouillat G, *et al.*: **Biomarkers related to bullous pemphigoid activity and outcome.** *Exp Dermatol.* 2017; **26**(12): 1240–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
64. **F** Riani M, Le Jan S, Plée J, *et al.*: **Bullous pemphigoid outcome is associated with CXCL10-induced matrix metalloproteinase 9 secretion from monocytes**

- and neutrophils but not lymphocytes. *J Allergy Clin Immunol.* 2017; **139**(3): 863–872.e3.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
65. **F** Joly P, Roujeau JC, Benichou J, *et al.*: **A comparison of oral and topical corticosteroids in patients with bullous pemphigoid.** *N Engl J Med.* 2002; **346**(5): 321–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 66. **F** Joly P, Roujeau JC, Benichou J, *et al.*: **A comparison of two regimens of topical corticosteroids in the treatment of patients with bullous pemphigoid: a multicenter randomized study.** *J Invest Dermatol.* 2009; **129**(7): 1681–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 67. Venning VA, Taghipour K, Mohd Mustapa MF, *et al.*: **British Association of Dermatologists' guidelines for the management of bullous pemphigoid 2012.** *Br J Dermatol.* 2012; **167**(6): 1200–14.
[PubMed Abstract](#) | [Publisher Full Text](#)
 68. **F** Williams HC, Wojnarowska F, Kirtschig G, *et al.*: **Doxycycline versus prednisolone as an initial treatment strategy for bullous pemphigoid: a pragmatic, non-inferiority, randomised controlled trial.** *Lancet.* 2017; **389**(10079): 1630–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 69. **F** Amagai M, Ikeda S, Hashimoto T, *et al.*: **A randomized double-blind trial of intravenous immunoglobulin for bullous pemphigoid.** *J Dermatol Sci.* 2017; **85**(2): 77–84.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 70. Kamaguchi M, Iwata H, Mori Y, *et al.*: **Anti-idiotypic Antibodies against BP-IgG Prevent Type XVII Collagen Depletion.** *Front Immunol.* 2017; **8**: 1669.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 71. **F** Cho YT, Chu CY, Wang LF: **First-line combination therapy with rituximab and corticosteroids provides a high complete remission rate in moderate-to-severe bullous pemphigoid.** *Br J Dermatol.* 2015; **173**(1): 302–4..
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 72. **F** Lamberts A, Euverman HI, Terra JB, *et al.*: **Effectiveness and Safety of Rituximab in Recalcitrant Pemphigoid Diseases.** *Front Immunol.* 2018; **9**: 248.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 73. **F** Yayli S, Pelivani N, Beltraminelli H, *et al.*: **Detection of linear IgE deposits in bullous pemphigoid and mucous membrane pemphigoid: a useful clue for diagnosis.** *Br J Dermatol.* 2011; **165**(5): 1133–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 74. Moriuchi R, Nishie W, Ujiie H, *et al.*: **In vivo analysis of IgE autoantibodies in bullous pemphigoid: a study of 100 cases.** *J Dermatol Sci.* 2015; **78**(1): 21–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
 75. **F** Van Beek N, Lüttmann N, Huebner F, *et al.*: **Correlation of Serum Levels of IgE Autoantibodies Against BP180 With Bullous Pemphigoid Disease Activity.** *JAMA Dermatol.* 2017; **153**(1): 30–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 76. **F** Freire PC, Muñoz CH, Stingl G: **IgE autoreactivity in bullous pemphigoid: eosinophils and mast cells as major targets of pathogenic immune reactants.** *Br J Dermatol.* 2017; **177**(6): 1644–53.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 77. **F** Fairley JA, Baum CL, Brandt DS, *et al.*: **Pathogenicity of IgE in autoimmunity: successful treatment of bullous pemphigoid with omalizumab.** *J Allergy Clin Immunol.* 2009; **123**(3): 704–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 78. **F** Balakirski G, Alkhateeb A, Merk HF, *et al.*: **Successful treatment of bullous pemphigoid with omalizumab as corticosteroid-sparing agent: report of two cases and review of literature.** *J Eur Acad Dermatol Venereol.* 2016; **30**(10): 1778–82.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. Soner Uzun

Department of Dermatology and Venereology, Akdeniz University, Antalya, Turkey

Competing Interests: Soner Uzun has published with Jun Yamagami once in the last 3 years on an international consensus paper on the diagnosis and management of pemphigus (doi: <http://dx.doi.org/10.1016/j.jaad.2018.02.021>).

2. Ralf Ludwig

Department of Dermatology, University of Lübeck, Lübeck, Germany

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research