REVIEW

Recent advances in understanding grasslands [version 1; peer review: 2 approved]

Carly J. Stevens

Lancaster Environment Centre, Lancaster University, Lancaster, UK

V1 First published: 30 Aug 2018, 7(F1000 Faculty Rev):1363
<https://doi.org/10.12688/f1000research.15050.1>

Latest published: 30 Aug 2018, 7(F1000 Faculty Rev):1363
<https://doi.org/10.12688/f1000research.15050.1>

Abstract

Grasslands are a vitally important ecosystem, supporting a wide range of ecosystem services and high levels of biodiversity. As a consequence, they have long been a focus for ecologists, playing host to some of the world's longest-running ecological experiments and providing the inspiration for many long-standing theories and debates. Because the field of grassland ecology is broad, encompassing many areas of ecology, this article picks some areas of particular debate and development to look at recent advances. The areas include relationships between diversity and productivity, ecosystem stability and ecosystem service provision, global change threats from nutrient addition, invasive species, climate change, and plant soil interactions.

Keywords

Climate change, diversity, ecosystem services, ecosystem stability, grassland ecology, invasive species, nutrient addition, plant-soil interactions.

Open Peer Review

Approval Status

	1	2
version 1
30 Aug 2018		

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Eric Allan**, University of Bern, Bern, Switzerland
2. **Till Kleinebecker**, Gießen University, Gießen, Germany

Any comments on the article can be found at the end of the article.

Corresponding author: Carly J. Stevens (c.stevens@lancaster.ac.uk)

Author roles: Stevens CJ: Conceptualization, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2018 Stevens CJ. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Stevens CJ. **Recent advances in understanding grasslands [version 1; peer review: 2 approved]**
F1000Research 2018, 7(F1000 Faculty Rev):1363 <https://doi.org/10.12688/f1000research.15050.1>

First published: 30 Aug 2018, 7(F1000 Faculty Rev):1363 <https://doi.org/10.12688/f1000research.15050.1>

Introduction

Grasslands cover 40% of the world's terrestrial surface¹, and they are found on all continents except Antarctica, in a wide range of climates, and on a wide range of soil types. They can vary in their species richness from monocultures up to botanically species-rich habitats which support diverse animal communities. Grasslands are typically dominated by grasses (*Poaceae*) and other grass-like plants. Some grasslands occur naturally, whilst others must be maintained by active management such as cutting or grazing. They are also incredibly important to mankind, providing many different services. Unsurprisingly, given their importance, extent, and variation, grasslands have been a focus for many ecologists and the home of many ecological theories, some of which remain intensely debated. In this article, I will highlight some of the recent advances in understanding grasslands. The field of grassland ecology is broad and developing rapidly. A Web of Science search with grasslands as a key word reveals almost 9,000 papers published between January 2016 and May 2018. As a consequence, this is not an exhaustive review but rather focusses on some key causes and consequences of biodiversity declines in grasslands, picking out some areas of key developments, controversies, and findings that I believe have helped us to advance our understanding of how these complex, intriguing, and beautiful ecosystems work.

Variation in diversity between grasslands

Productivity–diversity relationships

Grasslands across the world vary hugely in both physical and biological characteristics, and explaining relationships between them has led to much discussion. One of the fiercest ecological debates of recent decades has concerned the relationship between productivity and diversity. The humpbacked model² predicts that in extreme environments only a few stress-tolerant species survive and diversity is low. Both productivity and diversity increase until, when resource levels are high, diversity declines again, most likely due to competition or species pools. Grasslands account for as much as one-third of the net primary production on land³, and many of the original studies exploring the relationship between productivity and diversity originated in grasslands^{2,4}. In 2011, Adler *et al.*⁵ published a paper in *Science* that reignited the debate. Using a global dataset of 48 grasslands on five continents that were part of the Nutrient Network (NutNet), they found no consistent relationship between productivity and richness. Since then, we have seen publications that fall on both sides of the debate. Recently, Fraser *et al.*⁶ collected data from 30 sites on six continents and performed a similar analysis to that employed by Adler *et al.*⁵. They found strong support for the humpbacked model, with 19 of 28 sites showing significant concave-down quadratic relationships between plant species richness and productivity⁶. They identified a number of hypotheses for why they found much stronger support for the humpbacked model than did Adler *et al.* Whilst the debate regarding whether there is a single hypothesis that can explain the relationship between plant diversity and productivity will likely rage for a long time, most authors can agree that there is a need to develop our understanding of the mechanisms that underlie the

relationships identified. Grace *et al.*⁷ recently made progress toward this objective by showing that integrative modelling, considering multiple potential drivers of both richness and diversity, has substantially higher explanatory power than bivariate analyses, arguing for more integration of ideas and simultaneous tests of their combined implications.

Threats to grasslands

Land-use change

Globally, land-use change is a major driver of community change and habitat loss. The impacts of land-use change can be seen across trophic groups. A recent paper has taken a functional trait approach to examine shifts in invertebrate communities in response to land-use intensification. Working across 124 grasslands of differing intensities of land-use in Germany, Simons *et al.*⁸ collected data on a range of traits in insect and spider species to demonstrate that higher-intensity landscapes favoured smaller, more mobile, and less-specialised species. The collection of functional trait data for invertebrates is labour intensive, but the authors argue it is essential for understanding the impacts of land-use change on invertebrates.

Response to nutrient additions

One of the most challenging issues we currently face is the extent to which we have perturbed nutrient cycles and the impact this is having on the environment. High levels of nutrient addition have long been recognised to reduce species richness. We expect that nutrient enrichment results in a switch from belowground competition for nutrients to aboveground competition for light. Because some plants are taller, they receive more light per unit size than do smaller plants, thus precipitating competitive exclusion. Up to now, there has been limited evidence to conclusively demonstrate this mechanism, but a recent study demonstrated that an increase in light asymmetry is the main cause of species loss under nutrient enrichment⁹. DeMalach *et al.*⁹ used a combination of light measurements through the grassland canopy and plant height in fertilised and unfertilised grasslands to calculate light asymmetry and determine the competitive effect, demonstrating that it is an increase in the rate of light decay through the canopy rather than an increase in canopy height that is responsible for the competitive effect of grasses on forbs.

Another recent advance has been in the increased recognition of multiple factors limiting production in grasslands. Two recent papers from the NutNet^{10–12} have demonstrated that, contrary to popular opinion, where nitrogen (or nitrogen and phosphorus) is deemed a key determinant of aboveground net primary productivity, other nutrients are important in determining production, and not only were many grasslands limited by multiple nutrients¹⁰ but the number of added nutrients predicted diversity loss. Adding nutrients reduced niche dimensionality, increased productivity, and increased compositional turnover¹¹. Nutrient addition is clearly a considerable threat to grassland biodiversity, yet in many parts of the world it does not receive sufficient attention in policy or research, meaning there are many knowledge gaps we need to address.

Invasive species

There are many different mechanisms that have been identified for the success and spread of invasive species in grasslands and other habitats. A recent paper by Broadbent *et al.*¹³ highlighted the potential importance of root competition in the interactions between invasive and native grasses in New Zealand. This small-scale study only investigated the relationship between three species but demonstrated the importance of belowground competition, a mechanism that has received very little attention, in driving their interactions and highlights this as an area in need of future research. In contrast, a large meta-analysis was performed by Liu *et al.*¹⁴ to test whether invasive species benefit more from global environmental change than do native species. They compiled a database of published studies that gave performance measures for 74 invasive plant species and 117 native plant species in response to global change drivers. They found that invasive species responded more strongly to carbon dioxide enrichment and elevated temperature (and did not respond significantly to nitrogen deposition and increased precipitation), indicating that the problems caused by invasive plant species are likely to get worse under a changing climate. This study suggested that drought may be detrimental for invasive species, as did a seedbank study in Californian annual grasslands, which found that seeds of exotic annual grasses declined whilst native annual forbs increased¹⁵. Invasive species have very large impacts on grassland communities in some parts of the world, meaning that understanding and predicting these impacts is a priority.

Climate change

Long-term experiments are critically important in ecology¹⁶, and as grassland ecologists we are lucky to host some of the longest-running experiments in the world, including the 150-year-old Park Grass Experiment¹⁷. For climate change research, long-term experiments are especially valuable, as it takes time for plant species to respond. One of the longest-running climate change experiments is the Buxton Climate Change Impacts Laboratory (BCCIL) in northern England¹⁸. Recent research at BCCIL has demonstrated that climate change (warming and rainfall manipulation) has rapid direct impacts on the soil microbial community but also indirect impacts mediated by changes in plant species composition, which occur over longer time scales¹⁹.

Shifts in species composition are likely as a result of climate change but do not necessarily result in changes to ecosystem stability²⁰. Also, using a long-term investigation, Reich *et al.*²¹ showed that whilst in the first 12 years of carbon dioxide enrichment C₃ plant biomass increased markedly, C₄ plant biomass did not. This is expected, since C₄ plants are thought to be less limited by carbon dioxide; however, in the latter 8 years of the experiment, the responses switched, with biomass depressed in C₃ plants but not in C₄ plants. Fay *et al.*²² demonstrated that carbon dioxide concentrations have a strong effect on flowering in four of the five grassland species they investigated. They utilised contrasting soil conditions to demonstrate that impacts were mediated by productivity and nutrient status. The impacts of climate change will be felt globally, but

we are only just beginning to understand likely impacts. In the future, we need more research to understand not only likely impacts of climate change but also how climate change and other global change drivers are likely to interact; long-term experiments will be key to achieving this.

Effects of biodiversity loss

Diversity–stability relationships

One of the main arguments for biodiversity conservation is that more-diverse communities will be more stable and better able to resist perturbation. A number of mechanisms have been suggested for this relationship between diversity and stability, including asynchrony (productivity of one species increases, compensating for declines in another species), portfolio effects (statistical averaging of fluctuations in species properties), and functional redundancy (species loss is compensated by other species fulfilling a similar function). A recent study of 2,671 species from 300 plots, across three regions in Germany, indicated that asynchrony was the primary driver of stability rather than diversity alone²³. Evidence for this is mixed, but several recent papers, including two synthesis studies combining results from multiple experiments in grasslands, have provided strong evidence in support of the argument that diversity begets stability. Hautier *et al.*²⁴ used results from 12 multi-year experiments to show that changes in biodiversity caused by a range of environmental change drivers were a major factor in determining the impact on stability, whilst Isbell *et al.*²⁵ used data from 46 experiments that manipulated grassland diversity to show that diversity increased resistance to climate events. Further results from the NutNet collaborative experiment have shown that eutrophication weakens the relationship between diversity and stability. We would expect this to occur as a result of diversity losses, but in this case it was actually due to an increase in temporal variability of productivity²⁶.

Diversity–service provision relationships

Another often-cited negative consequence of the loss of biodiversity is that more-diverse systems support more ecosystem services. Many of the early papers in this field were conducted in grasslands, and now there is a move towards testing these relationships in the “real-world” in natural or managed grasslands. Allan *et al.*²⁷ provided strong evidence to support this, using 150 grassland plots spread across three regions of Germany as part of the Biodiversity Exploratories experiment. Their results show that diversity loss and functional composition change, caused by land-use intensification, is just as important as the land-use intensification itself in terms of its impact on ecosystem service delivery. It is not just richness that is important though: Winfree *et al.*²⁸ compared composition, richness, and abundance of pollinators to determine their relative contributions to pollination services and found that the abundance of dominant species was the most important factor whereas richness was relatively unimportant because most of the species were not responsible for service delivery. Whilst this is interesting evidence in favour of considering the composition of communities, this study was focussed on one trophic group and one ecosystem service; other recent studies have demonstrated the need for multiple trophic groups to

be considered. Results also from the Biodiversity Exploratories project demonstrated that high richness in multiple trophic groups had stronger positive effects on ecosystem service provision than did richness in any single group. This was particularly true for cultural and regulating services²⁹ and for the provision of multiple ecosystem services³⁰.

Conclusions

Grassland ecology, and the broader field of ecology, are rapidly moving fields. Developing analytical and statistical techniques combined with innovative approaches and co-ordinated networks are allowing us to address questions which we were not previously able to. However, many questions remain, and there are rarely, if ever, definitive answers to questions in ecology. Luckily, in much of grassland ecology, there is a willingness to embrace change and accept that rules are there to be broken. There are, however, unprecedented threats to

grassland habitats through climate change, nutrient deposition, invasive species, and habitat loss, to name a few, so the need to understand impacts and protect valuable habitats is more pressing than ever.

Abbreviations

BCCIL, Buxton Climate Change Impacts Laboratory; NutNet, Nutrient Network

Grant information

The author(s) declared that no grants were involved in supporting this work.

Acknowledgements

I would like to acknowledge the helpful input of reviewers Eric Allan, Till Kleinebecker, and Jin-Sheng He.

References

- Blair J, Nippert J, Briggs J: **Grassland Ecology**. In *Ecology and the Environment* (ed R. Monson) (Springer) 2014; 389–423.
[Publisher Full Text](#)
- Grime JP: **Competitive Exclusion in Herbaceous Vegetation**. *Nature*. 1973; **242**: 344–7.
[Publisher Full Text](#)
- Vitousek PM: **Grassland ecology: Complexity of nutrient constraints**. *Nat Plants*. 2015; **1**: 15098.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Tilman D: **Resource competition and community structure**. *Monogr Popul Biol*. 1982; **17**: 1–296.
[PubMed Abstract](#)
- Adler PB, Seabloom EW, Borer ET, *et al.*: **Productivity is a poor predictor of plant species richness**. *Science*. 2011; **333**(6050): 1750–3.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Fraser LH, Pither J, Jentsch A, *et al.*: **Plant ecology. Worldwide evidence of a unimodal relationship between productivity and plant species richness**. *Science*. 2015; **349**(6245): 302–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Grace JB, Anderson TM, Seabloom EW, *et al.*: **Integrative modelling reveals mechanisms linking productivity and plant species richness**. *Nature*. 2016; **529**(7586): 390–3.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Simons NK, Weisser WW, Gossner MM: **Multi-taxa approach shows consistent shifts in arthropod functional traits along grassland land-use intensity gradient**. *Ecology*. 2016; **97**(3): 754–64.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- DeMalach N, Zaady E, Kadmon R: **Light asymmetry explains the effect of nutrient enrichment on grassland diversity**. *Ecol Lett*. 2017; **20**(1): 60–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Fay PA, Prober SM, Harpole WS, *et al.*: **Grassland productivity limited by multiple nutrients**. *Nat Plants*. 2015; **1**: 15080.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Harpole WS, Sullivan LL, Lind EM, *et al.*: **Addition of multiple limiting resources reduces grassland diversity**. *Nature*. 2016; **537**(7618): 93–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Borer ET, Grace JB, Harpole WS, *et al.*: **A decade of insights into grassland ecosystem responses to global environmental change**. *Nat Ecol Evol*. 2017; **1**(5): 118.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Broadbent A, Stevens CJ, Peltzer DA, *et al.*: **Belowground competition drives invasive plant impact on native species regardless of nitrogen availability**. *Oecologia*. 2018; **186**(2): 577–87.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Liu Y, Oduor AMO, Zhang Z, *et al.*: **Do invasive alien plants benefit more from global environmental change than native plants?** *Glob Chang Biol*. 2017; **23**(8): 3363–70.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- LaForgia ML, Spasojevic MJ, Case EJ, *et al.*: **Seed banks of native forbs, but not exotic grasses, increase during extreme drought**. *Ecology*. 2018; **99**(4): 896–903.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Silvertown J, Tallowin J, Stevens C, *et al.*: **Environmental myopia: a diagnosis and a remedy**. *Trends Ecol Evol*. 2010; **25**(10): 556–61.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Silvertown J, Poulton P, Johnston E, *et al.*: **The Park Grass Experiment 1856–2006: Its contribution to ecology**. *J Ecol*. 2006; **94**(4): 801–14.
[Publisher Full Text](#)
- Grime JP, Brown VK, Thompson K, *et al.*: **The response of two contrasting limestone grasslands to simulated climate change**. *Science*. 2000; **289**(5480): 762–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Sayer EJ, Oliver AE, Fridley JD, *et al.*: **Links between soil microbial communities and plant traits in a species-rich grassland under long-term climate change**. *Ecol Evol*. 2017; **7**(3): 855–62.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Liu H, Mi Z, Lin L, *et al.*: **Shifting plant species composition in response to climate change stabilizes grassland primary production**. *Proc Natl Acad Sci U S A*. 2018; **115**(16): 4051–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Reich PB, Hobbie SE, Lee TD, *et al.*: **Unexpected reversal of C₃ versus C₄ grass response to elevated CO₂ during a 20-year field experiment**. *Science*. 2018; **360**(6386): 317–20.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Fay PA, Aspinwall MJ, Collins HP, *et al.*: **Flowering in grassland predicted by CO₂ and resource effects on species aboveground biomass**. *Glob Chang Biol*. 2018; **24**(4): 1771–81.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Blüthgen N, Simons NK, Jung K, *et al.*: **Land use imperils plant and animal community stability through changes in asynchrony rather than diversity**. *Nat Commun*. 2016; **7**: 10697.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Hautier Y, Tilman D, Isbell F, *et al.*: **Plant ecology. Anthropogenic environmental changes affect ecosystem stability via biodiversity**. *Science*. 2015; **348**(6232): 336–40.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Isbell F, Craven D, Connolly J, *et al.*: **Biodiversity increases the resistance of ecosystem productivity to climate extremes**. *Nature*. 2015; **526**(7574): 574–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Hautier Y, Seabloom EW, Borer ET, *et al.*: **Eutrophication weakens stabilizing**

- effects of diversity in natural grasslands.** *Nature*. 2014; **508**(7497): 521–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
27. Allan E, Manning P, Alt F, *et al.*: **Land use intensification alters ecosystem multifunctionality via loss of biodiversity and changes to functional composition.** *Ecol Lett*. 2015; **18**(8): 834–43.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 28. Winfree R, Fox JW, Williams NM, *et al.*: **Abundance of common species, not species richness, drives delivery of a real-world ecosystem service.** *Ecol Lett*. 2015; **18**(7): 626–35.
[PubMed Abstract](#) | [Publisher Full Text](#)
 29. Millenium Ecosystem Assessment: **Ecosystems and Human Well-being: Synthesis.** (Island Press), 2005.
[Reference Source](#)
 30. Soliveres S, van der Plas F, Manning P, *et al.*: **Biodiversity at multiple trophic levels is needed for ecosystem multifunctionality.** *Nature*. 2016; **536**(7617): 456–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. Till Kleinebecker

Division of Landscape Ecology and Landscape Planning, Gießen University, Gießen, Germany

Competing Interests: No competing interests were disclosed.

2. Eric Allan

Institute of Plant Sciences, University of Bern, Bern, Switzerland

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research