Check for updates

RESEARCH ARTICLE

Soil is a key factor influencing gut microbiota and its effect is comparable to that exerted by diet for mice [version 1; peer review: 1 approved, 1 approved with reservations]

Dongrui Zhou^{1,2}, Zhimao Bai³, Honglin Zhang⁴, Na Li¹, Zhiyu Bai¹, Fudong Cheng⁵, Haitao Jiang⁴, Chuanbin Mao⁶, Xiao Sun⁵, Zuhong Lu⁵

¹Key Laboratory of Child Development and Learning Science (Ministry of Education), School of Biological Science & Medical Engineering, Southeast University, Nanjing, China

²Suzhou Research Institute, Southeast University, Suzhou, China

³Key Laboratory of Environmental Medicine Engineering of Ministry of Education, School of Public Health, Southeast University, Nanjing, China

⁴College of Food Science, Nanjing Xiaozhuang University, Nanjing, China

⁵State Key Laboratory of Bioelectronics, Southeast University, Nanjing, China

⁶Department of Chemistry and Biochemistry, University of Oklahoma, Norman, OK, USA

V1 First published: 01 Oct 2018, 7:1588
<https://doi.org/10.12688/f1000research.15297.1>

Latest published: 01 Oct 2018, 7:1588
<https://doi.org/10.12688/f1000research.15297.1>

Abstract

Exposure to an unsanitary environment increases the diversity and alters the composition of gut microbiota. To identify the key element in the unsanitary environment responsible for this phenomenon, we investigated the effect and the extent by which the soil in our environment influenced the composition of gut microbiota. Results show that adding unsterile or sterile soil to bedding, either before birth or after weaning, influences significantly the composition of mice gut microbiota. Specifically, unsterile soil increases the richness and biodiversity of gut microbiota. Interestingly, based on UniFrac distance analysis of 16S rRNA sequences, the impact of soil on gut microbiota is comparable to that exerted by diet. These findings provide a potential new strategy for intervening on the human gut microbial community and preventing disease.

Keywords

gut microbiota, hygiene hypothesis, factors influencing gut microbiota, 16S rRNA sequencing

Open Peer Review

Approval Status

	1	2
version 1
01 Oct 2018	view	view

1. **Xuegong Zhang** , Tsinghua University, Beijing, China

2. **Jose F. Garcia-Mazcorro** , Universidad Autónoma de Nuevo León, San Nicolás de los Garza, Mexico

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Dongrui Zhou (junbai1013@seu.edu.cn)

Author roles: **Zhou D:** Formal Analysis, Funding Acquisition, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing; **Bai Z:** Data Curation, Formal Analysis, Software; **Zhang H:** Methodology; **Li N:** Methodology; **Bai Z:** Methodology; **Cheng F:** Software; **Jiang H:** Methodology; **Mao C:** Writing – Review & Editing; **Sun X:** Funding Acquisition; **Lu Z:** Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by The National Natural Science Foundation of China (Grants No. 31770540 and 61472078).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2018 Zhou D *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Zhou D, Bai Z, Zhang H *et al.* **Soil is a key factor influencing gut microbiota and its effect is comparable to that exerted by diet for mice [version 1; peer review: 1 approved, 1 approved with reservations]** F1000Research 2018, 7:1588 <https://doi.org/10.12688/f1000research.15297.1>

First published: 01 Oct 2018, 7:1588 <https://doi.org/10.12688/f1000research.15297.1>

Introduction

The relationship between human gut microbiota and certain diseases has become increasingly apparent^{1,2}. The main contributing factors are delivery mode³⁻⁵, age^{6,7}, antibiotic treatment^{4,6}, diet, and the living environment. They affect gut microbiota to different degrees. Among these factors, diet is the most widely studied one, as it modulates the composition and then changes the function of the microbial community in humans⁸⁻¹⁵. The microbiota of malnourished children was shown to impair growth and cause metabolic abnormalities in the brain and other organs of recipient gnotobiotic mice^{16,17}. By altering the gut microbiota, diet also contributes to other chronic illnesses, such as obesity^{1,2}, cardiovascular diseases¹⁸, and autism¹⁹. The human gut microbiota responds rapidly to dietary changes^{8,20}, even though long-term dietary habits remain the dominant force in determining the composition of an individual's gut microbiota^{10,15,21,22}.

Following what is known as the hygiene hypothesis, an overly clean modern life leads to immune dysfunction and more allergic diseases^{23,24}. Many studies have shown large differences in the human gut microbiome between populations in rural areas across the world and those in Europe or America^{7,9,25}. Our early research suggested that the cleanness of a living environment substantially altered the composition of gut microbiota in mice²⁶. After analyzing the elements in bedding material, it was first proposed that microbes might play a key role in changing the composition of mouse gut microbiota²⁷. However, a follow-up study showed that adding microbes to the bedding had a limited effect on the composition of the dominant genera²⁸. Thus, the unsanitary environment factors that act on gut microbial communities remain to be identified.

Unsanitary environments are rich in surface soil. This hosts a large number of microbes and is rich in nutrients. At present, it is unclear what effect the soil in our environment exerts on the composition of our gut microbiota, to what extent it causes such alterations, and what the time-scale of any changes may be. To understand these interrelated issues, we raised C57 mice on four different beddings. Groups were designed to explore the effect of sterile and unsterile soil, the extent by which soil compared with normal diet or high-fat diet (formula shown in [Supplementary Table 1](#)), as well as the timing of exposure to soil (before birth or after weaning) on the gut bacterial community ([Supplementary Figure 1](#)).

Results

Soil-associated changes in gut microbiota at phylum level

We analyzed the bacterial community composition using MiSeq sequencing targeted to the 16S rRNA gene in up to 180 fecal samples ([Supplementary Table 2](#)). For each sample, we PCR-amplified the V4 hypervariable region using the 515F-907R primer set²⁹. The sequencing data set consisted of 10,937,871 high-quality, classifiable 16S rRNA gene sequences, with at least 45,816 sequences per sample ([Supplementary Table 2](#)).

On the basis of the obtained sequences, we discovered that sterile and unsterile soil significantly altered gut bacterial

community composition ([Figure 1](#) and [Supplementary Table 3](#)) in a diet-dependent manner. Mice fed a normal diet and living on sterile soil after weaning exhibited significantly more Bacteroidetes ($P < 0.001$) and fewer Firmicutes ($P < 0.001$) ([Figure 1A](#) and [Supplementary Table 3a](#)). Unsterile soil added before birth also significantly increased the number of Bacteroidetes ($P = 0.032$). Instead, mice fed a high-fat diet and raised on bedding with unsterile soil added before birth or after weaning showed significantly more Actinobacteria ($P < 0.001$) ([Figure 1B](#) and [Supplementary Table 3b](#)).

Microbes in the environment were reported to affect the colonization of the intestinal microflora in newborns³⁰. Here, we found that microbes and soil in the bedding could change the composition of gut microbiota after mice were weaned. When mice were fed a normal diet and lived on bedding with unsterile soil, they consistently showed more Bacteroidetes ($P < 0.001$) and fewer Firmicutes ($P < 0.001$) ([Figure 1E](#) and [Supplementary Table 4a](#)) during the first 3 weeks. After the third week, the abundance of these two phyla returned to the level from the previous week. When fed a high-fat diet and provided with bedding containing unsterile soil, mice showed an increased abundance of Actinobacteria on the third week of treatment ($P = 0.030$), becoming highest on the fourth week ($P < 0.001$) ([Figure 1F](#) and [Supplementary Table 4b](#)). In summary, we observed that the gut microbiome could respond rapidly to alterations in the living environment and diet, potentially facilitating adaptation to a variety of lifestyles.

Soil-associated changes in gut microbial diversity

We further discovered that unsterile soil could increase microbial diversity, particularly for mice on a normal diet, whether it was added before birth or after weaning ([Figure 2](#), [Supplementary Figure 3](#), and [Supplementary Table 5](#), [Supplementary Table 6](#)). By comparison, microbial diversity was unaffected by sterile soil in mice fed a normal diet and decreased in those on a high-fat diet ([Supplementary Figure 3](#) and [Supplementary Table 5](#)). Earlier, we found that merely adding environmental microbes to the bedding could increase diversity but had little effect on the composition of dominant bacteria in mice gut microbiota²⁸. Hence, microbes appear to contribute mainly to microbial diversity, whereas soil may alter microbial community structure.

Soil-associated changes in gut microbiota differ between male and female mice

An epidemiological survey has shown that immune system diseases, such as asthma and wheeze, display a skewed sex bias towards males³¹. Here, we found that abundance of the phyla Bacteroidetes and Firmicutes was related to gender, except in mice with sterile bedding with unsterile soil added prior to birth (DE group) exposed to unsterile soil bedding from birth ([Supplementary Figure 4](#), [Supplementary Figure 5](#) and [Supplementary Table 7](#)). Bacteroidetes were more abundant in males than in females of the mice raised on specific-pathogen-free bedding (SPF group) fed a normal diet ($P = 0.019$). By contrast, Firmicutes were more abundant in males than in females in the mice raised on sterile grade murine bedding with dirty unsterile soil (6:11 (w/w)) added after weaning (SD group) on a high-fat

Figure 1. Changes in the relative abundance of Actinobacteria, Bacteroidetes, and Firmicutes in fecal samples from mice treated with different living beddings and fed a normal or high-fat diet. Values from all available fecal samples were averaged ($n = 8, 9$, or 10 per treatment). (A,B) Relative abundance of bacteria on the eighth week in fecal samples of mice raised on specific-pathogen-free grade (SPF) sterile bedding (control), sterile bedding with sterile soil added after weaning (SS), sterile bedding with dirty unsterile soil added after weaning (SD), and sterile bedding with unsterile soil added prior to birth (DE) fed (A) a normal diet (ND) or (B) a high-fat (HF) diet. (C–F) Relative bacterial abundance in samples collected on the fourth (W4), fifth (W5), seventh (W7), and eighth (W8) week from mice of the (C) SPF-ND group, (D) SPF-HF group, (E) SD-ND group, and (F) SD-HF group.

Figure 2. Representation of α diversity in gut microbiota based on fecal samples collected from specific-pathogen-free (SPF) (blue curve) and sterile bedding with dirty unsterile soil added after weaning (SD) (red curve) groups over the fourth to the eighth week of age. (A–C) SPF and SD groups fed a normal diet: (A) operational taxonomic units (OTUs), (B) Chao 1 estimates, and (C) Shannon index. (D–F) SPF and SD groups fed a high-fat diet: (D) OTUs, (E) Chao 1 estimates, and (F) Shannon index. Values from all available fecal samples were averaged ($n = 8–10$; $*P < 0.05$, $**P < 0.01$, based on a two-tailed least significant difference test).

diet ($P = 0.042$) (Supplementary Figure 5 and Supplementary Table 7).

Soil-associated changes in gut microbiota at operational taxonomic unit (OTU) level

Random Forests is a supervised machine-learning technique that can classify a sample by estimating the importance of OTUs at species level according to their relative abundance and sample probability³². We found that distinct community signatures existed between any two treatments (Supplementary Table 8–Supplementary Table 10). For mice on a high-fat diet, there were 79 predictive species-level OTUs between the (SPF) and DE groups (baseline error = 0.44, cross-validation error = 0 ± 0), of which 63 were overrepresented in the DE group. Moreover, out of these 63, 25 and 20 were assigned to the classes Actinobacteria and Bacilli, respectively (Supplementary Table 8b). A comparison between SPF and SD groups gave similar results (Supplementary Table 8a); however, the decrease in beneficial bacteria was greater in the mice raised on sterile grade murine bedding with sterile soil (6:11 (w/w)) added after weaning (SS group) than in the SPF group (Supplementary Table 8c).

We further analyzed differences in the composition of gut bacterial communities between any two groups of mice on a normal diet (Supplementary Table 9). Compared to the SPF group, DE and SD groups presented more Actinobacteria and Bacilli, whereas the SS group showed more Bacteroidia (Supplementary Table 9). Notably, bacteria of the Bacteroidales S24–7 family were more prevalent among SPF mice fed a high-fat diet than in those fed a normal diet. Instead, the latter had more bacteria of the families Lachnospiraceae and Ruminococcaceae (Supplementary Table 10a). The DE group fed a high-fat diet showed more Actinobacteria and Bacilli than animals fed a normal diet. On the contrary, the latter had more Bacteroidales S24–7 and Lachnospiraceae (Supplementary Table 10c). Thus, unsterile/sterile soil bedding affected the structure of mice gut microbial communities in a diet-dependent manner. The effect varied depending on whether i) mice were raised on bedding with soil added before birth or after weaning, ii) the bedding contained sterile or unsterile soil, and iii) mice were fed a normal or high-fat diet.

Gut microbial community profiles identify soil as a key determinant of changes to the mouse gut microbiome

We used UniFrac distances to analyze the 16S rRNA datasets and measure similarities among microbial communities. The distance between mice exposed to two different living environments and fed a high-fat diet was similar to that between mice using the same bedding, and fed either a normal or a high-fat diet (Figure 3 and Supplementary Table 11). The distance between SPF groups fed a normal or a high-fat diet did not differ significantly from that between SPF and SS, and between SD and DE mice fed a high-fat diet. Indeed, these groups were among the least distant. Distances between mice maintained on different beddings and a high-fat diet were greater than those between SPF groups fed different diets. Thus, soil appears to influence the composition of the gut microbial community to the same extent as diet. In addition, a change in living environment

Figure 3. Differences in gut microbial communities between treatments. For mice fed a normal diet, the shortest UniFrac distance was between specific-pathogen-free (SPF) and sterile bedding with unsterile soil added prior to birth (DE) groups, and the longest one was between sterile bedding with dirty unsterile soil added after weaning (SD) and sterile bedding with sterile soil added after weaning (SS) groups; comparisons between SPF and SD, and between DE and SS showed no significant differences. For mice fed a high-fat diet, the shortest distances were SPF–SS and SD–DE, with no significant difference between them; the longest distances were SD–SS and DE–SS; SPF–SD was significantly longer than SPF–DE ($n = 8–10$; $*P < 0.05$, $**P < 0.01$, based on a two-tailed least significant difference test).

also had an important effect on gut microbiota. When fed a normal diet, microbiota in the gut of SPF and DE mice were more similar than those of other groups (Figure 3 and Table 11).

Unsupervised clustering using PCoA of UniFrac distance matrices confirmed that living environments and diets explained the variation in the data set (Figure 4 and Supplementary Figure 6, Supplementary Figure 7). When fed the same diet, the gut microbial communities of mice from one living environment clustered apart from those of mice subjected to other treatments (Figure 4A, B). For two different living environments, the two high-fat diet groups showed a similar transfer distance as mice maintained on the same bedding but fed two different diets (Figure 4C, D and Supplementary Figure 6). For the SPF group fed a normal diet, almost no differences were observed among

Figure 4. Principal coordinates analysis (PCoA) of unweighted UniFrac distances for fecal microbiota sampled on the eighth week from mice on four types of bedding. Analysis was based on the Illumina bacterial 16S rRNA gene data set (V4 region). Mice from the four groups were fed a (A) normal or (B) high-fat diet. (C) Mice in the sterile bedding with dirty unsterile soil added after weaning (SD) and specific-pathogen-free (SPF) groups were fed a normal or high-fat diet. (D) Mice of the sterile bedding with sterile soil added after weaning (SS) and sterile bedding with unsterile soil added prior to birth (DE) groups were fed a normal or high-fat diet.

fecal samples collected at four different time points (Supplementary Figure 7A). These results confirm that the living environment exerts a great influence on the composition of gut microbiota. It should be noted that we did not find significant clustering with respect to cage or gender.

Discussion

In the present study, we identified differences in gut microbiota between mice living on sterile or unsterile soil and fed a normal or a high-fat diet. Sterile and unsterile soil substantially altered the composition of gut microbiota, with unsterile soil also increasing bacterial diversity. In addition, we demonstrated that unsterile soil added before birth or after weaning altered the gut microbiome, irrespective of whether mice were fed a normal or a high-fat diet. Hence, we suggest that soil is one of the key factors influencing gut microbiota and that its effect is comparable to that exerted by diet.

Mice have the habit of coprophagy and fur combing, through which they may swallow some soil. As a result of agriculture

and animal husbandry, our ancestors were also in a close and daily contact with soil. This practice is still common among infants (age 12–14 months), who crawl on grounds rich in soil and animal feces. During this period, they use their mouths to feel the world around them and experience a rapid development of the immune and nervous systems. Our results may explain in part some epidemic diseases occurring in modern life.

We show that sterile soil present in the surrounding environment has a role in shaping the composition and structure of gut microbiota. At birth, the gut is sterile and ambient microbes play a key role in shaping the bacterial community^{3–6,30,33}. Our early research showed that microbes in the environment had a limited capacity to influence the composition and structure of dominant bacterial communities in adult mice²⁸. Here, we show that sterile soil in the bedding significantly altered the composition and structure of gut microbial communities in weaned mice, but had little effect on their diversity and richness. By contrast, unsterile soil changed microbiota composition and structure, as well as its diversity and richness. Together, these studies

indicate that a healthy gut microbiota benefits from a close contact with soil and the microbes in it. It is possible that unsterile soil promotes the growth of certain bacteria, while inhibiting others, especially in mice fed a high-fat diet.

The most obvious effect of unsterile soil on the intestinal microflora was the predominance of the phylum Actinobacteria in mice fed a high-fat diet. Vatanen *et al.*³⁴ studied the development of gut microbiota in infants from birth until the age of 3 years and found that early-onset autoimmune diseases were common in Finland and Estonia but were less prevalent in Russia. Their results showed that Actinobacteria were more abundant among infants under 12 months in Russia than in Finland and Estonia; however, there was no difference after the age of 1 year³⁴. This might be explained by small infants having a diet rich in protein and fat, and those in Russia being in closer contact with unsterile soil.

The results also indicate how the timing of exposure to unsterile soil influences the formation of bacterial communities. Some studies reported that early environmental exposure exerted a sustained influence on the acquired gut microbiota, and that the early microbiota shaped the later one^{35–37}. Our results reveal that microbiota differed depending on whether mice were exposed to unsterile soil early or after weaning. It remains to be determined whether the difference between gut microbiota of mice that were exposed to soil before birth or after weaning influences the animals' health.

Our synergetic assessment of host soil and diet provides additional insight into the effect of diet on gut microbiota, and the possible effects on colorectal cancer and immune system diseases. We speculate that differences in bacterial flora mirror a different metabolite composition. Epidemiological studies indicate that red and processed meat intake is associated with an increased risk of colorectal cancer^{38,39} and diseases of the immune system⁴⁰. However, in pastoral areas, such as Tibet, where the local diet is rich in beef and mutton, mortality from cancer and childhood asthma is the lowest in China^{41,42}. Compared with a modern western lifestyle, the residents of Tibet are in close daily contact with soil. To this end, unsterile soil may alter microbiota composition and structure, increasing its diversity and richness, and affecting the metabolic processing of red meats. This, in turn, may enhance human immunity and resistance to chronic diseases; however, this mechanism needs to be confirmed via further research.

Our results provide further evidence in support of the hygiene hypothesis. In developed countries, human living environments and lifestyle have changed immensely during the last few decades. People spend ~92% of their time indoors⁴³ and there is very little soil in their living environment. At the same time, many microbes appear to have been lost from the modern human body^{7,9,25}, while the burden of chronic inflammatory diseases, including atopic diseases, has increased dramatically. Our results suggest that an important cause of these diseases is represented by the lack of contact with soil and soil microorganisms.

The cleanness of our living environment and diet could be readily modified. This approach may provide a new simple route to intervene on the human gut microbial community, as well as design specific diets aimed at disease prevention. A few open questions nevertheless remain. Further research will determine how the soil from the host environment alters human gut microbiota, whether through swallowing, body mucosae, or by other means. Similarly, it remains to be investigated whether different types of soil or soil microbes have different effects on intestinal microecology.

Methods

Animals and groups

We purchased 6-week-old male (n=1) and female (n=3) C57 mice from B & K Universal (Shanghai, China) for breeding. One male was mated with three females. After the females became pregnant, two of them were shifted to a separate cage. The newborn mice were weaned and when they were 4 weeks old they were moved to separate cages.

We selected 8 males and 24 females from among the weaned mice to carry out further experiments. Next, two males and six females were selected from different cages and transferred to room number 2. The cages were furnished with sterile grade murine bedding and unsterile soil (6:11 w/w bedding to soil). The remaining 6 males and 18 females were raised in room number 1, on sterile grade bedding. The bedding was changed once a week. The soil used in the experiments was from the top 10 cm of a farm ground with abundant goats, hens, and ducks.

When mice were 6 weeks old, they were bred with 1 male per 3 females. In room number 1, 30 males and 30 females, aged 23–37 days (weight, 8.6–21.4 g) the fourth week group, W4), were selected for experiments. They were randomly separated into three groups: the SPF group, raised on specific-pathogen-free bedding; the SD group, raised on sterile bedding with unsterile soil (6:11 (w/w)) added after weaning; and the SS group, raised on sterile bedding with sterile soil (6:11 (w/w)) after weaning. The SPF group served as the control. The randomization procedure was as follows. Considering gender matching, more than 80 mice were used in this experiment, and it was very difficult to breed so many mice in the same time. When we had about 20 mice with an age difference within about 10 days, we divided them into two groups, one fed with normal diet and the other with a high-fat diet. These 20 mice had been treated with same bedding. The other groups of 20 mice were treated with the second or third of type bedding. The breeding mice were uniformly distributed and randomly grouped. The SD group was transferred to room number 2 after weaning. Next, 10 males and 10 females, who were born in room number 2, aged 23–33 days (W4) were selected for experiments and raised on the same kind of bedding as before weaning (group DE). For all experiments, two or three mice were kept in a cage and the bedding was changed once a week.

For each group, 10 mice (five males and five females) were fed the same sterile grade commercial normal pellet diet as before

weaning, while 10 mice (five males and five females) were fed a sterile grade commercial high-fat pellet diet after weaning (Supplementary Table 1). All animals were fed the same sterile distilled water. They were allowed free access to water and food. All animals were housed in two specific-pathogen-free animal rooms with a 12-h light/12-h dark cycle at $24^{\circ}\text{C} \pm 2^{\circ}\text{C}$ and humidity of $40\% \pm 5\%$.

For SPF and SD groups, the first fecal samples were collected before mice were shifted to separate cages, and then on the fifth, seventh, and eighth week (Supplementary Figure 2). For DE and SS groups, fecal samples were collected on the eighth week (Supplementary Figure 2). After collection, fecal samples were put immediately into an ice box and stored at -80°C within 8 h.

Care and use of animals

Animal experiments were performed in strict accordance with the guidelines of the Animal Research Ethics Board of Southeast University. All experiments were approved by the Animal Care Research Advisory Committee of Southeast University and the National Institute of Biological Sciences (approval number: 2014063009). All efforts were made to alleviate the suffering of animals.

Specifically, mouse health was monitored every other day and body weight measurements were performed every week. Mouse health was assessed by observing changes in body weight or fecal shape, and external physical appearance. The animals would be euthanized to minimize pain and distress if they became severely ill over the experimental period, as when they showed one or more symptoms, such as 15–20% weight loss, diarrhea, loss of hair quality, pain (arched back and curled up posture), or listlessness for more than 1 week.

Euthanasia was performed as follows. A 1% (w/v) dose of sodium pentobarbital (50 mg/kg) was administered through an intraperitoneal injection using a hypodermic needle. When mice lost consciousness, they were killed by neck dislocation and death was confirmed. Two individuals were required to perform injections, one to hold the animal and the other to perform the procedure. Animals were not left unattended during the procedure. The ARRIVE reporting guidelines were followed during this study (Supplementary File 1)⁴⁴.

DNA extraction

Genomic DNA was extracted from fecal samples according to the protocol proposed by Zoetendal *et al.*⁴⁵. Each sample (40–60 mg) was weighed and put into 15 ml 1 × PBS (pH 7.4). Samples were re-suspended completely by vigorous shaking and subsequently centrifuged at $700 \times g$ at 4°C for 5 min. The supernatants were transferred to a 15-ml tube and centrifuged at $9000 \times g$ at 4°C for 5 min. The pellets were transferred to a 2-ml Eppendorf tube and re-suspended in 300 μl 10× TE buffer (10 mM Tris-HCl pH 8.0, and 1 mM EDTA pH 8.0). Lysozyme (100 μl , 200 mg/ml) was added to each tube, and incubated for 1 h at 37°C . Subsequently, 50 μl 10% SDS (w/v) and 20 μl proteinase K (20 mg/ml) were added to the samples and incubated for 2 h at 50°C . Next, 100 μl 5 M NaCl was added, followed by incubation at 65°C for 10 min. The samples were mixed

gently with an equal volume of Tris-Phenol (pH 8.0) for 5 min. The mixtures were centrifuged at $9000 \times g$ for 5 min. The upper layer was transferred to a new tube. This step was repeated twice, after which chloroform was used instead of Tris-Phenol to clean the samples one more time. Finally, a 1/10 volume of 3 M sodium acetate (pH 5.2) and two volumes of 95% ethanol were added. Samples were mixed gently and centrifuged at 4°C and $9000 \times g$ for 20 min. The precipitated genomic DNA was washed twice in 200 μl 70% ethanol. The dried DNA samples were re-suspended in 50 μl sterile double-distilled water. DNA quality was monitored by gel electrophoresis and spectrophotometry (260/280 nm). The samples were stored at -20°C .

MiSeq sequencing and data handling

The analysis of gut microbial communities was performed using MiSeq high-throughput sequencing of bacterial 16S ribosomal RNA (rRNA) genes. For each sample, the V4 hypervariable region was amplified using the 515F and 907R primer set, as described by Angenent *et al.*²⁹. PCR was performed with 300 ng microbial community DNA as a template, using 1 μl of Trans Start Fast Pfu Taq DNA Polymerase (TransGen Biotech, Beijing, China), 0.25 mM of each dNTP, 1× Trans Start Fast Pfu buffer, 0.2 μM forward primer, and 0.2 μM reverse primer, in a total reaction volume of 50 μl . The cycling conditions were: 94°C for 3 min followed by 27 cycles of 95°C for 45 s, 50°C for 45 s, and 72°C for 45 s, with a final extension at 72°C for 5 min. Each sample was amplified in duplicate, combined, purified, and then quantified using the QuantiFluor™-ST Handheld Fluorometer with UV/Blue Channels (Promega Corporation, Fitchburg, WI, USA). Sequencing of the PCR product libraries was performed on an Illumina MiSeq platform (Illumina Inc., San Diego, CA, USA) using 2 × 250 bp chemistry. We merged the raw paired-end reads using FLASH software version 1.2.7, set to a minimum overlap of 10 bp. Other parameters were set to default settings. PCR artifacts were removed by eliminating low-quality sequences using Trimmomatic version 0.30⁴⁶. Chimeric sequences were removed using Usearch version 7.1⁴⁷. We clustered the quality-checked sequences into *de novo* OTUs at a 97% similarity threshold using the QIIME software version 1.8.0 software package⁴⁸. The generated OTUs were classified using the RDP classifier software version 11.5 at a 70% confidence threshold for sequences longer than 200 bp⁴⁹.

Data analysis and statistical tests

All samples were rarefied down to 45,816 sequences per sample to prevent bias due to sampling depth. We used QIIME to calculate α diversity indices of Chao1 estimator and Shannon diversity/richness for each sample⁴⁸. We calculated the metrics of unweighted UniFrac distances between any two treatments and performed principal coordinates analysis (PCoA) through QIIME to examine dissimilarities in community composition. PCoA was used to compare groups of samples based on unweighted UniFrac distance metrics by plotting n samples in $(n - 1)$ -dimensional space.

Random Forests analysis in R version 3.2.3⁷ was performed with 500 trees and all default settings using 16S rRNA-based OTUs from the Illumina V4 data sets as described previously⁵⁰. We

used out-of-box (OOB) error to estimate the generalization error for all 16S rRNA comparisons involving all treatments. For each comparison, 100 relevant subsets of samples were extracted from the table of OTUs, and the average OOB error estimates and OTU importance estimates were calculated from subset samples. For a direct evaluation of the predictive strength of the OTUs, we compared generalization errors at various sequencing depths: the lowest observed depth of 45,816 sequences and at sequencing depths of 100, 1000, 10,000, and 40,000 reads per sample. The mean and standard deviation of the OOB error were estimated for each classification task using 100 independent rarefactions of the data. The expected 'baseline' error was obtained by a classifier that simply predicted the most common class label.

T- test and analysis of variance with post hoc Tukey's test, for comparison of two or more than two groups, respectively, were performed using SPSS software, version 18.0 (SPSS, Inc., Chicago, IL, USA).

Data availability

Metagenomic sequence data for each mouse are available from the Sequence Read Archive, accession number, PRJNA491246: <https://identifiers.org/insdc.sra/PRJNA491246>.

Grant information

This work was supported by The National Natural Science Foundation of China (Grants No. 31770540 and 61472078).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgments

We thank Liling Zhang, Zhencheng Xue, Shengqin Wang, Xiaoni Dai, Yue Hou, and Chunpeng He for technical assistance; and Xiaorong Wang for animal experiments.

Supplementary material

Supplementary Figure 1. Mice treatments.

Mice were raised on four types of bedding, each with two kinds of diet, resulting in four groups fed a normal diet (ND) and another four fed a high-fat diet (HF). The first group of mice was raised in room 1 in cages with sterile bedding of specific-pathogen-free grade (SPF) without soil (SPF-ND and SPF-HF) and served as controls. The second group was raised in cages with sterile bedding (same as the control) (SS-ND) and with sterile soil (6:11 (w/w) bedding to soil) added after weaning (SS-HF). The third group (SD-ND and SD-HF) was taken from room 1 to room 2 after weaning and then raised in cages with sterile bedding and unsterile soil (6:11 (w/w)). The last group (DE-ND and DE-HF) was born and raised in a cage with sterile bedding and unsterile soil (6:11 (w/w)).

[Click here to access the data.](#)

Supplementary Figure 2. Time-line indicating collection of fecal samples, bedding and diet conditions.

For the specific-pathogen-free, no soil (SPF) and sterile bedding, unsterile soil after weaning (SD) groups, fecal samples were collected four times. The first time was after mice were weaned at the age of 4 weeks (4W). For the SD group, the first sampling took place before addition of unsterile soil (and a high-fat diet). For the SPF-HF group, the first sampling occurred before exposure to a high-fat diet. The second, third, and fourth sampling times corresponded to the ages of five (5W), seven (7W), and eight (8W) weeks. For sterile bedding, unsterile soil (DE) and sterile bedding, sterile soil (SS) mice, samples were collected just once, at eight weeks. For the DE group, unsterile soil was added to the bedding before birth, for SD and SS mice it was added after weaning at the age of four weeks. All high-fat diet treatments were performed at the age of 4 weeks just after the mice were weaned.

[Click here to access the data.](#)

Supplementary Figure 3. Bacterial richness and diversity of fecal microbiota sampled on the eighth week: comparison between living bedding groups fed a normal or high-fat diet.

(A, B) Rarefaction curves of Chao1 estimators for mice fed (A) a normal or (B) a high-fat diet. (C, E, G) Mice fed a normal diet showing: (C) Chao1 estimators, (E) unique operational taxonomic units (OTUs), and (G) Shannon index. (D, F, H) Mice fed a high-fat diet showing: (D) Chao1 estimators, (F) unique OTUs, and (H) Shannon index. Values from all available samples were averaged. n = 8, 9, or 10 per treatment. *P < 0.05 and **P < 0.01 based on a two-tailed least significant difference test. SPF, sterile bedding/no soil; SD, sterile bedding/unsterile soil after weaning; DE, sterile bedding/unsterile soil before birth; SS, sterile bedding/sterile soil after weaning.

[Click here to access the data.](#)

Supplementary Figure 4. Changes in the relative abundance of the Bacteroidetes, Actinobacteria, and Firmicutes phyla for male and female mice fed a normal diet.

Values from all available samples were averaged (n = 3, 4, or 5 per treatment). (A) Female and (B) male mice of the sterile bedding/no soil (SPF), sterile bedding/sterile soil after weaning (SS), sterile bedding/unsterile soil after weaning (SD), and sterile bedding/unsterile soil before birth (DE) groups collected on the eighth week. (C to F) Samples collected on the fourth (W4), fifth (W5), seventh (W7), and eighth (W8) week: (C) female and (D) male mice of the SPF group; (E) female and (F) male mice of the SD group.

[Click here to access the data.](#)

Supplementary Figure 5. Changes in the relative abundance of the Bacteroidetes, Actinobacteria, and Firmicutes phyla for male and female mice fed a high-fat diet.

Values from all available samples were averaged (n was 3, 4, or 5 per treatment). (A) Female and (B) male mice of the sterile bedding/no soil (SPF), sterile bedding/sterile soil after weaning (SS), sterile bedding/unsterile soil after weaning (SD), and sterile bedding/unsterile soil before birth (DE) groups collected on the eighth week. (C to F) Samples collected on the fourth (W4), fifth (W5), seventh (W7), and eighth (W8) week: (C) female and (D) male mice of the SPF group; (E) female and (F) male mice of the SD group.

[Click here to access the data.](#)

Supplementary Figure 6. Principal coordinates analysis (PCoA) of unweighted UniFrac distances between fecal microbiota of mice from four living bedding conditions sampled on the eighth week.

The analysis was based on the V4 region of bacterial 16S rRNA gene sequencing data sets. The different treatments are color-coded. (A) sterile bedding/no soil (SPF) versus sterile bedding/unsterile soil before birth (DE) group; (B) SPF versus sterile bedding/sterile soil after weaning (SS) group; (C) sterile bedding/unsterile soil after weaning (SD) versus DE group; (D) SD versus SS group. All groups were fed either a normal diet (ND) or a high-fat diet (HF). n = 8, 9, or 10 per treatment.

[Click here to access the data.](#)

Supplementary Figure 7. PCoA of unweighted UniFrac distances between fecal microbiota of SPF and SD mice sampled over four weeks.

The analysis was based on the V4 region of bacterial 16S rRNA gene sequencing datasets. The different ages are color-coded. (A, B) sterile bedding/no soil (SPF) group fed a (A) normal or (B) high-fat diet; (C, D) SD group fed a (C) normal diet (CD) or (D) high-fat diet (HFD). n = 8, 9, or 10 per treatment.

[Click here to access the data.](#)

Supplementary Table 1. Diet formula.

[Click here to access the data.](#)

Supplementary Table 2. Sample information.

[Click here to access the data.](#)

Supplementary Table 3. Multiple phylum-level comparisons.

[Click here to access the data.](#)

Supplementary Table 4. Phylum abundance comparison.

[Click here to access the data.](#)

Supplementary Table 5. Multiple comparisons of diversity.

[Click here to access the data.](#)

Supplementary Table 6. Diversity index.

[Click here to access the data.](#)

Supplementary Table 7. Phylum abundance comparison between female and male mice.

[Click here to access the data.](#)

Supplementary Table 8. Random Forests for mice fed a high-fat diet.

[Click here to access the data.](#)

Supplementary Table 9. Random Forests for mice fed a normal diet.

[Click here to access the data.](#)

Supplementary Table 10. Random Forests for mice between diets.

[Click here to access the data.](#)

Supplementary Table 11. P-values of multiple comparisons for unweighted UniFrac distances.

[Click here to access the data.](#)

Supplementary File 1. Completed ARRIVE checklist.

[Click here to access the data.](#)

References

1. Ley RE, Turnbaugh PJ, Klein S, *et al.*: **Microbial ecology: human gut microbes associated with obesity.** *Nature.* 2006; **444**(7122): 1022–1023.
[PubMed Abstract](#) | [Publisher Full Text](#)
2. Turnbaugh PJ, Ley RE, Mahowald MA, *et al.*: **An obesity-associated gut microbiome with increased capacity for energy harvest.** *Nature.* 2006; **444**(7122): 1027–1031.
[PubMed Abstract](#) | [Publisher Full Text](#)
3. Dominguez-Bello MG, Costello EK, Contreras M, *et al.*: **Delivery mode shapes the acquisition and structure of the initial microbiota across multiple body habitats in newborns.** *Proc Natl Acad Sci U S A.* 2010; **107**(26): 11971–11975.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
4. Costello EK, Stagaman K, Dethlefsen L, *et al.*: **The application of ecological theory toward an understanding of the human microbiome.** *Science.* 2012; **336**(6086): 1255–1262.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
5. Bäckhed F, Roswall J, Peng Y, *et al.*: **Dynamics and Stabilization of the Human Gut Microbiome during the First Year of Life.** *Cell Host Microbe.* 2015; **17**(5): 690–703.
[PubMed Abstract](#) | [Publisher Full Text](#)
6. Koenig JE, Spor A, Scalfone N, *et al.*: **Succession of microbial consortia in the developing infant gut microbiome.** *Proc Natl Acad Sci U S A.* 2011; **108** Suppl 1: 4578–4585.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
7. Yatsunenko T, Rey FE, Manary MJ, *et al.*: **Human gut microbiome viewed across age and geography.** *Nature.* 2012; **486**(7402): 222–227.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
8. David LA, Maurice CF, Carmody RN, *et al.*: **Diet rapidly and reproducibly alters the human gut microbiome.** *Nature.* 2014; **505**(7484): 559–563.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
9. De Filippo C, Cavalieri D, Di Paola M, *et al.*: **Impact of diet in shaping gut microbiota revealed by a comparative study in children from Europe and rural Africa.** *Proc Natl Acad Sci U S A.* 2010; **107**(33): 14691–14696.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
10. Wu GD, Chen J, Hoffmann C, *et al.*: **Linking long-term dietary patterns with gut microbial enterotypes.** *Science.* 2011; **334**(6052): 105–108.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
11. Cotillard A, Kennedy SP, Kong LC, *et al.*: **Dietary intervention impact on gut microbial gene richness.** *Nature.* 2013; **500**(7464): 585–588.
[PubMed Abstract](#) | [Publisher Full Text](#)
12. Kovatcheva-Datchary P, Nilsson A, Akrami R, *et al.*: **Dietary Fiber-Induced Improvement in Glucose Metabolism Is Associated with Increased Abundance of *Prevotella*.** *Cell Metab.* 2015; **22**(6): 971–982.
[PubMed Abstract](#) | [Publisher Full Text](#)
13. Walker AW, Ince J, Duncan SH, *et al.*: **Dominant and diet-responsive groups of bacteria within the human colonic microbiota.** *ISME J.* 2011; **5**(2): 220–230.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
14. Ley RE, Hamady M, Lozupone C, *et al.*: **Evolution of mammals and their gut microbes.** *Science.* 2008; **320**(5883): 1647–1651.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
15. Muegge BD, Kuczynski J, Knights D, *et al.*: **Diet drives convergence in gut microbiome functions across mammalian phylogeny and within humans.** *Science.* 2011; **332**(6032): 970–4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
16. Charbonneau MR, O'Donnell D, Blanton LV, *et al.*: **Sialylated Milk Oligosaccharides Promote Microbiota-Dependent Growth in Models of Infant Undernutrition.** *Cell.* 2016; **164**(5): 859–871.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
17. Blanton LV, Charbonneau MR, Salih T, *et al.*: **Gut bacteria that prevent growth impairments transmitted by microbiota from malnourished children.** *Science.* 2016; **351**(6275): pii: aad3311.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
18. Vijay-Kumar M, Aitken JD, Carvalho FA, *et al.*: **Metabolic syndrome and altered gut microbiota in mice lacking Toll-like receptor 5.** *Science.* 2010; **328**(5975): 228–231.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
19. Buffington SA, Di Prisco GV, Auchtung TA, *et al.*: **Microbial Reconstitution Reverses Maternal Diet-Induced Social and Synaptic Deficits in Offspring.** *Cell.* 2016; **165**(7): 1762–1775.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
20. Faith JJ, McNulty NP, Rey FE, *et al.*: **Predicting a human gut microbiota's response to diet in gnotobiotic mice.** *Science.* 2011; **333**(6038): 101–104.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
21. Koeth RA, Wang Z, Levinson BS, *et al.*: **Intestinal microbiota metabolism of L-carnitine, a nutrient in red meat, promotes atherosclerosis.** *Nat Med.* 2013; **19**(5): 576–585.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
22. Wu GD, Compher C, Chen EZ, *et al.*: **Comparative metabolomics in vegans and omnivores reveal constraints on diet-dependent gut microbiota metabolite production.** *Gut.* 2016; **65**(1): 63–72.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
23. Ege MJ, Mayer M, Normand AC, *et al.*: **Exposure to environmental microorganisms and childhood asthma.** *N Engl J Med.* 2011; **364**(8): 701–709.
[PubMed Abstract](#) | [Publisher Full Text](#)
24. Stein MM, Hrusch CL, Gozdz J, *et al.*: **Innate Immunity and Asthma Risk in Amish and Hutterite Farm Children.** *N Engl J Med.* 2016; **375**(5): 411–421.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
25. Schnorr SL, Candela M, Rampelli S, *et al.*: **Gut microbiome of the Hadza hunter-gatherers.** *Nat Commun.* 2014; **5**: 3654.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
26. Zhou D, Zhang H, Bai Z, *et al.*: **Exposure to soil, house dust and decaying plants increases gut microbial diversity and decreases serum immunoglobulin E levels in BALB/c mice.** *Environ Microbiol.* 2016; **18**(5): 1326–1337.
[PubMed Abstract](#) | [Publisher Full Text](#)
27. Zhou D: **Impact of sanitary living environment on gut microbiota.** *Precision Medicine.* 2016; **2**: e1161.
[Publisher Full Text](#)
28. Bai Z, Zhang H, Li N, *et al.*: **Impact of Environmental Microbes on the Composition of the Gut Microbiota of Adult BALB/c Mice.** *PLoS One.* 2016; **11**(8): e0160568.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
29. Angenent LT, Kelley ST, St Amand A, *et al.*: **Molecular identification of potential pathogens in water and air of a hospital therapy pool.** *Proc Natl Acad Sci U S A.* 2005; **102**(13): 4860–4865.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
30. Penders J, Thijs C, Vink C, *et al.*: **Factors influencing the composition of the intestinal microbiota in early infancy.** *Pediatrics.* 2006; **118**(2): 511–521.
[PubMed Abstract](#) | [Publisher Full Text](#)
31. Whitacre CC: **Sex differences in autoimmune disease.** *Nat Immunol.* 2001; **2**(9): 777–780.
[PubMed Abstract](#) | [Publisher Full Text](#)
32. Knights D, Costello EK, Knight R: **Supervised classification of human microbiota.** *FEMS Microbiol Rev.* 2011; **35**(2): 343–359.
[PubMed Abstract](#) | [Publisher Full Text](#)
33. Adlerberth I, Lindberg E, Aberg N, *et al.*: **Reduced enterobacterial and increased staphylococcal colonization of the infantile bowel: an effect of hygienic lifestyle?** *Pediatr Res.* 2006; **59**(1): 96–101.
[PubMed Abstract](#) | [Publisher Full Text](#)
34. Vatanen T, Kostic AD, d'Hennezel E, *et al.*: **Variation in Microbiome LPS Immunogenicity Contributes to Autoimmunity in Humans.** *Cell.* 2016; **165**(4): 842–853.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
35. Merrifield CA, Lewis MC, Berger B, *et al.*: **Neonatal environment exerts a sustained influence on the development of the intestinal microbiota and metabolic phenotype.** *ISME J.* 2016; **10**(1): 145–157.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
36. Eggesbo M, Moen B, Peddada S, *et al.*: **Development of gut microbiota in infants not exposed to medical interventions.** *APMIS.* 2011; **119**(1): 17–35.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
37. Martin R, Makino H, Cetinyurek Yavuz A, *et al.*: **Early-Life Events, Including Mode of Delivery and Type of Feeding, Siblings and Gender, Shape the Developing Gut Microbiota.** *PLoS One.* 2016; **11**(6): e0158498.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
38. Bastide NM, Chenni F, Audebert M, *et al.*: **A central role for heme iron in colon carcinogenesis associated with red meat intake.** *Cancer Res.* 2015; **75**(5): 870–879.
[PubMed Abstract](#) | [Publisher Full Text](#)
39. Potera C: **Red Meat and Colorectal Cancer: Exploring the Potential HCA Connection.** *Environ Health Perspect.* 2016; **124**(10): A189.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
40. Kau AL, Ahern PP, Griffin NW, *et al.*: **Human nutrition, the gut microbiome and the immune system.** *Nature.* 2011; **474**(7351): 327–336.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
41. Zeng Y, Du J, Pu X, *et al.*: **Coevolution between Cancer Activities and Food Structure of Human Being from Southwest China.** *Biomed Res Int.* 2015; **2015**: 497934.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
42. Yangzong Y, Shi Z, Nafstad P, *et al.*: **The prevalence of childhood asthma in China: a systematic review.** *BMC Public Health.* 2012; **12**: 860.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
43. Ott W: **Human activity patterns: A review of the literature for estimating time spend indoors, outdoors and in transit.** Las Vegas. 1989.
44. Kilkenny C, Browne WJ, Cuthill IC, *et al.*: **Improving bioscience research reporting: the ARRIVE guidelines for reporting animal research.** *PLoS Biol.* 2010; **8**(6): e1000412.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
45. Zoetendal EG, Booijink CC, Klaassens ES, *et al.*: **Isolation of RNA from bacterial samples of the human gastrointestinal tract.** *Nat Protoc.* 2006; **1**(2): 954–959.
[PubMed Abstract](#) | [Publisher Full Text](#)
46. Bolger AM, Lohse M, Usadel B: **Trimmomatic: a flexible trimmer for Illumina sequence data.** *Bioinformatics.* 2014; **30**(15): 2114–2120.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
47. Edgar RC, Haas BJ, Clemente JC, *et al.*: **UCHIME improves sensitivity and**

- speed of chimera detection. *Bioinformatics*. 2011; **27**(16): 2194–2200.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
48. Wang Q, Garrity GM, Tiedje JM, *et al.*: Naive Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Appl Environ Microbiol*. 2007; **73**(16): 5261–5267.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
49. Caporaso JG, Kuczynski J, Stombaugh J, *et al.*: QIIME allows analysis of high-throughput community sequencing data. *Nat Methods*. 2010; **7**(5): 335–336.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
50. Liaw A, Wiener M: Classification and regression by randomForest. *R News*. 2002; **2**(3): 18–22.
[Reference Source](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 21 February 2020

<https://doi.org/10.5256/f1000research.16664.r59635>

© 2020 Garcia-Mazcorro J. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Jose F. Garcia-Mazcorro

Research Group Medical Eco-Biology, Faculty of Veterinary Medicine, Universidad Autónoma de Nuevo León, San Nicolás de los Garza, Mexico

This is a very interesting study with lots of potential for multiples areas in medicine and biomedical sciences. I have only a few comments that may improve the manuscript.

- Do not use microflora.
- Do you have any information about the soil characteristics?
- I am missing more details in the discussion regarding the mechanism behind the effect of soil in their immediate environment. The authors talk about coprophagy but do not mention anything about whether they observed this behavior in the animals. Also, if possible, it would not take much time and money to analyze the microbiota in the soil and to investigate whether there are any shared OTUs. In this regard, I hope the following paper is somehow useful for the authors¹. Is it possible that the soil changed some parameters and behavior of the immune system and therefore the effect on the microbiota could be indirect?
- Could you please add information about any OTUs that were shared among the animals?
- The clustering of samples in Fig. 4 is simply too strong. For me, such a strong clustering would only be explained by processing and sequencing replicates of the same samples. Please explain anything you can add to explain such a strong clustering.
- Please consider performing any additional analysis on the 16S sequences you think is appropriate (e.g. PICRUSt, BugBase, etc).
- The link provided to SRA seems not to be working.

References

1. Lankelma JM, Birnie E, Weehuizen TAF, Scicluna BP, et al.: The gut microbiota as a modulator of innate immunity during melioidosis. *PLoS Negl Trop Dis*. **11** (4): e0005548 [PubMed Abstract](#) | [Publisher Full Text](#)

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Partly

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Partly

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Microbial ecology, bioinformatics.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Reviewer Report 03 February 2020

<https://doi.org/10.5256/f1000research.16664.r38951>

© 2020 Zhang X. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Xuegong Zhang

Key Lab of Bioinformatics, Bioinformatics Division, Beijing National Research Centre for Information Science and Technology (BNRIST), Department of Automation, School of Life Sciences, Tsinghua University, Beijing, China

This is an interesting paper that shows the influence of bedding soil on the experimental mice's microbiome, especially the differential effects on mice with different diet. The experiment results

and statistical analyses are overall convincing, but there are a few minor points that the work can be further improved with more solid conclusions:

1. The paper only described the soils used "was from the top 10cm of a farm ground with abundant goats, hens and ducks". No further information about the procedures of sterilizing the soils were provided. If the authors can conduct a systematic analysis on the biological and chemical characteristics of sterile and unsterile soil samples, more concrete results might be observed.
2. Some of the observations reported in the paper was based on p-values that are not very small, like $p=0.042$ or 0.019 . It will be better if the authors also present the effect size of those tests and discuss the issue of multiple tests.
3. In beginning of the Introduction, the authors stated that "The main contributing factors are delivery mode, age, antibiotic treatment, diet, and the living environment". While those are the main factors that have been reported to affect gut microbiota, the statement was too strong and exclusive. Actually those factors only explain part of the variations, and there could be other important factors that have not been revealed yet.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Partly

If applicable, is the statistical analysis and its interpretation appropriate?

Partly

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Pattern Recognition, Machine Learning and Bioinformatics.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research