RESEARCH ARTICLE

Relationship between *Candida albicans* and *Streptococcus mutans* in early childhood caries, evaluated by quantitative PCR [version 1; peer review: 2 approved]

Endang W. Bachtiar , Boy M. Bachtiar

Oral Biology and Oral Science Research Center Faculty of Dentistry, Universitas Indonesia, Jakarta, 10430, Indonesia

V1 First published: 16 Oct 2018, 7:1645
<https://doi.org/10.12688/f1000research.16275.1>
 Latest published: 06 Dec 2018, 7:1645
<https://doi.org/10.12688/f1000research.16275.2>

Abstract

Background: The aim of this study was to analyze the synergistic relationship between *Candida albicans* and *Streptococcus mutans* in children with early childhood caries (ECC) experience.

Methods: Dental plaque and unstimulated saliva samples were taken from 30 subjects aged 3-5 years old, half with (n=15, dmft > 4) and half without (n=15) ECC. The abundance of *C. albicans* and *S. mutans* and relative to total bacteria load were quantify by real-time PCR (qPCR). This method was also employed to investigate the mRNA expression of glycosyltransferase (*gtfB*) gene in dental plaque. Student's t-test and Pearson's correlation were used to perform statistical analysis.

Results: Within the ECC group, the quantity of both microorganisms were higher in the saliva than in dental plaque. The ratio of *C. albicans* to total bacteria was higher in saliva than in plaque samples ($p < 0.05$). We observed the opposite for *S. mutans* ($p < 0.05$). The different value of *C. albicans* and *S. mutans* in saliva was positively correlated, and negatively correlated in dental plaque. Transcription level of *S. mutans gtfB* showed a positive correlation with *C. albicans* concentration in dental plaque.

Conclusion: *C. albicans* has a positive correlation with cariogenic traits of *S. mutans* in ECC-related biofilm of young children.

Keywords

Early childhood caries, C.albicans/ S. mutans, Saliva, Dental plaque, qPCR, Indonesian

Open Peer Review

Approval Status

	1	2
version 2 (revision) 06 Dec 2018		
version 1 16 Oct 2018	 view	 view
1. Zamirah Zainal-Abidin , National University of Malaysia (UKM), Kuala Lumpur, Malaysia		
2. Shahida Mohd-Said , National University of Malaysia, Kuala Lumpur, Malaysia		
Any reports and responses or comments on the article can be found at the end of the article.		

Corresponding author: Boy M. Bachtiar (boybachtiar@gmail.com)

Author roles: **Bachtiar EW:** Data Curation, Methodology, Project Administration, Software, Supervision, Validation, Writing – Review & Editing; **Bachtiar BM:** Conceptualization, Formal Analysis, Supervision, Validation, Visualization, Writing – Original Draft Preparation

Competing interests: No competing interests were disclosed.

Grant information: This study was supported by a grant from the Ministry of Research, Technology, and Higher Education, The Republic of Indonesia 2017.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2018 Bachtiar EW and Bachtiar BM. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. Data associated with the article are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

How to cite this article: Bachtiar EW and Bachtiar BM. **Relationship between *Candida albicans* and *Streptococcus mutans* in early childhood caries, evaluated by quantitative PCR [version 1; peer review: 2 approved]** F1000Research 2018, 7:1645 <https://doi.org/10.12688/f1000research.16275.1>

First published: 16 Oct 2018, 7:1645 <https://doi.org/10.12688/f1000research.16275.1>

Introduction

Early childhood caries (ECC) remain the most common childhood oral health problem, globally¹, and *Streptococcus mutans* has been known for its important role in ECC development^{2,3}. However, in recent years, *Candida albicans* has frequently been linked with its synergistic relationship with *S. mutans* in dental plaque recovered from children with ECC^{4,5}. Consequently, many studies using different methods have been conducted to identify, quantify, and explored the relationship of this fungus with *S. mutans*^{6–8}. However, a controversial report does exist, where *C. albicans* tends to decrease the cariogenic traits of *S. mutans* in *in vitro* dual-species biofilm⁹. Therefore, the main purpose of this study was to validate the synergistic relationships between *C. albicans* and *S. mutans*, when growing in caries-related biofilm. For this reason, we required preschool children with ECC experience, and we used qPCR since it is practice and reliable as a quantitative molecular tool of clinical oral samples¹⁰. The fungus and bacterium concentrations in saliva sample were used as control, and we compared the outcomes with those subjects noted as children with free caries (FC).

Methods

Subjects

Oral samples were collected from 30 required preschool children (male and female, 3–5 years old), in two different location located near (about 30 km) to Jakarta, the capital city of Indonesia. The diagnostic of ECC referred to the criteria provided by the American Academy of Pediatric Dentistry, as previously reported¹¹. The preschool children were recruited to get 15 subjects for each group. Thus, in this study, children were categorized into two different groups; children without any history of caries, including white-spot lesions, (caries free; CF group) and ECC group with decay-missing-filled teeth (dmft) index >4. To be included as subjects in this study, the children were required to be free of symptomatic oral candidiasis, have the absence of any medication therapy during the one month before this study, and have not worn any intraoral appliances. Before oral samples collection, written permission (informed consent) for children to participate was obtained from parents or guardians, according to the guidance provided by the Ethics Committee of Faculty of Dentistry, Universitas Indonesia.

Samples from supragingival plaque, obtained from the selected teeth deciduous (either molar or incisive) were isolated with sterile cotton rolls and pure cotton buds. Samples from the ECC group was obtained by gathering carious biofilm around the affected enamel¹², including dentin, as the fungus does not invade carious human dentin¹³. For the FC group, samples were obtained from enamel in clinically sound gingival areas. In each group, samples collected from molar or incisor, upper or lower teeth were not separated. Therefore, the obtained plaques were pooled to give a single sample for each subject and put in a microcentrifuge tube containing 1 ml PBS (pH 7.4). Unstimulated saliva was collected from all subjects on the same occasion and immediately after the plaque collection, by spitting the saliva into sterile Falcon plastic tubes. The minimum volume of collected saliva was 0.5 ml.

Saliva and plaque samples were immediately cold-transported to the laboratory. For saliva samples, after centrifugation, the sediments were washed three times with 0.5 ml sterile milli Q water between each centrifugation and kept in -80°C until use. Similarly, plaque samples from caries-free or those with ECC were cold-transported to the laboratory and processed as mentioned above, then stored at -80°C until use.

Quantification of *C. albicans*, *S. mutans*, and total bacteria by qPCR

Bacterial/fungal DNA was obtained by centrifugation each sample in the microtube, using Trizol reagent (Sigma-Aldrich, Dorset, UK). Extracted DNA sample was kept at -20°C after determining the concentration and the quality by Qubit assay reagents (Invitrogen, Carlsbad, CA). The genomic DNA samples were dissolved in Tris-EDTA (TE) buffer and kept in -20°C freeze until used. Further, the DNA samples were quantified through a qPCR reaction with universal primers for the 16S rRNA genes and *C. albicans*/*S. mutans*-specific primers as shown in Table 1. For PCR-quantification, each sample was run in triplicate on an ABI StepOnePlus Real-Time PCR System with SYBR Green PCR Master Mix (Applied Biosystems, Foster City, CA, USA) according to the manufacturer's protocol.

The PCR conditions were run in a final reaction volume of 10 µl, composed of 50 ng of sample DNA and 1 µM of each

Table 1. Primers used in this study.

Primer name	Sequences	References
<i>C. albicans</i>	Forward: 5'-CACGACGGAGTTTCACAAGA -3'	14
	Reverse: 5'-CGATGGAAGTTTGAGGCAAT-3'	
<i>S. mutans</i>	Forward: 5'-CCTACGGGAGGCAGCAGTAG-3'	14
	Reverse: 5'-CAACAGAGCTTTACGATCCGAAA-3'	
Universal bacterial 16S rDNA	Forward: 5'-GTGSTGCAYGGYTGTCTGCA-3'	15
	Reverse: 5'-ACGTCRTCCMCACCTTCCTC-3'	
<i>gtfB</i>	Forward: 5'-AGCAATGCAGCCAATCTACAAAT-3'	14
	Reverse: 5'-ACGAACTTTGCCGTTATTGTCA-3'	

primer (Table 1), with thermal cycling condition consisted of a 10 min initial denaturation at 95°C, followed by 40 cycles of 95°C for 15 s and 65°C for 60 s. The cycle threshold (Ct) were determined automatically by the instrument, and a dissociation curve of the amplified fragment set as follow; 95°C for 15 s, 60°C for 60 s, and 95°C for 15 s.

Estimating the amount of genomic DNA of both microorganisms tested was determined by constructing standard curves with r^2 values for both organisms tested as well as total bacteria (Figure 1A–C). To do this, we used a 10-fold serial dilution of extracted fungal and bacterial genomic DNA from overnight cultured of *C. albicans* ATCC 10231, *S. mutans* Xc, and *Escherichia coli* JM 107, respectively. The number (CFU/ml) assessed by plating culture dilutions on sabouraud agar, tryptone-yeast extract cysteine with sucrose and bacitracin (TYCSB) agar and Luria Bertani (LB) broth for *C. albicans*¹⁶, *S. mutans*¹⁷, and *E. coli*¹⁸, respectively. The same strains were used as positive control for qPCR. Therefore, quantification of *C. albicans* and *S. mutans* from plaques and saliva achieved by plotting the Ct values against the log of the respective standard curve. In this study, the ratio of *C. albicans* or *S. mutans* in the microbial community, in each sample, was determined as each microorganism proportion to total bacteria.

For both *C. albicans* and *S. mutans*, the detection limit by qPCR method was determined according to the limit of

quantification (LOQ), obtained by the highest dilution of the template of the standard curve. When the Ct value of samples was higher than the LOQ, they would be considered positive, but their melting curve profile should be the same as those of the standards included when running the qPCR.

qPCR analysis of *S. mutans* *gtfB* in dental plaque samples

RNA isolation, purification, and reverse transcription of cDNA were performed similarly those in the previous study¹⁹. Platinum SYBR Green qPCR SuperMix-UDG (Invitrogen Life Technologies, Carlsbad, CA, USA), passive reference (ROX, Invitrogen), and *S. mutans* *gtfB* primers (Table 1), as well as 1 µg of cDNA, were used to quantify the cDNA, and non-transcribed RNA samples were used as control for genomic DNA contamination. The qPCR reaction was run on a similar machine as stated above with cycling conditions consisted of a 10 min initial denaturation at 95°C followed by 40 PCR cycles of 15 s at 95°C, and 60°C for 1 min. The formula of fold change $2^{-\Delta\Delta C_t}$ was used to calculate *S. mutans* *gtfB* gene expression that was normalized to the 16S rRNA, a well-established housekeeping gene²⁰, and *gtfB* expressed in dental plaque of FC group was set to be the control.

Statistical analysis

The variables for quantification, proportion, and the mean quantitative gene expression were assessed using Student's t-test, while Pearson's correlation two-tailed test was used to depict

Figure 1. Standard curve construction and melting temperature of qPCR. Standard curves of total bacteria (A), *C. albicans* (B) and *S. mutans* (C). Also shown are melt curve profiles and melting temperatures for total bacteria (D), *C. albicans* (E), and *S. mutans* (F).

a linear association. Microsoft Excel software was used to perform statistical analysis, and a p -value < 0.05 was considered significant.

Results

Quantitative levels of *C. albicans* and *S. mutans* and their proportion in saliva and dental plaque samples

Standard curves were used to determine the corresponding number of microorganism tested while melting peaks were used to assess the specificity of the amplicon using saliva and plaque samples (Figure 1D–F).

In general, this study showed that in all saliva and plaque samples, from either FC or ECC children, both *C. albicans* and *S. mutans* were present. The quantification (log DNA copies) and proportion (% to total bacteria) of *C. albicans* and *S. mutans* in saliva, as well as plaque samples, are presented in Figure 2. Comparatively, it was observed, in either sample tested, a significantly higher number of both microorganisms in ECC children was found more than in those of the FC children ($p < 0.05$). However, in either group, the quantitative level of *C. albicans*, in the saliva sample was found to be significantly lower than those of *S. mutans* ($p < 0.05$). When comparing plaque and saliva samples within ECC children, we observed that the load (log DNA copies) of either microorganism in plaque was significantly lower than that in saliva ($p < 0.05$, Figure 2A).

Furthermore, we compared the proportion of *C. albicans* and *S. mutans* DNA relative to total bacterial DNA in saliva and dental plaque samples, in FC and ECC children. Within ECC children, there was a significantly higher proportion ratio (Ca/Tb) of *C. albicans* in saliva samples (35.5%), than that in plaque samples (13.5%) ($p < 0.05$). For FC children the ratio was not statistically different (saliva, 8% and plaque, 5.3%) (Figure 2B).

Similar analysis was carried out for the *S. mutans* proportion ratio (Sm/Tb). The result showed a different trend, with a significantly higher proportion of *S. mutans* in plaque (99%) than that in the saliva (62%) of ECC children (Figure 2B). The result also showed the proportion of this bacterium to total bacteria in saliva and plaque samples showed a significant difference between ECC and FC children ($p < 0.05$). Interestingly, there was a trend within dental plaque sample in ECC children, *S. mutans* DNA increased most with increased of *C. albicans*' DNA (Figure 2B).

Association between the value of *C. albicans* and *S. mutans* in dental plaque and saliva of ECC children

We further evaluated the possible linear relationship of *C. albicans* and *S. mutans* load or their proportion and ECC experience in the subjects. Pearson correlations coefficient analysis revealed that the association between the numbers of these two microorganisms in saliva was moderate positively significant ($r = 0.1$, $p < 0.05$). On the contrary, a weak negative not substantial ($r = 0.03$, $p > 0.05$) between the decreasing number of *C. albicans* and the quantity of *S. mutans* in plaque samples observed in the ECC group (Figure 3A and B).

Quantification of *gtfB* gene transcription and its correlation with *C. albicans* and *S. mutans* amount in dental plaque

To confirm all the above results, we selected the *gtfB* gene, which has been reported to be mostly involved in the synergistic relationship between *C. albicans* and *S. mutans* in biofilm development⁷, and compared its expression in each dental plaque of children tested. The qPCR result showed that level of mRNA *gtfB* was induced approximately 4.5-fold in ECC-derived dental plaque samples, and it was a significant difference compared to transcription level of *gtfB* mRNA in dental plaque sampled from children with FC ($p < 0.05$) (Figure 4A). Also,

Figure 2. Mean and standard error of absolute numbers (A) and ratios (B) of *C. albicans* and *S. mutans* detected within the same saliva (Sa) and dental plaque (Dp) samples. CF, caries-free; ECC, early childhood caries. * $p < 0.05$.

Figure 3. Correlation between *C. albicans* and *S. mutans* loads in saliva (A) and in dental plaque (B), in the same subjects. Each circle depicts the value of *C. albicans* (Y-axis) and *S. mutans* (X-axis) in log CFU/ml for each subject.

Figure 4. *S. mutans gtfB* gene expression. (A) The fold regulation of *gtfB* cDNA that was normalized to the amount of cDNA of 16S rRNA. (B, C) the correlation between *gtfB* transcription rate and the amount of *C. albicans* and *S. mutans*, respectively.

gtfB transcription levels and the amount of *C. albicans* and *S. mutans* (CFU/ml) in dental plaque of children with ECC showed moderate ($r = 0.2$) and strong ($r = 0.9$) positive correlations, respectively, which was statistically significant ($p < 0.05$, Figure 3B and C).

Dataset 1. The raw data associated with this study

<https://dx.doi.org/10.5256/f1000research.16275.d221304>

Data are grouped according to the figure with which they are associated.

Discussion

Numerous studies^{7,8,21} have supported the idea that ECC may be better understood by focusing on the effect of the functional relationship between species within consortia instead of individual pathogens. This study focused on the relationship between *C. albicans* and *S. mutans*, as these oral microorganisms are frequently detected in the plaque of children with ECC^{12,22}. Since ECC may indicate a vast proliferation of the cariogenic microorganism, we sought to evaluate the extent to which the amount of *C. albicans* might correlate to the ECC experience. To do this, we used qPCR. This method enabled the quantification of the targeted microorganisms' genomic DNA from oral samples²³. Thus, we examined the similarities and differences by comparing the amount and ratio to total bacteria of each microorganism in saliva and dental plaque samples, and verified their correlation with the occurrence of ECC. In general, we observed that the oral cavities of preschool children in this study, with or without ECC, are colonized by yeast (*C. albicans*) and bacteria, as represented by *S. mutans*. Overall, the fungus presence was always simultaneously detected with *S. mutans*, although children with ECC had a higher amount of *C. albicans* and *S. mutans* in their oral cavity, compared to those children with FC. As expected, in addition to saliva, these oral microbiotas commonly exist together in an ECC-related biofilm.

Our observation is in accordance with results from other studies, which found that in addition to *S. mutans* as a specific cariogenic bacterium^{24,25}, *C. albicans* can be part of the dental lesion^{26,27}. Moreover, previous studies showed strong synergism when *C. albicans* and *S. mutans* co-existed in biofilm, suggesting that this co-existence enhanced their virulence^{7,8,28}. However, our data showed, the fungus was detected at lower levels in dental plaque, compared to *S. mutans*. This support the previous *in vitro* study²⁹, which found that the presence of *C. albicans* might favor the extensive colonization of *S. mutans* in dental biofilm.

The causes to generate site specificity bacteria proportion are believed to include local sucrose concentration in the oral cavity^{30,31}. In addition to sucrose³², many factors may link to the presence of *C. albicans* in children oral cavity. These include infection at birth, baby's feeding bottles, infected pacifiers, and carious teeth¹². We speculate that a high cariogenic diet might influence the interaction between *C. albicans* and *S. mutans* in these children tested. In turns, it becomes critical for ECC, since the presence of sucrose in the children oral cavity may lead to the ability of this species to grow within structured microbial biofilm^{33,34}. Further studies are therefore necessary.

The presence of yeast and bacteria is one of the local factors that contributes to the etiology of ECC⁷. To obtain an overall insight into the impact of simultaneous participation of *C. albicans* and *S. mutans* when detecting together in each sample tested, we compared the percent proportion of *C. albicans* or *S. mutans* relative to total bacteria. As expected, the proportion of *C. albicans* was higher in saliva than in carious plaque among children with ECC. Other studies have reported this phenomenon, where *Candida* species were frequently isolated more, qualitatively and quantitatively, from saliva than from dental plaque³⁵ and

subgingival samples³⁶. During in saliva, this fungus might act as a bridge for oral bacteria to adhere to a mucosal surface, a mechanism that may protect this bacterium from being removed by salivary flow and swallowing²¹. On the other side, biofilm formation is vital for *C. albicans* to survive as a pathogen, which involves attachment, colonization, and development of structural biofilm integrity composed of yeast and hypha^{37,38}. Our data illustrate that co-adhesion between *C. albicans* and *S. mutans* in cariogenic biofilm is one mechanism by which the fungus, in yeast form, survives in the oral cavity^{39,40}. Although the fungus morphology was not observed in this study, it has been reported by other studied that hypha morphology is not crucial for the *C. albicans*–*S. mutans* relationship when they grow in multispecies dental biofilms^{41–43}. Therefore, in addition to the morphology, both the number and proportion of salivary *C. albicans* influence the fungal–bacterial relationships, which further increases the risk of caries. Additionally, the flushing effect of saliva, as part of innate defense mechanism, might contribute to decrease *Candida* adherence to the oral surface⁴⁴, including tooth surface. Thus, the quality and quantity of saliva have an essential role in maintaining fungus behavior, as commensal or pathogen⁴⁵.

We observed that the proportion of *S. mutans* was higher in dental plaque than in saliva, and there was a tendency for the percentage of *C. albicans* to be lower in carious plaque, where the proportion of *S. mutans* increased. This observation indicates that *S. mutans* has an active role in orchestrating the development of cariogenic biofilms^{46,47}. This species has an essential part in attenuating the virulence of the fungus⁴⁸ by interfering with the fungus transition, from yeast to hypha form when these oral microflorae interact and grow in biofilm^{49,50}. This result further supported by the data of correlation analysis, in which the fungus-bacterium concentration in dental plaque sample showed a negative association, although a positive correlation was found in saliva sediment. This suggests that the ECC rate may not be connected to the quantity of *C. albicans* involves.

One of the mediators for the synergistic relationship between *C. albicans* and *S. mutans* is the streptococcal GtfB enzyme^{7,32}. Our finding indicates that cariogenic biofilm developed in ECC children accompanied by the increased transcription level of *gtfB* mRNA and enhance of *S. mutans* growth in dental plaque derived from children with ECC. In a clinical situation, this observation is relevant, since GtfB is the enzyme that synthesizes glucan polymers from sucrose⁶. Thus, result of this study suggests that a high sucrose concentration, which is critical for the development of dental caries, might exist in the children oral cavity. Future studies are thus recommended.

Collectively, this study indicates that although the proportion of *C. albicans* was less in ECC-associated biofilm, it may support facilitation of a fungal-bacterial synergistic relationship. Yeast cells could be used by *S. mutans* to promote fitness and the bacterial survival, as shown by enhanced transcription level of *gtfB*, which reflects that more extracellular polysaccharides were produced to promote the fungal-bacterium relationship in caries-related biofilm⁷.

The results of the present study cannot explain the reason for such an association. However, at least this study provides information on ECC experience among preschool children. The presence of *C. albicans* in dental plaque and saliva could be merely an indicator of oral health conditions and the high carbohydrate intake among the young children selected in this study, which might confer a survival advantage for *C. albicans* and it is favorable for ECC development. More studies are needed.

There is some limitation in this study. First, the primary disadvantage of qPCR used in this study is its inability to separate and quantify the viable from nonviable cells. This technique may result in false positives or an overly high estimation, as all DNA extracted from live or dead *C. albicans*, or *S. mutans* cell will be amplified. Since the number of viable cells is especially significant for diagnosing and monitoring disease, adding cell viability information to qPCR-based diagnostics should be considered. Second, the number of children involved in this study was small (15 subjects per group) because of difficulty in sample collection, primarily to obtain the plaque on the dentin surface.

Conclusions

This study shows that *C. albicans* contributes to increasing concentration of *S. mutans* by inducing the expression of *gtfB*

mRNA in ECC-related biofilm. Therefore, results from this study would be useful as a starting point to consider *C. albicans*, as a potential target in prevention programs to reduce the high rates of ECC in individuals or groups of young children.

Data availability

Dataset 1. The raw data associated with this study. Data are grouped according to the figure with which they are associated. DOI: <https://doi.org/10.5256/f1000research.16275.d221304>⁵¹.

Grant information

This study was supported by a grant from the Ministry of Research, Technology, and Higher Education, The Republic of Indonesia 2017.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgments

We thank Anissa, Vivi, and Asti for the laboratory work at Oral Science Research Centre, Faculty of Dentistry, Universitas Indonesia. We also thank for Arif, Tika, Inge, and Hanli for collecting oral samples.

References

- Petersen PE, Bourgeois D, Ogawa H, *et al.*: **The global burden of oral diseases and risks to oral health.** *Bull World Health Organ.* 2005; **83**(9): 661–669.
[PubMed Abstract](#) | [Free Full Text](#)
- Tanzer JM, Baranowski LK, Rogers JD, *et al.*: **Oral colonization and cariogenicity of *Streptococcus gordonii* in specific pathogen-free TAN:SPFOM(OM)BR rats consuming starch or sucrose diets.** *Arch Oral Biol.* 2001; **46**(4): 323–333.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Koga T, Oho T, Shimazaki Y, *et al.*: **Immunization against dental caries.** *Vaccine.* 2002; **20**(16): 2027–2044.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Raja M, Hannan A, Ali K: **Association of oral candidal carriage with dental caries in children.** *Caries Res.* 2010; **44**(3): 272–276.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Ellepola K, Liu Y, Cao T, *et al.*: **Bacterial GtfB Augments *Candida albicans* Accumulation in Cross-Kingdom Biofilms.** *J Dent Res.* 2017; **96**(10): 1129–1135.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kuramitsu HK: **Virulence properties of oral bacteria: impact of molecular biology.** *Curr Issues Mol Biol.* 2001; **3**(2): 35–36.
[PubMed Abstract](#)
- Falsetta ML, Klein MI, Colonne PM, *et al.*: **Symbiotic relationship between *Streptococcus mutans* and *Candida albicans* synergizes virulence of plaque biofilms in vivo.** *Infect Immun.* 2014; **82**(5): 1968–1981.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Koo H, Bowen WH: ***Candida albicans* and *Streptococcus mutans*: a potential synergistic alliance to cause virulent tooth decay in children.** *Future Microbiol.* 2014; **9**(12): 1295–1297.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Willems HM, Kos K, Jabra-Rizk MA, *et al.*: ***Candida albicans* in oral biofilms could prevent caries.** *Pathog Dis.* 2016; **74**(5): pii: ftw039.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lyons SR, Griffen AL, Leys EJ: **Quantitative real-time PCR for *Porphyromonas gingivalis* and total bacteria.** *J Clin Microbiol.* 2000; **38**(6): 2362–2365.
[PubMed Abstract](#) | [Free Full Text](#)
- Drury TF, Horowitz AM, Ismail AI, *et al.*: **Diagnosing and reporting early childhood caries for research purposes. A report of a workshop sponsored by the National Institute of Dental and Craniofacial Research, the Health Resources and Services Administration, and the Health Care Financing Administration.** *J Public Health Dent.* 1999; **59**(3): 192–197.
[PubMed Abstract](#) | [Publisher Full Text](#)
- de Carvalho FG, Silva DS, Hebling J, *et al.*: **Presence of mutans streptococci and *Candida* spp. in dental plaque/dentine of carious teeth and early childhood caries.** *Arch Oral Biol.* 2006; **51**(11): 1024–1028.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Maijala M, Rautemaa R, Järvensivu A, *et al.*: ***Candida albicans* does not invade carious human dentine.** *Oral Dis.* 2007; **13**(3): 279–284.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Feldman M, Ginsburg I, Al-Quntar A, *et al.*: **Thiazolidinedione-8 Alters Symbiotic Relationship in *C. albicans*-*S. mutans* Dual Species Biofilm.** *Front Microbiol.* 2016; **7**: 140.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Maeda H, Fujimoto C, Haruki Y, *et al.*: **Quantitative real-time PCR using TaqMan and SYBR Green for *Actinobacillus actinomycetemcomitans*, *Porphyromonas gingivalis*, *Prevotella intermedia*, *tetQ* gene and total bacteria.** *FEMS Immunol Med Microbiol.* 2003; **39**(1): 81–86.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Bachtir EW, Bachtir BM, Jarosz LM, *et al.*: **Al-2 of *Aggregatibacter actinomycetemcomitans* inhibits *Candida albicans* biofilm formation.** *Front Cell Infect Microbiol.* 2014; **4**: 94.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Van Palenstein Helderma WH, Ijsseldijk M, Huis in 't Veld JH: **A selective medium for the two major subgroups of the bacterium *Streptococcus mutans* isolated from human dental plaque and saliva.** *Arch Oral Biol.* 1983; **28**(7): 599–603.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Bachtir BM, Coloe PJ, Fry BN: **Knockout mutagenesis of the *kpsE* gene of *Campylobacter jejuni* 81116 and its involvement in bacterium-host interactions.** *FEMS Immunol Med Microbiol.* 2007; **49**(1): 149–154.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Bachtir BM, Bachtir EW: **Proinflammatory MG-63 cells response infection with *Enterococcus faecalis* cps2 evaluated by the expression of TLR-2, IL-1 β , and iNOS mRNA.** *BMC Res Notes.* 2017; **10**(1): 401.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Stipp RN, Gonçalves RB, Höfling JF, *et al.*: **Transcriptional analysis of *gtfB*, *gtfC*, and *gpbB* and their putative response regulators in several isolates of**

- Streptococcus mutans*. Oral Microbiol Immunol. 2008; 23(6): 466–473.**
[PubMed Abstract](#) | [Publisher Full Text](#)
21. Krom BP, Kidwai S, Ten Cate JM: ***Candida* and other fungal species: forgotten players of healthy oral microbiota.** *J Dent Res.* 2014; 93(5): 445–451.
[PubMed Abstract](#) | [Publisher Full Text](#)
 22. Qiu R, Li W, Lin Y, *et al.*: **Genotypic diversity and cariogenicity of *Candida albicans* from children with early childhood caries and caries-free children.** *BMC oral health.* 2015; 15(1): 144.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 23. Kirakodu SS, Govindaswami M, Novak MJ, *et al.*: **Optimizing qPCR for the Quantification of Periodontal Pathogens in a Complex Plaque Biofilm.** *Open Dent J.* 2008; 2: 49–55.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 24. Bowen WH, Koo H: **Biology of *Streptococcus mutans*-derived glucosyltransferases: role in extracellular matrix formation of cariogenic biofilms.** *Caries Res.* 2011; 45(1): 69–86.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 25. Krzyściak W, Jurczak A, Kościelniak D, *et al.*: **The virulence of *Streptococcus mutans* and the ability to form biofilms.** *Eur J Clin Microbiol Infect Dis.* 2014; 33(4): 499–515.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 26. Xiao J, Moon Y, Li L, *et al.*: ***Candida albicans* Carriage in Children with Severe Early Childhood Caries (S-ECC) and Maternal Relatedness.** *PLoS One.* 2016; 11(10): e0164242.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 27. Yang XQ, Zhang Q, Lu LY, *et al.*: **Genotypic distribution of *Candida albicans* in dental biofilm of Chinese children associated with severe early childhood caries.** *Arch Oral Biol.* 2012; 57(8): 1048–1053.
[PubMed Abstract](#) | [Publisher Full Text](#)
 28. Sztajer H, Szafranski SP, Tomasch J, *et al.*: **Cross-feeding and interkingdom communication in dual-species biofilms of *Streptococcus mutans* and *Candida albicans*.** *ISME J.* 2014; 8(11): 2256–2271.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 29. Barbieri DdS'AV VV, Fraiz FC, Lavoranti OJ, *et al.*: **Analysis of the *in vitro* adherence of *Streptococcus mutans* and *Candida albicans*.** *Braz J Microbiol.* 2007; 38(4): 624–631.
[Publisher Full Text](#)
 30. Dawes C, MacPherson LM: **The distribution of saliva and sucrose around the mouth during the use of chewing gum and the implications for the site-specificity of caries and calculus deposition.** *J Dent Res.* 1993; 72(5): 852–857.
[PubMed Abstract](#) | [Publisher Full Text](#)
 31. Hallett KB, O'Rourke PK: **Early childhood caries and infant feeding practice.** *Community Dent Health.* 2002; 19(4): 237–242.
[PubMed Abstract](#)
 32. Gregoire S, Xiao J, Silva BB, *et al.*: **Role of glucosyltransferase B in interactions of *Candida albicans* with *Streptococcus mutans* and with an experimental pellicle on hydroxyapatite surfaces.** *Appl Environ Microbiol.* 2011; 77(18): 6357–6367.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 33. Uppuluri P, Pierce CG, López-Ribot JL: ***Candida albicans* biofilm formation and its clinical consequences.** *Future Microbiol.* 2009; 4(10): 1235–1237.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 34. Metwalli KH, Khan SA, Krom BP, *et al.*: ***Streptococcus mutans*, *Candida albicans*, and the human mouth: a sticky situation.** *PLoS Pathog.* 2013; 9(10): e1003616.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 35. Salam MA, Senpuku H, Nomura Y, *et al.*: **Isolation of opportunistic pathogens in dental plaque, saliva and tonsil samples from elderly.** *Jpn J Infect Dis.* 2001; 54(5): 193–195.
[PubMed Abstract](#)
 36. Jabra-Rizk MA, Ferreira SM, Sabet M, *et al.*: **Recovery of *Candida dubliniensis* and other yeasts from human immunodeficiency virus-associated periodontal lesions.** *J Clin Microbiol.* 2001; 39(12): 4520–4522.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 37. Finkel JS, Mitchell AP: **Genetic control of *Candida albicans* biofilm development.** *Nat Rev Microbiol.* 2011; 9(2): 109–118.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 38. Ramage G, Saville SP, Thomas DP, *et al.*: ***Candida* biofilms: an update.** *Eukaryot Cell.* 2005; 4(4): 633–638.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 39. Nikawa H, Egusa H, Makihiro S, *et al.*: **Alteration of the coadherence of *Candida albicans* with oral bacteria by dietary sugars.** *Oral Microbiol Immunol.* 2001; 16(5): 279–283.
[PubMed Abstract](#) | [Publisher Full Text](#)
 40. Thein ZM, Samaranyake YH, Samaranyake LP: **Effect of oral bacteria on growth and survival of *Candida albicans* biofilms.** *Arch Oral Biol.* 2006; 51(8): 672–680.
[PubMed Abstract](#) | [Publisher Full Text](#)
 41. Hwang G, Liu Y, Kim D, *et al.*: ***Candida albicans* mannans mediate *Streptococcus mutans* exoenzyme GtfB binding to modulate cross-kingdom biofilm development *in vivo*.** *PLoS Pathog.* 2017; 13(6): e1006407.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 42. Pereira-Cenci T, Deng DM, Kraneveld EA, *et al.*: **The effect of *Streptococcus mutans* and *Candida glabrata* on *Candida albicans* biofilms formed on different surfaces.** *Arch Oral Biol.* 2008; 53(8): 755–764.
[PubMed Abstract](#) | [Publisher Full Text](#)
 43. Joyner PM, Liu J, Zhang Z, *et al.*: **Mutanobactin A from the human oral pathogen *Streptococcus mutans* is a cross-kingdom regulator of the yeast-mycelium transition.** *Org Biomol Chem.* 2010; 8(24): 5486–5489.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 44. Ramage G, Vandewalle K, Wickes BL, *et al.*: **Characteristics of biofilm formation by *Candida albicans*.** *Rev Iberoam Micol.* 2001; 18(4): 163–170.
[PubMed Abstract](#)
 45. Epstein JB, Truelove EL, Izutzu KT: **Oral candidiasis: pathogenesis and host defense.** *Rev Infect Dis.* 1984; 6(1): 96–106.
[PubMed Abstract](#) | [Publisher Full Text](#)
 46. Kanasi E, Johansson I, Lu SC, *et al.*: **Microbial risk markers for childhood caries in pediatricians' offices.** *J Dent Res.* 2010; 89(4): 378–383.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 47. Tanner AC, Kent RL Jr, Holgerson PL, *et al.*: **Microbiota of severe early childhood caries before and after therapy.** *J Dent Res.* 2011; 90(11): 1298–1305.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 48. Barbosa JO, Rossoni RD, Vilela SF, *et al.*: ***Streptococcus mutans* Can Modulate Biofilm Formation and Attenuate the Virulence of *Candida albicans*.** *PLoS One.* 2016; 11(3): e0150457.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 49. Jarosz LM, Deng DM, van der Mei HC, *et al.*: ***Streptococcus mutans* competence-stimulating peptide inhibits *Candida albicans* hypha formation.** *Eukaryot Cell.* 2009; 8(11): 1658–1664.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 50. Vilchez R, Lemme A, Ballhausen B, *et al.*: ***Streptococcus mutans* inhibits *Candida albicans* hyphal formation by the fatty acid signaling molecule *trans*-2-decenoic acid (SDSF).** *ChemBiochem.* 2010; 11(11): 1552–1562.
[PubMed Abstract](#) | [Publisher Full Text](#)
 51. Bachtar EW, Bachtar BM: **Dataset 1 in: *Candida albicans* and *Streptococcus mutans* relationship in early childhood caries, evaluated by quantitative real time-PCR.** *F1000Research.* 2018.
<http://www.doi.org/10.5256/f1000research.16275.d221304>

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 06 November 2018

<https://doi.org/10.5256/f1000research.17775.r39489>

© 2018 Mohd-Said S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Shahida Mohd-Said

Centre for Restorative Dentistry, Faculty of Dentistry, National University of Malaysia, Kuala Lumpur, Malaysia

This paper reports an interesting and relevant study on ECC in children. The methods and contents of the papers are current, relevant and well-structured. However, some improvements could benefit the presentation including checking on grammar and typos, some revisions to sentence construction, highlights on the significance of this study, how the findings can be important to update of knowledge and current management of ECC, and comparison of data from Indonesia with other populations to perhaps illustrate novelty and significance to current available data.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

I cannot comment. A qualified statistician is required.

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 08 Nov 2018

Endang Bachtiar, Universitas Indonesia, Jakarta, Indonesia

Dear reviewer,

I am pleased to resubmit for publication the revised version of **Relationship between *Candida albicans* and *Streptococcus mutans* in early childhood caries, evaluated by quantitative PCR**.

We appreciate all the insightful comments provided by the reviewer. Based on the reviewer's guidance, includes a number of positive changes, we endeavoured to improve the fit of the paper with the journal.

We hope that these revisions improve the paper such that the editor and reviewer now deem it worthy of publication in F1000 Research. Thank you for taking the time to help us improve the paper.

Sincerely,

Authors: Endang W Bachtiar and Boy M Bachtiar

Responses to reviewer #1:

We have fixed all grammar, typos, and some revisions to sentence construction. Furthermore all responses to the suggestions have been added in the manuscript in the yellow highlight sentences.

Competing Interests: No competing interests

Reviewer Report 31 October 2018

<https://doi.org/10.5256/f1000research.17775.r39491>

© 2018 Zainal-Abidin Z. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Zamirah Zainal-Abidin

Centre for Craniofacial Diagnostics and Biosciences, Faculty of Dentistry, National University of Malaysia (UKM), Kuala Lumpur, Malaysia

Methods:

1. The authors did not mention how the examiners were calibrated, in order to reduce the

inter- and intra-examiner variability when examining the preschool children, before grouping them into the early childhood caries and caries free groups.

2. The authors may have been able to collect more information on the oral hygiene habits or dietary intake of the subjects, before taking the oral samples.

Results:

1. In the caption for Figure 2, the symbol $*p<0.05$ should be indicated clearly. For Figure 2A, it is to show that there is a statistically-significant difference between the *C. albicans* in saliva and dental plaque in the ECC group, and a statistically-significant difference between the *S. mutans* in saliva and dental plaque in the ECC group. The use of top square bracket ($\square$) with an asterisk would be better to indicate which parameters are statistically compared.
2. For Figure 4A, the use of top square bracket ($\square$) with an asterisk would be better to indicate the parameters which are statistically compared.

Discussion:

1. In paragraphs 3 and 7, the authors speculated that a high cariogenic/high sucrose concentration diet might influence the interaction between *C. albicans* and *S. mutans* in the tested subjects. The authors may have been able to draw a better suggestion if the information on the oral hygiene habits or diet is obtained during the recruitment of the subjects (See my comment in Methods (2)).
2. In paragraphs 3, 7 and 9, the authors could have suggested what further studies are necessary to address the suggested theories/findings.

General comments:

1. The paper has a few grammar and language issues, in the Abstract and Methods sections.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Partly

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 08 Nov 2018

Endang Bachtiar, Universitas Indonesia, Jakarta, Indonesia

Dear editor and reviewer,

I am pleased to resubmit for publication the revised version of **Relationship between *Candida albicans* and *Streptococcus mutans* in early childhood caries, evaluated by quantitative PCR.**

We appreciate all the insightful comments provided by the reviewer. Based on the reviewer's guidance, includes a number of positive changes, we endeavoured to improve the fit of the paper with the journal.

We hope that these revisions improve the paper such that the editor and reviewer now deem it worthy of publication in F1000 Research. Thank you for taking the time to help us improve the paper.

Sincerely,

Authors : 1. Endang W Bachtiar
2. Boy M Bachtiar

Author's response to reviewer comments
Comment provided by Reviewer #1

Methods:

Reviewer's comment:

The authors did not mention how the examiners were calibrated, in order to reduce the inter- and intra-examiner variability when examining the preschool children, before grouping them into the early childhood caries and caries-free groups.

Author's response:

We thank the reviewer for this correction. The problem has been fixed. As suggested by the reviewer, we have added sentence within material and method as follow: "two weeks prior to collect to clinical samples, the examiners were calibrated and trained by providing with the manual describing study protocol and guidance regarding the examination of preschool children. Therefore, only those trained-examiners evaluated the preschool children".

Reviewer's comment:

The authors may have been able to collect more information on the oral hygiene habits or dietary intake of the subjects, before taking the oral samples.

Author's response:

In this study, we used a questionnaire regarding the parent's/ guardian's perception concerning children's general and oral health, the risk factor for caries, dietary intake, and access to dental care. We noticed, in general, most of the responders believed that dental decay is a natural phenomenon. However, considering the complexity of the issue studied in this study, it is not the authors' intent to include oral hygiene habits or dietary intake of the subjects (preschool children). Therefore, the related-data were not included in the current study.

Reviewer's comment:

In the caption for Figure 2, the symbol $*p<0.05$ should be indicated clearly. For Figure 2A, it is to show that there is a statistically significant difference between the *C. albicans* in saliva and dental plaque in the ECC group, and a statistically significant difference between the *S. mutans* in saliva and dental plaque in the ECC group. The use of top square bracket ($\square$) with an asterisk would be better to indicate which parameters are statistically compared.

For Figure 4A, the use of top square bracket ($\square$) with an asterisk would be better to indicate the parameters which are statistically compared.

Author's response:

According to the suggestions provided by Reviewer#1, we have corrected the old figures. The top square bracket has been added in Fig. 2A to indicate a statistically significant difference as suggested. For the Fig. 4A, we have added the top square bracket with an asterisk to indicate a significant difference between the parameters compared. Thank you.

Discussion:

Reviewer's comment:

In paragraphs 3 and 7, the authors speculated that a high cariogenic/high sucrose concentration diet might influence the interaction between *C. albicans* and *S. mutans* in the tested subjects. The authors may have been able to draw a better suggestion if the information on the oral hygiene habits or diet is obtained during the recruitment of the subjects (See my comment in Method (2)).

Author's response:

As mentioned above (response to the reviewer's comment in methods (2); it is not the authors' intent to include the oral hygiene habits or diet of the preschool children in this study. This is because the information regarding oral hygiene habit or diet, provided by the

accompanying guardian are difficult to be understood. Therefore, it is hard to interpret oral hygiene habit or diet data.

Reviewer's comment:

In paragraphs 3, 7 and 9, the authors could have suggested what further studies are necessary to address the suggested theories/findings.

Author's responses:

According to reviewer suggestion, we have added a better suggestion at the end of paragraph-3, as follow; "Further studies to obtain data regarding cariogenic diet are therefore necessary". Additionally, we have added a sentence at the end of paragraph 7, as follow: "More studies regarding the involvement of gtfB expression, and how it relates to oral health conditions as well as cariogenic intake in Indonesian preschool children, are needed.

At the end of paragraph 9 (conclusion), as suggested by the reviewer, we added sentences; "Moreover, since the information regarding oral hygiene habit or diet, provided by the accompanying guardian are difficult to be understood, future studies involving preschool children may wish to involve examiners who are more experience in dealing with such parents/guardians when planning studies". Thank you.

Dear Editor,

We are pleased to inform you that; 1/ for all the revision, as suggested by the reviewer, were showed in yellow-highlighting words, 2/ we revised the figure 2 and 4 to address the reviewer's comments. Although the Fig. 3 is not included in reviewer's comment, we revised the font, from the original (Arial) to new version (Times New Roman). Thus, all founds depicted in figures are similar. Thank you.

Competing Interests: No competing interests

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research