ANTIBODY VALIDATION ARTICLE

Comparative study of commercially available and homemade anti-VAMP7 antibodies using CRISPR/Cas9-depleted HeLa cells and VAMP7 knockout mice [version 1; peer review: 1 approved, 1 approved with reservations]

Agathe Verraes^{1*}, Beatrice Cholley^{2*}, Thierry Galli^{1,2}, Sebastien Nola^{1,2}

¹Membrane Traffic in Health and Disease, INSERM ERL U950, Univ Paris Diderot, Sorbonne Paris Cité, Institut Jacques Monod, CNRS UMR 7592, Paris, France, 75013, France

²Membrane Traffic in Healthy & Diseased Brain, Université Paris Descartes, Paris, 75014, France

* Equal contributors

V1 First published: 16 Oct 2018, 7:1649
<https://doi.org/10.12688/f1000research.15707.1>
 Latest published: 07 Feb 2019, 7:1649
<https://doi.org/10.12688/f1000research.15707.2>

Abstract

VAMP7 (vesicle-associated membrane protein) belongs to the intracellular membrane fusion SNARE (Soluble N-ethylmaleimide-sensitive factor attachment protein receptors) protein family. In this study, we used CRISPR/Cas9 genome editing technology to generate VAMP7 knockout (KO) human HeLa cells and mouse KO brain extracts in order to test the specificity and the background of a set of commercially available and homemade anti-VAMP7 antibodies. We propose a simple profiling method to analyze western blotting and immunocytochemistry staining profiles and determine the extent of the antibodies' specificity. Using this method, we were able to rank the performance of a set of available antibodies and further showed an optimized procedure for VAMP7 immunoprecipitation, which we validated using wild-type and KO mouse brain extracts.

Keywords

VAMP7, SNARE, monoclonal, polyclonal, CRISPR/Cas9, KO, immunoprecipitation

This article is included in the **Antibody** Validations gateway.

Open Peer Review

Approval Status

	1	2
version 2 (revision) 07 Feb 2019	 view	
version 1 16 Oct 2018	 view	 view

1. **Alison H. Banham** , University of Oxford, Oxford, UK

2. **Marc G. Coppolino**, University of Guelph, Guelph, Canada

Olivia Grafinger , University of Guelph, Guelph, Canada

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Thierry Galli (thierry.galli@inserm.fr)

Author roles: **Verraes A:** Investigation, Writing – Original Draft Preparation; **Cholley B:** Investigation, Methodology, Validation, Writing – Original Draft Preparation; **Galli T:** Conceptualization, Funding Acquisition, Investigation, Project Administration, Supervision, Writing – Review & Editing; **Nola S:** Conceptualization, Formal Analysis, Investigation, Project Administration, Resources, Software, Validation, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: Work in our group was funded by grants from Association Française contre les Myopathies (Research Grant 16612), the French National Research Agency (NeuroImmunoSynapse ANR-13-BSV2-0018-02; MetDePaDi ANR-16-CE16-0012), the Ecole des Neurosciences de Paris (ENP), the Fondation pour la Recherche Médicale (FRM), Prix Coup d'Élan pour la recherche française of the Fondation Bettencourt Schueller, awards of the Association Robert Debré pour la Recherche Médicale to T.G.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2018 Verraes A *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. Data associated with the article are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

How to cite this article: Verraes A, Cholley B, Galli T and Nola S. **Comparative study of commercially available and homemade anti-VAMP7 antibodies using CRISPR/Cas9-depleted HeLa cells and VAMP7 knockout mice [version 1; peer review: 1 approved, 1 approved with reservations]** F1000Research 2018, 7:1649 <https://doi.org/10.12688/f1000research.15707.1>

First published: 16 Oct 2018, 7:1649 <https://doi.org/10.12688/f1000research.15707.1>

Introduction

Intracellular membrane fusion in the secretory and endocytic pathways relies on SNARE proteins (soluble N-ethylmaleimide-sensitive factor attachment protein receptors) for membrane fusion events. In order to allow apposition and fusion between two membranes, vesicular (v)- and target (t)-SNARE form a so-called trans-SNARE complex, or SNAREpin. VAMP7 (vesicle associated membrane protein 7) is a clostridial neurotoxin-insensitive v-SNARE that belongs to the “Longin” subfamily (as opposed to the short “Brevins”, like VAMPs 1-3): it encompasses an amino-terminal extension, the Longin domain, which acts as an auto-regulatory domain¹. VAMP7 mediates Golgi-derived, late endosomal and lysosomal and autophagosomal related membrane fusion events^{2,3} and co-localizes to a large extent with the tetraspanin CD63^{4,5}. VAMP7 is involved in exocytosis in several cell types⁶, including neurons⁷, in neurotransmitter basal release^{8,9} and specific brain circuits and functions¹⁰. VAMP7 exocytosis was shown to be regulated by an integrin-, FAK-, and Src-dependent mechanism in developing neurons¹¹ and its transport to the cell periphery by VARP, Rab21 and Kif5⁷, while retrograde transport depends on LRRK1¹². In non-neuronal cells, VAMP7 secretory vesicles release compounds such as ATP^{13,14} and interleukin-12⁴. In addition, VAMP7 regulates trafficking of membrane proteins, including the tetraspanin CD82¹⁵ and the cold-sensing channel TRPM8¹⁶. VAMP7 plays an essential role in cell migration and invasion^{17–19}. VAMP7 also contributes to the regulation of membrane composition of sphingolipids and GPI-anchored proteins²⁰.

At present date, several antibodies against VAMP7 are commercially available. However, not all of them have been extensively characterized and many reported studies have been conducted using exogenous expression. This is too little information regarding the sensitivity of these antibodies, and may limit their use for super-resolution imaging or proximity ligation assay, which require detection of endogenous proteins.

In this survey, we took advantage of the genome editing CRISPR/Cas9 technology to generate VAMP7-knockout (KO) human HeLa cells. This genetically modified cell line allowed us to test the specificity and background of available commercially or homemade VAMP7 antibodies. Here we compared four mouse monoclonal and four rabbit polyclonal antibodies by western blotting and immunofluorescence using standard protocols. We analyzed the data using a simple profiling of both western blots and immunocytochemistry images to extract a specificity index. We further characterized the best ones by immunoprecipitation assays using VAMP7 constructs from different origins and wild type or VAMP7 knockout mouse tissues.

Material and methods

Cell culture

HeLa and Cos-7 cells (ATCC CCL-2 and CRL-1651, respectively) were maintained at 37°C and 5% CO₂ in a humidified incubator, and grown in Dulbecco's modified Eagle's medium (DMEM), supplemented with 10% fetal calf serum (FCS), 100 units/ml penicillin, and 100 µg/ml streptomycin (Gibco, Thermo Fisher Scientific). Cells were regularly split using Trypsin-EDTA to maintain exponential growth. Transfection

of cells was performed using Lipofectamine 2000 according to the manufacturer's instructions. All culture media reagents were from Thermo Fisher Scientific.

Generation of VAMP7-depleted cells

Inactivation of VAMP7 was achieved using CRISPR RNA-CAS9 guide constructs based on a previously published protocol²¹. Briefly, sequences for sgRNA were preferentially chosen within the first exon region of VAMP7 genomic gene, with the help of the “RGEN Cas designer”²² web-based tool (<http://rgenome.net/cas-designer/>). To limit off-targets, an oligo sequence with ≤2 putative mismatches throughout the whole genome or an ‘out of frame’ score <66 were excluded. The sgRNA target sequences used are: 5'-caccgAACAGCAAAAAGAATCGCCA-3' (forward) and 5'-aaacTGGCGATTCTTTTGTCTGTTc-3' (reverse). Oligonucleotides (10 mM) were heated at 95°C for 5 min and annealed by ramping down the temperature from 95°C to 25°C at 5°C min⁻¹. Annealed primers were ligated into pSpCas9(BB)-2A-Puro (PX459) vector (Addgene) using the BbsI sites. After validation by sequencing, the targeting constructs were transfected into HeLa cells following a previously described protocol. An empty pSpCas9(BB)-2A-Puro (PX459) vector was used to generate “control” cells. At 24 h post-transfection, cells were diluted 1/10 and transfected ones were selected by 1 µg/ml puromycin addition for 72 h. The selected populations were then seeded into a 96-well plate at 1 cell per well. Clones derived from single cells were amplified and screened for deficiency by immunoblotting.

Plasmids

The human and rat GFP-VAMP7 constructs are the same as those that have been described previously²³. Plasmid containing mouse VAMP7 cDNA was a kind gift from Maurizio D'Esposito (IGCB, Naples, Italy). Mouse VAMP7 was amplified by PCR and cloned into pEGFP-C3 (Clontech) using HindIII / BamHI restriction sites.

Mouse colony husbandry and care

The wild-type (WT) and VAMP7 knockout (KO) littermate male cohort was established at the Mouse Clinical Institute animal facility as previously described¹⁰. They had a mixed 129/Sv-C57BL/6 genetic background. They were weaned at 4 weeks and housed two to six per M.I.C.E. cage by sex and litter regardless of the genotype under standard conditions and maintained in a room with controlled temperature (21–22°C) under a 12 h light/dark cycle (light on at 7:00 A.M.), with food (standard chow diet, Safe D04) and water available *ad libitum*. All experiments were performed in accordance with the European Communities Council Directive regarding the care and use of animals for experimental procedures (2010/63/UE) and were approved by the local ethical committee (CEEA40-Comité d'Ethique Buffon). Mice were euthanized by cervical dislocation. All efforts were made to ameliorate the suffering of animals and to reduce their number per experiment (1 animal cortex per condition for immunoprecipitation experiment).

Cortex isolation

The cortex from WT and VAMP7 KO 8 weeks-old C57bl/6 mice were isolated according to a previously published protocol¹⁰. Cortex were dissociated by pipette trituration in 500 µl TSE

(50 mM Tris-HCl, pH 8.0, 150 mM NaCl, 1 mM EDTA) supplemented with 1% Triton X-100 and cOmplete protease inhibitor tablets (Roche Applied Science) and volume was adjusted in order to get ~2 mg/ml protein final concentration (considering 10 mg of tissue is equivalent to ~1 mg final protein). Lysis was performed by incubation under agitation at 4°C for 30min. After clarification by 16,000 x g centrifugation for 30 min, protein concentration of the supernatant was estimated using Protein Assay Dye Reagent Concentrate (Bio-Rad Laboratories) and immunoprecipitation was carried out (see below).

Affinity purification

The TG50 (for “Thierry Galli #50”) serum raised against VAMP7 was generated by Covalab (Villeurbanne, France; animal house registration number C21 464 04 EA) using immunization of New Zealand white rabbit with GST-VAMP7 (1-188 aa, as previously described for TG11²⁴ and TG18²⁵) and then purified by affinity chromatography. Briefly, serum was clarified and de-lipidated by high-speed centrifugation (70,000 rpm) and applied on a 6xHis-VAMP7 (coiled-coil 1-188 aa) covalently cross-linked HITrap-NHS column (GE Healthcare) overnight at 4°C using a peristaltic pump (0.3 ml/min). We washed the column with filtered and degassed PBS (20 ml, 1 ml/min).

Specific antibodies were eluted using 200 mM Glycine/HCl pH 2.2, and collected in tubes containing neutralizing buffer (TBS 10X, 1X final). For each fraction, protein concentration was quantified by optical density (280 nm) measurement.

Immunoblotting

References for all tested antibodies and reagents used for immunoblotting are listed in [Table 1](#) and [Table 2](#), respectively. Cells were washed in cold phosphate buffered saline (PBS) and lysed 20 min in TSE (50mM Tris-HCl, pH 8.0, 150mM NaCl, 1 mM EDTA) supplemented with 1% Triton X-100 and cOmplete protease inhibitor tablets (Roche Applied Science). Lysates were clarified by centrifugation 30 min at 16,000 x g, and protein concentration was estimated using Protein Assay Dye Reagent Concentrate (Bio-Rad Laboratories). Following heat denaturation at 95°C during 5 min, proteins were separated by 15% SDS-PAGE and transferred to a 0.45-µm nitrocellulose filter (Amersham, GE Healthcare) at 40 mA overnight. Membrane was blocked with 5% low-fat milk in TBS buffer for 20 min at room temperature and probed with indicated primary antibodies diluted in 5% skimmed milk in TBS-T, overnight at 4°C. Following primary antibody incubation, the filter was washed 3x 5 min in TBS-T at room temperature, and probed with

Table 1. Details of primary and secondary antibodies.

Antibody	Origin	Manufacturer	Catalog number	RRID	WB dilution	IF dilution
Anti-VAMP7 (D4D5J)	Rabbit	Cell Signaling Technology	14811	Not available	1:800	
Anti-VAMP7 (D8Y1R)	Rabbit	Cell Signaling Technology	13876	Not available		1:200
Anti-VAMP7	Rabbit	Sigma-Aldrich	SAB3500844	Not available	1:800	1:200
Anti-VAMP7	Rabbit	Synaptic Systems	232 003	AB_2212953	1:800	1:500
Anti-VAMP7 (TG50)	Rabbit	TG lab (affinity purified)	Not relevant	Not relevant	1:800	1:500
Anti-VAMP7 (TG50)	Rabbit	Covalab (Protein A purified)	pab01031-P	Not available	Not tested	Not tested
Anti-VAMP7	Mouse	Creative Diagnostics	CABT-37960MH	AB_2358849	1:800	1:200
Anti-VAMP7	Mouse	R&D Systems	MAB6117	AB_10571669	1:800	1:200
Anti-VAMP7 (158.2)	Mouse	TG lab (original clone)	Not relevant	Not relevant	1:800	1:200
Anti-VAMP7 (158.2)	Mouse	Synaptic Systems (recloned 158.2)	232 011	AB_2619947	1:800	1:200
Anti-α-tubulin	Mouse	Sigma-Aldrich	T6074	AB_477582	1:800	1:200
Anti-GFP	Rabbit	Abcam	ab6673	AB_305643		
Anti-GFP	Mouse	Roche	11814460001	AB_390913		
Normal IgG	Rabbit	Sigma-Aldrich	I-5006	AB_1163659		
Anti-mouse IgG HRP	Goat	Jackson Immuno Research	115-035-008	AB_2313585	1:10000	
Anti-rabbit IgG HRP	Goat	Jackson Immuno Research	111-035-008	AB_2337937	1:10000	
Anti-mouse IgG IRDye 800	Donkey	Rockland	610-731-124	AB_220145	1:5000	
Anti-rabbit IgG IRDye 800	Donkey	Rockland	611-731-127	AB_220157	1:5000	
Anti-mouse AF 488	Goat	Invitrogen	A-11029	AB_2534088		1:500
Anti-rabbit AF 488	Goat	Invitrogen	A-11034	AB_2576217		1:500

Table 2. Details of reagents used for immunoblotting.

Process	Reagent	Manufacturer	Catalog number	Concentration/ composition
Sample preparation	Sample reducing agent LDS Sample Buffer	Invitrogen Invitrogen	NP0009 NP0008	10X 4X
Protein ladder	SeeBlue Plus2 Prestained standard	Invitrogen	LC5925	1X
SDS-PAGE Running Buffer	Tris base Glycine SDS	Euromedex Euromedex Euromedex	200923-A 261286405C EU0660	25mM 190mM 0.1% w/v
Protein blotting membrane	0.45µm nitrocellulose transfer membrane			
Wash buffer/ diluent for blocking and antibody (TBS-T)	Tris base NaCl Tween-20	Euromedex VWR Euromedex	200923-A 27810.295 EU0660	20 mM 150 mM 0.1% w/v
Blocking buffer	Milk powder TBS-T			5% w/v 1X
SDS-PAGE Transfer Buffer	Tris base Glycine Ethanol	Euromedex Euromedex VWR	200923-A 261286405C 20824.365	25mM 190mM 20% w/v

HRP- or fluorescently-labeled secondary antibodies diluted in TBS-T for 1 h at room temperature. The filter was then washed 3× 5 min in TBS-T at room temperature. All incubation and washes were performed with gentle rocking. Proteins were detected by enhanced chemiluminescence (ThermoFisher Scientific) and imaged using ImageQuant LAS-4000 (Fujitsu Life Sciences), or scanned in an Odyssey infrared imaging system (Li-Cor).

Immunofluorescence staining

The following immunofluorescence staining protocol was performed, with all steps carried out at room temperature unless stated otherwise (see Table 3 for reagent details). HeLa cells were washed once in PBS, fixed with 4% paraformaldehyde in PBS for 20 min, quenched for 20 min with 50 mM NH₄Cl in PBS and permeabilized by treatment with 0.3% Triton X-100 in PBS for 4 min. After blocking with 10% FCS + 0.3% Triton in PBS for 30 min, cells were incubated overnight with the primary antibodies diluted in 3% FCS + 0.3% Triton in PBS at 4°C. After several washes with 3% FCS, 0.3% Triton in PBS, cells were then incubated with the secondary antibodies in 3% FCS, 0.3% Triton in PBS for 1 h, then washed several times in 0.3% Triton in PBS. Coverslips were partially dried and mounted in Prolong medium (Invitrogen), and then left to set overnight. Fluorescence microscopy and imaging were performed using an upright epifluorescence microscope (DMRA2, Leica Microsystem) equipped with a CMOS camera (Orca Flash 4.0 LT, Hamamatsu) and a HCX PL APO 100x/1.40-0.70 oil CS oil-immersion Leica objective. Dilution of primary antibodies (Table 3) was formerly optimized to get relatively equivalent signal intensity in the WT cells for the same microscope settings (time of exposure, binning, objective, etc.), allowing direct comparison of signal between antibodies.

Immunoprecipitation

Transfected Cos7 cells were washed once in PBS 1X then lysed as described in the “immunoblotting” section. Immunoprecipitation experiments were carried out as followed (see Table 4 for reagent details). Briefly, 1 mg of protein extract was submitted to immunoprecipitation by incubation overnight at 4°C with 2.5 µg of antibodies that were pre-coupled with 25 µl magnetic beads (Dynabeads M-280, Invitrogen). Beads were then extensively washed with TSE-1% Triton and beads resuspended in 2X Laemmli buffer. Samples were loaded on 4–12% Bis-Tris NuPAGE (ThermoFisher Scientific) or RunBlue SDS (Expedeon) gels with manufacturer-recommended electrophoresis buffer, processed for western blotting using HRP-coupled secondary antibodies and enhanced chemiluminescence (ThermoFisher Scientific).

For immunoprecipitation of endogenous VAMP7, 1 mg of mouse cortex extracts (see “Cortex isolation” section) were submitted to immunoprecipitation as for cell extracts, excepted that 5 µg of antibodies, 40 µl magnetic beads (Dynabeads M-280, Invitrogen), fluorescent secondary antibodies and an Odyssey infrared imaging system (LI-COR, Lincoln, Nebraska, USA) were used.

Quantitative analysis

All quantification analyses was performed using ImageJ (1.49n) and data were computed in Microsoft Excel.

For western blotting signal quantification, a 20-pixel-wide straight line vertically crossing each lane was drawn to generate an intensity profile (see Figure 1A and B, left panel). Local background correction was performed by manually drawing a segmented line under the peaks representing the bands detected by western blotting (Figure B, right panel). The VAMP7

Table 3. Details of reagents used for immunofluorescence.

Process	Reagent	Manufacturer	Catalog number	Concentration/Composition
Washing	PBS	Home-made		1x
Fixing	Paraformaldehyde	Prolabo	PRO-28794.295	4% in PBS 1x
Permeabilizing, Washing (PBS-T)	PBS Triton X100	Home-made Merck	9410	1x 0.3% w/v
Blocking	Fetal Bovine Serum (FBS) PBS-T	Thermofisher	10270106	10% v/v
Antibody diluent	Fetal Bovine Serum (FBS) PBS-T	Thermofisher	10270106	3% v/v
Nuclear staining	DAPI (4',6-diamidino-2-phenylindole)	Invitrogen	D3571	0.2µg/ml in PBS
Mounting	Prolong antifade	Invitrogen	P36930	1X

Table 4. Details of reagents used for immunoprecipitation.

Process	Reagent	Manufacturer	Catalog number	Concentration/Composition
Magnetic Coupling	Dynabeads M-280 Sheep anti-mouse IgG Dynabeads M-280 Sheep anti-rabbit IgG	Invitrogen Invitrogen	11201D 11203D	10mg/ml 10mg/ml
Washing (TSE-T)	Tris HCl pH8.0 NaCl EDTA Triton X100	Sigma VWR Eurobio Merck	T2694 27810.295 GAUEDT0064 9410	50 mM 150 mM 1 mM 1% w/v
Sample preparation	Sample reducing agent LDS Sample Buffer	Invitrogen Invitrogen	NP0009 NP0008	10X 4X

band was defined as the ~25 kDa band that would disappear or diminish in intensity in the KO extract compared with the WT extract. Areas under all peaks and the VAMP7 one, shown in grey and blue, respectively (Figure 1B, right panel), were measured. In order to estimate the signal-to-noise ratio of each antibody, taking into account the intensity of the band of interest over the intra-lane non-specific ones and the specificity of the signal in the control condition versus the KO ones, a so-called “WB specificity index” was calculated for each lane of the western blot, using the following equation:

$$\text{WB specificity index} = \left\{ \frac{\text{Area}_{(\text{peak of interest})}}{\text{Area}_{(\text{all peaks})}} \right\}_{\text{control}} - \left\{ \frac{\text{Area}_{(\text{peak of interest})}}{\text{Area}_{(\text{all peaks})}} \right\}_{\text{VAMP7 KO}}$$

For immunocytochemistry, rather than a raw intensity comparison between control and KO conditions, we chose to rely on the subcellular distribution of VAMP7 to reflect the correct staining pattern of the tested antibodies better. This was estimated using a previously described method with slight modifications²⁶, for each antibody, in control and VAMP7 KO cells. A 20-pixel-wide band (Figure 2B) was drawn from the nucleus (defined from the edge of DAPI staining) to the cell leading edge and going through the perinuclear region pointing to the most elongated part of the cell. Length of this band was normalized by binning in order to obtain a 10-pixel-long line from which

a fluorescence intensity profile was generated. Profile from control and KO cells were plotted in function of the distance from nucleus to periphery (i.e. 0–100% on the x axis, Figure 2B, lower panel). A so-called “IF specificity index” was calculated by measuring the area between the control and the KO intensity profiles, which can be computed by the following equation:

$$\text{IF specificity index} = \int_0^{100} \text{Intensity}_{\text{control}} - \int_0^{100} \text{Intensity}_{\text{VAMP7 KO}}$$

Antibodies with a high score exhibit both a more intense signal in the perinuclear region than the peripheral one in the control and a low signal distribution for VAMP7 KO condition compared to control. Poor score corresponds to either a correct distribution of VAMP7 signal in the control but a very high background in the KO condition, or an unspecific/random distribution of VAMP7 signal in the control, or a combination of both cases.

Results

Tools development and description of antibodies tested in this study

In order to characterize and compare a set of commercially available and homemade (i.e from “Thierry Galli’s lab”, hereafter referred as “TG lab”) anti-VAMP7 antibodies (Table 1), we first generated VAMP7 knockout cells, using CRISPR/Cas9 engineering²¹, as described in the methods section. HeLa cells

Figure 1. Comparative immunoblot analysis of endogenous VAMP7 level of expression in control or CRISPR/Cas9-modified VAMP7-depleted HeLa cells. (A) Immunoblot performed on lysates from control (Ctrl) and VAMP7 knockout (KO) HeLa cells. An equal amount of total protein extracts from each condition was run in replicates. Following transfer on nitrocellulose and blocking, membrane was sliced and each piece was probed with indicated anti-VAMP7 antibodies (expected size: ~25 kDa). Time of exposure for each condition is provided. For loading control, membrane was washed and incubated with anti- α -tubulin antibody (expected size: ~50 kDa). **(B)** Example of quantification of western blotting signal from **(A)** (see dotted red line). Intensity profile (right panel) was generated from a 20-pixel-wide straight line (yellow, left panel) across each lane. On the intensity profile (right panel), areas corresponding to the VAMP7 signal and the non-specific bands are shown in blue and gray, respectively (see Methods section for details). **(C)** Quantification of each antibody tested by western blotting shown in **(A)**. The “specificity index” represents the signal-to-noise ratio of the antibodies, reflecting the intensity of the VAMP7 band amongst the overall signal per lane (including non-specific bands) and its specificity in the control condition compared to KO (see Methods section for details). AP, affinity purified.

Figure 2. Comparative immunostaining of VAMP7 endogenous level of expression in control or depleted HeLa cells. (A) Control and VAMP7 knockout (KO) HeLa cells were grown on glass coverslips, fixed in paraformaldehyde, blocked and immunostained with the indicated mouse or rabbit anti-VAMP7 antibodies (green) and DAPI (blue). KO condition allows the estimation of background signal. Samples were imaged with an epifluorescence microscope using a 100X objective. Bars, 25 μ m. **(B)** Example of quantification of immunofluorescence signal from **(A)**. Distribution of VAMP7 fluorescence was measured across a 20-pixel-wide straight line (yellow) drawn from the border of the nucleus towards the plasma membrane in control and VAMP7 KO cells (left panel). Intensity profiles were plotted on the same graph (right panel) and area between control and VAMP7 KO curve (blue) was measured in order to generate a "specificity index" (see Methods section). **(C)** Quantification of each antibody tested by IF shown in **(A)**. The "specificity index" reflects both pattern of distribution of VAMP7 across the cell and signal-to-noise ratio given by the tested antibodies (see Methods section for details). AP, affinity purified.

were chosen as they express VAMP7 endogenously in a detectable amount by western blotting and immunocytochemistry¹⁵.

Several of the tested antibodies were described in the provider's datasheet to only work for immunofluorescence detection (e.g. Cell Signaling, catalogue number D81Y1R) or western blotting (e.g. Cell Signaling, catalogue number D4D5J) and were used accordingly. Our lab generated two antibodies, the mouse monoclonal "158.2"²⁷ and the rabbit polyclonal "TG50", which are commercially available from Synaptic Systems and Covalab (TG50 as protein A purified serum), respectively. Only the in-house affinity-purified (AP) version of the TG50 antibody was included in this study because we wanted an affinity-purified serum as best possible positive control.

Characterization of antibodies by western blotting

We compared four monoclonal and four polyclonal rabbit antibodies by western blotting using control or VAMP7-KO HeLa cell extracts (Figure 1). We used an anti-tubulin antibody as loading control. All antibodies tested in the described conditions (Table 2) were sensitive enough to detect a prominent band at the expected molecular weight (~25 kDa) in the control condition. This band was absent in the VAMP7-KO cell lines in all cases. However, some non-specific bands were visible for all the tested antibodies in both control and VAMP7-KO cell lysates, particularly with polyclonal Synaptic Systems (catalogue number 232 003) and Sigma-Aldrich (catalogue number T6074) antibodies. Therefore, all the tested antibodies showed a signal that was specific for VAMP7, but they also showed variable background bands. As assessed by intensity profile analysis (Figure 1B) and our western blotting specificity index (Figure 1C), the TG lab (TG50) antibody showed the best signal-to-noise ratio using this Western blotting conditions, which may not be surprising, because it had been affinity-purified.

Characterization of antibodies by immunofluorescence

In order to better characterize these antibodies (Table 1), we performed immunostaining in control and VAMP7 KO HeLa cells. For this assay, the rabbit antibody from Cell Signalling Technology clone D8Y1R was used instead of the D4D5J clone, according to the manufacturer's recommendations. To compare the specificity of the antibodies, we adjusted their dilution (Table 3) to get relatively equivalent signal intensity in the WT cells with the same acquisition time on the microscope. According to this assay, the mouse antibodies from Creative Diagnostics (CABT-37960MH), Synaptic Systems (158.2 –232 011), TG lab (158.2) and the rabbit antibodies Synaptic Systems (232 003) and TG lab (TG50) stained perinuclear membrane structures and vesicles dispersed in the cytoplasm, a typical and already described localization pattern for VAMP7 in HeLa cells^{10,15,28}. However, in the WT HeLa cells, the R&D Systems (MAB6117) antibody gave a homogenous signal, which spread into the nucleus, the Cell Signaling (14811) antibody, seemed to also stain perinuclear ER-like structures and the Sigma-Aldrich (T6074) antibody exhibited a diffuse cytoplasmic pattern with an absence of vesicular staining. All the tested antibodies seemed to show an overall lower-intensity signal in the VAMP7 KO cells

compared to control. According to the "secondary-only" condition that reveals the internal background signal of the experiment (Figure 2A, right panel) and the distribution analysis we conducted (Figure 2B, see Methods for details) to compute the IF specificity index (Figure 2C), the most convincing signal-to-noise ratio was observed with the Creative Diagnostics (CABT-37960MH), Synaptic Systems (232 003) and TG lab (158.2 and TG50) antibodies.

Characterization of antibodies by immunoprecipitation

Taken together, immunoblot and immunofluorescence assays suggest that the homemade anti-VAMP7 antibodies (158.2 and TG50) showed the best endogenous signal-to-noise ratio in HeLa cells. To check the inter-species specificity of these antibodies, we carried out immunoprecipitation assays in Cos cells overexpressing mouse, rat or human GFP-tagged VAMP7 constructs (Figure 3A). We used 158.2 and TG50 for immunoprecipitation and immunoblot and species-specific IgG and anti-GFP antibodies as negative and positive controls for immunoprecipitation, respectively (Table 4). Both 158.2 and TG50 antibodies exhibited a sharp ~50 kDa band in all immunoprecipitation lanes, demonstrating a relatively equivalent ability to precipitate either mouse, rat or human VAMP7, while a very faint signal was observed in the negative control IgG IP lanes.

We next wondered whether or not these antibodies could immunoprecipitate endogenous VAMP7 from tissue and test background signal in KO tissue. To this aim, we chose to perform immunoprecipitation on lysates from WT or VAMP7 KO mouse cortex using the rabbit TG50 antibody because it performed the best in previous assays and to observe immunoprecipitated VAMP7 with the mouse 158.2 antibody (Figure 3B). Although 158.2 antibody showed stronger background and multiple bands compared to the pattern seen in HeLa cells (Figure 1), a clear band at the expected molecular weight (~25 kDa) was present for the WT immunoprecipitation condition but not VAMP7 KO, demonstrating the specificity and low background noise of the TG50 antibody for immunoprecipitation of endogenous VAMP7 in mouse tissue extracts. Altogether, we showed here that both 158.2 and TG50 antibodies were able to immunoprecipitate VAMP7 from different species and that immunoprecipitation of endogenous VAMP7 could be performed in mouse tissue extracts using TG50 for IP and 158.2 for subsequent western blotting.

Dataset 1. Raw images of experimental replicates for Figure 1, immunoblotting experiments

<https://dx.doi.org/10.5256/f1000research.15707.d221360>

This dataset includes uncropped blots for all experimental replicates that are represented in Figure 1. Treatments and immunoblot methods were performed as outlined in Figure 1. Blots were probed with indicated anti-VAMP7 antibodies and anti- α -tubulin antibodies was used as a loading control. (A) Dataset used for Figure 1, with cropped regions in red dashed line. (B) Additional set of raw images of a replicate experiment. Quantification as performed in Figure 1 is shown in lower panel. Note that although signal intensity and background are different within these two replicates, the relative performance of the different tested antibodies remained the same.

A

Transfection	GFP-VAMP7 (mouse)					GFP-VAMP7 (rat)					GFP-VAMP7 (human)				
IP antibody	158.2	IgGM	TG50	IgGR	GFP	158.2	IgGM	TG50	IgGR	GFP	158.2	IgGM	TG50	IgGR	GFP
IP beads specificity	anti-mouse		anti-rabbit			anti-mouse		anti-rabbit			anti-mouse		anti-rabbit		
WB (primary Ab)	TG50 (AP)		158.2			TG50 (AP)		158.2			TG50 (AP)		158.2		
WB (IRDye 800 secondary Ab)	anti-rabbit		anti-mouse			anti-rabbit		anti-mouse			anti-rabbit		anti-mouse		
	SN	IP	SN	IP	SN	IP	SN	IP	SN	IP	SN	IP	SN	IP	SN
kDa															
49		*		°		*		°		*		°		*	
39															
28															

B

Cortex lysate	WT		VAMP7 KO	
IP antibody	TG50	IgGR	TG50	IgGR
IP beads	anti-mouse IgG			
WB (primary Ab)	158.2			
WB (secondary Ab)	anti-mouse IgG IRDye 800			
	IN	SN	IP	SN
kDa				
62				
49			*	
39				
28				

Figure 3. Comparison of available anti-VAMP7 antibodies for exogenous and endogenous immunoprecipitation. (A) Cos cells overexpressing GFP-tagged mouse, rat or human VAMP7 constructs were lysed and VAMP7 was immunoprecipitated using indicated specific antibodies ("IP Antibody"). Normal isotype IgG ("IgGM" or "IgGR") and GFP antibody were used as negative and positive control, respectively. Supernatants after immunoprecipitation (SN) and immunoprecipitates (IP) were probed with indicated VAMP7 antibodies (WB primary Ab). *VAMP7 band of interest (expected size of GFP-VAMP7 constructs: ~50 kDa). °Absence of band at VAMP7 corresponding size. **(B)** VAMP7 was immunoprecipitated from wild-type ("WT") and VAMP7 knock-out (KO) mouse cortex lysates using rabbit anti-VAMP7 TG50 antibody and detected with mouse anti-VAMP7 158.2. Normal isotype IgG (IgM or IgGR) were used as negative control. Inputs (IN), supernatants after immunoprecipitation (SN) and immunoprecipitates (IP) were probed with mouse anti-VAMP7 158.2 antibody. *VAMP7 band of interest (expected size of GFP-VAMP7 constructs: ~50 kDa). °Absence of band at expected size. ~heavy and light chains of the antibody used for immunoprecipitation. AP = affinity purified.

Dataset 2. Raw images of additional experimental replicates for Figure 2, immunofluorescence experiments

<https://dx.doi.org/10.5256/f1000research.15707.d221361>

This dataset includes additional images from experimental replicates of the images presented in Figure 2. Immunofluorescence staining methods and quantification (lower panel) were performed as described for Figure 2. Images were taken at 40× objective. Bar, 15µm.

Dataset 3. Raw images of immunoprecipitation experiments for Figure 3, immunoprecipitation

<https://dx.doi.org/10.5256/f1000research.15707.d221362>

Uncropped data from Figure 3 (A) and replicate (C) for VAMP7 immunoprecipitation from Cos-7 cell lysate overexpressing GFP-tagged mouse, rat or human VAMP7 constructs. Uncropped immunoblotting data from Figure 3 (B) and replicate (D) for VAMP7 immunoprecipitation from WT and VAMP7 KO mouse cortex extracts. Antibodies used for immunoprecipitation and subsequent immunoblotting are indicated. Red dashed lines show GFP-VAMP7 protein and cropped region, respectively. IN=Input (50 µg in A and C, 100 µg in B and D); SN = supernatant after immunoprecipitation; IP = immunoprecipitate; * = GFP-VAMP7; ° = Absence of band at GFP-VAMP7 size (~50 kDa); ~: immunoglobulins.

Discussion

In this antibody survey, we used genome-edited VAMP7 KO HeLa cells to compare several commercially available and home-made antibodies using standard methods for western blotting and immunocytochemistry. Apart from poor reactivity/quality, high background observed in western blotting or non-specific signal of some tested antibodies in IF could be due to non-fully optimized technical procedures. For example, different blocking agent could be used, such as bovine serum albumin for immunoblot or immunofluorescence assays, or cells could have been fixed differently (methanol, glutaraldehyde). We also cannot totally exclude some batch effect for the poor signal observed with some commercial antibodies. Furthermore, we voluntarily restricted this survey to the newly generated VAMP7-depleted HeLa cell line generated in this study and tissues from KO mice. We chose to evaluate the background signal of these antibodies in KO cells, criteria of choice that is both crucial for any assay and not always convincingly characterized. Conducting the same study in a different cell type or species might have led to different conclusions regarding background and specificity. Although it might have reduced the overall background, particularly in the VAMP7 KO immunofluorescence conditions, we chose to use epifluorescence microscopy rather than confocal imaging in order to give a global overview of the signal obtained with these antibodies and because many studies still largely rely on wide field microscopy. With these words of caution and limitations, we conclude that 158.2 and TG50 antibodies appeared as the best performers in our assays.

The datasheet provided with the Synaptic Systems (158.2 –232 011) antibody indicates that it is specific for rat and mouse. Here we provide evidence that it is also able to specifically recognize VAMP7 in human HeLa cells, both in immunoblot and immunofluorescence assays. This is in good agreement with

the fact that human, rat and mouse VAMP7 protein sequences share more than 94% identity (alignment with www.uniprot.org website). MAb 158.2 also immunoprecipitated mouse, rat and human GFP-tagged VAMP7 (Figure 3A). Altogether, 158.2 thus appeared as a suitable antibody in all species and for all applications. However, this antibody may not be very sensitive, as it did not allow for the endogenous detection of low amounts of the protein, particularly in tissues (Figure 3). More generally, further comparative study should be conducted to formally assess the efficiency of this set of antibodies in non-human cell lines or tissues, particularly in immunocytochemistry. However, immunoprecipitation using TG50 (protein A purified serum available from Covalab, ref. pab01031-P) and 158.2 detection appeared as a valid strategy for specific isolation of the endogenous VAMP7.

Finally, we proposed here an easy profile comparison of WT and KO western blotting and immunocytochemistry signals to rank the quality of antibodies directed against membrane-associated proteins as a decision-making tool for more complex studies.

Data availability

Dataset 1. Raw images of experimental replicates for Figure 1, immunoblotting experiments. This dataset includes uncropped blots for all experimental replicates that are represented in Figure 1. Treatments and immunoblot methods were performed as outlined in Figure 1. Blots were probed with indicated anti-VAMP7 antibodies and anti-α-tubulin antibodies was used as a loading control. (A) Dataset used for Figure 1, with cropped regions in red dashed line. (B) Additional set of raw images of a replicate experiment. Quantification as performed in Figure 1 is shown in lower panel. Note that although signal intensity and background are different within these two replicates, the relative performance of the different tested antibodies remained the same. DOI: <https://doi.org/10.5256/f1000research.15707.d221360>²⁹.

Dataset 2. Raw images of additional experimental replicates for Figure 2, immunofluorescence experiments. This dataset includes additional images from experimental replicates of the images presented in Figure 2. Immunofluorescence staining methods and quantification (lower panel) were performed as described for Figure 2. Images were taken at 40× objective. Bar, 15µm. DOI: <https://doi.org/10.5256/f1000research.15707.d221361>³⁰.

Dataset 3. Raw images of immunoprecipitation experiments for Figure 3, immunoprecipitation. Uncropped data from Figure 3 (A) and replicate (C) for VAMP7 immunoprecipitation from Cos-7 cell lysate overexpressing GFP-tagged mouse, rat or human VAMP7 constructs. Uncropped immunoblotting data from Figure 3 (B) and replicate (D) for VAMP7 immunoprecipitation from WT and VAMP7 KO mouse cortex extracts. Antibodies used for immunoprecipitation and subsequent immunoblotting are indicated. Red dashed lines show GFP-VAMP7 protein and cropped region, respectively. IN=Input (50 µg in A and C, 100 µg in B and D); SN = supernatant after immunoprecipitation; IP = immunoprecipitate; * = GFP-VAMP7; ° = Absence

of band at GFP-VAMP7 size (~50 kDa); ~: immunoglobulins. DOI: <https://doi.org/10.5256/f1000research.15707.d221362>³¹.

Grant information

Work in our group was funded by grants from Association Française contre les Myopathies (Research Grant 16612), the French National Research Agency (NeuroImmunoSynapse ANR-13-BSV2-0018-02; MetDePaDi ANR-16-CE16-0012), the Ecole des Neurosciences de Paris (ENP), the Fondation pour la Recherche Médicale (FRM), Prix Coup d'Élan pour la recherche

française of the Fondation Bettencourt Schueller, awards of the Association Robert Debré pour la Recherche Médicale to T.G.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgements

We thank L. Danglot and C. Vannier for helpful discussion. We are grateful to Dr Maurizio D'Esposito (IGCB, Naples, Italy) for mouse VAMP7 plasmid.

References

- Daste F, Galli T, Tareste D: **Structure and function of longin SNAREs.** *J Cell Sci.* 2015; **128**(23): 4263–4272.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Proux-Gillardeaux V, Rudge R, Galli T: **The tetanus neurotoxin-sensitive and insensitive routes to and from the plasma membrane: fast and slow pathways?** *Traffic.* 2005; **6**(5): 366–373.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Wang Y, Li L, Hou C, *et al.*: **SNARE-mediated membrane fusion in autophagy.** *Semin Cell Dev Biol.* 2016; **60**: 97–104.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Chiaruttini G, Piperno GM, Jouve M, *et al.*: **The SNARE VAMP7 Regulates Exocytic Trafficking of Interleukin-12 in Dendritic Cells.** *Cell Rep.* 2016; **14**(11): 2624–2636.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Coco S, Raposo G, Martinez S, *et al.*: **Subcellular localization of tetanus neurotoxin-insensitive vesicle-associated membrane protein (VAMP)/VAMP7 in neuronal cells: evidence for a novel membrane compartment.** *J Neurosci.* 1999; **19**(22): 9803–9812.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Chaineau M, Danglot L, Galli T: **Multiple roles of the vesicular-SNARE TI-VAMP in post-Golgi and endosomal trafficking.** *FEBS Lett.* 2009; **583**(23): 3817–3826.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Burgo A, Proux-Gillardeaux V, Sotirakis E, *et al.*: **A molecular network for the transport of the TI-VAMP/VAMP7 vesicles from cell center to periphery.** *Dev Cell.* 2012; **23**(1): 166–180.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Scheuber A, Rudge R, Danglot L, *et al.*: **Loss of AP-3 function affects spontaneous and evoked release at hippocampal mossy fiber synapses.** *Proc Natl Acad Sci USA.* 2006; **103**(44): 16562–16567.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Hua Z, Leal-Ortiz S, Foss SM, *et al.*: **v-SNARE composition distinguishes synaptic vesicle pools.** *Neuron.* 2011; **71**(3): 474–487.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Danglot L, Zylbersztein K, Petkovic M, *et al.*: **Absence of TI-VAMP/Vamp7 leads to increased anxiety in mice.** *J Neurosci.* 2012; **32**(6): 1962–1968.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Gupton SL, Gertler FB: **Integrin signaling switches the cytoskeletal and exocytic machinery that drives neuritogenesis.** *Dev Cell.* 2010; **18**(5): 725–736.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Wang G, Nola S, Bovio S, *et al.*: **Biomechanical Control of Lysosomal Secretion Via the VAMP7 Hub: A Tug-of-War between VARP and LRRK1.** *iScience.* 2018; **4**: 127–143.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Verderio C, Cagnoli C, Bergami M, *et al.*: **TI-VAMP/VAMP7 is the SNARE of secretory lysosomes contributing to ATP secretion from astrocytes.** *Biol Cell.* 2012; **104**(4): 213–228.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Fader CM, Aguilera MO, Colombo ML: **ATP is released from autophagic vesicles to the extracellular space in a VAMP7-dependent manner.** *Autophagy.* 2012; **8**(12): 1741–1756.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Danglot L, Chaineau M, Dahan M, *et al.*: **Role of TI-VAMP and CD82 in EGFR cell-surface dynamics and signaling.** *J Cell Sci.* 2010; **123**(Pt 5): 723–735.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Ghosh D, Pinto S, Danglot L, *et al.*: **VAMP7 regulates constitutive membrane incorporation of the cold-activated channel TRPM8.** *Nat Commun.* 2016; **7**: 10489.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Williams KC, Coppolino MG: **Phosphorylation of membrane type 1-matrix metalloproteinase (MT1-MMP) and its vesicle-associated membrane protein 7 (VAMP7)-dependent trafficking facilitate cell invasion and migration.** *J Biol Chem.* 2011; **286**(50): 43405–43416.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Proux-Gillardeaux V, Raposo G, Irinopoulou T, *et al.*: **Expression of the Longin domain of TI-VAMP impairs lysosomal secretion and epithelial cell migration.** *Biol Cell.* 2007; **99**(5): 261–271.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Steffen A, Le Dez G, Poincloux R, *et al.*: **MT1-MMP-dependent invasion is regulated by TI-VAMP/VAMP7.** *Curr Biol.* 2008; **18**(12): 926–931.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Molino D, Nola S, Lam SM, *et al.*: **Role of tetanus neurotoxin insensitive vesicle-associated membrane protein in membrane domains transport and homeostasis.** *Cell Logist.* 2015; **5**(1): e1025182.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ran FA, Hsu PD, Wright J, *et al.*: **Genome engineering using the CRISPR-Cas9 system.** *Nat Protoc.* 2013; **8**(11): 2281–2308.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Park J, Bae S, Kim JS: **Cas-Designer: a web-based tool for choice of CRISPR-Cas9 target sites.** *Bioinformatics.* 2015; **31**(24): 4014–4016.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Martinez-Arca S, Alberts P, Zahraoui A, *et al.*: **Role of tetanus neurotoxin insensitive vesicle-associated membrane protein (TI-VAMP) in vesicular transport mediating neurite outgrowth.** *J Cell Biol.* 2000; **149**(4): 889–900.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Galli T, Zahraoui A, Vaidyanathan VV, *et al.*: **A novel tetanus neurotoxin-insensitive vesicle-associated membrane protein in SNARE complexes of the apical plasma membrane of epithelial cells.** *Mol Biol Cell.* 1998; **9**(6): 1437–1448.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Martinez-Arca S, Proux-Gillardeaux V, Alberts P, *et al.*: **Ectopic expression of syntaxin 1 in the ER redirects TI-VAMP- and cellubrevin-containing vesicles.** *J Cell Sci.* 2003; **116**(Pt 13): 2805–2816.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kuster A, Nola S, Dingli F, *et al.*: **The Q-soluble N-Ethylmaleimide-sensitive Factor Attachment Protein Receptor (Q-SNARE) SNAP-47 Regulates Trafficking of Selected Vesicle-associated Membrane Proteins (VAMPs).** *J Biol Chem.* 2015; **290**(47): 28056–28069.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Muzerelle A, Alberts P, Martinez-Arca S, *et al.*: **Tetanus neurotoxin-insensitive vesicle-associated membrane protein localizes to a presynaptic membrane compartment in selected terminal subsets of the rat brain.** *Neuroscience.* 2003; **122**(1): 59–75.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Advani RJ, Yang B, Prekeris R, *et al.*: **VAMP-7 mediates vesicular transport from endosomes to lysosomes.** *J Cell Biol.* 1999; **146**(4): 765–776.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Verraes A, Cholley B, Galli T, *et al.*: **Dataset 1 in: Comparative study of commercially available and homemade anti-VAMP7 antibodies using CRISPR/Cas9-depleted HeLa cells and VAMP7 knockout mice.** *F1000Research.* 2018. <http://www.doi.org/10.5256/f1000research.15707.d221360>
- Verraes A, Cholley B, Galli T, *et al.*: **Dataset 2 in: Comparative study of commercially available and homemade anti-VAMP7 antibodies using CRISPR/Cas9-depleted HeLa cells and VAMP7 knockout mice.** *F1000Research.* 2018. <http://www.doi.org/10.5256/f1000research.15707.d221361>
- Verraes A, Cholley B, Galli T, *et al.*: **Dataset 3 in: Comparative study of commercially available and homemade anti-VAMP7 antibodies using CRISPR/Cas9-depleted HeLa cells and VAMP7 knockout mice.** *F1000Research.* 2018. <http://www.doi.org/10.5256/f1000research.15707.d221362>

Open Peer Review

Current Peer Review Status: ? ✓

Version 1

Reviewer Report 27 November 2018

<https://doi.org/10.5256/f1000research.17141.r40504>

© 2018 Coppolino M et al. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Marc G. Coppolino

Department of Molecular and Cellular Biology, University of Guelph, Guelph, ON, Canada

Olivia Grafinger

Department of Molecular and Cellular Biology, University of Guelph, Guelph, Ontario, Canada

This study assessed the performance of four monoclonal and four polyclonal antibodies to VAMP7. Six commercially available antibodies, and two antibodies generated by the researchers, were compared by Western blot, immunocytochemistry and immunoprecipitation. Lysates from HeLa and HeLa-VAMP7 knockout cells, and extracts from wildtype and VAMP7 knockout mouse brain tissue were used to examine specificity and background of the antibodies. The findings indicate some of the antibodies outperform comparators in blotting, immunocytochemistry, with reduced background and higher specificity, and perform well in immunoprecipitations. This work offers careful characterization of VAMP7 antibodies for use by other researchers in making decisions about antibody use for analytical studies of this protein.

The study is well conceived and executed. The approaches used are suitable, and the description of work is adequately detailed. Data are clearly presented, and for the most part conclusions are reasonable.

There are a few points that should be addressed to strengthen the study:

1. For the Western blot analyses, the antibody concentrations used (ug/ml) should be provided to allow more accurate comparison of antibody performance.
2. In the IF specificity index graph (Fig. 2C), it appears that the authors only measured the intensity profile in one cell for each antibody. These results would be more convincing if the intensity profiles of multiple cells were shown, as there can be variations in staining patterns that can arise from different cell morphologies, etc.
3. It is not clear why some of the antibodies, which performed reasonably well in Western blot and immunocytochemistry (Creative Diagnostics, Synaptic Systems), were not analysed in immunoprecipitations. An explanation for this, or information on their performance in this

regard, would be useful.

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Are sufficient details of materials, methods and analysis provided to allow replication by others?

Partly

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

We confirm that we have read this submission and believe that we have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 26 October 2018

<https://doi.org/10.5256/f1000research.17141.r39566>

© 2018 Banham A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Alison H. Banham

Nuffield Division of Clinical Laboratory Sciences, Radcliffe Department of Medicine, John Radcliffe Hospital, University of Oxford, Oxford, UK

This study has compared the specificity of a panel of antibodies against VAMP7, a member of the SNARE protein family. Using CRISPR/Cas9 to create a HeLa-VAMP7 knockout cell line together with knockout mouse tissues the authors effectively demonstrate that the panel of antibodies are able to recognise the endogenous VAMP7 protein, the majority performing effectively in both WB and IF. Image analysis was performed to profile the signal-to-noise index in both techniques. While this does provide quantitative data with which to prioritise the antibodies, I reached the same conclusions without needing the profiling data. This approach may indeed have utility in other projects but perhaps consideration should be given to incorporating a measure for antibody sensitivity. Antibodies with similar specificity and different levels of sensitivity may have a similar specificity index score but may not be equally useful. Two antibodies were also validated for their ability to IP recombinant orthologous VAMP7 proteins and one for IP of the endogenous protein. This is a useful comparative study that will be of interest to other researchers sourcing antibodies to further characterise this molecule.

Comments:

1) P3: For clarity I would suggest rephrasing 'Invalidation' of VAMP7 as inactivation, knockout or deletion.

2) P5 and P9 bottom left column: Readers are referred to Table 3 for dilutions of primary antibodies yet this information seems to be provided in Table 1.

3) While the image analysis provides quantitative information, the best antibodies identified using this approach were also easily identifiable by looking at the WB and IF images presented. With regard to the WB data in Figure 1, if just using the WB specificity index then there potentially appeared to be relatively little to choose between the two best antibodies. However, by eye, TG50 appears to have much greater sensitivity for VAMP7 detection, yet this distinction is not conveyed by a score which focuses on signal-to-noise.

4) In Figure 2 – there is clearly significant heterogeneity in the distribution of VAMP7 both within and between individual HeLa cells. It is not clear from the methods whether data from individual cells or a large number were used to generate the IF specificity index and this information should be provided. The data would be more robust if multiple cells were sampled.

5) Figure 2 indicates Cell Signaling 14811 performed poorly, however on the supplier's site this antibody (D4D5J) is not recommended for IF and it is clearly stated by the authors in the results that an alternative (D8Y1R) was used. This might just be an error in labelling on the figure? However, the text describing these IF data also refers to the antibody being used as 14811 (which is D4D5J). This needs to be clarified and it would be helpful within the manuscript and figures to consistently refer to either the clone names or the catalogue number when describing an antibody.

6) In the discussion the authors correctly comment that not using individually optimised conditions for each antibody in IF may disadvantage some of the reagents. I think it would have been more helpful to other researchers wanting to know which antibody works best for IF to be able to compare images obtained using an optimal staining protocol, such as that determined by each manufacturer.

7) In the discussion the authors comment on the sensitivity of 158.2 being insufficient to detect low levels of the endogenous VAMP7 protein by WB. When TG50 seemed to be more sensitive in WB analysis then it would perhaps have been the more obvious choice for the WB detection antibody. This point could be added to the discussion.

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Are sufficient details of materials, methods and analysis provided to allow replication by others?

Partly

Are all the source data underlying the results available to ensure full reproducibility?

Partly

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research