REVIEW

The role of NAFLD in cardiometabolic disease: an update

[version 1; peer review: 2 approved]

Sarah Faasse, Hayley Braun, Miriam Vos

Health Sciences Research Building, Emory University, 1760 Haygood Drive, W-450, Atlanta, GA 30322, USA

V1 First published: 09 Feb 2018, 7(F1000 Faculty Rev):170
<https://doi.org/10.12688/f1000research.12028.1>

Latest published: 09 Feb 2018, 7(F1000 Faculty Rev):170
<https://doi.org/10.12688/f1000research.12028.1>

Abstract

Non-alcoholic fatty liver disease (NAFLD) is the most common cause of chronic liver disease in the world, yet the complex pathogenesis remains to be fully elucidated. The prevalence of NAFLD has risen precipitously in recent years and is now a leading indication for liver transplantation. New waitlist registrants with non-alcoholic steatohepatitis-induced cirrhosis increased by 170% from 2004 to 2013. In addition, patients with NAFLD are at increased risk of both cardiovascular disease and type II diabetes. In this update, recent studies contributing to the understanding of the place of NAFLD in cardiometabolic disease will be discussed.

Keywords

non-alcoholic fatty liver disease, cardiometabolic disease, type 2 diabetes mellitus

Open Peer Review

Approval Status

	1	2
version 1 09 Feb 2018		

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Giovanni Targher**, University and Azienda Ospedaliera Universitaria Integrata of Verona, Verona, Italy
2. **Nicola Santoro**, Yale School of Medicine, New Haven, USA

Any comments on the article can be found at the end of the article.

Corresponding author: Miriam Vos (mvos@emory.edu)

Author roles: **Faasse S:** Conceptualization, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Braun H:** Conceptualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Vos M:** Conceptualization, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: SF and HB declare that they have no competing interests. MBV has NAFLD-related research support or in-kind research services from Resonance Health, Nutrition Science Initiative, AMRA, Siemens, Shire, Target PharmaSolutions, Labcorp, and Perspectum and serves as an NAFLD consultant for Allergan, AMRA, Boehringer Ingelheim, Bristol-Myers Squibb, Immuron, Intercept, Shire, and Target PharmaSolutions.

Grant information: This work was supported through a grant provided by the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK) R21 HD0809056-01.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2018 Faasse S *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Faasse S, Braun H and Vos M. **The role of NAFLD in cardiometabolic disease: an update [version 1; peer review: 2 approved]** F1000Research 2018, 7(F1000 Faculty Rev):170 <https://doi.org/10.12688/f1000research.12028.1>

First published: 09 Feb 2018, 7(F1000 Faculty Rev):170 <https://doi.org/10.12688/f1000research.12028.1>

Introduction

Non-alcoholic fatty liver disease (NAFLD), now the most common liver disease in the world, is defined as the accumulation of fat in the liver beyond 5% of total liver weight in the absence of excessive alcohol use and other pathogenic sources such as viral hepatitis, Wilson disease, alpha-1-antitrypsin deficiency, autoimmune hepatitis, or other metabolic diseases¹⁻⁴. The estimated prevalence of NAFLD in the US is now 19%, and worldwide prevalence estimates are between 25.24 and 32.5%^{5,6}. The marked increase in NAFLD prevalence has occurred concomitantly with increases in rates of obesity and the metabolic syndrome (MS) worldwide. Obesity itself is heterogenic, and a greater understanding of the relevance of body fat distribution patterns to NAFLD has developed recently as a growing number of studies using magnetic resonance imaging (MRI) to measure fat distribution have been published⁷⁻⁹. The previous mainstay was dual-energy x-ray absorptiometry, useful for visceral fat¹⁰, muscle mass, and total adiposity, but it misses hepatic fat. Body mass index (BMI) can both underestimate disease, as in the case of metabolically obese, normal-weight individuals¹¹, and overestimate disease, as in the case of obese (BMI of more than 30 kg/m²) but metabolically healthy people.

People in the metabolically unhealthy category, regardless of BMI, tend to have increased visceral adipose tissue (VAT)¹². VAT is metabolically active and has been thought to be complicit in the dysmetabolic syndrome¹³. Individuals with a higher proportion of VAT are more likely to have NAFLD¹⁴ and a higher risk of associated dysmetabolic disease^{15,16}. This leads to questions of which drives which: hepatic fat leading to VAT, or vice versa? There are two studies that demonstrate in both adults and adolescents that intrahepatic fat, rather than VAT, is most closely associated with insulin resistance (IR), suggesting that intrahepatic fat is the primary driver of metabolic dysfunction in the centrally located adiposity phenotype^{7,8}. Within the liver, *de novo* lipogenesis (DNL) is the primary dysfunction of lipid metabolism leading to steatosis accumulation. In healthy metabolism, DNL is not a major contributor of lipids in the liver. However, in NAFLD, DNL is abnormally upregulated¹⁷, and it increases further in the setting of weight gain¹⁸. In NAFLD, approximately one fourth of the fat in the liver is directly from DNL¹⁹.

Amino acid metabolism is another important area of disturbance in NAFLD, and newer techniques such as high-throughput, high-resolution metabolomics have advanced studies in this area. Specifically, levels of branched chain amino acids (leucine, isoleucine, and valine) and aromatic amino acids (tyrosine) have been found to be altered in NAFLD^{20,21}. Notably, a similar amino acid profile was also found to predict risk of future type 2 diabetes mellitus (T2DM)²². In pediatric patients with NAFLD, tyrosine metabolism was found to be most significantly altered in adolescent patients, and a recent study shows that elevated levels of tyrosine, glutamate, and glutamate/(serine+glycine) ratio (that is, GSG index) were associated with IR and that alanine, glutamate, isoleucine, and valine were increased in non-obese patients with NAFLD^{21,23}. More studies are needed to understand whether dysregulated amino acids represent promising biomarkers or future therapeutic targets for NAFLD.

Genetic factors

As is typical of many chronic diseases, NAFLD is a genetic condition with an environmental trigger that induces the expression of the phenotype. In the case of NAFLD, this trigger appears to be, at a minimum, a hypercaloric condition. The most established genotype complicit in liver steatosis is a mutation of the rs738409 allele of patatin-like phospholipase domain (PNPLA3)²⁴. The mutation is a single-nucleotide polymorphism (SNP) of codon 148, resulting in a missense mutation of isoleucine to methionine (I148M)²⁵. The homozygous variant is most commonly found in Hispanics. A recent article found that adiposity potentiates the effect of this missense mutation, resulting in a higher risk for development of NAFLD as BMI increases in patients with this particular variant²⁶. Interestingly, another variant allele of PNPLA3 (rs6006460) that results in a serine-to-isoleucine (S453I) missense appears to have a hepatoprotective effect, associated with lower intrahepatic fat content, and is most commonly found in African-Americans, an ethnic group that has a relatively low prevalence of NAFLD²⁷. Other SNPs have been identified in the pathogenesis of NAFLD, glucokinase regulatory protein (*GCKR*) rs1260326²⁸, and transmembrane 6 superfamily member 2 (*TM6SF2*) rs58542926²⁹, and when assessed with PNPLA3 rs6006460, these three SNPs seem to have an additive effect on determining intrahepatic fat content²⁹.

Prenatal exposure

It is vital to consider NAFLD within the life span of cardiometabolic disease as it is not an isolated disease (Figure 1)³⁰. Some important studies have emerged that suggest prenatal origins for NAFLD. Friedman *et al.* conducted an elegant set of experiments in rhesus monkeys and demonstrated that a high-fat diet in insulin-resistant pregnant monkeys predicted fatty liver in the offspring, regardless of the diet (healthy or not) given to the infant monkeys^{31,32}. Limited but intriguing evidence in humans supports this finding. Two recent MRI-based studies found that higher maternal BMI positively correlates with both higher postnatal infant adipose tissue and intrahepatic fat content, suggesting that the burden of intrahepatic fat begins *in utero*^{33,34}. Similarly, infants born to obese mothers with gestational diabetes (GDM) had, on average, 68% more hepatic fat content by MRI quantification than those born to normal-weight women³⁴. An autopsy study of stillborn babies showed that histopathologic hepatic steatosis was significantly more prevalent in children whose mothers had maternal GDM (79%) compared with controls (17%) ($P < 0.001$)³⁵. Together, these studies suggest that maternal IR may program the liver during development and drive hepatocyte function toward steatosis. More information is needed regarding maternal environmental exposures and the effect on NAFLD predisposition in children.

Non-alcoholic fatty liver disease in childhood and adolescence

Moving forward to early childhood, there are currently little data on the status of the liver in children who are one to seven years old, but markers of MS can be seen in preschool-aged children. Prenatal and perinatal exposures of GDM and high birth weight are both established risk factors for MS in childhood³⁶. In addition, both high and low birth weight are correlated not only

Figure 1. Temporality of risk factors, comorbidities, and potential consequences of non-alcoholic fatty liver disease. These findings suggest a common exposure or complementary or additive relationship between non-alcoholic fatty liver disease and other metabolic diseases. BMI, body mass index; NASH, non-alcoholic steatohepatitis.

with NAFLD but also with increased risk of more advanced liver disease³⁷. Some evidence for NAFLD in preschool-aged children within high risk groups also exists. In a study mostly of Hispanic children in an obesity clinic, about one fourth of children starting at preschool age had elevated alanine aminotransferase (ALT) for age, and this proportion rose to nearly half by middle-school age³⁸.

NAFLD in most children is diagnosed in the peripubertal period. During puberty, IR transiently increases^{39,40}, and children who are obese or have existing IR have a greater increase in IR during puberty, which may contribute to greater pathology in the liver⁴¹. Histologically, children in puberty can display greater hepatic steatosis and more severe portal inflammation than adults, whereas lobular inflammation and ballooning appear milder in children compared with adults⁴².

Non-alcoholic fatty liver disease as a risk factor of cardiometabolic disease

The increase in IR and worsening of NAFLD can be somewhat of a “chicken or the egg” question. Few longitudinal studies exist in children to help answer this, but in a study of Japanese adults, after adjustment for other confounding factors, patients with fatty liver were more likely (hazard ratio (HR) 1.49) to become dysmetabolic compared with their counterparts without fatty liver⁴³. This supports the view of NAFLD as a driver of ongoing IR, visceral adiposity, and metabolic dysfunction^{15,44}.

There is convincing evidence that NAFLD is an independent risk factor for cardiovascular disease events even after adjustment for conventional risk factors (odds ratio 4.12, 95% confidence interval (CI) 1.58 to 10.75, $P = 0.004$)⁴⁵. Significantly more patients with NAFLD had either coronary artery disease (7.5% versus 1.4%) or stroke (0.9% versus 0.2%) than non-NAFLD control patients⁴⁶. Data suggest that cardiac intima-media thickness (CIMT), an early predictor of atherosclerosis, and Framingham risk score significantly increase across quartiles of the fatty liver index (an accurate predictor of hepatic steatosis on ultrasonography)^{47,48}. In addition, patients who developed hepatic steatosis showed an increase in CIMT whereas those

without did not see a significant change⁴⁷. In regard to cardiac dysfunction risk, patients with NAFLD, particularly those with non-alcoholic steatohepatitis, have worsened right ventricular function compared with non-NAFLD counterparts⁴⁹. Coupled with type 2 diabetes, the risk for left ventricular dysfunction is also higher: patients with both T2DM and NAFLD have greater left ventricular diastolic dysfunction than those with T2DM alone⁵⁰.

There is also evidence that NAFLD may increase risk of T2DM^{44,51}. One study found the prevalence of T2DM to be higher in patients with NAFLD when compared with their obese, non-NAFLD counterparts (6.5% versus <1%). Notably, girls with NAFLD were more likely to have T2DM than boys with NAFLD, despite an overall higher prevalence of NAFLD in boys⁵². A longitudinal cohort study found that patients with only MS had a significantly higher risk of developing T2DM than those with NAFLD and no MS but that patients with both NAFLD and MS had the greatest risk of developing T2DM, almost four times greater than those with only NAFLD and two times greater than those with only MS. In addition, among those with MS, NAFLD significantly increased risk of T2DM (HR 1.93, 95% CI 1.38–2.71)⁵³. These authors hypothesized that the absence of an independent association between NAFLD and T2DM in those patients without MS may be due to differences in the severity of NAFLD⁵³. This theory is further supported by an independent study that found that NAFLD severity was associated with increased T2DM incidence even after controlling for MS and other predictors of T2DM⁵⁴.

Conclusions

In summary, the intertwined stories of NAFLD and cardiometabolic dysregulation appear far more complex than previously hypothesized. In light of recent evidence suggesting that NAFLD may develop *in utero* and that NAFLD increases risk of both cardiovascular disease and T2DM, we contend that it is important to assess and treat NAFLD in childhood in order to prevent progression to these diseases. It is no longer possible to view NAFLD solely as a liver disease resulting from metabolic dysregulation; rather, it may be a causative factor in the spiral

toward T2DM and cardiovascular events. The typical NAFLD clinical assessment is currently confined to liver pathology (steatosis, fibrosis, and screening for consequential cirrhosis and hepatocellular carcinoma), but this growing body of evidence emphasizes the importance of a broader patient assessment and demonstrates that NAFLD is a disease with systemic pathophysiological consequences.

The practical applications of these conclusions are several. More knowledge is needed regarding hepatic fat in infants and young children and to understand whether hepatic steatosis disappears and reappears or is sustained from birth. This will require the development of a better and safer non-invasive measurement of fat in very young children. Furthermore, children in groups known to be at high risk for NAFLD should be studied longitudinally so that the onset of the liver disease can be better understood. This will be critical for developing a public health approach to prevent NAFLD and to test interventions that interrupt its course at the onset. Lastly, emerging evidence now suggests that, once children develop NAFLD, they are at high risk for the development of T2DM. This needs to be confirmed in

a large-scale, long-term longitudinal study of pediatric NAFLD, and if it is confirmed, this population could be given priority for T2DM prevention programs.

Competing interests

SF and HB declare that they have no competing interests. MBV has NAFLD-related research support or in-kind research services from Resonance Health, Nutrition Science Initiative, AMRA, Siemens, Shire, Target PharmaSolutions, Labcorp, and Perspectum and serves as an NAFLD consultant for Allergan, AMRA, Boehringer Ingelheim, Bristol-Myers Squibb, Immuron, Intercept, Shire, and Target PharmaSolutions.

Grant information

This work was supported through a grant provided by the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK) R21 HD0809056-01.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- Schwimmer JB, Deutsch R, Kahen T, *et al.*: **Prevalence of fatty liver in children and adolescents.** *Pediatrics*. 2006; **118**(4): 1388–93.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Welsh JA, Karpen S, Vos MB: **Increasing prevalence of nonalcoholic fatty liver disease among United States adolescents, 1988–1994 to 2007–2010.** *J Pediatr*. 2013; **162**(3): 496–500.e1.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Wong RJ, Aguilar M, Cheung R, *et al.*: **Nonalcoholic steatohepatitis is the second leading etiology of liver disease among adults awaiting liver transplantation in the United States.** *Gastroenterology*. 2015; **148**(3): 547–55.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Uppal V, Mansoor S, Furuya KN: **Pediatric Non-alcoholic Fatty Liver Disease.** *Curr Gastroenterol Rep*. 2016; **18**(5): 24.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Lazo M, Hernaez R, Eberhardt MS, *et al.*: **Prevalence of nonalcoholic fatty liver disease in the United States: the Third National Health and Nutrition Examination Survey, 1988–1994.** *Am J Epidemiol*. 2013; **178**(1): 38–45.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Younossi ZM, Koenig AB, Abdelatif D, *et al.*: **Global epidemiology of nonalcoholic fatty liver disease-Meta-analytic assessment of prevalence, incidence, and outcomes.** *Hepatology*. 2016; **64**(1): 73–84.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- D'Adamo E, Cali AM, Weiss R, *et al.*: **Central role of fatty liver in the pathogenesis of insulin resistance in obese adolescents.** *Diabetes Care*. 2010; **33**(8): 1817–22.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Fabbri E, Magkos F, Mohammed BS, *et al.*: **Intrahepatic fat, not visceral fat, is linked with metabolic complications of obesity.** *Proc Natl Acad Sci U S A*. 2009; **106**(36): 15430–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Baum T, Cordes C, Diekmeyer M, *et al.*: **MR-based assessment of body fat distribution and characteristics.** *Eur J Radiol*. 2016; **85**(8): 1512–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Kaul S, Rothney MP, Peters DM, *et al.*: **Dual-energy X-ray absorptiometry for quantification of visceral fat.** *Obesity (Silver Spring)*. 2012; **20**(6): 1313–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ruderman NB, Schneider SH, Berchtold P: **The “metabolically-obese,” normal-weight individual.** *Am J Clin Nutr*. 1981; **34**(8): 1617–21.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Neeland IJ, Ayers CR, Rohatgi AK, *et al.*: **Associations of visceral and abdominal subcutaneous adipose tissue with markers of cardiac and metabolic risk in obese adults.** *Obesity (Silver Spring)*. 2013; **21**(9): E439–47.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Fontana L, Eagon JC, Trujillo ME, *et al.*: **Visceral fat adipokine secretion is associated with systemic inflammation in obese humans.** *Diabetes*. 2007; **56**(4): 1010–3.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Caprio S, Perry R, Kursawe R: **Adolescent Obesity and Insulin Resistance: Roles of Ectopic Fat Accumulation and Adipose Inflammation.** *Gastroenterology*. 2017; **152**(7): 1638–46.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Ko YH, Wong TC, Hsu YY, *et al.*: **The Correlation Between Body Fat, Visceral Fat, and Nonalcoholic Fatty Liver Disease.** *Metab Syndr Relat Disord*. 2017; **15**(6): 304–11.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Lee S, Kuk JL: **Visceral fat is associated with the racial differences in liver fat between black and white adolescent boys with obesity.** *Pediatr Diabetes*. 2017; **18**(7): 660–663.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Lambert JE, Ramos-Roman MA, Browning JD, *et al.*: **Increased de novo lipogenesis is a distinct characteristic of individuals with nonalcoholic fatty liver disease.** *Gastroenterology*. 2014; **146**(3): 726–35.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Fabbri E, Tiemann Luecking C, Love-Gregory L, *et al.*: **Physiological Mechanisms of Weight Gain-Induced Steatosis in People With Obesity.** *Gastroenterology*. 2016; **150**(1): 79–81.e2.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Donnelly KL, Smith CI, Schwarzenberg SJ, *et al.*: **Sources of fatty acids stored in liver and secreted via lipoproteins in patients with nonalcoholic fatty liver disease.** *J Clin Invest*. 2005; **115**(5): 1343–51.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Lake AD, Novak P, Shipkova P, *et al.*: **Branched chain amino acid metabolism profiles in progressive human nonalcoholic fatty liver disease.** *Amino Acids*. 2015; **47**(3): 603–15.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Jin R, Banton S, Tran VT, *et al.*: **Amino Acid Metabolism is Altered in Adolescents with Nonalcoholic Fatty Liver Disease-An Untargeted, High Resolution Metabolomics Study.** *J Pediatr*. 2016; **172**: 14–19.e5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Wang TJ, Larson MG, Vasan RS, *et al.*: **Metabolite profiles and the risk of developing diabetes.** *Nat Med*. 2011; **17**(4): 448–53.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)

23. **F** Gaggini M, Carli F, Rosso C, *et al.*: **Altered amino acid concentrations in NAFLD: Impact of obesity and insulin resistance.** *Hepatology*. 2018; **67**(1): 145–158.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
24. **F** Arslanow A, Stokes CS, Weber SN, *et al.*: **The common PNPLA3 variant p.I148M is associated with liver fat contents as quantified by controlled attenuation parameter (CAP).** *Liver Int*. 2016; **36**(3): 418–26.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
25. Liu W, Baker RD, Bhatia T, *et al.*: **Pathogenesis of nonalcoholic steatohepatitis.** *Cell Mol Life Sci*. 2016; **73**(10): 1969–87.
[PubMed Abstract](#) | [Publisher Full Text](#)
26. **F** Stender S, Kozlitina J, Nordestgaard BG, *et al.*: **Adiposity amplifies the genetic risk of fatty liver disease conferred by multiple loci.** *Nat Genet*. 2017; **49**(6): 842–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
27. **F** Romeo S, Kozlitina J, Xing C, *et al.*: **Genetic variation in PNPLA3 confers susceptibility to nonalcoholic fatty liver disease.** *Nat Genet*. 2008; **40**(12): 1461–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
28. Santoro N, Zhang CK, Zhao H, *et al.*: **Variant in the glucokinase regulatory protein (GCKR) gene is associated with fatty liver in obese children and adolescents.** *Hepatology*. 2012; **55**(3): 781–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
29. **F** Goffredo M, Caprio S, Feldstein AE, *et al.*: **Role of TM6SF2 rs58542926 in the pathogenesis of nonalcoholic pediatric fatty liver disease: A multiethnic study.** *Hepatology*. 2016; **63**(1): 117–25.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
30. Cioffi CE, Welsh JA, Cleeton RL, *et al.*: **Natural History of NAFLD Diagnosed in Childhood: A Single-Center Study.** *Children (Basel)*. 2017; **4**(5): pii: E34.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
31. Thorn SR, Baquero KC, Newsom SA, *et al.*: **Early life exposure to maternal insulin resistance has persistent effects on hepatic NAFLD in juvenile nonhuman primates.** *Diabetes*. 2014; **63**(8): 2702–13.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
32. **F** Wesolowski SR, Kasmi KC, Jonscher KR, *et al.*: **Developmental origins of NAFLD: a womb with a clue.** *Nat Rev Gastroenterol Hepatol*. 2017; **14**(2): 81–96.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
33. Modi N, Murgasova D, Ruager-Martin R, *et al.*: **The influence of maternal body mass index on infant adiposity and hepatic lipid content.** *Pediatr Res*. 2011; **70**(3): 287–91.
[PubMed Abstract](#) | [Publisher Full Text](#)
34. Brumbaugh DE, Teare P, Cree-Green M, *et al.*: **Intrahepatic fat is increased in the neonatal offspring of obese women with gestational diabetes.** *J Pediatr*. 2013; **162**(5): 930–6.e1.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
35. **F** Patel KR, White FV, Deutsch GH: **Hepatic steatosis is prevalent in stillborns delivered to women with diabetes mellitus.** *J Pediatr Gastroenterol Nutr*. 2015; **60**(2): 152–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
36. Boney CM, Verma A, Tucker R, *et al.*: **Metabolic syndrome in childhood: association with birth weight, maternal obesity, and gestational diabetes mellitus.** *Pediatrics*. 2005; **115**(3): e290–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
37. **F** Newton KP, Feldman HS, Chambers CD, *et al.*: **Low and High Birth Weights Are Risk Factors for Nonalcoholic Fatty Liver Disease in Children.** *J Pediatr*. 2017; **187**: 141–146.e1.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
38. Beacher DR, Ariza AJ, Fishbein MH, *et al.*: **Screening for elevated risk of liver disease in preschool children (aged 2-5 years) being seen for obesity management.** *SAGE Open Med*. 2014; **2**: 2050312114555211.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
39. Guzzaloni G, Grugni G, Mazzilli G, *et al.*: **Comparison between beta-cell function and insulin resistance indexes in prepubertal and pubertal obese children.** *Metabolism*. 2002; **51**(8): 1011–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
40. Moran A, Jacobs DR Jr, Steinberger J, *et al.*: **Insulin resistance during puberty: results from clamp studies in 357 children.** *Diabetes*. 1999; **48**(10): 2039–44.
[PubMed Abstract](#) | [Publisher Full Text](#)
41. Cree-Green M, Triolo TM, Nadeau KJ: **Etiology of insulin resistance in youth with type 2 diabetes.** *Curr Diab Rep*. 2013; **13**(1): 81–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
42. Takahashi Y, Inui A, Fujisawa T, *et al.*: **Histopathological characteristics of non-alcoholic fatty liver disease in children: Comparison with adult cases.** *Hepatol Res*. 2011; **41**(11): 1066–74.
[PubMed Abstract](#) | [Publisher Full Text](#)
43. **F** Hashimoto Y, Hamaguchi M, Fukuda T, *et al.*: **Fatty liver as a risk factor for progression from metabolically healthy to metabolically abnormal in non-overweight individuals.** *Endocrine*. 2017; **57**(1): 89–97.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
44. **F** Adams LA, Anstee QM, Tilg H, *et al.*: **Non-alcoholic fatty liver disease and its relationship with cardiovascular disease and other extrahepatic diseases.** *Gut*. 2017; **66**(6): 1138–53.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
45. Hamaguchi M, Kojima T, Takeda N, *et al.*: **Nonalcoholic fatty liver disease is a novel predictor of cardiovascular disease.** *World J Gastroenterol*. 2007; **13**(10): 1579–84.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
46. **F** Mahfood Haddad T, Hamdeh S, Kamnathareddy A, *et al.*: **Nonalcoholic fatty liver disease and the risk of clinical cardiovascular events: A systematic review and meta-analysis.** *Diabetes Metab Syndr*. 2017; **11** Suppl 1: S209–S216.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
47. **F** Pais R, Giral P, Khan JF, *et al.*: **Fatty liver is an independent predictor of early carotid atherosclerosis.** *J Hepatol*. 2016; **65**(1): 95–102.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
48. **F** Targher G, Byrne CD, Lonardo A, *et al.*: **Non-alcoholic fatty liver disease and risk of incident cardiovascular disease: A meta-analysis.** *J Hepatol*. 2016; **65**(3): 589–600.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
49. **F** Sunbul M, Kivrak T, Durmus E, *et al.*: **Nonalcoholic Steatohepatitis Score is an Independent Predictor of Right Ventricular Dysfunction in Patients with Nonalcoholic Fatty Liver Disease.** *Cardiovasc Ther*. 2015; **33**(5): 294–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
50. **F** Mantovani A, Pernigo M, Bergamini C, *et al.*: **Nonalcoholic Fatty Liver Disease Is Independently Associated with Early Left Ventricular Diastolic Dysfunction in Patients with Type 2 Diabetes.** *PLoS One*. 2015; **10**(8): e0135329.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
51. **F** Targher G, Marchesini G, Byrne CD: **Risk of type 2 diabetes in patients with non-alcoholic fatty liver disease: Causal association or epiphenomenon?** *Diabetes Metab*. 2016; **42**(3): 142–56.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
52. **F** Newton KP, Hou J, Crimmins NA, *et al.*: **Prevalence of Prediabetes and Type 2 Diabetes in Children With Nonalcoholic Fatty Liver Disease.** *JAMA Pediatr*. 2016; **170**(10): e161971.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
53. **F** Bae JC, Kim SK, Han JM, *et al.*: **Additive effect of non-alcoholic fatty liver disease on the development of diabetes in individuals with metabolic syndrome.** *Diabetes Res Clin Pract*. 2017; **129**: 136–43.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
54. Park SK, Seo MH, Shin HC, *et al.*: **Clinical availability of nonalcoholic fatty liver disease as an early predictor of type 2 diabetes mellitus in Korean men: 5-year prospective cohort study.** *Hepatology*. 2013; **57**(4): 1378–83.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. **Nicola Santoro**

Department of Pediatrics, Yale School of Medicine, New Haven, CT, USA

Competing Interests: No competing interests were disclosed.

2. **Giovanni Targher**

Division of Endocrinology, Diabetes and Metabolism, Department of Medicine, University and Azienda Ospedaliera Universitaria Integrata of Verona, Verona, Italy

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research