RESEARCH ARTICLE

Intraocular pressure elevation precedes a phagocytosis decline in a model of pigmentary glaucoma [version 1; peer review: 1 not approved]

Yalong Dang ¹, Susannah Waxman¹, Chao Wang ^{1,2}, Priyal Shah ¹, Ralitsa T. Loewen¹, Nils A. Loewen ¹

¹Department of Ophthalmology, University of Pittsburgh School of Medicine, Pittsburgh, PA, 15213, USA

²Department of Ophthalmology, Xiangya Hospital of Central South University, Changsha, Hunan, 410008, China

V1 First published: 12 Feb 2018, 7:174
<https://doi.org/10.12688/f1000research.13797.1>

Latest published: 09 Apr 2018, 7:174
<https://doi.org/10.12688/f1000research.13797.2>

Abstract

Background: Outflow regulation and phagocytosis are key functions of the trabecular meshwork (TM), but it is not clear how the two are related in secondary open angle glaucomas characterized by an increased particle load. We hypothesized that diminished TM phagocytosis is not the primary cause of early ocular hypertension and recreated pigment dispersion in a porcine *ex vivo* model.

Methods: Sixteen porcine anterior chamber cultures received a continuous infusion of pigment granules (Pg), while 16 additional anterior chambers served as controls (C). Pressure transducers recorded the intraocular pressure (IOP). The phagocytic capacity of the trabecular meshwork was determined by fluorescent microspheres.

Results: The baseline IOPs in Pg and C were similar ($P=0.82$). A significant IOP elevation occurred in Pg at 48, 120, and 180 hours (all $P < 0.01$, compared to baseline). The pigment did not cause a reduction in TM phagocytosis at 48 hours, when the earliest IOP elevation occurred, but at 120 hours onward ($P=0.001$ compared to C). This reduction did not result in an additional IOP increase at 120 or 180 hours compared to the first IOP elevation at 48 hours ($P>0.05$).

Conclusions: In this porcine model of pigmentary glaucoma, an IOP elevation occurs much earlier than when phagocytosis fails, suggesting that two separate mechanisms might be at work.

Keywords

Pigment dispersion glaucoma, aqueous outflow, trabecular meshwork, intraocular pressure, phagocytosis

Open Peer Review

Approval Status

	1	2
version 2		
(revision)		
09 Apr 2018	view	view
version 1		
12 Feb 2018	view	

1. **Sergio C Saccà** , RCCS San Martino
University Hospital, Genoa, Italy

2. **Don S. Minckler**, University of California,
Irvine, USA

Any reports and responses or comments on the article can be found at the end of the article.

This article is included in the **Eye Health** gateway.

Corresponding author: Nils A. Loewen (loewen.nils@gmail.com)

Author roles: **Dang Y:** Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Waxman S:** Data Curation, Investigation, Validation, Writing – Original Draft Preparation; **Wang C:** Data Curation, Formal Analysis, Investigation, Methodology, Validation; **Shah P:** Data Curation, Investigation, Methodology, Writing – Review & Editing; **Loewen RT:** Formal Analysis, Investigation, Methodology; **Loewen NA:** Conceptualization, Data Curation, Formal Analysis, Funding Acquisition, Investigation, Methodology, Project Administration, Resources, Supervision, Validation, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: NIH CORE Grant P30 EY08098 to the Department of Ophthalmology, from the Eye and Ear Foundation of Pittsburgh, and from an unrestricted grant from Research to Prevent Blindness, New York, NY; National Eye Institute K08EY022737 (NAL); Initiative to Cure Glaucoma of the Eye and Ear Foundation of Pittsburgh (NAL); Research to Prevent Blindness, Departmental Grant (NAL); the Wiegand Fellowship of the Eye and Ear Foundation (YD); an unrestricted grant from the Third Xiangya Hospital of Central South University for studying at the University of Pittsburgh (CW).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2018 Dang Y *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. Data associated with the article are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

How to cite this article: Dang Y, Waxman S, Wang C *et al.* **Intraocular pressure elevation precedes a phagocytosis decline in a model of pigmentary glaucoma [version 1; peer review: 1 not approved]** F1000Research 2018, 7:174 <https://doi.org/10.12688/f1000research.13797.1>

First published: 12 Feb 2018, 7:174 <https://doi.org/10.12688/f1000research.13797.1>

Introduction

The conventional outflow is guarded by the trabecular meshwork (TM), a complex three dimensional, layered tissue that contains variable amounts of extracellular matrix (ECM)¹. The aqueous passes into Schlemm's canal by paracytosis or giant vacuoles². Failure to maintain a normal cytoskeleton and homeostasis of aqueous outflow can cause ocular hypertension¹. For instance, pigment dispersion³ and corticosteroids can alter the actin cytoskeleton and cause TM cell contraction resulting in an elevation of intraocular pressure (IOP)^{3,4}. Conversely, relaxing the cytoskeleton, for instance by using a Rho kinase inhibitor, can reverse these effects^{5,6}.

Phagocytosis of debris is another key function of TM cells². However, its direct and short-term effects on IOP regulation remain poorly understood¹. Chronic exposure to pigment⁷, erythrocyte-derived ghost cells⁸, inflammatory cells⁹, photoreceptor outer segments¹⁰, lens and pseudoexfoliation material^{11,12} can lead to secondary glaucomas.

We recently developed an *ex vivo* pigmentary glaucoma (PG) model that recreates the IOP elevation, stress fiber formation, and phagocytosis reduction characteristics of human PG³. A gene expression analysis indicated an activation of the RhoA signaling pathway, and a downstream effect of tight junction formation negatively regulated by RhoA-mediated actin cytoskeletal reorganization³. In the current study, we hypothesized that ocular hypertension is the result of a reorganization of the actin cytoskeleton and occurs before phagocytosis declines.

Methods

Pig eye perfusion culture and pigmentary glaucoma model

This study was conducted in accordance with the Association for Research in Vision and Ophthalmology [Statement for the Use of Animals in Ophthalmic and Vision Research](#). Because no live vertebrate animals were used and pig eyes were acquired from a local abattoir (Thoma Meat Market, Saxonburg, PA), no Institutional Animal Care and Use approval was required.

Thirty-two porcine eyes were cultured within 2 hours of enucleation. Extraocular tissues were removed, and the eyes were decontaminated with 5% povidone-iodine solution (CAT# 3955-16, United States Pharmacopeia, Rockville, MD) for two minutes and washed three times in phosphate buffered saline (PBS). Posterior segments, lenses, and irises were removed and the anterior segments with intact TM mounted in the perfusion system as previously described^{3,13,14}. We used the same method to generate pigment granules as recently described in a model of pigmentary glaucoma (PG)³. Briefly, pigment granules were produced by subjecting the iris to freeze-thaw and resuspension washing before dilution of the stock to a final concentration of 1.67×10^7 particles/ml. Eyes in the pigment dispersion group were continuously perfused with pigment added to the culture medium for up to 180 hours (Pg) and compared to controls (C). The perfusate consisted of Dulbecco's modified Eagle media (DMEM, SH30284, HyClone, GE Healthcare, UK) supplemented with 1% FBS and 1% antibiotics (15240062, Thermo Fisher Scientific, Waltham, MA) at a constant rate of 3 μ l/min using

a microinfusion pump (PHD 22/2000; Harvard Apparatus, Holliston, MA). IOP was measured intracamerally by a pressure transducer (SP844; MEMSCAP, Skoppum, Norway) and recorded at two-minute intervals (LabChart, ADInstruments, Colorado Springs, CO). Baseline IOPs were obtained after IOP stabilization for 48 hours.

TM phagocytosis

The *in situ* TM phagocytosis was measured using an epifluorescence microscope after microsphere perfusion. In brief, a suspension of 0.5 μ m carboxylate-modified yellow-green fluorescent microspheres¹⁵ (CAT# F8813, Thermo Fisher, Waltham, MA) at 5×10^8 particles/ml was added to the perfusate at 48, 120, and 180 hours and perfused for 24 hours. The eyes were removed from their perfusion dishes, washed three times with pre-warmed PBS, secured again in the perfusion dishes, and placed upside down for imaging. The TM, visualized from the underside of the transparent perfusion dish, was photographed and measured by acquiring the images with a camera and epifluorescence equipped dissecting fluorescence microscope (SZX16, Olympus, Tokyo, Japan) at a 680x510 pixel resolution and a 200 ms exposure. The mean fluorescence intensity was quantified by [ImageJ](#) (Version 1.50i, NIH) as previously described¹⁶ at 48, 120, and 180 hours by measuring the fluorescence intensity in the TM.

To validate that the microspheres were phagocytosed by TM cells, the TM was dissected and digested with collagenase type IA (C9891, Sigma Aldrich, St. Louis, MO) at 2mg/ml and 1% FBS for 30 min at room temperature. The cells were filtered with a 70-micron cell strainer and resuspended in 0.5 ml of PBS. The percentage of TM cells that had ingested fluorescent microspheres was determined using flow cytometry.

To get a more accurate visualization of the phagocytosed microbeads, we used confocal microscopy. TM cells were seeded into the wells of a six-well plate and fixed with 4% PFA. The cell membranes were labeled with *Lycopersicon esculentum* agglutinin (TL; Texas red-conjugated; #TL-1176, Vector Laboratories, Inc., Burlingame, CA) at room temperature for 1 hour. The cell nuclei were counterstained with DAPI (D1306, Thermo Fisher Scientific, Waltham, MA). Photos and 3D videos were taken using an upright laser scanning confocal at 400x magnification (BX61, Olympus, Tokyo, Japan).

Histology

After the TM phagocytosis assay, the anterior segments were fixed with 4% PFA for 24 hours, washed three times with PBS, dehydrated in 70% ethanol, and embedded in paraffin. Sections were cut to a thickness of 5 μ m and stained with hematoxylin and eosin (H&E).

Statistics

Data were presented as the mean $\pm$ standard error and analyzed by [PASW Statistics 18](#) (SPSS Inc., Chicago, IL). The baseline IOP was compared to the other time points of the same eye using a paired *t*-test. Other quantitative data were analyzed by one-way ANOVA. A *p* value ≤ 0.05 was considered statistically significant.

Results

In H&E stained tissue sections, normal TM (Figure 1A) presented as a sparsely pigmented (red arrowheads), multilayered, porous tissue with Schlemm's canal-like segments within the aqueous plexus at the outer layer (black arrows). Pigment granules were seen phagocytosed by trabecular meshwork cells, particularly in the uveal TM, at 48, 120, and 180 hours (Figure 1B, C and D) but were not dense enough to physically obstruct any part of the conventional outflow system.

Baseline IOP in Pg was comparable to C (12.2 ± 0.9 mmHg vs. 11.9 ± 0.9 mmHg, $P=0.82$). Pigment dispersion caused a significant IOP elevation at 48, 120, and 180 hours (19.5 ± 1.4 mmHg, 20.2 ± 1.4 mmHg and 22.8 ± 0.8 mmHg, $P=0.001$, $P<0.001$ and $P=0.002$, compared to baseline) while IOPs in C remained steady (13.1 ± 1.1 mmHg, 12.0 ± 0.9 mmHg and 14.0 ± 1.5 mmHg, all p values >0.05 , compared to baseline) (Figure 2A).

By inverting the perfusion dishes and washing away the microspheres in the intertrabecular spaces, the TM phagocytosis was visualized and quantified under an upright dissecting fluorescence microscope. Pigment did not cause any change of phagocytosis during early ocular hypertension at 48 hours (Figure 2Bi-ii, $96.3 \pm 5.0\%$ compared to the control, $P=0.723$), but did cause a reduction at the later phases of 120 hours (Figure 2Biii-iv, $58.3 \pm 2.3\%$, $P=0.001$) and 180 hours (Figure 2Bv-vi, $62.5 \pm 5.1\%$, $P=0.026$). However, the declining phagocytosis did not result in further elevation of IOP at 120 and 180 hours compared to the initial IOP elevation at 48 hours (20.2 ± 1.4 mmHg and 22.8 ± 0.8 mmHg versus 19.5 ± 1.4 mmHg, both $P>0.05$).

The microsphere ingestion by TM cells was further assessed by flow cytometry and confocal microscopy. 28.1% of TM cells had

phagocytosed microbeads in a normal perfusion eye (Figure 3A) and the confocal microscopy confirmed them as being located within the cells (Figure 3B) aided by tomato lectin-stained cell membranes and DAPI-stained nuclei. Confocal imaging showed clusters of green fluorescent microspheres within the intracellular space with no microspheres in the intercellular space. The 3D video also suggested the microspheres were in fact phagocytized and not merely on top of or below them since the microbeads were in the same z plane as the cells (Supplementary Video 1).

Dataset 1. Raw unedited images of Figure 1

<http://dx.doi.org/10.5256/f1000research.13797.d192088>

They are representative of 17 slides for histology.

Dataset 2. Raw unedited images of Figure 2B

<http://dx.doi.org/10.5256/f1000research.13797.d192089>

They are representative of 31 pictures for phagocytosis measurement.

Dataset 3. Raw unedited images of Figure 3B

<http://dx.doi.org/10.5256/f1000research.13797.d192090>

Dataset 4. The FACS output file for Figure 3A

<http://dx.doi.org/10.5256/f1000research.13797.d192091>

Dataset 5. The raw IOP and phagocytosis measurements at all time points

<http://dx.doi.org/10.5256/f1000research.13797.d192092>

Figure 1. Histology. Normal trabecular meshwork (TM) (A) was a multilayer, strainer-like structure with few pigment deposits (red arrowheads). *Ex vivo* perfusion with pigment granules at $1.67 \times 10^7/\text{ml}$ caused significant TM pigmentation at 48 hours (B), 120 hours (C) and 180 hours (D). No apparent occlusion of the outflow tract was found.

Figure 2. Reduction of intraocular pressure and TM phagocytic activity by pigment dispersion. Baseline IOPs in the pigment group (n=16) and the control (n=16) are comparable (12.2 ± 0.9 mmHg vs. 11.9 ± 0.9 mmHg, $P=0.82$). Pigment caused a significant IOP elevation at 48 hours and onward (all $P < 0.05$) while the IOP in the control group showed no significant difference to baseline at any time point (all $P > 0.05$) when compared to the baseline (**A**). TM phagocytosis was visualized *in situ*. The mean fluorescence intensity in the TM region was quantified by NIH ImageJ. TM phagocytosis in the pigment group was comparable to the control at 48 hours ($P=0.723$), (**Bi–ii**) but showed sharp decreases at 120 hours (**Biii–iv**) and 180 hours ($P=0.001$ and $P=0.026$, respectively) (**Bv–vi**).

Figure 3. Validation of TM phagocytosis by flow cytometry and confocal microscopy. To further confirm that microspheres were phagocytosed, we digested a normal sample TM tissue into single cell suspension and sent for flow cytometry. The results suggested that 28.1% of the TM cells were actively phagocytic (**A**). We then seeded these cells into a six well plate to form monolayer. After labeling them with tomato lectin, the confocal imaging showed that clusters of green fluorescent microspheres were located in the intracellular but not in the intercellular space (**B**).

Discussion

Phagocytosis is a defining feature of TM cells¹⁷ and plays a central but poorly understood role in the pathogenesis of several types of secondary glaucoma that include pigment, erythrocytes and ghost cells, inflammatory cells, photoreceptor outer segments, lens and pseudoexfoliation material^{13,18,19}. Although TM phagocytosis can remove particles from the aqueous humor²⁰, the direct and short-term effects on outflow regulation remain insufficiently explained¹. In this study, we measured IOP and TM phagocytic activity in the presence of pigment granules at different time points and found IOP was significantly elevated as early as 48 hours after exposure to pigment granules. This was contrasted by a phagocytic activity in Pg not different from C before the decrease at 120 and 180 hours. A worsening decline of TM phagocytosis at 120 and 180 hours did not result in a further increase of IOP. This suggests that reduction in phagocytosis is a downstream and secondary effect of actin cytoskeletal reorganization.

Pigment treatment has previously been shown to cause ocular hypertension in part by reorganizing the TM actin cytoskeleton and not by physical obstruction of the outflow tract^{7,21}. We have recently reported that long, thick, and continuous TM actin bundles emerge as early as 24 hours after pigment exposure³ and replicate this observation in the present study. Histological characteristics of pigment dispersion in porcine eyes matched those seen in samples from pigmentary glaucoma patients^{21–23} showing that pigment particles were taken up by TM cells.

In summary, the results indicate the IOP elevation caused by pigment dispersion is not the direct result of a physical obstruction of outflow or a chronically overwhelmed phagocytosis. The reduction in phagocytosis considerably lags the evolving hypertension supporting the notion that these cytoskeletal changes occur early on and are separate from the impact of pigment on canonical phagocytosis pathways³.

Supplementary material

Supplementary Video 1. Visualization of microsphere ingestion by a 3D reconstruction with confocal microscopy. We took a series of z-stack confocal microscopy images to reconstruct a 3D video, showing that the fluorescent microspheres were neither on the top nor below, but phagocytized by the TM cells.

[Click here to access the data.](#)

Data availability

All the raw data generated or analyzed in this study are included in following datasets.

Dataset 1. Raw unedited images of [Figure 1](#). They are representative of 17 slides for histology. [10.5256/f1000research.13797.d192088²⁴](#)

Dataset 2. Raw unedited images of [Figure 2B](#). They are representative of 31 pictures for phagocytosis measurement. [10.5256/f1000research.13797.d192089²⁵](#)

Dataset 3. Raw unedited images of [Figure 3B](#). [10.5256/f1000research.13797.d192090²⁶](#)

Dataset 4. The FACS output file for [Figure 3A](#). [10.5256/f1000research.13797.d192091²⁷](#)

Dataset 5. The raw IOP and phagocytosis measurements at all time points. [10.5256/f1000research.13797.d192092²⁸](#)

Competing interests

No competing interests were disclosed.

Grant information

NIH CORE Grant P30 EY08098 to the Department of Ophthalmology, from the Eye and Ear Foundation of Pittsburgh, and from an unrestricted grant from Research to Prevent Blindness, New York, NY; National Eye Institute K08EY022737 (NAL); Initiative to Cure Glaucoma of the Eye and Ear Foundation of Pittsburgh (NAL); Research to Prevent Blindness, Departmental Grant (NAL); the Wiegand Fellowship of the Eye and Ear Foundation (YD); an unrestricted grant from the Third Xiangya Hospital of Central South University for studying at the University of Pittsburgh (CW).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

1. Llobet A, Gasull X, Gual A: **Understanding trabecular meshwork physiology: a key to the control of intraocular pressure?** *News Physiol Sci.* 2003; **18**(5): 205–9. [PubMed Abstract](#) | [Publisher Full Text](#)
2. Tamm ER: **The trabecular meshwork outflow pathways: structural and functional aspects.** *Exp Eye Res.* 2009; **88**(4): 648–55. [PubMed Abstract](#) | [Publisher Full Text](#)
3. Dang Y, Waxman S, Wang C, *et al.*: **Trabecular Meshwork Failure In A Model Of Pigmentary Glaucoma [Internet].** *bioRxiv.* 2017; 118448, [cited 2017 Apr 21]. [Reference Source](#)
4. Pattabiraman PP, Rao PV: **Mechanistic basis of Rho GTPase-induced extracellular matrix synthesis in trabecular meshwork cells.** *Am J Physiol Cell Physiol.* 2010; **298**(3): C749–63. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
5. Tanihara H, Inoue T, Yamamoto T, *et al.*: **Intra-ocular pressure-lowering effects**

- of a Rho kinase inhibitor, ripasudil (K-115), over 24 hours in primary open-angle glaucoma and ocular hypertension: a randomized, open-label, crossover study. *Acta Ophthalmol.* 2015; **93**(4): e254–60.
[PubMed Abstract](#) | [Publisher Full Text](#)
6. Tanihara H, Inoue T, Yamamoto T, *et al.*: Phase 1 clinical trials of a selective Rho kinase inhibitor, K-115. *JAMA Ophthalmol.* 2013; **131**(10): 1288–95.
[PubMed Abstract](#) | [Publisher Full Text](#)
 7. Epstein DL, Freddo TF, Anderson PJ, *et al.*: Experimental obstruction to aqueous outflow by pigment particles in living monkeys. *Invest Ophthalmol Vis Sci.* 1986; **27**(3): 387–95.
[PubMed Abstract](#)
 8. Campbell DG, Simmons RJ, Grant WM: Ghost cells as a cause of glaucoma. *Am J Ophthalmol.* 1976; **81**(4): 441–50.
[PubMed Abstract](#) | [Publisher Full Text](#)
 9. Moorthy RS, Mermoud A, Baerveldt G, *et al.*: Glaucoma associated with uveitis. *Surv Ophthalmol.* 1997; **41**(5): 361–94.
[PubMed Abstract](#) | [Publisher Full Text](#)
 10. Callender D, Jay JL, Barrie T: Schwartz-Matsuo syndrome: atypical presentation as acute open angle glaucoma. *Br J Ophthalmol.* 1997; **81**(7): 608–608.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 11. Lee RK: The molecular pathophysiology of pseudoexfoliation glaucoma. *Curr Opin Ophthalmol.* 2008; **19**(2): 95–101.
[PubMed Abstract](#) | [Publisher Full Text](#)
 12. Ritch R, Schlötzer-Schrehardt U, Konstas AG: Why is glaucoma associated with exfoliation syndrome? *Prog Retin Eye Res.* 2003; **22**(3): 253–75.
[PubMed Abstract](#) | [Publisher Full Text](#)
 13. Loewen RT, Roy P, Park DB, *et al.*: A Porcine Anterior Segment Perfusion and Transduction Model With Direct Visualization of the Trabecular Meshwork. *Invest Ophthalmol Vis Sci.* 2016; **57**(3): 1338–44.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 14. Dang Y, Waxman S, Wang C, *et al.*: Freeze-thaw decellularization of the trabecular meshwork in an ex vivo eye perfusion model. *PeerJ Preprints.* 2017; **5**: [cited 2017 Jan 24]. Report No.: e2736v1.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 15. Loewen RT, Brown EN, Scott G, *et al.*: Quantification of Focal Outflow Enhancement Using Differential Canalograms. *Invest Ophthalmol Vis Sci.* 2016; **57**(6): 2831–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 16. Dang Y, Waxman S, Wang C, *et al.*: Rapid learning curve assessment in an ex vivo training system for microincisional glaucoma surgery. *PeerJ Preprints.* 2017; [cited 2017 Jan 25]. Report No. 5: e2745v1.
[Publisher Full Text](#)
 17. Buller C, Johnson DH, Tschumper RC: Human trabecular meshwork phagocytosis. Observations in an organ culture system. *Invest Ophthalmol Vis Sci.* 1990; **31**(10): 2156–63.
[PubMed Abstract](#)
 18. Liton PB, Lin Y, Gonzalez P, *et al.*: Potential role of lysosomal dysfunction in the pathogenesis of primary open angle glaucoma. *Autophagy.* 2009; **5**(1): 122–4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 19. Zhang X, Ognibene CM, Clark AF, *et al.*: Dexamethasone inhibition of trabecular meshwork cell phagocytosis and its modulation by glucocorticoid receptor beta. *Exp Eye Res.* 2007; **84**(2): 275–84.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 20. Abu-Hassan DW, Acott TS, Kelley MJ: The Trabecular Meshwork: A Basic Review of Form and Function. *J Ocul Biol.* 2014; **2**(1): pii: http://fulltextarticles.avenonline.org/JOCB-2334-2838-02-0017.html.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 21. Gottanka J, Johnson DH, Grehn F, *et al.*: Histologic findings in pigment dispersion syndrome and pigmentary glaucoma. *J Glaucoma.* 2006; **15**(2): 142–51.
[PubMed Abstract](#) | [Publisher Full Text](#)
 22. Kupfer C, Kuwabara T, Kaiser-Kupfer M: The histopathology of pigmentary dispersion syndrome with glaucoma. *Am J Ophthalmol.* 1975; **80**(5): 857–62.
[PubMed Abstract](#) | [Publisher Full Text](#)
 23. Alvarado JA, Murphy CG: Outflow obstruction in pigmentary and primary open angle glaucoma. *Arch Ophthalmol.* 1992; **110**(12): 1769–78.
[PubMed Abstract](#) | [Publisher Full Text](#)
 24. Dang Y, Waxman S, Wang C, *et al.*: Dataset 1 in: Intraocular pressure elevation precedes a phagocytosis decline in a model of pigmentary glaucoma. *F1000Research.* 2018.
[Data Source](#)
 25. Dang Y, Waxman S, Wang C, *et al.*: Dataset 2 in: Intraocular pressure elevation precedes a phagocytosis decline in a model of pigmentary glaucoma. *F1000Research.* 2018.
[Data Source](#)
 26. Dang Y, Waxman S, Wang C, *et al.*: Dataset 3 in: Intraocular pressure elevation precedes a phagocytosis decline in a model of pigmentary glaucoma. *F1000Research.* 2018.
[Data Source](#)
 27. Dang Y, Waxman S, Wang C, *et al.*: Dataset 4 in: Intraocular pressure elevation precedes a phagocytosis decline in a model of pigmentary glaucoma. *F1000Research.* 2018.
[Data Source](#)
 28. Dang Y, Waxman S, Wang C, *et al.*: Dataset 5 in: Intraocular pressure elevation precedes a phagocytosis decline in a model of pigmentary glaucoma. *F1000Research.* 2018.
[Data Source](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 26 March 2018

<https://doi.org/10.5256/f1000research.14999.r32109>

© 2018 Saccà S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Sergio C Saccà

Department of Neuroscience and Sense Organs, RCCS San Martino University Hospital, Genoa, Italy

I read the article titled: Intraocular pressure elevation precedes a phagocytosis decline in a model of pigmentary glaucoma by Dang *et al.* The title of the article is very stimulating but the content has several things that should be corrected. The authors state that in the conventional way of outflow the aqueous passes into Schlemm's canal by paracytosis or giant vacuoles. This is a partial view that does not correspond to the truth. I invite the authors to read Saccà *et al.* The Outflow Pathway: A Tissue With Morphological and Functional Unity¹. Then the effects of the inhibitors of the Rho kinase inhibitor may have the ability to modify the metabolism of endothelial cells in the node in which the authors think it is open to question. These drugs have the ability to block TM motility by exposing more cells to aqueous humor. This is the reason why we initially witness a decline in IOP then this worsens because the exposed cells are blocked and are more exposed to apoptosis (to read Saccà *et al.* from DNA damage to functional changes of the trabecular meshwork in aging and glaucoma²). Also the hypothesis formulated by the authors to explain the pigmentary glaucoma is partial in the Trabecoloto the so-called pores have a very low meaning and above all the pigment does not obstruct anything. We do not work like sinks - I am amazed that today we can still believe that TM is a porous tissue, and above all, how it is possible to think that outflow is a passive phenomenon. In pigmentary glaucoma there are 2 fundamental pathogenic moments. The first concerns the back sheet of the iris, where the pigment comes from (a defect that involves the loss of the pigment itself) the second point concerns autophagy which is not able to ensure adequate cell homeostasis. So these cells first enter suffering and then die and hence glaucoma is why the cells do not work, not because it obstructs something. There are cells, not tubes. The cytoskeleton in all this has little to do with it: pigment granules are only a metabolic burden that in the case of pigment dispersal syndrome is absorbed by a functioning autophagy, while in pigment glaucoma, autophagy does not work. Finally, I remember that the changes in the cytoskeleton occur in all types of glaucoma. I do not believe that this article can be indexed in this form and needs a remake of both the introduction and the discussion and the results should be reviewed in the light of a more modern interpretation of the physio-pathogenical events.

References

1. Saccà SC, Gandolfi S, Bagnis A, Manni G, et al.: The Outflow Pathway: A Tissue With Morphological and Functional Unity. *J Cell Physiol.* 2016; **231** (9): 1876-93 [PubMed Abstract](#) | [Publisher Full Text](#)
2. Saccà SC, Gandolfi S, Bagnis A, Manni G, et al.: From DNA damage to functional changes of the trabecular meshwork in aging and glaucoma. *Ageing Res Rev.* 2016; **29**: 26-41 [PubMed Abstract](#) | [Publisher Full Text](#)

Is the work clearly and accurately presented and does it cite the current literature?

No

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

No

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Glaucoma pathogenesis and Molecular biology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to state that I do not consider it to be of an acceptable scientific standard, for reasons outlined above.

Author Response 31 Mar 2018

Nils Loewen, University of Pittsburgh School of Medicine, Pittsburgh, USA

Reviewer 1:

I read the article titled: Intraocular pressure elevation precedes a phagocytosis decline in a model of pigmentary glaucoma by Dang et al. The title of the article is very stimulating but the content has several things that should be corrected.

Authors: We thank the reviewer, Professor Saccà, for having taken the time to carefully review our manuscript. We hope that we are able to address all of them with extensive manuscript changes as requested.

Reviewer 1:

1) The authors state that in the conventional way of outflow the aqueous passes into Schlemm's canal by paracytosis or giant vacuoles. This is a partial view that does not correspond to the truth. I invite the authors to read Saccà et al. The Outflow Pathway: A Tissue With Morphological and Functional Unity¹.

Authors: Thank you for bringing this review article to our attention. We are pleased to use this publication and your other article as additional references. We have modified the Introduction to include more comprehensive details about conventional outflow better aligned with Saccà et al. We would like to change the less common phrase “paracytosis” with “paracellular passage through pores” to distinguish it from the active process of pathogen intrusion through epithelia e.g. by haemophilus.¹ Overall, it appears our manuscript matches Saccà et al well who state that the main mechanisms of aqueous outflow are “a paracellular route [...] and a transcellular pathway”. We have added to the Introduction the description of a transcellular pathway through intracellular pores² with inducible pore density that depends on outflow demand³.

In the reworded Introduction we now state: “The aqueous passes from the anterior chamber into Schlemm's canal (SC) by entering first the uveal TM (UTM), the corneoscleral TM (CTM) and finally the juxtacanalicular TM (JCT)⁴. The JCT contains proteoglycans and hyaluronans and presents the aqueous humor with an increasingly tighter fluid passageway towards the SC in a process referred to as funneling⁵. The aqueous eventually passes the inner wall of SC endothelium mainly by two different mechanisms, a paracellular route in between endothelial cells and a transcellular route⁶ consisting of intracellular pores and giant vacuoles that are time and pressure dependent^{7–9}. Failure of the TM and the inner wall of SC endothelial cells to maintain homeostasis and a normal cytoskeleton and can cause ocular hypertension¹⁰. For instance, pigment dispersion¹¹ and corticosteroids can increase and contract actin stress fibers and result in an elevation of the intraocular pressure (IOP)^{11,12}. Conversely, relaxing the cytoskeleton with Rho kinase inhibitors can reverse these effects^{13,14}. Phagocytosis of debris is another function of TM cells⁸. A chronic phagocytosis demand in the form of pigment¹⁵, erythrocyte-derived ghost cells¹⁶, inflammatory cells¹⁷, photoreceptor outer segments¹⁸, lens and pseudoexfoliation material^{19,20} can all lead to secondary glaucomas even though the amount of material itself is unlikely to cause a physical outflow obstruction. Although these glaucomas make for a sizable fraction of open angle glaucomas, it was difficult to study the cellular mechanism that leads to an IOP elevation.”

2) Then the effects of the inhibitors of the Rho kinase inhibitor may have the ability to modify the metabolism of endothelial cells in the node in which the authors think it is open to question. These drugs have the ability to block TM motility by exposing more cells to aqueous humor. This is the reason why we initially witness a decline in IOP then this worsens because the exposed cells are blocked and are more exposed to apoptosis (to read Saccà et al. from DNA damage to functional changes of the trabecular meshwork in aging and glaucoma²).

Authors: We completely agree with the Reviewer. We would like to point out that - consistent with Saccà et al - we wrote that “we hypothesized that ocular hypertension is the result of a reorganization of the actin cytoskeleton and occurs before phagocytosis declines.” We did not intend to describe outflow through the TM as sink-like. We hope the expanded Introduction makes it more obvious that this is a paracellular and transcellular process.

In addition, we specifically stated before that “material itself is unlikely to cause a physical outflow obstruction” (Introduction), that granules “were not dense enough to physically obstruct any part of the conventional outflow system” (Results) and discuss that this had already been shown before: “Pigment treatment has previously been shown to cause ocular hypertension in part by reorganizing the TM actin cytoskeleton and not by physical obstruction of the outflow tract^{15,21}.” (Discussion).

We now clarify in the Discussion that “Past investigations suggested that the outflow obstruction was, in fact, physical^{15,22} but newer studies indicated that ocular hypertension is in part caused indirectly by reorganization of the TM actin cytoskeleton²³. Consistent with Zhou et al.²³, we recently reported that long, thick, and continuous TM actin bundles emerge as early as 24 hours after pigment exposure¹¹ and replicate this observation in the present study.”

3) Also the hypothesis formulated by the authors to explain the pigmentary glaucoma is partial in the Trabecolator the so-called pores have a very low meaning and above all the pigment does not obstruct anything. We do not work like sinks - I am amazed that today we can still believe that TM is a porous tissue, and above all, how it is possible to think that outflow is a passive phenomenon.

Authors: We did not state anything about an outflow obstruction or a location in our hypothesis. We wrote, “In the current study, we hypothesized that ocular hypertension is the result of a reorganization of the actin cytoskeleton and occurs before phagocytosis declines.” As we detail in our Response to 3), we never wrote that the TM functions as a sink. On the contrary, we elaborate multiple times that there is no physical obstruction as detailed in our Response to 2). Please see our response above where we hope to clarify this point in the rewritten manuscript.

4) In pigmentary glaucoma there are 2 fundamental pathogenic moments. The first concerns the back sheet of the iris, where the pigment comes from (a defect that involves the loss of the pigment itself) the second point concerns autophagy which is not able to ensure adequate cell homeostasis. So these cells first enter suffering and then die and hence glaucoma is why the cells do not work, not because it obstructs something. There are cells, not tubes.

Authors: We completely agree as detailed in 1)-3) and have expanded our manuscript to make this more obvious.

5) The cytoskeleton in all this has little to do with it: pigment granules are only a metabolic burden that in the case of pigment dispersal syndrome is absorbed by a functioning autophagy, while in pigment glaucoma, autophagy does not work. Finally, I remember that the changes in the cytoskeleton occur in all types of glaucoma. I do not believe that this article can be indexed in this form and needs a remake of both the introduction and the discussion and the results should be reviewed in the light of a more modern interpretation of the physio-pathogenical events.

Authors: We are unsure why the Reviewer states that cytoskeletal changes have nothing to do with an IOP elevation after a phagocytosis challenge. This is rather well established: Zhou et al found that 4 h after phagocytosis, the cytoskeletal structure in trabecular meshwork cells was disrupted²³. As we detail in the Introduction “pigment dispersion¹¹ and corticosteroids can increase and contract actin stress fibers and result in an elevation of

the intraocular pressure (IOP)^{11,12}. Conversely, relaxing the cytoskeleton with Rho kinase inhibitors can reverse these effects^{13,14}."

The new finding described in this manuscript is that the IOP rises when the cytoskeleton changes occur as a result from pigment exposure but before phagocytosis starts to fail. This has not been observed before because an appropriate ex vivo anterior segment model of pigment dispersion did not exist.

Our recent studies^{11,24,25} show that ex vivo perfused pig eyes experience reduced outflow in response to continuous exposure to pigment at a concentration far lower (10,000-fold) than that used in previous bolus experiments²⁶ and that severe cytoskeletal changes are associated with this. Conversely, we show that rho-kinase inhibitors normalize the cytoskeleton and the phagocytosis²⁴. TM regulates aqueous outflow by changing its cytoskeleton, stiffness, cell adhesion, migration, contraction, and phagocytosis^{10,27}, in which ROCK signaling plays a central role^{28,29}. There is a close interaction between actin cytoskeleton reorganization, cell or extracellular matrix stiffness and aqueous outflow facility^{30,31}. Only a small proportion of stress fiber formation was found in the normal TM culture while pigment exposure increased it by 2.06-fold ¹¹.

Lastly, we caution not to confuse TM cell phagocytosis of pigmented debris with autophagy. To address the Reviewer's concern we have added to the Discussion his own reference³² and state that "An increased pigmentary debris may interfere with many intracellular functions, including important autophagy functions that can cause a gradual deterioration of TM cell function^{32,33}."

References used by Reviewer 1

1. Saccà SC, Gandolfi S, Bagnis A, Manni G, Damonte G, Traverso CE, Izzotti A: The Outflow Pathway: A Tissue With Morphological and Functional Unity. *J Cell Physiol*. 2016; 231 (9): 1876-93 [PubMed Abstract](#) | [Publisher Full Text](#)
2. Saccà SC, Gandolfi S, Bagnis A, Manni G, Damonte G, Traverso CE, Izzotti A: From DNA damage to functional changes of the trabecular meshwork in aging and glaucoma. *Ageing Res Rev*. 2016; 29: 26-41 [PubMed Abstract](#) | [Publisher Full Text](#)

References used by the Authors in this Reply

1. van Schilfgaarde M, van Alphen L, Eijk P, Everts V, Dankert J. Paracytosis of *Haemophilus influenzae* through cell layers of NCI-H292 lung epithelial cells. *Infect Immun*. 1995 Dec;63(12):4729-4737. PMID: PMC173678
2. Johnson M, Erickson K. Mechanisms and routes of aqueous humor drainage. *Principles and Practice of Ophthalmology*. Saunders Philadelphia; 2000;4:2577-2595.
3. Braakman ST, Pedrigi RM, Read AT, Smith JAE, Stamer WD, Ethier CR, Overby DR. Biomechanical strain as a trigger for pore formation in Schlemm's canal endothelial cells. *Exp Eye Res*. 2014 Oct;127:224-235. PMID: PMC4175173
4. Saccà SC, Gandolfi S, Bagnis A, Manni G, Damonte G, Traverso CE, Izzotti A. The Outflow Pathway: A Tissue With Morphological and Functional Unity. *J Cell Physiol*. 2016 Sep;231(9):1876-1893. PMID: 26754581
5. Ethier CR, Coloma FM, Sit AJ, Johnson M. Two pore types in the inner-wall endothelium of Schlemm's canal. *Invest Ophthalmol Vis Sci*. 1998;39:2041-2048.
6. Alvarado JA, Betanzos A, Franse-Carman L, Chen J, González-Mariscal L. Endothelia of Schlemm's canal and trabecular meshwork: distinct molecular, functional, and anatomic

- features. *Am J Physiol Cell Physiol*. 2004 Mar;286(3):C621–34. PMID: 14613887
7. Johnstone MA, Grant WM. Pressure-dependent changes in structures of the aqueous outflow system of human and monkey eyes. *Am J Ophthalmol*. 1973 Mar;75(3):365–383. PMID: 4633234
8. Tamm ER. The trabecular meshwork outflow pathways: structural and functional aspects. *Exp Eye Res*. 2009 Apr;88(4):648–655. PMID: 19239914
9. Braakman ST, Daniel Stamer W, Overby DR. A fluorescent permeability assay for Schlemm's canal endothelial cells in response to stretch. *Invest Ophthalmol Vis Sci*. The Association for Research in Vision and Ophthalmology; 2014 Apr 30;55(13):5983–5983.
10. Llobet A, Gasull X, Gual A. Understanding trabecular meshwork physiology: a key to the control of intraocular pressure? *News Physiol Sci*. 2003 Oct;18:205–209. PMID: 14500801
11. Dang Y, Waxman S, Wang C, Loewen RT, Sun M, Loewen N. A porcine ex vivo model of pigmentary glaucoma. *Sci Rep [Internet]*. 2018 Mar 26;8. Available from: <http://dx.doi.org/10.1038/s41598-018-23861-x>
12. Pattabiraman PP, Rao PV. Mechanistic basis of Rho GTPase-induced extracellular matrix synthesis in trabecular meshwork cells. *Am J Physiol Cell Physiol*. 2010 Mar;298(3):C749–63. PMID: PMC2838580
13. Tanihara H, Inoue T, Yamamoto T, Kuwayama Y, Abe H, Suganami H, Araie M, K-115 Clinical Study Group. Intra-ocular pressure-lowering effects of a Rho kinase inhibitor, ripasudil (K-115), over 24 hours in primary open-angle glaucoma and ocular hypertension: a randomized, open-label, crossover study. *Acta Ophthalmol*. 2015 Jun;93(4):e254–60. PMID: 25487877
14. Tanihara H, Inoue T, Yamamoto T, Kuwayama Y, Abe H, Araie M, K-115 Clinical Study Group. Phase 1 clinical trials of a selective Rho kinase inhibitor, K-115. *JAMA Ophthalmol*. 2013 Oct;131(10):1288–1295. PMID: 23787820
15. Epstein DL, Freddo TF, Anderson PJ, Patterson MM, Bassett-Chu S. Experimental obstruction to aqueous outflow by pigment particles in living monkeys. *Invest Ophthalmol Vis Sci*. 1986 Mar;27(3):387–395. PMID: 3949467
16. Campbell DG, Simmons RJ, Grant WM. Ghost cells as a cause of glaucoma. *Am J Ophthalmol*. 1976 Apr;81(4):441–450. PMID: 1266922
17. Moorthy RS, Mermoud A, Baerveldt G, Minckler DS, Lee PP, Rao NA. Glaucoma associated with uveitis. *Surv Ophthalmol*. 1997 Mar;41(5):361–394. PMID: 9163835
18. Callender D, Jay JL, Barrie T. Schwartz–Matsuo syndrome: atypical presentation as acute open angle glaucoma. *Br J Ophthalmol*. BMJ Publishing Group Ltd; 1997 Jul 1;81(7):608–608.
19. Lee RK. The molecular pathophysiology of pseudoexfoliation glaucoma. *Curr Opin Ophthalmol*. 2008 Mar;19(2):95–101. PMID: 18301281
20. Ritch R, Schlötzer-Schrehardt U, Konstas AGP. Why is glaucoma associated with exfoliation syndrome? *Prog Retin Eye Res*. 2003 May;22(3):253–275. PMID: 12852486
21. Gottanka J, Johnson DH, Grehn F, Lütjen-Drecoll E. Histologic findings in pigment dispersion syndrome and pigmentary glaucoma. *J Glaucoma*. 2006 Apr;15(2):142–151. PMID: 16633228
22. Alvarado JA, Murphy CG. Outflow obstruction in pigmentary and primary open angle glaucoma. *Arch Ophthalmol*. 1992 Dec;110(12):1769–1778. PMID: 1463421
23. Zhou L, Li Y, Beatrice YJ. Alteration of cytoskeletal structure, integrin distribution, and migratory activity by phagocytic challenge in cells from an ocular tissue—The trabecular meshwork. *In Vitro CellDevBiol-Animal*. Springer-Verlag; 1999 Mar 1;35(3):144–149.
24. Dang Y, Wang C, Shah P, Waxman S, Loewen RT, Loewen NA. Ocular Hypotension, Actin

Stress Fiber Disruption and Phagocytosis Increase by RKI-1447, a Rho-Kinase Inhibitor [Internet]. 2018 [cited 2018 Feb 26]. Available from:

<http://dx.doi.org/10.20944/preprints201802.0026.v1>

25. Wang C, Dang Y, Loewen R, Waxman S, Shah P, Xia X, Loewen NA. Impact of Pigment Dispersion on Trabecular Meshwork Cells. ResearchGate preprint [Internet]. 2018 Jan 25 [cited 2018 Jan 25]; Available from:

https://www.researchgate.net/publication/322627138_Impact_of_Pigment_Dispersion_on_Trabecular_Meshwork

26. Epstein DL, Fredro TF, Anderson PJ, Patterson MM, Bassett-Chu S. Experimental obstruction to aqueous outflow by pigment particles in living monkeys. Invest Ophthalmol Vis Sci. 1986 Mar;27(3):387–395. PMID: 3949467

27. Abu-Hassan DW, Acott TS, Kelley MJ. The Trabecular Meshwork: A Basic Review of Form and Function. J Ocul Biol Dis Infor [Internet]. 2014 May;2(1). Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/25356439> PMCID: PMC4209746

28. Nakajima E, Nakajima T, Minagawa Y, Shearer TR, Azuma M. Contribution of ROCK in contraction of trabecular meshwork: proposed mechanism for regulating aqueous outflow in monkey and human eyes. J Pharm Sci. 2005 Apr;94(4):701–708. PMID: 15682386

29. Wang J, Liu X, Zhong Y. Rho/Rho-associated kinase pathway in glaucoma (Review). Int J Oncol. 2013 Nov;43(5):1357–1367. PMID: 24042317

30. Doornaert B, Leblond V, Planus E, Galiacy S, Laurent VM, Gras G, Isabey D, Lafuma C. Time course of actin cytoskeleton stiffness and matrix adhesion molecules in human bronchial epithelial cell cultures. Exp Cell Res. 2003 Jul 15;287(2):199–208. PMID: 12837276

31. Wang K, Read AT, Sulchek T, Ethier CR. Trabecular meshwork stiffness in glaucoma. Exp Eye Res. 2017 May 1;158:3–12.

32. Saccà SC, Gandolfi S, Bagnis A, Manni G, Damonte G, Traverso CE, Izzotti A. From DNA damage to functional changes of the trabecular meshwork in aging and glaucoma. Ageing Res Rev. 2016 Aug;29:26–41. PMID: 27242026

33. Liton PB, Lin Y, Gonzalez P, Epstein DL. Potential role of lysosomal dysfunction in the pathogenesis of primary open angle glaucoma. Autophagy. 2009 Jan;5(1):122–124. PMCID: PMC2745819

Competing Interests: No competing interest

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research