RESEARCH ARTICLE

REVISED Prevalence and associated risk factors of HIV in prisons in Balochistan, Pakistan: a cross-sectional study [version 2; peer review: 1 approved, 2 approved with reservations]

Muhammad Dawood Khan ¹, Ahmad Wali², Razia Fatima ³, Aashifa Yaqoob³, Shoaib Aziz ²

¹Department of Health, Provincial AIDS Control Program, Quetta, Balochistan, 87300, Pakistan

²Department of Health, Provincial TB Control Program, Quetta, Balochistan, 87300, Pakistan

³Ministry of National Health Services, Regulation and Coordination, National TB Control Program, Islamabad, Pakistan

v2 First published: 20 Nov 2018, 7:1821
<https://doi.org/10.12688/f1000research.16994.1>
 Latest published: 15 Mar 2019, 7:1821
<https://doi.org/10.12688/f1000research.16994.2>

Abstract

Background: The prevalence of HIV is 0.1% in Pakistan, with epidemics in high-risk groups. The infection is on verge of transmission from key populations to the general population through people who inject drugs and sexual transmission. Prisoners are vulnerable to infectious diseases like HIV. This study was conducted in four prisons in Balochistan, Pakistan to determine the prevalence of HIV and associated risk factors.

Methods: This cross sectional study was conducted from March to June 2018, in the prisons of Balochistan. WHO-approved rapid diagnostic kits were used for determining the prevalence of HIV and structured interviews were conducted for the assessment of risk factors.

Results: Out of 2084 screened prisoners, 33 (1.6%) were found to be positive. A subset of 104 interviews was analyzed for risk factors of HIV. Among HIV-infected prisoners 68.8% (OR 4.48; 95% CI 1.41-14.2) had extramarital sex, 43.8% (OR 2.09 95% CI 0.69-6.28) had a homosexual experience, and 50% had a history of needle sharing (OR 43; 95% CI 7.77-237). About 94% (OR 16.42; 95% CI 2.09-129.81) of prisoners had a history of drug addiction of any type while 50% (OR 13; 95% CI 2.82-60.01) of HIV-infected had a history of using injectable drugs. Around 75% of HIV-infected prisoners had spent 1-5 years in prison, and 25% had spent more than 10 years.

Conclusion: The high prevalence of HIV in prisons of Balochistan demands that preventive and treatment strategies should be designed and implemented carefully, allowing early diagnosis and treatment initiation to minimize the spread of infection among the prisons and ultimately their onward transmission into the community.

Open Peer Review

Approval Status

	1	2	3
version 2 (revision) 15 Mar 2019		 view	
		↑	
version 1 20 Nov 2018	 view	 view	 view

1. **Quaid Saeed** , National AIDS Control Programme, Islamabad, Pakistan
2. **Pamela Valera**, Rutgers University School of Public Health, New Brunswick, USA
3. **Laurent Getaz** , Geneva University Hospitals & University of Geneva, Geneva, Switzerland
Geneva University Hospitals & University of Geneva, Geneva, Switzerland

Any reports and responses or comments on the article can be found at the end of the article.

Keywords

Prison, HIV, AIDS, prevalence, risk-factors.

This article is included in the **TDR** gateway.

Corresponding author: Muhammad Dawood Khan (dkachakzai@gmail.com)

Author roles: **Khan MD:** Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Project Administration, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Wali A:** Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Supervision, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Fatima R:** Formal Analysis, Funding Acquisition, Methodology, Resources, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing; **Yaqoob A:** Formal Analysis, Investigation, Methodology, Supervision, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Aziz S:** Investigation, Methodology, Writing – Original Draft Preparation

Competing interests: No competing interests were disclosed.

Grant information: This research was conducted through the Structured Operational Research and Training Initiative (SORT IT), a global partnership led by the Special Programme for Research and Training in Tropical Diseases at the World Health Organization (WHO/TDR). The training model is based on a course developed jointly by the International Union Against Tuberculosis and Lung Disease (The Union, Paris, France) and Médecins Sans Frontières (MSF, Geneva, Switzerland). The specific SORT IT programme that resulted in this publication was implemented by the National Tuberculosis Control Programme of Pakistan, through the support of the Global Fund to Fight AIDS, Tuberculosis and Malaria (The Global Fund, Geneva, Switzerland). The publication fee was covered by the Special Programme for Research and Training in Tropical Diseases at the World Health Organization (WHO/TDR).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2019 World Health Organisation. This is an open access article distributed under the terms of the **Creative Commons Attribution IGO License**, which permits copying, adaptation and distribution in any medium or format for any purpose, provided the original work is properly cited, a link is provided to the license, and any changes made are indicated. Any such copying, adaptation and distribution must not in any way suggest that World Health Organisation endorses you or your use.

How to cite this article: Khan MD, Wali A, Fatima R *et al.* **Prevalence and associated risk factors of HIV in prisons in Balochistan, Pakistan: a cross-sectional study [version 2; peer review: 1 approved, 2 approved with reservations]** F1000Research 2019, 7:1821 <https://doi.org/10.12688/f1000research.16994.2>

First published: 20 Nov 2018, 7:1821 <https://doi.org/10.12688/f1000research.16994.1>

REVISED Amendments from Version 1

As per feedback of the reviewer I have removed grammatical errors and expanded the conclusion.

See referee reports

Introduction

Globally; about 36.7 million people were living with HIV. HIV infection causes AIDS and is epidemic throughout the world. In 2016; about 1.8 million were reported as newly infected and 1 million died from AIDS related illnesses in the same period (<http://www.unaids.org>). In Pakistan, an estimated 0.13 million peoples are infected with HIV, out of which only 17% are registered with the AIDS Control programme and 54% of total registered people currently living with HIV/AIDS are getting treatment from antiretroviral therapy (ART) centres (<http://www.nacp.gov.pk>). The prevalence rate of HIV is 0.1% in Pakistan (<http://www.unaids.org>). AIDS was limited and epidemic among key populations over the decades in Pakistan, but the latest evidence suggests that there is a shift from key populations through transmission via needle-sharing by injecting drug users (IDUs) to sexual transmission via multiple sexual partners, and further transmission to their intimate partners is increasing¹. According to Integrated Biological and Behavioural Survey (IBBS), HIV prevalence was 38.4% among people who inject drugs (PWIDs), about 7.1% among transgender (TG), 3.5% among men who have sex with men and 2.2% among female sex workers¹⁻³.

Prisons are the easily approachable venues for health interventions but unfortunately, the prisoners are among the most marginalized and restricted populations where access to health services, interventions and surveys are very limited⁴. Prisoners are one of the most vulnerable population for acquiring infectious diseases especially HIV because people from diverse background, communities and risk factors are kept together for varying time⁵. The prevalence of HIV is alarmingly increasing in PWIDs and evidence through IBBS indicated that 38.5% of PWIDs were arrested in past 12 months which makes the other prisoners also vulnerable as PWIDs have access to drugs in the prisons². A study of HIV-positive prisoners in five jails in Sindh, Pakistan indicates that 12.3% of jail prisoners have ever used drugs by injection in the jails⁶.

The revised Pakistan AIDS Strategy III, developed in 2017 by the National AIDS Control Program (NACP) to guide Pakistan's overall national response for HIV and AIDS till 2020, through focused interventions with set targets, costs, roles and responsibilities has emphasized on adopting precision targeting (including prisoners) to achieve the required level of impact on the epidemic¹. The Balochistan AIDS Control Program (BACP) conducted awareness and screening of prisoners in Central Prison Gaddani in 2017 and found that 27 (6.85%) out of 394 prisoners were positive for HIV. Most of the studies from the country have found that HIV is prevalent at about 2% in prisoners which is much higher than the prevalence in the general population^{4,5,7-13}.

Availability of limited data across the different prisons of the province and high prevalence of HIV found in central jail Gaddani warranted evidence to be produced by conducting a detailed study in other prisons of Balochistan as well. The study aimed to assess the prevalence and risk factors of HIV in prisoners of four major prisons of Balochistan.

Methods

Study design and setting

A descriptive cross-sectional study was carried out from March to June 2018 in four major prisons of Balochistan i.e. Quetta, Gaddani, Mach and Loralai. Balochistan is the most widely spread and less populated province of Pakistan with the population of 12.3 million (<http://www.pbs.gov.pk>). Area wise it is the largest province and covers 347,190 km² and comprises of 33 districts (<http://www.balochistan.gov.pk>). The province is bordered by Afghanistan in the north and north-west, Iran in the south-west, Khyber Pakhtunkhwa province in the north-east and Punjab and Sindh provinces in the east.

BACP is one of the vertical programs of the health department, providing preventive and curative services to high-risk population in the province. The program has two treatment centres (ART Centres) located in one of the largest tertiary care hospital (Bolan Medical Complex Hospital) at provincial capital Quetta and District Headquarter (DHQ) hospital of Kech, in the city of Turbat. The program is also running 30 screening centres in DHQ Hospitals. Patients with HIV are referred to ART centres for registration and treatment provision from all the screening centres and other health care facilities of the province.

Specific settings. There are 11 prisons in Balochistan of which three are central prisons and the remaining eight are district jails¹⁴. Jails are meant to keep the prisoners who are under trial or imprisoned for short durations, whereas prisons are meant to keep the prisoners who are imprisoned for long duration. In Balochistan the jail may serve as prison and prison may serve as jail due to limited capacity for prisoners and long distances from the trial court. The four prisons were selected on the basis of number of prisoners and type of jail, as these prisons cater the majority of the prisoners from Balochistan.

The provincial program started 6-month-long service delivery packages (SDPs) in four selected prisons of Balochistan from March to August 2018. In these SDPs, screening as well as treatment and preventive services were provided to jail prisoners.

Study population/participants

The study included all the prisoners enrolled at the time of screening irrespective of their status or duration of imprisonment, age, sex, marital status, education status, history of drug addiction, place of origin and employment status. Prisoners were screened for HIV after obtaining written informed consent. Blood specimen collection and screening were conducted in the health facility of jails. Interviews were conducted for assessment of risk factors with HIV. The prisoners found positive on screening were linked with treatment centres in order to commence treatment procedures.

Data collection methods

Data was collected by a team of trained data collectors that was already involved in screening and provision of SDPs in the mentioned prisons.

Screening was conducted through World Health Organization (WHO)-approved rapid diagnostic kits for HIV Screening (Alere Determine HIV-1/2 Ag/Ab Combo, with sensitivity >99%) provided by the Balochistan AIDS Control Program. The HIV status of HIV-positive prisoners was confirmed by initial confirmatory rapid diagnostic kits (Uni-Gold HIV with Specificity > 99%) and followed by confirmatory rapid diagnostic kits (SD Bioline HIV-1/2 3.0 with Sensitivity > 99%, Specificity > 98%)¹⁵.

The risk factors identification for HIV were assessed through structured a questionnaire developed, pretested and validated by the United Nations Office on Drugs and Crime (UNODC), available on OSF¹⁹. The interviews primarily were not part of initial agreement between BACP and partner organization delivering the SDPs, but keeping in view the large number of HIV-positive prisoners found in Gaddani prison, they were added to assess the risk factors for HIV. For this purpose, interviews were conducted in specially dedicated visits of the prisons. A total of 150 prisoners were approached to take part in the structured interview, out of which 135 prisoners gave their written informed consent to this aspect of the study. Out of 135 interviews conducted 104 were selected for final analysis while the remaining 31 were discarded due to lack of required information and incomplete filling. A total of 16 out of 104 (15.3%) interviewed prisoners were positive for HIV.

The screening and interview processes were randomly monitored by principal investigator to ensure the quality of data collection.

Data variables collected

The outcome variables of the study included the total number of prisoners screened for HIV and the number and proportion of prisoners found positive for HIV. The presumptive exposure variables assessed were age, sex, education and occupation status of prisoners, type of imprisonment and time spent in imprisonment, ever been arrested for drug related offense (Y/N), marital status and extramarital sex status including homosexual sex status (Y/N), sharing needles, razors and blades (Y/N), history of drug addiction (Y/N) and injectable drug addiction (Y/N) and history of blood transfusion, surgery, dental procedure and tattooing (Y/N).

Statistical analysis

Data was double-entered into Epi-Data software (version 3.1) and analysed in Epi-Analysis (version 2.2.2.183 for analysis, Epi-Data Association, Odense, Denmark). Descriptive statistics were used to describe the data. Odds ratio (OR) with 95% confidence interval (CI) was calculated for possible presumptive variables.

Ethical approval

The SDPs had already commenced in the selected prisons before the start of this study by BACP, with the permission of the Health Department of Balochistan and Home Department

Government of Balochistan. Screening, health education sessions and collection of data was routinely done for the SDPs so the program data was utilized. The principal investigator was also the monitor of SDPs from BACP. Study and ethical approval was obtained from BACP (No. BACP: 01/2018-432) to collect and analyse the program data for the research purpose.

Results

Initial screening and demographic information

A total of 2084 prisoners were screened, of whom 33 were found to be positive for HIV (Figure 1). All the HIV-positive prisoners were approached for assessment of risk factors but only 16 gave informed written consent for the interview.

The median age of the prisoners was 25 years (range 13–80 years, 94.23% in 15–45 years age group) years and 53 (50.9%) of them were married. The majority of them were uneducated (36.53%) or had been in education for less than 5 years (16.34%). Among them, 15% were unemployed and 8% were working on daily wages before the imprisonment. Around 93% of the prisoners were from Balochistan, 4% from other provinces and 3% from other countries (Table 1).

Around 75% HIV infected prisoners were convicted, 75% had spent less than five years, 25% had spent more than ten years in prisons. 22% of total interviewed prisoners had been ever arrested for drug related offense (Table 2).

Information surrounding HIV risk factors

Extramarital sex was reported by 11 (68.8%) out of 16 HIV-positive prisoners (OR 4.48; 95% CI 1.41–14.2). Homosexuality was reported by 7 (43.8%) HIV-positive prisoners (OR 2.09; 95% CI 0.69–6.28). About 50% of HIV positive prisoners had history of needle sharing (OR 43; 95% CI 7.77–237.87). Almost 94% of HIV positive prisoners had history of drug addiction of any type, of whom 50% had history of injectable drugs (OR 13; 95% CI 2.82–60.01). History of sharing razorblades and blood transfusion was 9.6% in HIV-positive prisoners; 12.5% had gone through any surgical procedure, 8.6% through any dental procedure and 6.7% had history of tattooing in the past 5 years (Table 3).

Discussion

This study was conducted in four major jails in Balochistan, Pakistan, and revealed that the HIV prevalence was 1.6%, which is much higher than the prevalence rate in general population in Pakistan³. A previous study conducted in Quetta jail indicates that the HIV prevalence was 0.5%¹⁶. A limitation of that study was generalizability to the whole province as the sample size was very small (only 200 prisoners were screened) and the prisoners of Quetta jail, although large in number, were mostly under trial, so the long-term sentenced prisoners were excluded from being assessed. The study findings are consistent with studies conducted in other provinces where the HIV prevalence ranged from 1 to 2.9%^{4,5,7,9,11,13}.

Based on our study findings, the majority of the HIV-positive prisoners belonged to the 15–45 years age group. The same age range was also found to be significant in other studies conducted

Figure 1. Flow diagram of prisoners screened and interviewed for HIV in prisons in Balochistan. *Test 1 = Alere Determine HIV-1/2 Ag/Ab Combo for detection of Antigen/Antibody, **Test 2 = Uni-Gold HIV, **Test 3 = SD Bioline HIV-1/2 3.0.

in Pakistan^{5,11,13}. Education status was also found as an important factor in HIV risk assessment, as a large majority of HIV positive prisoners (62.5%) were not educated or had been educated for less than 5 years, as found by another study conducted in Lahore¹¹.

The prisoners who were single had a higher prevalence of HIV, as 62.5% of HIV-positive prisoners were unmarried. The findings are consistent with a study conducted in different jails of six cities in Sindh province¹³. Imprisonment status and duration of imprisonment was significant as 81% of HIV-positive prisoners were convicted and 25% prisoners had spent more than 10 years in prisons.

A substantial proportion of HIV-infected prisoners were involved in behaviour considered to be presumptive risk factors associated with HIV; 68.8% were involved in extramarital sex and 43.8% had engaged in homosexual practices. A study of

prisoners in Karachi found involvement of prisoners in homosexuality and sexual encounters with females to be 21.3% and 45.9%, respectively⁴. Moreover, drug addiction, especially injectable drugs involving the sharing of needles, was also found to be present in significant majority of the HIV infected prisoners. Risk of acquiring of HIV increases in prisons as studies shows that 38.5% of PWID were arrested in past 12 months and around 12.3% of arrested PWID had access to injectable drugs in prisons^{2,17}.

The young age of the prisoners, being less educated, having low socioeconomic status, addiction to injectable drugs, having multiple sex partners, sharing needles and involvement in drug related offense were notable findings of the study and consistent with studies conducted elsewhere in Pakistan^{4,5,7-13}.

Prison settings highly favour the spread of infectious disease as the prisoners from different sociodemographic backgrounds

Table 1. Sociodemographic characteristics of prisoners interviewed for assessment of risk factors for HIV.

Variable	HIV positive, n (%)	HIV negative, n (%)	Total, n (%)
Total	16 (15.4)	88 (84.6)	104
Age, years			
<15	0	1 (0.96)	1 (0.96)
15–45	16 (100)	82 (78.8)	98 (94.2)
45–65	00	02 (1.92)	02 (1.92)
65 and above	00	03 (2.88)	03 (2.88)
Sex			
Male	16 (100)	88 (100)	104 (100)
Place of origin			
Within province	16 (100)	81 (77.8)	97 (93.2)
Out of province	00	04 (3.84)	04 (3.84)
Out of country	00	03 (2.88)	03 (2.88)
Education			
Not educated	09 (56.3)	29 (33.0)	38 (36.5)
<5 years	01 (6.25)	16 (18.1)	17 (16.3)
6–10 years	02 (12.5)	20 (22.7)	22 (21.2)
11–14 years	04 (25.0)	20 (22.7)	24 (23.1)
>14 years	00	02 (2.27)	02 (1.92)
Religious	00	01 (1.13)	01 (0.96)
Occupation			
Unemployed	02 (12.5)	14 (15.9)	16 (15.3)
Employed*	12 (75.0)	68 (77.3)	80 (76.9)
Daily wages	02 (12.5)	06 (6.81)	08 (7.69)
Marital status			
Unmarried	10 (62.5)	41 (46.6)	51 (49.0)
Married	06 (37.5)	47 (53.4)	53 (51.0)

*Employed includes prisoners who have worked both in public, private sector and/or self-employed.

like low literacy levels and having pre-imprisonment risk factors such as drug use, extramarital sex encounters etc. are forced to live together in confined space without having recreational facilities⁴. The prisoners are vulnerable to acquiring new infections in prisons due to overcrowding, poor medical facilities (especially diagnostic investigations) and the unavailability of harm reduction approaches such as the use of condoms or needle exchange program¹⁸. All these factors may not only lead to greater chances of spread of infections like HIV among the jails but can also be a major concern among the general population due to frequent flow of prisoners into the community⁴.

Aside from the many risk factors responsible for spread of infections in prison settings, prisons are still favourable venues for the screening and treatment of HIV, as the prisoners are confined to limited vicinity and easily approachable for diagnosis, treatment and health education. If the prisoners are approached within

the prisons for diagnosis and treatment, this can help halt the spread of HIV to the general population and vice versa⁴.

The level of health services provided in prisons plays an important role in dealing with the associations of HIV-infected persons with the community. In a resource-constrained country like Pakistan, the chain of transmission can be broken by introducing urgent prevention efforts, including health education, initial screening at the time of entry of prisoners in prisons, routine periodic screening of all the prisoners and treatment of infected prisoners in each prison so that its spread within the prisons and to the community after release of the prisoner could be minimized⁴. The presence of HIV-infected prisoners warrants that information and education programs are made accessible in the prisons to stop the spread of infectious diseases in the prisons, with particular emphasis on transmission through injectable drug use and high-risk sexual

Table 2. Imprisonment status of prisoners in prisons setting of Balochistan evaluated during March-June 2018 (n=104).

Variable	HIV positive, n (%)	HIV negative, n (%)	Total, n (%)
Total	16 (15.4)	88 (84.6)	104
Type of imprisonment			
Under trial	03 (18.8)	26 (29.5)	29 (27.8)
Convicted	13 (81.3)	62 (70.5)	75 (72.1)
Time imprisoned, years			
< 1	08 (50)	37 (42.0)	45 (43.2)
1–5	04 (25)	34 (38.6)	38 (36.5)
6 – 10	00	12 (13.6)	12 (11.5)
> 10	04 (25)	05 (5.7)	09 (8.65)
Ever arrested for drug-related offense			
No	10 (62.5)	71 (80.7)	81 (77.8)
Yes	06 (37.5)	17 (19.3)	23 (22.1)

Table 3. Risk factors associated with prevalence of HIV in prisons setting of Balochistan. 2018.

Variable	HIV positive, n (%)	HIV negative, n (%)	OR	95% CI
Total	16 (15.4)	88 (84.6)		
Extramarital sex status	11 (68.8)	27 (30.7)	4.48	1.41–14.2
Homosexual sex status	07 (43.8)	22 (25)	2.09	0.69–6.28
Sharing needles	08 (50)	02 (2.3)	43	7.77–237.87
Sharing razors and blades	03 (18.8)	07 (8)	2.67	0.61–11.66
History of drug addiction	15 (93.7)	42 (47.7)	16.42	2.09–129.81
Type of drug addiction				
Injectable drugs	08 (50)	03 (7.1)	13	2.82–60.01
Other drugs	08 (50)	39 (44.3)		
History of blood transfusion	00 (00)	10 (11.3)	00	
History of surgery	02 (12.5)	11 (12.5)	01	0.2–5.01
History of dental procedure	00 (00)	09 (10)	00	
History of tattooing	02 (12.5)	05 (5.68)	2.37	0.42–13.44

OR, odds ratio; CI, confidence interval.

activities. Training the health care providers to identify and treat those suspected of being, or known to be infected with HIV and the provision of testing services may help in early diagnosis of infections and their treatment to reduce the spread of infection.

Being a comprehensive study covering four major jails of Balochistan province the study is strengthened by universal sampling for screening in which all the prisoners present during the

study period were screened for HIV. The risk factors may vary in those prisoners who were not interviewed.

Conclusion

The high prevalence of HIV in prisons in Balochistan, compared with the non-prison population, indicates that preventive strategies should be designed and implemented carefully to minimize the spread of infection among the prisons and ultimately their linkages to the community. Early diagnosis of

HIV infection through screening, initiation of treatment, addressing the stigma universally attached with the disease, peer based, risk reduction education program and linking HIV-positive inmates to the community care after release are effective ways of preventing the spread of infection.

Data availability

Underlying data

Raw data for the present study, including demographic information and answers to the questionnaire, is available on OSF, DOI: <https://doi.org/10.17605/OSF.IO/8Z4EJ>¹⁹.

Extended data

The questionnaire used in this study is available on OSF, DOI: <https://doi.org/10.17605/OSF.IO/8Z4EJ>¹⁹.

Grant information

This research was conducted through the Structured Operational Research and Training Initiative (SORT IT), a global partnership led by the Special Programme for Research and Training

in Tropical Diseases at the World Health Organization (WHO/TDR). The training model is based on a course developed jointly by the International Union Against Tuberculosis and Lung Disease (The Union, Paris, France) and Médecins Sans Frontières (MSF, Geneva, Switzerland). The specific SORT IT programme that resulted in this publication was implemented by the National Tuberculosis Control Programme of Pakistan, through the support of the Global Fund to Fight AIDS, Tuberculosis and Malaria (The Global Fund, Geneva, Switzerland). The publication fee was covered by the Special Programme for Research and Training in Tropical Diseases at the World Health Organization (WHO/TDR).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgements

The Provincial AIDS Control Program, Balochistan, through its program Manager Dr. Noor Muhammad Qazi, supported the research by allowing the principal investigator to utilize the program data and resources.

References

- National AIDS Control Program Pakistan: **Pakistan AIDS Strategy III (2015-2021)**. 2017.
- Pakistan NACP 2nd Generation HIV Surveillance in Pakistan Round 5: **Integrated Biological & Behavioral Surveillance in Pakistan**. 2016.
[Reference Source](#)
- Pakistan NACP: **AIDS Epidemic Modelling Exercise for Pakistan 2017**. 2017.
[Reference Source](#)
- Kazi AM, Shah SA, Jenkins CA, *et al.*: **Risk factors and prevalence of tuberculosis, human immunodeficiency virus, syphilis, hepatitis B virus, and hepatitis C virus among prisoners in Pakistan**. *Int J Infect Dis*. 2010; **14** Suppl 3: e60–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ikram N, Firdus R, Tariq Baig PA: **Screening of Jail Inmates for Hepatitis B,C and HIV Infections**. *J Rawalpindi Med Coll*. 2011; **15**(2): 79–81.
- Bergstrom A, Achakzai B, Furqan S, *et al.*: **Drug-related HIV epidemic in Pakistan: a review of current situation and response and the way forward beyond 2015**. *Harm Reduct J*. 2015; **12**(1): 43.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Nafees M, Qasim A, Jaffer G, *et al.*: **HIV infection, HIV/HCV and HIV/HBV co-infections among jail inmates of Lahore**. *Pak J Med Sci*. 2011; **27**(4): 837–41.
[Reference Source](#)
- Ali M, Nadeem M, Numan M, *et al.*: **Thirty years of HIV in Pakistan: a systematic review of prevalence and current scenario**. *Future Virol*. 2017; **12**(10): 609–23.
[Publisher Full Text](#)
- Waheed U, Satti HS, Arshad M, *et al.*: **Epidemiology of HIV/AIDS and syphilis among high risk groups in Pakistan**. *Pak J Zool*. 2017; **49**(5): 1829–34.
[Publisher Full Text](#)
- Shah SSA, Ali M, Ahmad M, *et al.*: **Screening of jail inmates for HIV and tuberculosis**. *Pak J Med Health Sci*. 2013; **7**(1): 172–5.
[Reference Source](#)
- Manzoor S, Tahir Z, Anjum A: **Prevalence of HIV and Tuberculosis Among Jail Inmates in Lahore - Pakistan**. *Biomedica*. 2009; **25**: 36–41.
[Reference Source](#)
- Iqbal KJ, Arshad J, Muhammad A, *et al.*: **Health status and imprisonment profile of jail inmates of district jail Rahim Yar Khan, Pakistan**. *Explor Anim Med Res*. 2012; **2**(2): 146–50.
[Reference Source](#)
- Safdar S, Mehmood A, Abbas SQ: **Short Report Prevalence of HIV / AIDS among jail inmates in Sindh**. *J Pak Med Assoc*. 2009; **59**(2): 111–112.
[Reference Source](#)
- Qureshi H, Iqbal R: **Review of Health System in Prisons of Punjab, Pakistan**. 2012.
[Reference Source](#)
- Pakistan Country Strategy for HIV Testing and Counselling based on Situation and Response Analysis**. 2012.
[Reference Source](#)
- Kakar N: **Prevalence of human immunodeficiency virus in the inmates of district Jail Quetta, Balochistan**. *Pure Appl Biol*. 2017; **6**(1): 164–170.
[Publisher Full Text](#)
- Akhtar S, Luby SP, Rahbar MH, *et al.*: **HIV/AIDS knowledge, attitudes and beliefs based prediction models for practices in prison inmates, Sindh, Pakistan**. *Southeast Asian J Trop Med Public Health*. 2001; **32**(2): 351–61.
[PubMed Abstract](#)
- Bick JA: **Infection Control in Jails and Prisons**. *Clin Infect Dis*. 2007; **45**(8): 1047–55.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Khan MD: **Prevalence and Associated Risk Factors of HIV in Prisons in Balochistan, Pakistan: A Cross-Sectional Study**. *OSF Web*. 2018.

Open Peer Review

Current Peer Review Status:

Version 2

Reviewer Report 23 April 2019

<https://doi.org/10.5256/f1000research.20272.r45799>

© 2019 Valera P. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Pamela Valera

Rutgers University School of Public Health, New Brunswick, NJ, USA

Revise the abstract - Introduction - the prevalence of HIV is very small, but how this is written it sounds more fear mongering given the results that only 1.6% out of 2,084 were HIV+. I would focus on the facts only. The conclusion states - high prevalence compared to whom, please revise?

Replace "homosexual experience" with "same-sex experience" in the abstract and entire manuscript.

Grammatical errors have not been addressed throughout the paper. For example: "*Prisons are the easily approachable venues for health interventions but unfortunately, the prisoners are among the most marginalized and restricted populations where access to health services, interventions and surveys are very limited*". This should say - "Prisons are an important venue for health interventions as inmates are among the most marginalized and restricted populations where access to health services, interventions, and survey research are very limited". I am unsure that prisons are easily approachable venues for health interventions.

Describe IRB procedures.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Prison Health, HIV/AIDS, and Prison Education.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Version 1

Reviewer Report 18 February 2019

<https://doi.org/10.5256/f1000research.18580.r42871>

© 2019 Getaz L. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Laurent Getaz

¹ Division of Prison Health, Geneva University Hospitals & University of Geneva, Geneva, Switzerland

² Division of Tropical and Humanitarian Medicine, Geneva University Hospitals & University of Geneva, Geneva, Switzerland

This article describes the prevalence of and risk factors associated with HIV in four prisons in Pakistan. These data are very important for prioritizing programs that are essential for HIV control in this vulnerable population.

I propose a major revision of this manuscript.

Methods section:

- How was the HIV screening test proposed in the study population? Was the test optional or mandatory? If the test was not mandatory, what was the participation rate?
- How did you calculate the sample size, in particular for the number of participants who responded to the questionnaire?
- Please clarify the HIV diagnostic algorithm. For participants with a positive "Alere Determine HIV test" result, the HIV status "was confirmed by initial confirmatory rapid diagnostic kits (Uni-Gold) and followed by confirmatory rapid diagnostic kits (SD Bioline)". In the case of a positive "Alere determine test" result and negative "Uni-Gold test" result, did you process the SD Bioline test? How did you manage discordant results?

Discussion section:

- "Education status was also found as an important factor in HIV risk assessment, as a large majority of HIV positive prisoners (62.5%) were not educated or had been educated for less than 5 years". This statement is not true. In the results section, you should provide the results of a bivariate analysis to estimate the association between education level and HIV prevalence. The level of education is not associated with HIV (62.5% versus 51.1%, $p=0.4$, $OR=1.6$ (0.5-5.1)).
- "The prisoners who were single had a higher prevalence of HIV, as 62.5% of HIV-positive prisoners were unmarried". In the results section, you must also provide the results of a bivariate analysis to estimate the association between this variable and HIV. Marital status is not associated with infection (62.5% versus 46.6%, $p=0.24$).
- I believe an epidemiologist should be involved as a coauthor to validate the interpretation of the results. You should discuss the result in which a statistically significant association is identified in the study population and put it in perspective with the data from the literature. And when a generally recognized risk factor is not associated with infection in the study population, you should still discuss it. For example, in your study, "extramarital sex status" ($OR=4.48$, $IC95\%$ 1-4-14.2) is associated with HIV but not "homosexual sex status" ($OR=2.1$, $IC95\%$ 0.7-6.3). Also, you cannot state "A substantial proportion of HIV-

infected prisoners were involved in behaviour considered to be presumptive risk factors associated with HIV; 68.8% were involved in extramarital sex and 43.8% had engaged in homosexual practices".

- I suggest you describe the limitations of the study. What potential biases do you suspect, and what are their expected impacts on the results?

The manuscript should be proofread by a native English speaker for spelling, grammar and punctuation, as it contains many errors throughout. Many phrases contain errors, which makes reading the manuscript difficult.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Partly

Are sufficient details of methods and analysis provided to allow replication by others?

Partly

If applicable, is the statistical analysis and its interpretation appropriate?

Partly

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Tropical medicine / Internal medicine / Public health / Health in detention

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Reviewer Report 11 February 2019

<https://doi.org/10.5256/f1000research.18580.r42869>

© 2019 Valera P. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Pamela Valera

Rutgers University School of Public Health, New Brunswick, NJ, USA

There is an urgent public health need to understand HIV/AIDS disease transmission globally especially in areas where war and conflict continues and resources to address communicable diseases are limited.

This paper is the first to my knowledge to shed light on a growing HIV epidemic especially in correctional facilities in different prisons and jails in Pakistan. The study design provides opportunities for future research to understand HIV/AIDS transmission in correctional facilities. However, I would have much appreciated more literature on the characteristics of offenders prior to incarceration.

The findings while not generalizable were instrumental in helping readers understand high-risk behaviors among people who inject drugs and people who engage in same-sex behaviors. Perhaps, the next step of this study might be to conduct a qualitative study exploring the lived experience of inmates living with HIV/AIDS in remote correctional settings in Pakistan

The conclusion points researchers in the right direction. We have a limited understanding of HIV prevalence and HIV risk-taking behaviors of offenders in Pakistan and other areas. We need to do more in understanding HIV prevalence in correctional settings, but also discuss opportunities for prevention intervention programs. Perhaps, a section on implications or future research in the conclusion section should be included.

Here are potential articles to consider for this section:

- Scott, D. P., Harzke, A. J., Mizwa, M. B., Pugh, M., & Ross, M. - Evaluation of an HIV peer education program in Texas prisons¹.
- Solomon, L., Montague, B. T., Beckwith, C. G., Baillargeon, J., Costa, M., Dumont, D., Rich, J. D. - Survey finds that many prisons and jails have room to improve HIV testing and coordination of postrelease treatment².
- Valera, P., Chang, Y., & Lian, Z. - HIV risk inside US prisons: A systematic review of risk reduction interventions conducted in US prisons³.

Lastly, the manuscript would greatly benefit from proof-reading and addressing grammatical errors throughout the article.

References

1. Scott D, Harzke A, Mizwa M, Pugh M, et al.: Evaluation of an HIV Peer Education Program in Texas Prisons. *Journal of Correctional Health Care*. 2004; **10** (2): 151-173 [Publisher Full Text](#)
2. Solomon L, Montague BT, Beckwith CG, Baillargeon J, et al.: Survey finds that many prisons and jails have room to improve HIV testing and coordination of postrelease treatment. *Health Aff (Millwood)*. 2014; **33** (3): 434-42 [PubMed Abstract](#) | [Publisher Full Text](#)
3. Valera P, Chang Y, Lian Z: HIV risk inside U.S. prisons: a systematic review of risk reduction interventions conducted in U.S. prisons. *AIDS Care*. **29** (8): 943-952 [PubMed Abstract](#) | [Publisher Full Text](#)

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Prison Health, HIV/AIDS, and Prison Education

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 05 Mar 2019

Muhammad Khan, Provincial AIDS Control Program, Quetta, Pakistan

Dear Pamela Valera

Thank you very much for your valuable feedback. I have tried to remove the grammatical mistakes from the manuscript and amend the conclusion section based on the references provided by you.

Competing Interests: No competing interest disclosed.

Reviewer Report 27 December 2018

<https://doi.org/10.5256/f1000research.18580.r40898>

© 2018 Saeed Q. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Quaid Saeed

National AIDS Control Programme, Islamabad, Pakistan

The work quite clearly cites the current and latest literature regarding the HIV disease transmission dynamics in Pakistan and also cites other such studies conducting in different

prisons of Pakistan. The study design is appropriate and the study has been conducted in the most difficult and remote prison in Pakistan giving good insight on conditions prevalent in such remote prison settings.

The study design was appropriate and a good number of prison inmates (2084) were selected. As with other similar studies the prevalence found was more than the one in the general population which warrants that urgent preventive measures should be taken to control spread of this infection in this closed population. In Pakistan, outbreaks of HIV have been reported in different jails of the country in the past. Sensational reports came in the media regarding outbreak of HIV in different prisons but no scientific studies were carried out to determine the reasons of this spread. This study is a good effort to highlight the prevalent conditions and risks for spread of HIV in prison settings.

The study outlines the risk factors associated with HIV transmission in this population as part of the general population but does not shed ample light on prevailing high-risk behaviours inside the prisons that may have led to transmission of this disease in the prisons. Thus, most of the readers would be interested whether high risk behaviours are prevalent in these prisons and whether drugs and injectable equipment is available in the prisons. Having said that it is not easy to divulge upon these areas since most prison authorities deny prevalence of such practices in their jurisdiction and do not allow such questions to be asked and explored by investigators. It is therefore conceivable that the of this study questionnaire must have been cleared and vetted by the jail authorities and therefore such information could not be explored during this study.

As far as the methods of analysis was concerned the details provided were quite useful for other investigators to replicate such a study. Details were provided on statistical tools applied and used.

The conclusion drawn are in line with the findings of the study. We lack data on prevalent behaviours of prison inmates in different prisons in Pakistan and such similar studies are needed to advocate and start prevention interventions in these settings in the country.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research