REVIEW

Voltage-gated calcium channel $\alpha_2\delta$ subunits: an assessment of proposed novel roles [version 1; peer review: 2 approved]

Annette C. Dolphin

Department of Neuroscience, Physiology and Pharmacology, University College London, Gower Street, London, WC1E 6BT, UK

V1 First published: 21 Nov 2018, 7(F1000 Faculty Rev):1830
<https://doi.org/10.12688/f1000research.16104.1>

Latest published: 21 Nov 2018, 7(F1000 Faculty Rev):1830
<https://doi.org/10.12688/f1000research.16104.1>

Abstract

Voltage-gated calcium (Ca_v) channels are associated with β and $\alpha_2\delta$ auxiliary subunits. This review will concentrate on the function of the $\alpha_2\delta$ protein family, which has four members. The canonical role for $\alpha_2\delta$ subunits is to convey a variety of properties on the Ca_v1 and Ca_v2 channels, increasing the density of these channels in the plasma membrane and also enhancing their function. More recently, a diverse spectrum of non-canonical interactions for $\alpha_2\delta$ proteins has been proposed, some of which involve competition with calcium channels for $\alpha_2\delta$ or increase $\alpha_2\delta$ trafficking and others which mediate roles completely unrelated to their calcium channel function. The novel roles for $\alpha_2\delta$ proteins which will be discussed here include association with low-density lipoprotein receptor-related protein 1 (LRP1), thrombospondins, α -neurexins, prion proteins, large conductance (big) potassium (BK) channels, and *N*-methyl-d-aspartate (NMDA) receptors.

Keywords

calcium channel, alpha2delta, interaction

Open Peer Review

Approval Status

	1	2
version 1 21 Nov 2018		

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Emilio Carbone**, Lab of Cellular Physiology and Molecular Neuroscience, Torino, Italy
2. **Jutta Engel**, Saarland University, School of Medicine, Homburg, Germany

Any comments on the article can be found at the end of the article.

Corresponding author: Annette C. Dolphin (a.dolphin@ucl.ac.uk)

Author roles: Dolphin AC: Conceptualization, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: My own work referenced in this study has been funded by Wellcome Trust and MRC.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2018 Dolphin AC. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Dolphin AC. **Voltage-gated calcium channel $\alpha_2\delta$ subunits: an assessment of proposed novel roles [version 1; peer review: 2 approved]** F1000Research 2018, 7(F1000 Faculty Rev):1830 <https://doi.org/10.12688/f1000research.16104.1>

First published: 21 Nov 2018, 7(F1000 Faculty Rev):1830 <https://doi.org/10.12688/f1000research.16104.1>

Introduction

Voltage-gated calcium (Ca_v) channels are ubiquitously present in excitable cells and are essential for their function. They can be divided into three classes (Ca_v1 – 3). All except the Ca_v3 (T type) channels are associated with several auxiliary subunits—termed $\alpha_2\delta$ and β —together with an additional γ subunit in skeletal muscle^{1,2} (Figure 1). One of these subunits, $\alpha_2\delta$, conveys a variety of properties on the channels but recently has also been reported to have distinct effects on both other ion channels and other biological processes. These novel aspects of $\alpha_2\delta$ function are the subject of this review. This topic is important, as $\alpha_2\delta$ -1 is the therapeutic target of the $\alpha_2\delta$ ligand (gabapentinoid) class of drugs^{3,4}, which are widely prescribed for several indications, including many types of neuropathic pain.

The $\alpha_2\delta$ subunits have a well-established canonical role to influence the trafficking and function of the Ca_v1 and Ca_v2 channels, increasing the density of these channels on the plasma membrane⁵. They also direct trafficking of the channels to specific subcellular sites, including neuronal processes^{5,6}. In addition, the $\alpha_2\delta$ subunits increase Ca_v function by influencing the biophysical properties of the calcium currents^{7–10}, over and above their effect on trafficking⁶.

More recently, $\alpha_2\delta$ -1 proteins have been proposed to have non-classic functions of two types: (a) additional functions related to calcium channels, either to link the calcium channel complexes to other proteins or to influence calcium channel function, and (b) roles not associated with calcium channel function.

For (a), I will discuss several topics, including the association of $\alpha_2\delta$ proteins with α -neurexins to influence synaptic transmission^{11,12}. The $\alpha_2\delta$ -1 protein has also been found to interact potentially with large conductance (big) potassium (BK)

channels¹³, a process which it has been suggested influences calcium channel function by sequestering the $\alpha_2\delta$ subunits. For (b), I will discuss novel roles associated with the association of $\alpha_2\delta$ with thrombospondins (TSPs), an interaction which has been found to influence synaptogenesis in some systems¹⁴. I will also discuss the proposed association of $\alpha_2\delta$ with *N*-methyl-D-aspartate (NMDA) receptors¹⁵ (Figure 2). It is possible that the gabapentinoid drugs also act by influencing these various novel targets.

Topology, domain structure, and biochemical properties of $\alpha_2\delta$ proteins

The $\alpha_2\delta$ subunit was first identified as two proteins— α_2 and δ —co-purifying as integral constituents of the calcium channel complex present in skeletal muscle T-tubules^{16–18}. It was found that $\alpha_2\delta$ is encoded by a single gene and is subsequently processed into α_2 and δ ^{17,18}. Four mammalian $\alpha_2\delta$ genes have been cloned (*CACNA2D1–4*)^{16,19–21}.

All the $\alpha_2\delta$ proteins have highly related topology^{22,23}, with an N-terminal signal sequence, indicating that the N-terminus is extracellular (Figure 3). The hydrophobic C-terminus of $\alpha_2\delta$, and its behavior as an integral membrane protein, led to its being categorized as a transmembrane protein^{17,18}. However, it was subsequently identified to have a strongly predicted glycosylphosphatidylinositol (GPI)-anchor ω -site²⁴. Indeed, multiple pieces of experimental evidence indicate that $\alpha_2\delta$ -1, $\alpha_2\delta$ -2, and $\alpha_2\delta$ -3 (and probably $\alpha_2\delta$ -4 by prediction) are GPI-anchored^{24–26}.

The $\alpha_2\delta$ subunit genes encode a single precursor protein, which is post-translationally proteolytically processed into two polypeptides. The folding of $\alpha_2\delta$ in the endoplasmic reticulum involves the formation of multiple disulfide bonds both within and between the α_2 and δ moieties, so that, despite their cleavage, the α_2 and δ polypeptides remain disulfide-bonded

Figure 1. The subunit structure of voltage-gated calcium channels of the Ca_v1 and Ca_v2 family. The Ca_v α_1 subunit with 24 transmembrane segments and the intracellular β and the extracellular $\alpha_2\delta$ subunits are shown. The γ subunit (γ_1) is associated with $\text{Ca}_v1.1$ only and is not depicted.

Figure 2. Summary of $\alpha_2\delta$ interactions with other proteins. The various ion channels and other proteins with which $\alpha_2\delta$ subunits have been found to interact are shown. BK, large conductance (big) potassium; LRP1, low-density lipoprotein receptor-related protein 1; NMDA, *N*-methyl-D-aspartate; TSP, thrombospondin.

Figure 3. The post-translational processing of $\alpha_2\delta$ subunits. The hydrophobic N-terminal signal sequence is a signal for the polypeptide to co-translationally pass through the membrane of the endoplasmic reticulum (ER). This signal sequence is cleaved off. The glycosylphosphatidylinositol (GPI) anchor is added in the ER by an endopeptidase transamidase, which cleaves the C-terminal signal peptide at the ω -site and adds a pre-formed GPI lipid anchor. Multiple disulfide bonds are formed as the protein folds in the ER, and N-glycosylation occurs at multiple sites. Mature glycosylation is then completed in the Golgi complex, and it is likely that proteolytic cleavage of $\alpha_2\delta$ also occurs here [27](#). The GPI anchor can also be modified during trafficking.

together^{17,18}. The role for the proteolytic cleavage between α_2 and δ has been shown to be key to the mature function of these proteins^{6,28}, and $\text{Ca}_v2.2$ associates to a greater extent with the mature cleaved form of $\alpha_2\delta$ -1 than with the uncleaved form²⁸.

A von Willebrand factor A (VWA) domain is present in the α_2 moiety of all $\alpha_2\delta$ proteins^{29,30}; these widespread domains are generally involved in extracellular protein–protein interactions. A key motif in VWA domains is the metal ion-dependent adhesion site (MIDAS), which involves coordination of the divalent cation by a ring of up to five polar or charged residues²⁹. $\alpha_2\delta$ -1 and $\alpha_2\delta$ -2 have a “perfect” MIDAS site³⁰, whereas $\alpha_2\delta$ -3 and $\alpha_2\delta$ -4 have a missing polar residue²⁹. The $\alpha_2\delta$ subunits also contain multiple Cache domains^{22,31,32}, which have homology to domains found in bacterial chemotaxis receptors.

A recent cryo-electron microscopic structural study of the skeletal muscle calcium channel complex provided detailed information on the structure of $\alpha_2\delta$ -1, confirmed the topology of $\alpha_2\delta$ subunits, and identified the interaction sites between $\alpha_2\delta$ and $\text{Ca}_v1.1$ ³², reinforcing the importance of the VWA domain interaction, previously identified³⁰, and also providing evidence for C-terminal GPI anchoring rather than a transmembrane segment associated with $\alpha_2\delta$ -1. The study also identified four sites of disulfide bonding between α_2 and δ , one of which was found previously by mutagenesis³³.

The complex biochemistry of $\alpha_2\delta$ proteins represents a challenge for their study, and it is important to be aware of their distinct biochemical characteristics in terms of their multiple glycosylation sites and disulfide bonds, proteolytic cleavage into α_2 and δ , and GPI anchoring (Figure 3). All of these properties might be inadvertently disrupted by the placement of epitope tags or production of mutants, to the detriment of their function^{6,24,26,33}. Furthermore, as elegantly shown very recently with respect to $\alpha_2\delta$ proteins¹², co-immunoprecipitation experiments require multiple controls to be sure of the specificity of any interaction, and additional experiments are needed to determine whether any association is direct. This is particularly true when potential binding partners are co-expressed in transfected cells, where elevated concentrations may result in aberrant interactions being detected.

Properties of $\alpha_2\delta$ as a voltage-gated calcium channel subunit

For the Ca_v1 and Ca_v2 channels, $\alpha_2\delta$ universally augments expressed calcium current density^{7–9,30}. The $\alpha_2\delta$ subunits also have effects on both kinetic and voltage-dependent properties of the channels, including activation and inactivation. In general, there is a negative shift in the voltage dependence of steady-state inactivation^{30,34}. In some cases, there is also a hyperpolarization of the voltage dependence of activation, particularly for $\text{Ca}_v1.2$. Here, it has been shown that $\alpha_2\delta$ -1 mediates a negative shift in voltage-sensor movement in response to depolarization³⁵. There is also an increase in activation and inactivation kinetics^{36,37}, although these effects depend on the particular $\alpha1$, β , and $\alpha_2\delta$ subunit used (for a recent review, see 10). Results from co-expression studies (which inevitably

lack many components of the native environment) are reinforced by parallel experiments in more intact systems, including using tissues from $\alpha_2\delta$ knockout mice^{20,38–42} and small interfering RNA (siRNA) knockdown of $\alpha_2\delta$ -1 in skeletal muscle cells⁴³ or cardiac myocytes⁴⁴.

Role for $\alpha_2\delta$ -1 in calcium channel trafficking

The effect of $\alpha_2\delta$ subunits to increase calcium current density can be partially explained by an increase in the trafficking of the channels to augment the amount on the cell surface⁵. The exact mechanism whereby $\alpha_2\delta$ increases the density of Ca_v channels in the plasma membrane is still unclear. There was no effect of $\alpha_2\delta$ -1 to reduce the internalization of $\text{Ca}_v2.2$ ⁵, indicating that the effect is likely to be on forward trafficking. Furthermore, the trafficking of $\alpha_2\delta$ itself is blocked by a dominant-negative rab11 construct, suggesting the involvement of the recycling endosomes⁴⁵.

The VWA domain within the α_2 moiety of $\alpha_2\delta$ is important for both trafficking of $\alpha_2\delta$ and its associated effect on Ca_v channel trafficking and function^{5,30,46,47}. Furthermore, the presence of alternatively spliced exon 37a in the proximal C-terminus of $\text{Ca}_v2.2$, which is a minor splice variant expressed particularly in certain DRG neurons^{48,49}, increases $\text{Ca}_v2.2$ currents⁴⁸ and also increases its cell surface density via binding to adaptor proteins⁵⁰. We found that this increase was lost in the absence of $\alpha_2\delta$ subunits, suggesting that this auxiliary subunit promotes particular steps in the forward trafficking process⁵⁰.

Proteomic study of Ca_v2 calcium channels

A comprehensive study of the Ca_v2 channel proteome was performed by using antibodies against $\text{Ca}_v2.1$ or $\text{Ca}_v2.2$, together with antibodies against β subunits, and cataloguing the associated proteins⁵¹. Many proteins were found to be part of this complex, although such studies do not indicate whether the interaction is direct or indirect. In contrast to initial purification studies of N-type channels⁵², and rather surprisingly to many in the field, the interaction of the channels with $\alpha_2\delta$ proteins was found to be much less than 1:1; indeed, it depended on the mildness of the detergent used to solubilize the membranes, resulting in more or less $\alpha_2\delta$ associated with the complex. Since we found that $\alpha_2\delta$ subunits are present in lipid raft fractions⁵³ and subsequently identified that they are GPI-anchored²⁴, this supports the possibility that there is a rather mobile interaction between the $\alpha1$ and $\alpha_2\delta$ subunits^{54,55} or that this interaction is more labile to disruption. Certainly, it also points to a pool of $\alpha_2\delta$ which is not associated with calcium channels, which has also been identified by studies of calcium channel membrane mobility⁵⁴.

Importance of studies in knockout mouse models for elucidating potential novel roles for $\alpha_2\delta$ subunits

The genetic ablation of particular $\alpha_2\delta$ subunits has been found to affect neuronal and synaptic morphology in several systems^{55–57}, pointing to roles for $\alpha_2\delta$ that may or may not involve calcium channels^{22,59}. Knockout mice have been generated for $\alpha_2\delta$ -1³⁸, $\alpha_2\delta$ -2²⁰, $\alpha_2\delta$ -3⁶⁰, and $\alpha_2\delta$ -4⁴¹. These have led to important findings regarding both calcium channel function in

specific tissues and potential roles for the $\alpha_2\delta$ proteins in neuronal and synaptic morphology and in physiological functions, especially in tissues such as cochlear hair cells⁴², spiral ganglion neurons⁵⁷, retinal photoreceptor cells⁵⁸, and Purkinje neurons^{20,56}, where one subtype of $\alpha_2\delta$ predominates. However, complementary approaches are also required to elucidate the mechanisms of such effects.

Importance of $\alpha_2\delta$ in disease states

Neuropathic pain. *Cacna2d1*, encoding $\alpha_2\delta$ -1, is one of many genes whose expression is altered in experimental animals as a result of damage to sensory nerves, which may lead to chronic neuropathic pain. There is a consistent elevation of $\alpha_2\delta$ -1 mRNA and protein^{61–66} in every damaged DRG neuron^{39,62}. Furthermore, we have shown that, in $\alpha_2\delta$ -1 knockout mice³⁸, there is a marked reduction in baseline responses to mechanical and cold stimulation, and a very retarded hyperalgesic response to sciatic nerve injury, in comparison with wild-type littermate mice³⁹.

Other diseases. *CACNA2D1* mutations in humans have been identified to cause cardiac dysfunction, including short QT syndrome⁶⁷ and Brugada syndrome⁶⁸. *Cacna2d1* knockout also resulted in a cardiovascular phenotype in mice involving a reduction in basal ventricular cardiac contractility and lower calcium current in ventricular myocytes³⁸. *CACNA2D2* mutations in both humans and mice result in a recessive phenotype including epilepsy and ataxia^{20,56,69–73}, as well as a hearing deficit, related to aberrant trans-synaptic channel organization⁴². Furthermore, developmentally associated upregulation of $\alpha_2\delta$ -2 expression suppressed axon regeneration in adult spinal cord, although the mechanism remains unclear⁷⁴. *Cacna2d3* knockout mice have a hearing deficit⁵⁷ and a central pain phenotype^{60,75}. Finally, *CACNA2D4* mutations in both humans and mice are associated with night blindness^{76,77} and retinal degeneration⁵⁸.

Mechanism of action of gabapentinoid drugs which bind to $\alpha_2\delta$ -1 and $\alpha_2\delta$ -2

The $\alpha_2\delta$ subunits are the target for gabapentinoid drugs⁷⁸, which bind to both $\alpha_2\delta$ -1 and $\alpha_2\delta$ -2 with similar affinity⁷⁹. However, from studies of mice with mutations in the gabapentin binding site within either $\alpha_2\delta$ -1 or $\alpha_2\delta$ -2, it was concluded that their therapeutic target both in alleviation of neuropathic pain and in epilepsy is $\alpha_2\delta$ -1^{4,80}. We have found, from *in vitro* experiments, that incubation with gabapentin lowers the amount of $\alpha_2\delta$ -1 and $\alpha_2\delta$ -2 on the cell surface^{5,45,81} by inhibiting their rab11-dependent recycling to the cell surface⁴⁵. *In vivo*, chronic administration of pregabalin to sensory nerve-injured rats reduced the elevation in the dorsal horn of pre-synaptic $\alpha_2\delta$ -1, interpreted as being due to inhibition of trafficking⁶². Thus, gabapentin is likely to influence the function of the other proteins to which these $\alpha_2\delta$ proteins have now been found to bind.

For the relevant Ca_v channels, we have also extensively examined the effects of gabapentin. They were initially found to have only small effects on calcium currents when applied acutely⁸². We found that longer-term incubation of cultured cells with gabapentin produced a clear reduction of calcium currents, both in

transfected cells, when $\alpha_2\delta$ -1 or $\alpha_2\delta$ -2 was co-expressed, and in DRG neurons^{45,81,83}. We also observed a corresponding reduction in the expression of Ca_v2 α 1 subunits on the cell surface^{5,45}.

Other interaction partners for $\alpha_2\delta$ proteins related to their function as calcium channel subunits

Several studies in recent years have provided evidence for novel interactions of proteins with $\alpha_2\delta$ subunits; such interactions then impinge on the function of the calcium channel complex. These interactions may be involved positively in the trafficking of $\alpha_2\delta$ proteins (for example, low-density lipoprotein [LDL] receptor-related protein 1, LRP1)²⁷. By contrast, in several studies, the binding partners have been found to sequester $\alpha_2\delta$ proteins, limiting their access to the Ca_v channels, thus reducing both the function and the plasma membrane localization of calcium channels. This mechanism has been proposed for α -neurexins¹¹ and for BK channels¹³ as well as pathologically for a mutant form of prion protein (PrP)⁸⁴. These will all be considered in turn.

Trafficking of $\alpha_2\delta$ -1 by the multifunctional transport protein LRP1

The LRP family represents a large group of ligand-binding and trafficking proteins, including the LDL receptor and LRP1–6. They are multifunctional, multi-domain receptors, interacting with many protein ligands via their ligand-binding domains, mediating both forward trafficking and endocytosis of these ligands⁸⁵. They are also involved as co-receptors, affecting intracellular cell signaling processes^{86,87}.

LRP1 is a ubiquitous membrane protein with four ligand-binding domains (Figure 4a) and is involved in forward trafficking of proteins, including several TSPs^{88–92}, PrP⁹³, and NMDA receptors⁹⁴. LRP1 is also involved in clathrin-dependent endocytosis^{85,95}. It is present in synapses⁹⁴ and is implicated in neurite outgrowth⁹⁶. Whether different LRP proteins bind to overlapping sets of protein ligands is unclear, but LRP5/6 are also involved in Wnt signaling⁸⁷.

We recently showed that LRP1 binds to $\alpha_2\delta$ -1²⁷ and the same is true for $\alpha_2\delta$ -2 and $\alpha_2\delta$ -3 (Ivan Kadurin and Annette Dolphin, preliminary results). For $\alpha_2\delta$ -1, we showed this interaction is direct, involving the VWA domain of $\alpha_2\delta$ -1 and LRP1 ligand-binding domains II and IV (Figure 4a)²⁷. The association is modulated by the LRP chaperone, receptor-associated protein (RAP), which is required for the correct folding of all LRP proteins and for their trafficking out of the endoplasmic reticulum^{97,98}. We found that the LRP1/RAP combination increases mature glycosylation, proteolytic processing, and cell-surface expression of $\alpha_2\delta$ -1 and also increases plasma membrane expression and function of $\text{Ca}_v2.2$ when co-expressed with $\alpha_2\delta$ -1²⁷. Since LRP1 is able to bind more than one ligand at different sites⁹⁹, it is possible that it forms a bridge between $\alpha_2\delta$ -1 and other proteins, such as TSPs.

Sequestration of $\alpha_2\delta$ -3 by interaction with α -neurexins

There are three vertebrate neurexin genes, and each can form α - and β -neurexins from different promoters. The α -neurexins have been found to be important for coupling calcium channels

Figure 4. Protein domains involved in novel $\alpha_2\delta$ interactions. (a) Interaction of $\alpha_2\delta$ -1 (and $\alpha_2\delta$ -2/3) with the ligand-binding repeats II and IV of low-density lipoprotein receptor-related protein 1 (LRP1) (red). Other domains in LRP1 are epithelial growth factor (EGF)-like repeats (orange) and β -propeller domains (cyan)²⁷. i/c, intracellular; TM, transmembrane. (b) Interaction of neurexin 1 α with $\alpha_2\delta$ -3, via its laminin-like globular (LG) repeats (L, green) 1 and 5. E, EGF-like repeat (orange). Neurexin 1 α is cleaved by a disintegrin and metalloprotease 10 (ADAM 10) (arrow) to have the observed effects on synaptic transmission, but it is not clear whether this is required for the interaction with $\alpha_2\delta$ -3¹¹. (c) Interaction of the extracellular N-terminus of large conductance (big) potassium (BK) α subunits with $\alpha_2\delta$ -1. The three blue arrows indicate the three alternative N-terminal translation initiation sites, the third being the most commonly used¹³. S0 is the additional transmembrane domain (red). (d) Interaction of $\alpha_2\delta$ -1 von Willebrand factor A (VWA) domain with the EGF-like domains (black bars) of both pentameric (left) and trimeric (right) thrombospondins (TSPs)¹⁴. (e) Interaction of a C-terminal region of $\alpha_2\delta$ -1 beyond its GPI-anchor site (dashed orange/white region) with the N-methyl-D-aspartate (NMDA) receptor GluN1, GluN2A, and GluN2B subunits¹⁵.

to synaptic transmission¹⁰⁰. Whereas in mammalian synapses the neurexins are pre-synaptic and bind to post-synaptic neuroligins, in *Caenorhabditis elegans* this polarity is reversed at many synapses. It has been found in the worm that post-synaptic neurexin 1 α at the neuromuscular junction binds, via its laminin-like globular 1 (LG1) domain, to pre-synaptic unc-36 (similar to $\alpha_2\delta$ -3), thus decreasing its availability to bind to the pre-synaptic unc-2 (a Ca_v2 -like channel) that mediates neurotransmitter release¹¹. This was found to reduce synaptic transmission, an effect which required a proteolytically cleaved fragment of neurexin, shed from the post-synaptic plasma membrane (Figure 4b). In transfected cells, mouse neurexin 1 α was found to bind $\alpha_2\delta$ -3 and to decrease $\text{Ca}_v2.2$ current, whereas there was no effect on $\text{Ca}_v2.2$ currents in the presence of $\alpha_2\delta$ -1 or $\alpha_2\delta$ -2¹¹. An attractive suggestion is that this type of pre- to post-synaptic interaction may contribute to trans-synaptic nanoscale organization¹⁰¹. However, in view of recent results described below, it will be important in the future to identify the site of selective interaction on the $\alpha_2\delta$ -3 protein of the LG1 domain (and LG5 in the mouse)¹¹ of neurexin 1 α .

In contrast, a more recent article has identified positive effects of neurexin 1 α in the presence of $\alpha_2\delta$ -1 (but not $\alpha_2\delta$ -3) on pre-synaptic Ca^{2+} transients in hippocampal neurons and in parallel on $\text{Ca}_v2.1$ calcium currents¹². Importantly, very carefully done experiments, designed to detect an interaction of neurexin 1 α with $\alpha_2\delta$ -1 or $\alpha_2\delta$ -3, failed to find a specific association between the two proteins, as every protein tested (α -neurexin, neuroligin, and two forms of cadherin) was pulled down with $\alpha_2\delta$ -1 (and also $\alpha_2\delta$ -3 co-immunoprecipitated with neurexin 1 α). The authors concluded that neurexin 1 α does not form stable complexes with $\alpha_2\delta$ subunits but nevertheless influences their function. Their results also provide a warning that $\alpha_2\delta$ proteins may be rather prone to co-immunoprecipitation artefacts.

Sequestration of $\alpha_2\delta$ -1 by interaction with BK channels

A recent study has identified that BK α subunits bind to $\alpha_2\delta$ -1 subunits via the BK N-terminus¹³, and the authors suggest that this interaction sequesters $\alpha_2\delta$ -1 from Ca_v channels. BK channels are important mediators of cell excitability, as they respond to both

TSPs are extracellular matrix proteins which bind to a very large number of proteins, 83 being so far identified for TSP-1¹¹³; consequently, they have many functions^{114–116}. In the brain, they are produced by astrocytes and promote neurite outgrowth¹¹⁷, including the formation of silent excitatory synapses, lacking post-synaptic receptors¹¹⁸. It was then hypothesized that post-synaptic $\alpha_2\delta$ -1 could be the sought-after post-synaptic binding partner of TSPs to mediate synaptogenesis, independent of any effects on calcium channels. This was first tested using co-immunoprecipitation to determine whether TSPs or individual domains of TSPs interacted with C-terminally tagged $\alpha_2\delta$ -1¹⁴. An interaction which involved a key synaptogenic epithelial growth factor (EGF)-like domain was found (Figure 4d). As a note of caution, C-terminal tagging may interfere with trafficking of $\alpha_2\delta$ -1 by disrupting the GPI anchor^{24,26}. Nevertheless, gabapentin was found to inhibit the interaction between $\alpha_2\delta$ -1 and the EGF-like domain of TSP-2 and

to disrupt synaptogenesis. Furthermore, *in vivo*, gabapentin was found to disrupt whisker barrel plasticity following whisker removal in some of the mice examined¹⁴.

TSP-4 is upregulated in rodent models of neuropathic pain¹¹⁹. Since $\alpha_2\delta$ -1 is also upregulated in DRGs following peripheral sensory nerve injury, several studies have investigated whether an interaction between these two proteins is important in neuropathic pain or the effect of gabapentin. Interestingly, in a recent article, it was suggested that pre-synaptic, rather than post-synaptic, $\alpha_2\delta$ -1 may be a synaptogenic binding partner for TSP-4 in the spinal cord¹²⁰.

We found (using overexpressed proteins) that TSP-4 modestly reduced the affinity for ³H-gabapentin binding to $\alpha_2\delta$ -1, although the effect on ³H-gabapentin binding was not reproduced with the TSP-4 synaptogenic EGF-like domain. Furthermore, we found only very weak and unreliable co-immunoprecipitation of the two proteins, which again could not be reproduced with the synaptogenic EGF-like domain of TSP-4¹²¹. We also could not demonstrate any interaction between $\alpha_2\delta$ -1 and TSP-4 on the cell surface of transfected cells, suggesting that the association between these two proteins to disrupt ³H-gabapentin binding is occurring intracellularly following co-transfection, when the two proteins are juxtaposed at high concentration¹²¹.

Nevertheless, there is evidence from other studies that $\alpha_2\delta$ subunits are important for synaptic morphology in several different systems^{57,58,122,123}. Whether the role for $\alpha_2\delta$ in calcium channel localization and function is responsible for these morphological changes has not always been investigated. However, $\alpha_2\delta$ was shown to increase pre-synaptic localization of the relevant $\alpha 1$ subunit in *Drosophila* neuromuscular junction synapses¹²⁴ as well as in retinal⁵⁸ and hippocampal⁶ synapses.

$\alpha_2\delta$ -1 as an NMDA receptor trafficking protein

It was recently shown that overexpression of $\alpha_2\delta$ -1 administered intrathecally into the spinal cord potentiates pre-synaptic and post-synaptic NMDA receptor activity, and it was further shown that $\alpha_2\delta$ -1 interacted with NMDA receptors, both in spinal cord and in overexpression studies¹⁵. The interaction was apparently specific for $\alpha_2\delta$ -1, as it did not occur with $\alpha_2\delta$ -2 or $\alpha_2\delta$ -3. The authors identified the site of interaction as the C-terminus of $\alpha_2\delta$ -1, surprisingly after the C-terminal GPI-anchor cleavage site (Figure 4e). This was determined using chimeras assembled from the different isoforms, swapping isoforms either between $\alpha 2$ and δ or with the C-terminus of δ ⁶. However, it is important to note that such chimeras may have disrupted the primary sequences involved in proteolytic cleavage between α_2 and δ , a process which is important for function^{6,28}, or it might have affected the sequences involved in GPI anchoring²⁴. Nevertheless, this result suggests either that a transmembrane version of $\alpha_2\delta$ -1 may be interacting with NMDA receptors, initially in the endoplasmic reticulum, or that the NMDA receptor interacts with the C-terminal peptide of $\alpha_2\delta$ -1 that is cleaved off during GPI-anchor attachment¹²⁵.

The GluN1, GluN2A, and GluN2B subunits of NMDA receptors were found to interact with $\alpha_2\delta$ -1, presumably via the

transmembrane or intracellular domains of these subunits, since the identified interaction is with the C-terminus of $\alpha_2\delta$ -1¹⁵. The C-termini of these NMDA receptors are rather different in both sequence and function^{126–128}, and determining the interaction site will be a key next step. It is of interest that $\alpha_2\delta$ -1 has not been previously detected in proteomic studies of post-synaptic densities¹²⁹. In contrast, other calcium channel subunits ($\text{Ca}_v1.2$, $\text{Ca}_v2.3$, and β) were identified. Another recent study also did not detect $\alpha_2\delta$ -1 when purifying NMDA receptors from mouse brain¹²⁸, although $\alpha_2\delta$ -1 is widely expressed in most brain regions^{130,131}. Therefore, it would be important to determine whether this interaction is for some reason observed only in the spinal cord. One possible reason is that it might be indirect (for example, via a scaffolding protein expressed in the spinal cord, interacting with both $\alpha_2\delta$ -1 and NMDA receptors).

Conclusions and future directions

The $\alpha_2\delta$ subunits are important auxiliary subunits of the Ca_v1 and Ca_v2 voltage-gated calcium channels. They play key roles in trafficking of these channels, both to the plasma membrane and to specific subcellular domains, and they have marked effects on the activation and other biophysical properties of these channels, indicating their importance as subunits of the channel complex rather than purely as chaperones. However, recent evidence suggests that they may bind to other proteins, and one role for such additional interactions could be to sequester particular $\alpha_2\delta$ subunits at specific sites away from the calcium channels in a dynamic manner and thus reduce calcium channel function. Evidence also suggests that $\alpha_2\delta$ proteins may independently influence other channels and also affect other functions of neurons. All of these novel functions will need to be critically explored in the future to evaluate further their physiological, pathological, and pharmacological relevance. Furthermore, the roles for novel $\alpha_2\delta$ -like protein, *Cachd1*, which enhances both T-type channels¹³² and N-type channels¹³³ as well as competes with $\alpha_2\delta$ -1¹³³, will be explored further in the future.

Abbreviations

BK, large conductance (big) potassium; EGF, epithelial growth factor; GPI, glycosylphosphatidylinositol; LDL, low-density lipoprotein; LG, laminin-like globular; LRP, low-density lipoprotein receptor-related protein; MIDAS, metal ion-dependent adhesion site; NMDA, *N*-methyl-D-aspartate; PrP, prion protein; RAP, receptor-associated protein; TSP, thrombospondin; VWA, von Willebrand factor A.

Grant information

My own work referenced in this study has been funded by Wellcome Trust and MRC.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgements

I would like to thank Mike Shipston, Mark Farrant, and Seth Grant for sharing their invaluable expertise with me as well as members of my group (particularly Laurent Ferron and Ivan Kadurin) for discussions.

References

1. Takahashi M, Seagar MJ, Jones JF, *et al.*: **Subunit structure of dihydropyridine-sensitive calcium channels from skeletal muscle.** *Proc Natl Acad Sci U S A*. 1987; **84**(15): 5478–82.
[PubMed Abstract](#) | [Publisher Full Text](#)
2. Leung AT, Imagawa T, Campbell KP: **Structural characterization of the 1,4-dihydropyridine receptor of the voltage-dependent Ca²⁺ channel from rabbit skeletal muscle. Evidence for two distinct high molecular weight subunits.** *J Biol Chem*. 1987; **262**(17): 7943–6.
[PubMed Abstract](#)
3. Brown JP, Gee NS: **Cloning and deletion mutagenesis of the alpha₂ delta calcium channel subunit from porcine cerebral cortex. Expression of a soluble form of the protein that retains [³H]gabapentin binding activity** *J Biol Chem*. 1998; **273**(39): 25458–65.
[PubMed Abstract](#) | [Publisher Full Text](#)
4. Field MJ, Cox PJ, Stott E, *et al.*: **Identification of the alpha₂-delta-1 subunit of voltage-dependent calcium channels as a molecular target for pain mediating the analgesic actions of pregabalin.** *Proc Natl Acad Sci U S A*. 2006; **103**(46): 17537–42.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
5. Cassidy JS, Ferron L, Kadurin I, *et al.*: **Functional exofacially tagged N-type calcium channels elucidate the interaction with auxiliary alpha₂delta-1 subunits.** *Proc Natl Acad Sci U S A*. 2014; **111**(24): 8979–84.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
6. Kadurin I, Ferron L, Rothwell SW, *et al.*: **Proteolytic maturation of alpha₂delta represents a checkpoint for activation and neuronal trafficking of latent calcium channels.** *eLife*. 2016; **5**: pii: e21143.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
7. Gurnett CA, De Waard M, Campbell KP: **Dual function of the voltage-dependent Ca²⁺ channel alpha₂delta subunit in current stimulation and subunit interaction.** *Neuron*. 1996; **16**(2): 431–40.
[PubMed Abstract](#) | [Publisher Full Text](#)
8. Shistik E, Ivanina T, Puri T, *et al.*: **Ca²⁺ current enhancement by alpha 2/delta and beta subunits in Xenopus oocytes: contribution of changes in channel gating and alpha 1 protein level.** *J Physiol*. 1995; **489**(Pt 1): 55–62.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
9. Mikami A, Imoto K, Tanabe T, *et al.*: **Primary structure and functional expression of the cardiac dihydropyridine-sensitive calcium channel.** *Nature*. 1989; **340**(6230): 230–3.
[PubMed Abstract](#) | [Publisher Full Text](#)
10. Dolphin AC: **Voltage-gated calcium channels and their auxiliary subunits: physiology and pathophysiology and pharmacology.** *J Physiol*. 2016; **594**(19): 5369–90.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
11. Tong XJ, López-Soto EJ, Li L, *et al.*: **Retrograde Synaptic Inhibition Is Mediated by alpha-Neurexin Binding to the alpha₂delta Subunits of N-Type Calcium Channels.** *Neuron*. 2017; **95**(2): 326–340.e5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
12. Brockhaus J, Schreitmüller M, Repetto D, *et al.*: **alpha-Neurexins Together with alpha₂delta-1 Auxiliary Subunits Regulate Ca²⁺ Influx through Ca_v2.1 Channels.** *J Neurosci*. 2018; **38**(38): 8277–94.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
13. Zhang FX, Gadotti VM, Souza IA, *et al.*: **BK Potassium Channels Suppress Cav₂delta Subunit Function to Reduce Inflammatory and Neuropathic Pain.** *Cell Rep*. 2018; **22**(8): 1956–64.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
14. Eroglu C, Allen NJ, Susman MW, *et al.*: **Gabapentin receptor alpha2delta-1 is a neuronal thrombospondin receptor responsible for excitatory CNS synaptogenesis.** *Cell*. 2009; **139**(2): 380–92.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
15. Chen J, Li L, Chen SR, *et al.*: **The alpha₂delta-1-NMDA Receptor Complex Is Critically Involved in Neuropathic Pain Development and Gabapentin Therapeutic Actions.** *Cell Rep*. 2018; **22**(9): 2307–21.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
16. Ellis SB, Williams ME, Ways NR, *et al.*: **Sequence and expression of mRNAs encoding the alpha 1 and alpha 2 subunits of a DHP-sensitive calcium channel.** *Science*. 1988; **241**(4873): 1661–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
17. Jay SD, Sharp AH, Kahl SD, *et al.*: **Structural characterization of the dihydropyridine-sensitive calcium channel alpha 2-subunit and the associated delta peptides.** *J Biol Chem*. 1991; **266**(5): 3287–93.
[PubMed Abstract](#)
18. De Jongh KS, Warner C, Catterall WA: **Subunits of purified calcium channels. Alpha 2 and delta are encoded by the same gene.** *J Biol Chem*. 1990; **265**(25): 14738–41.
[PubMed Abstract](#)
19. Klugbauer N, Lacinová L, Marais E, *et al.*: **Molecular diversity of the calcium channel alpha₂delta subunit.** *J Neurosci*. 1999; **19**(2): 684–91.
[PubMed Abstract](#) | [Publisher Full Text](#)
20. Barclay J, Balaguero N, Mione M, *et al.*: **Ducky mouse phenotype of epilepsy and ataxia is associated with mutations in the Cacna2d2 gene and decreased calcium channel current in cerebellar Purkinje cells.** *J Neurosci*. 2001; **21**(6): 6095–104.
[PubMed Abstract](#) | [Publisher Full Text](#)
21. Qin N, Yagel S, Momplaisir ML, *et al.*: **Molecular cloning and characterization of the human voltage-gated calcium channel alpha₂delta₁ subunit.** *Mol Pharmacol*. 2002; **62**(3): 485–96.
[PubMed Abstract](#) | [Publisher Full Text](#)
22. Dolphin AC: **Calcium channel auxiliary alpha₂delta and beta subunits: trafficking and one step beyond.** *Nat Rev Neurosci*. 2012; **13**(8): 542–55.
[PubMed Abstract](#) | [Publisher Full Text](#)
23. Davies A, Hendrich J, Van Minh AT, *et al.*: **Functional biology of the alpha₂delta subunits of voltage-gated calcium channels.** *Trends Pharmacol Sci*. 2007; **28**(5): 220–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
24. Davies A, Kadurin I, Alvarez-Laviada A, *et al.*: **The alpha₂delta subunits of voltage-gated calcium channels form GPI-anchored proteins, a posttranslational modification essential for function.** *Proc Natl Acad Sci U S A*. 2010; **107**(4): 1654–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
25. Alvarez-Laviada A, Kadurin I, Senatore A, *et al.*: **The inhibition of functional expression of calcium channels by prion protein demonstrates competition with alpha₂delta for GPI-anchoring pathways.** *Biochem J*. 2014; **458**(2): 365–74.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
26. Kadurin I, Alvarez-Laviada A, Ng SF, *et al.*: **Calcium currents are enhanced by alpha₂delta-1 lacking its membrane anchor.** *J Biol Chem*. 2012; **287**(40): 33554–66.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
27. Kadurin I, Rothwell SW, Lana B, *et al.*: **LRP1 influences trafficking of N-type calcium channels via interaction with the auxiliary alpha₂delta-1 subunit.** *Sci Rep*. 2017; **7**: 43802.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
28. Ferron L, Kadurin I, Dolphin AC: **Proteolytic maturation of alpha₂delta controls the probability of synaptic vesicular release.** *eLife*. 2018; **7**: pii: e37507.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
29. Whittaker CA, Hynes RO: **Distribution and evolution of von Willebrand/integrin A domains: widely dispersed domains with roles in cell adhesion and elsewhere.** *Mol Biol Cell*. 2002; **13**(10): 3369–87.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
30. Cantí C, Nieto-Rostro M, Foucault I, *et al.*: **The metal-ion-dependent adhesion site in the Von Willebrand factor-A domain of alpha₂delta subunits is key to trafficking voltage-gated Ca²⁺ channels.** *Proc Natl Acad Sci U S A*. 2005; **102**(32): 11230–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
31. Anantharaman V, Aravind L: **Cache - a signaling domain common to animal Ca²⁺-channel subunits and a class of prokaryotic chemotaxis receptors.** *Trends Biochem Sci*. 2000; **25**(11): 535–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
32. Wu J, Yan Z, Li Z, *et al.*: **Structure of the voltage-gated calcium channel Ca_v1.1 at 3.6 Å resolution.** *Nature*. 2016; **537**(7619): 191–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
33. Calderón-Rivera A, Andrade A, Hernández-Hernández O, *et al.*: **Identification of a disulfide bridge essential for structure and function of the voltage-gated Ca²⁺ channel alpha₂delta-1 auxiliary subunit.** *Cell Calcium*. 2012; **51**(1): 22–30.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
34. Felix R, Gurnett CA, De Waard M, *et al.*: **Dissection of functional domains of the voltage-dependent Ca²⁺ channel alpha₂delta subunit.** *J Neurosci*. 1997; **17**(18): 6884–91.
[PubMed Abstract](#) | [Publisher Full Text](#)
35. Savalli N, Pantazis A, Sigg D, *et al.*: **The alpha₂delta-1 subunit remodels Ca_v1.2 voltage sensors and allows Ca²⁺ influx at physiological membrane potentials.** *J Gen Physiol*. 2016; **148**(2): 147–59.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
36. Cantí C, Davies A, Dolphin A: **Calcium Channel alpha2delta Subunits: Structure, Functions and Target Site for Drugs.** *Curr Neuropharmacol*. 2003; **1**(3): 209–17.
[Publisher Full Text](#)
37. Qin N, Olcese R, Stefani E, *et al.*: **Modulation of human neuronal alpha 1E-type calcium channel by alpha 2 delta-subunit.** *Am J Physiol*. 1998; **274**(5 Pt 1): C1324–31.
[PubMed Abstract](#) | [Publisher Full Text](#)
38. Fuller-Bicer GA, Varadi G, Koch SE, *et al.*: **Targeted disruption of the voltage-dependent calcium channel alpha₂delta-1-subunit.** *Am J Physiol Heart Circ Physiol*. 2009; **297**(1): H117–24.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
39. Patel R, Bauer CS, Nieto-Rostro M, *et al.*: **alpha₂delta-1 gene deletion affects somatosensory neuron function and delays mechanical hypersensitivity in response to peripheral nerve damage.** *J Neurosci*. 2013; **33**(42): 16412–26.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**

40. Mastrolia V, Flucher SM, Obermair GJ, *et al.*: Loss of $\alpha_2\delta$ -1 Calcium Channel Subunit Function Increases the Susceptibility for Diabetes. *Diabetes*. 2017; 66(4): 897–907.
[PubMed Abstract](#) | [Publisher Full Text](#)
41. Wang Y, Fehlhaber KE, Sarria I, *et al.*: The Auxiliary Calcium Channel Subunit $\alpha_2\delta$ Is Required for Axonal Elaboration, Synaptic Transmission, and Wiring of Rod Photoreceptors. *Neuron*. 2017; 93(6): 1359–1374.e6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
42. Fell B, Eckrich S, Blum K, *et al.*: $\alpha_2\delta$ Controls the Function and Trans-Synaptic Coupling of $\text{Ca}_v1.3$ Channels in Mouse Inner Hair Cells and Is Essential for Normal Hearing. *J Neurosci*. 2016; 36(43): 11024–36.
[PubMed Abstract](#) | [Publisher Full Text](#)
43. Obermair GJ, Kugler G, Baumgartner S, *et al.*: The Ca^{2+} channel $\alpha_2\delta$ -1 subunit determines Ca^{2+} current kinetics in skeletal muscle but not targeting of $\alpha_1\text{S}$ or excitation-contraction coupling. *J Biol Chem*. 2005; 280(3): 2229–37.
[PubMed Abstract](#) | [Publisher Full Text](#)
44. Tuluc P, Kern G, Obermair GJ, *et al.*: Computer modeling of siRNA knockdown effects indicates an essential role of the Ca^{2+} channel $\alpha_2\delta$ -1 subunit in cardiac excitation-contraction coupling. *Proc Natl Acad Sci U S A*. 2007; 104(26): 11091–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
45. Tran-Van-Minh A, Dolphin AC: The $\alpha_2\delta$ ligand gabapentin inhibits the Rab11-dependent recycling of the calcium channel subunit $\alpha_2\delta$ -2. *J Neurosci*. 2010; 30(38): 12856–67.
[PubMed Abstract](#) | [Publisher Full Text](#)
46. Bourdin B, Briot J, Tétrault MP, *et al.*: Negatively charged residues in the first extracellular loop of the L-type $\text{Ca}_v1.2$ channel anchor the interaction with the $\text{Ca}_v\alpha_2\delta$ auxiliary subunit. *J Biol Chem*. 2017; 292(42): 17236–49.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
47. Hoppa MB, Lana B, Margas W, *et al.*: $\alpha_2\delta$ expression sets presynaptic calcium channel abundance and release probability. *Nature*. 2012; 486(7401): 122–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
48. Castiglioni AJ, Raingo J, Lipscombe D: Alternative splicing in the C-terminus of $\text{CaV}2.2$ controls expression and gating of N-type calcium channels. *J Physiol*. 2006; 576(Pt 1): 119–34.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
49. Altier C, Dale CS, Kisilevsky AE, *et al.*: Differential role of N-type calcium channel splice isoforms in pain. *J Neurosci*. 2007; 27(24): 6363–73.
[PubMed Abstract](#) | [Publisher Full Text](#)
50. Macabuag N, Dolphin AC: Alternative Splicing in $\text{Ca}_v2.2$ Regulates Neuronal Trafficking via Adaptor Protein Complex-1 Adaptor Protein Motifs. *J Neurosci*. 2015; 35(43): 14636–52.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
51. Müller CS, Haupt A, Bildl W, *et al.*: Quantitative proteomics of the $\text{CaV}2$ channel nano-environments in the mammalian brain. *Proc Natl Acad Sci U S A*. 2010; 107(34): 14950–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
52. Witcher DR, De Waard M, Sakamoto J, *et al.*: Subunit identification and reconstitution of the N-type Ca^{2+} channel complex purified from brain. *Science*. 1993; 261(5120): 486–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
53. Davies A, Douglas L, Hendrich J, *et al.*: The calcium channel $\alpha_2\delta$ -2 subunit partitions with $\text{Ca}_v2.1$ into lipid rafts in cerebellum: implications for localization and function. *J Neurosci*. 2006; 26(34): 8748–57.
[PubMed Abstract](#) | [Publisher Full Text](#)
54. Voigt A, Freund R, Heck J, *et al.*: Dynamic association of calcium channel subunits at the cellular membrane. *Neurophotonics*. 2016; 3(4): 041809.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
55. Obermair GJ, Tuluc P, Flucher BE: Auxiliary Ca^{2+} channel subunits: lessons learned from muscle. *Curr Opin Pharmacol*. 2008; 8(3): 311–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
56. Brodbeck J, Davies A, Courtney JM, *et al.*: The ducky mutation in *Cacna2d2* results in altered Purkinje cell morphology and is associated with the expression of a truncated $\alpha_2\delta$ protein with abnormal function. *J Biol Chem*. 2002; 277(10): 7684–93.
[PubMed Abstract](#) | [Publisher Full Text](#)
57. Pirone A, Kurt S, Zuccotti A, *et al.*: $\alpha_3\delta$ is essential for normal structure and function of auditory nerve synapses and is a novel candidate for auditory processing disorders. *J Neurosci*. 2014; 34(2): 434–45.
[PubMed Abstract](#) | [Publisher Full Text](#)
58. Kerov V, Laird JG, Joiner ML, *et al.*: $\alpha_2\delta$ -4 Is Required for the Molecular and Structural Organization of Rod and Cone Photoreceptor Synapses. *J Neurosci*. 2018; 38(27): 6145–60.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
59. Geisler S, Schöpp CL, Obermair GJ: Emerging evidence for specific neuronal functions of auxiliary calcium channel $\alpha_2\delta$ subunits. *Gen Physiol Biophys*. 2015; 34(2): 105–18.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
60. Neely GG, Hess A, Costigan M, *et al.*: A genome-wide *Drosophila* screen for heat nociception identifies $\alpha_2\delta$ as an evolutionarily conserved pain gene. *Cell*. 2010; 143(4): 628–38.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
61. Newton RA, Bingham S, Case PC, *et al.*: Dorsal root ganglion neurons show increased expression of the calcium channel $\alpha_2\delta$ -1 subunit following partial sciatic nerve injury. *Brain Res Mol Brain Res*. 2001; 95(1–2): 1–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
62. Bauer CS, Nieto-Rostro M, Rahman W, *et al.*: The increased trafficking of the calcium channel subunit $\alpha_2\delta$ -1 to presynaptic terminals in neuropathic pain is inhibited by the $\alpha_2\delta$ ligand pregabalin. *J Neurosci*. 2009; 29(13): 4076–88.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
63. Wang H, Sun H, Della Penna K, *et al.*: Chronic neuropathic pain is accompanied by global changes in gene expression and shares pathobiology with neurodegenerative diseases. *Neuroscience*. 2002; 114(3): 529–46.
[PubMed Abstract](#) | [Publisher Full Text](#)
64. Xiao HS, Huang QH, Zhang FX, *et al.*: Identification of gene expression profile of dorsal root ganglion in the rat peripheral axotomy model of neuropathic pain. *Proc Natl Acad Sci U S A*. 2002; 99(12): 8360–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
65. Perkins JR, Antunes-Martins A, Calvo M, *et al.*: A comparison of RNA-seq and exon arrays for whole genome transcription profiling of the L5 spinal nerve transection model of neuropathic pain in the rat. *Mol Pain*. 2014; 10: 7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
66. Luo ZD, Chaplan SR, Higuera ES, *et al.*: Upregulation of dorsal root ganglion ($\alpha_2\delta$) calcium channel subunit and its correlation with allodynia in spinal nerve-injured rats. *J Neurosci*. 2001; 21(6): 1868–75.
[PubMed Abstract](#) | [Publisher Full Text](#)
67. Templin C, Ghadri JR, Rougier JS, *et al.*: Identification of a novel loss-of-function calcium channel gene mutation in short QT syndrome (SQTS6). *Eur Heart J*. 2011; 32(9): 1077–88.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
68. Burashnikov E, Pfeiffer R, Barajas-Martinez H, *et al.*: Mutations in the cardiac L-type calcium channel associated with inherited J-wave syndromes and sudden cardiac death. *Heart Rhythm*. 2010; 7(12): 1872–82.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
69. Donato R, Page KM, Koch D, *et al.*: The ducky² mutation in *Cacna2d2* results in reduced spontaneous Purkinje cell activity and altered gene expression. *J Neurosci*. 2006; 26(48): 12576–86.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
70. Brill J, Klocke R, Paul D, *et al.*: *entla*, a novel epileptic and ataxic *Cacna2d2* mutant of the mouse. *J Biol Chem*. 2004; 279(8): 7322–30.
[PubMed Abstract](#) | [Publisher Full Text](#)
71. Ivanov SV, Ward JM, Tessarollo L, *et al.*: Cerebellar ataxia, seizures, premature death, and cardiac abnormalities in mice with targeted disruption of the *Cacna2d2* gene. *Am J Pathol*. 2004; 165(3): 1007–18.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
72. Pippucci T, Parmeggiani A, Palombo F, *et al.*: A novel null homozygous mutation confirms *CACNA2D2* as a gene mutated in epileptic encephalopathy. *PLoS One*. 2013; 8(12): e82154.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
73. Edvardson S, Oz S, Abulhijab FA, *et al.*: Early infantile epileptic encephalopathy associated with a high voltage gated calcium channelopathy. *J Med Genet*. 2013; 50(2): 118–23.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
74. Tedeschi A, Dupraz S, Laskowski CJ, *et al.*: The Calcium Channel Subunit $\alpha_2\delta$ Suppresses Axon Regeneration in the Adult CNS. *Neuron*. 2016; 92(2): 419–34.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
75. Landmann J, Richter F, Oros-Peusquens AM, *et al.*: Neuroanatomy of pain-deficiency and cross-modal activation in calcium channel subunit (*CACN*) $\alpha_2\delta$ knockout mice. *Brain Struct Funct*. 2018; 223(1): 111–30.
[PubMed Abstract](#) | [Publisher Full Text](#)
76. Wycisk KA, Budde B, Feil S, *et al.*: Structural and functional abnormalities of retinal ribbon synapses due to *Cacna2d4* mutation. *Invest Ophthalmol Vis Sci*. 2006; 47(8): 3523–30.
[PubMed Abstract](#) | [Publisher Full Text](#)
77. Wycisk KA, Zeitz C, Feil S, *et al.*: Mutation in the auxiliary calcium-channel subunit *CACNA2D4* causes autosomal recessive cone dystrophy. *Am J Hum Genet*. 2006; 79(5): 973–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
78. Gee NS, Brown JP, Dissanayake VU, *et al.*: The novel anticonvulsant drug, gabapentin (Neurontin), binds to the $\alpha_2\delta$ subunit of a calcium channel. *J Biol Chem*. 1996; 271(10): 5768–76.
[PubMed Abstract](#) | [Publisher Full Text](#)
79. Su T, Gong CH, Hang J, *et al.*: Human $\alpha_2\delta$ subunit of calcium channel: a novel gabapentin binding protein in brain. *Society for Neuroscience. Abstracts*. 2000 2000; 26: 40.20.
80. Lotarski S, Hain H, Peterson J, *et al.*: Anticonvulsant activity of pregabalin in the maximal electroshock-induced seizure assay in $\alpha_2\delta_1$ (R217A) and $\alpha_2\delta_2$ (R279A) mouse mutants. *Epilepsy Res*. 2014; 108(5): 833–42.
[PubMed Abstract](#) | [Publisher Full Text](#)
81. Hendrich J, Van Minh AT, Heblisch F, *et al.*: Pharmacological disruption of

- calcium channel trafficking by the $\alpha_2\delta_1$ ligand gabapentin. *Proc Natl Acad Sci U S A*. 2008; 105(9): 3628–33.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
82. Martin DJ, McClelland D, Herd MB, *et al.*: Gabapentin-mediated inhibition of voltage-activated Ca^{2+} channel currents in cultured sensory neurones is dependent on culture conditions and channel subunit expression. *Neuropharmacology*. 2002; 42(3): 353–66.
[PubMed Abstract](#) | [Publisher Full Text](#)
 83. Hebllich F, Tran Van Minh A, Hendrich J, *et al.*: Time course and specificity of the pharmacological disruption of the trafficking of voltage-gated calcium channels by gabapentin. *Channels (Austin)*. 2008; 2(1): 4–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
 84. **F** Senatore A, Colleoni S, Verderio C, *et al.*: Mutant PrP suppresses glutamatergic neurotransmission in cerebellar granule neurons by impairing membrane delivery of VGCC $\alpha_2\delta_1$ Subunit. *Neuron*. 2012; 74(2): 300–13.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 85. Lillis AP, Van Duyn LB, Murphy-Ullrich JE, *et al.*: LDL receptor-related protein 1: unique tissue-specific functions revealed by selective gene knockout studies. *Physiol Rev*. 2008; 88(3): 887–918.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 86. Orr AW, Pedraza CE, Pallero MA, *et al.*: Low density lipoprotein receptor-related protein is a calreticulin coreceptor that signals focal adhesion disassembly. *J Cell Biol*. 2003; 161(6): 1179–89.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 87. Chen S, Bubeck D, MacDonald BT, *et al.*: Structural and functional studies of LRP6 ectodomain reveal a platform for Wnt signaling. *Dev Cell*. 2011; 21(5): 848–61.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 88. Pallero MA, Elzie CA, Chen J, *et al.*: Thrombospondin 1 binding to calreticulin-LRP1 signals resistance to anoikis. *FASEB J*. 2008; 22(11): 3968–79.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 89. Wang S, Herndon ME, Ranganathan S, *et al.*: Internalization but not binding of thrombospondin-1 to low density lipoprotein receptor-related protein-1 requires heparan sulfate proteoglycans. *J Cell Biochem*. 2004; 91(4): 766–76.
[PubMed Abstract](#) | [Publisher Full Text](#)
 90. Mikhailenko I, Kounnas MZ, Strickland DK: Low density lipoprotein receptor-related protein/alpha 2-macroglobulin receptor mediates the cellular internalization and degradation of thrombospondin. A process facilitated by cell-surface proteoglycans. *J Biol Chem*. 1995; 270(16): 9543–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
 91. Strickland DK, Kounnas MZ, Argraves WS: LDL receptor-related protein: a multiligand receptor for lipoprotein and proteinase catabolism. *FASEB J*. 1995; 9(10): 890–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
 92. Godyna S, Liu G, Popa I, *et al.*: Identification of the low density lipoprotein receptor-related protein (LRP) as an endocytic receptor for thrombospondin-1. *J Cell Biol*. 1995; 129(5): 1403–10.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 93. **F** Parkyn CJ, Vermeulen EG, Mootoosamy RC, *et al.*: LRP1 controls biosynthetic and endocytic trafficking of neuronal prion protein. *J Cell Sci*. 2008; 121(Pt 6): 773–83.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 94. Nakajima C, Kulik A, Frotscher M, *et al.*: Low density lipoprotein receptor-related protein 1 (LRP1) modulates N-methyl-D-aspartate (NMDA) receptor-dependent intracellular signaling and NMDA-induced regulation of postsynaptic protein complexes. *J Biol Chem*. 2013; 288(30): 21909–23.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 95. Fuentealba RA, Liu Q, Zhang J, *et al.*: Low-density lipoprotein receptor-related protein 1 (LRP1) mediates neuronal Abeta42 uptake and lysosomal trafficking. *PLoS One*. 2010; 5(7): e11884.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 96. Yoon C, Van Niekirk EA, Henry K, *et al.*: Low-density lipoprotein receptor-related protein 1 (LRP1)-dependent cell signaling promotes axonal regeneration. *J Biol Chem*. 2013; 288(37): 26557–68.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 97. Jensen JK, Dolmer K, Schar C, *et al.*: Receptor-associated protein (RAP) has two high-affinity binding sites for the low-density lipoprotein receptor-related protein (LRP): consequences for the chaperone functions of RAP. *Biochem J*. 2009; 421(2): 273–82.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 98. Lee D, Walsh JD, Migliorini M, *et al.*: The structure of receptor-associated protein (RAP). *Protein Sci*. 2007; 16(8): 1628–40.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 99. Willnow TE, Goldstein JL, Orth K, *et al.*: Low density lipoprotein receptor-related protein and gp330 bind similar ligands, including plasminogen activator-inhibitor complexes and lactoferrin, an inhibitor of chylomicron remnant clearance. *J Biol Chem*. 1992; 267(36): 26172–80.
[PubMed Abstract](#)
 100. **F** Missler M, Zhang W, Rohmann A, *et al.*: Alpha-neurexins couple Ca^{2+} channels to synaptic vesicle exocytosis. *Nature*. 2003; 423(6943): 939–48.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 101. Biederer T, Kaeser PS, Blanpied TA: Transcellular Nanoalignment of Synaptic Function. *Neuron*. 2017; 96(3): 680–96.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 102. Berkefeld H, Fakler B, Schulte U: Ca^{2+} -activated K^{+} channels: from protein complexes to function. *Physiol Rev*. 2010; 90(4): 1437–59.
[PubMed Abstract](#) | [Publisher Full Text](#)
 103. **F** Shipton MJ, Tian L: Posttranscriptional and Posttranslational Regulation of BK Channels. *Int Rev Neurobiol*. 2016; 128: 91–126.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 104. Meera P, Wallner M, Song M, *et al.*: Large conductance voltage- and calcium-dependent K^{+} channel, a distinct member of voltage-dependent ion channels with seven N-terminal transmembrane segments (S0-S6), an extracellular N terminus, and an intracellular (S9-S10) C terminus. *Proc Natl Acad Sci U S A*. 1997; 94(25): 14066–71.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 105. Yan J, Olsen JV, Park KS, *et al.*: Profiling the phospho-status of the BK_{Ca} channel alpha subunit in rat brain reveals unexpected patterns and complexity. *Mol Cell Proteomics*. 2008; 7(11): 2188–98.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 106. **F** Berkefeld H, Sailer CA, Bildl W, *et al.*: BK_{Ca} -Cav channel complexes mediate rapid and localized Ca^{2+} -activated K^{+} signaling. *Science*. 2006; 314(5799): 615–20.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 107. Wallner M, Meera P, Toro L: Determinant for beta-subunit regulation in high-conductance voltage-activated and Ca^{2+} -sensitive K^{+} channels: an additional transmembrane region at the N terminus. *Proc Natl Acad Sci U S A*. 1996; 93(25): 14922–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 108. **F** Lorca RA, Ma X, England SK: The unique N-terminal sequence of the BK_{Ca} channel α -subunit determines its modulation by β -subunits. *PLoS One*. 2017; 12(7): e0182068.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 109. Chen L, Bi D, Tian L, *et al.*: Palmitoylation of the $\beta 4$ -subunit regulates surface expression of large conductance calcium-activated potassium channel splice variants. *J Biol Chem*. 2013; 288(18): 13136–44.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 110. Li Q, Fan F, Kwak HR, *et al.*: Molecular basis for differential modulation of BK channel voltage-dependent gating by auxiliary γ subunits. *J Gen Physiol*. 2015; 145(6): 543–54.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 111. Chen SR, Cai YQ, Pan HL: Plasticity and emerging role of BK_{Ca} channels in nociceptive control in neuropathic pain. *J Neurochem*. 2009; 110(1): 352–62.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 112. Wulf MA, Senatore A, Aguzzi A: The biological function of the cellular prion protein: an update. *BMC Biol*. 2017; 15(1): 34.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 113. Resovi A, Pinessi D, Chiorino G, *et al.*: Current understanding of the thrombospondin-1 interactome. *Matrix Biol*. 2014; 37: 83–91.
[PubMed Abstract](#) | [Publisher Full Text](#)
 114. Adams JC, Lawler J: The thrombospondins. *Cold Spring Harb Perspect Biol*. 2011; 3(10): a009712.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 115. Carlson CB, Lawler J, Mosher DF: Structures of thrombospondins. *Cell Mol Life Sci*. 2008; 65(5): 672–86.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 116. Kazeroonian S, Yee KO, Lawler J: Thrombospondins in cancer. *Cell Mol Life Sci*. 2008; 65(5): 700–12.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 117. Arber S, Caroni P: Thrombospondin-4, an extracellular matrix protein expressed in the developing and adult nervous system promotes neurite outgrowth. *J Cell Biol*. 1995; 131(4): 1083–94.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 118. **F** Christopherson KS, Ullian EM, Stokes CC, *et al.*: Thrombospondins are astrocyte-secreted proteins that promote CNS synaptogenesis. *Cell*. 2005; 120(3): 421–33.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 119. Pan B, Yu H, Park J, *et al.*: Painful nerve injury upregulates thrombospondin-4 expression in dorsal root ganglia. *J Neurosci Res*. 2015; 93(3): 443–53.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 120. Yu YP, Gong N, Kweon TD, *et al.*: Gabapentin prevents synaptogenesis between sensory and spinal cord neurons induced by thrombospondin-4 acting on pre-synaptic $\text{Ca}_v 2.2$ subunits and involving T-type Ca^{2+} channels. *Br J Pharmacol*. 2018; 175(12): 2348–61.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 121. Lana B, Page KM, Kadurin I, *et al.*: Thrombospondin-4 reduces binding affinity of [^3H]-gabapentin to calcium-channel $\alpha_2\delta_1$ -subunit but does not interact with $\alpha_2\delta_1$ on the cell-surface when co-expressed. *Sci Rep*. 2016; 6: 24531.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 122. Ly CV, Yao CK, Verstreken P, *et al.*: *straightjacket* is required for the synaptic stabilization of *cacophony*, a voltage-gated calcium channel alpha1 subunit. *J Cell Biol*. 2008; 181(1): 157–70.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 123. **F** Kurshan PT, Oztan A, Schwarz TL: Presynaptic $\alpha_2\delta_3$ is required for synaptic morphogenesis independent of its Ca^{2+} -channel functions. *Nat Neurosci*. 2009; 12(11): 1415–23.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)

124. Dickman DK, Kurshan PT, Schwarz TL: **Mutations in a *Drosophila* alpha,delta voltage-gated calcium channel subunit reveal a crucial synaptic function.** *J Neurosci.* 2008; **28**(1): 31–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
125. Hooper NM: **Determination of glycosyl-phosphatidylinositol membrane protein anchorage.** *Proteomics.* 2001; **1**(6): 748–55.
[PubMed Abstract](#) | [Publisher Full Text](#)
126. Ryan TJ, Kopanitsa MV, Indersmitten T, *et al.*: **Evolution of GluN2A/B cytoplasmic domains diversified vertebrate synaptic plasticity and behavior.** *Nat Neurosci.* 2013; **16**(1): 25–32.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
127. Frank RAW, Zhu F, Komiyama NH, *et al.*: **Hierarchical organization and genetically separable subfamilies of PSD95 postsynaptic supercomplexes.** *J Neurochem.* 2017; **142**(4): 504–11.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
128. Frank RA, Komiyama NH, Ryan TJ, *et al.*: **NMDA receptors are selectively partitioned into complexes and supercomplexes during synapse maturation.** *Nat Commun.* 2016; **7**: 11264.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
129. Peng J, Kim MJ, Cheng D, *et al.*: **Semiquantitative proteomic analysis of rat forebrain postsynaptic density fractions by mass spectrometry.** *J Biol Chem.* 2004; **279**(20): 21003–11.
[PubMed Abstract](#) | [Publisher Full Text](#)
130. Nieto-Rostro M, Sandhu G, Bauer CS, *et al.*: **Altered expression of the voltage-gated calcium channel subunit α - δ -1: a comparison between two experimental models of epilepsy and a sensory nerve ligation model of neuropathic pain.** *Neuroscience.* 2014; **283**: 124–37.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
131. Schlick B, Flucher BE, Obermair GJ: **Voltage-activated calcium channel expression profiles in mouse brain and cultured hippocampal neurons.** *Neuroscience.* 2010; **167**(3): 786–98.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
132. Cottrell GS, Soubrane CH, Hounshell JA, *et al.*: **CACHD1 is an α 2 δ -Like Protein That Modulates Ca_v3 Voltage-Gated Calcium Channel Activity.** *J Neurosci.* 2018; **38**(43): 9186–9201.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
133. Dahimene S, Page KM, Kadurin I, *et al.*: **The α 2 δ -like Protein Cachd1 Increases N-type Calcium Currents and Cell Surface Expression and Competes with α 2 δ -1.** *Cell Rep.* 2018; **25**(6): 1610–21.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. Jutta Engel

Department of Biophysics, Center for Integrative Physiology and Molecular Medicine (CIPMM), Saarland University, School of Medicine, Homburg, Germany

Competing Interests: No competing interests were disclosed.

2. Emilio Carbone

Department of Drug Science, Lab of Cellular Physiology and Molecular Neuroscience, Torino, Italy

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research