Check for updates

RESEARCH ARTICLE

Feasibility of incorporating mindfulness based mental health promotion to the pregnancy care program in Sri Lanka: a pilot study [version 1; peer review: 1 approved with reservations]

Thilini Agampodi, Subhashini Katumuluwa, Thulani Pattiyakumbura, Nilupulee Rankaduwa, Thushari Dissanayaka, Suneth Agampodi

Department of Community Medicine, Faculty of Medicine and Allied Sciences, Rajarata University of Sri Lanka, Saliyapura, 50008, Sri Lanka

V1 First published: 27 Nov 2018, 7:1850
<https://doi.org/10.12688/f1000research.17049.1>

Latest published: 04 Jan 2019, 7:1850
<https://doi.org/10.12688/f1000research.17049.2>

Abstract

Background: Though widely discussed, mindfulness-based interventions (MBI) to improve maternal mental health is limited by lack of studies with system incorporation. We evaluate the feasibility of incorporating a MBI program into routine antenatal care (ANC) in Sri Lanka.

Methods: MBI included learning mindfulness concepts, practicing mindfulness sitting/reclining meditation, performing mindful movements and practicing mindfulness in daily life. Feedback from the participants were obtained through an anonymous, self-administered, semi-structured questionnaire to determine the program's cultural appropriateness, usefulness, and feasibility.

Results: Participants reported that the training reduced the stress of their daily life, brought a sense of calmness to their mind and body, and improved their anger management. Participants felt strongly that this training would be very useful and a shortened version be included in the national ANC program.

Conclusions: A systematic assessment of impact of MBI is needed with system incorporation of the suggested programme.

Keywords

Maternal mental health; Mindfulness; Pregnancy; Antenatal care; Sri Lanka

Open Peer Review

Approval Status

	1	2
version 2		
(revision)
04 Jan 2019	view	view

version 1		
27 Nov 2018	view	

1. **Moraendage Wasantha Gunathunga**,
 University of Colombo, Colombo, Sri Lanka
Aruni Gallage, University of Colombo,
 Colombo, Sri Lanka

2. **Alys Einion-Waller** , Swansea University,
 Swansea, UK

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Thilini Agampodi (thilnichanchala@yahoo.com)

Author roles: **Agampodi T:** Conceptualization, Formal Analysis, Funding Acquisition, Investigation, Methodology, Project Administration, Resources, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing; **Katumuluwa S:** Data Curation, Formal Analysis, Investigation, Project Administration, Resources, Writing – Original Draft Preparation; **Pattiyakumbura T:** Data Curation, Formal Analysis, Investigation, Writing – Original Draft Preparation; **Rankaduwa N:** Data Curation, Investigation, Methodology, Writing – Original Draft Preparation; **Dissanayaka T:** Data Curation, Formal Analysis, Validation, Writing – Original Draft Preparation; **Agampodi S:** Conceptualization, Formal Analysis, Funding Acquisition, Methodology, Project Administration, Supervision, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This study was funded by the University grants of Rajarata University of Sri Lanka under the grant number RJT/RPandHDC/2015/FMAS/R/06.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2018 Agampodi T *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Agampodi T, Katumuluwa S, Pattiyakumbura T *et al.* **Feasibility of incorporating mindfulness based mental health promotion to the pregnancy care program in Sri Lanka: a pilot study [version 1; peer review: 1 approved with reservations]** F1000Research 2018, 7:1850 <https://doi.org/10.12688/f1000research.17049.1>

First published: 27 Nov 2018, 7:1850 <https://doi.org/10.12688/f1000research.17049.1>

Introduction

Pregnancy and childbirth represent a time of increased vulnerability, during which a woman is exposed to many physiological and psychosocial changes. This puts pregnant and postpartum women at increased risk of mental health problems. The prevalence of mental health disorders in pregnancy and postpartum period is much greater in low-income countries (LIC) and lower-middle-income countries (LMIC) (Fisher *et al.*, 2012) than in high income countries (HIC) (Hendrick *et al.*, 1998). Of the mental health problems experienced by pregnant and postpartum women, antenatal depression, anxiety, and post-partum depression are the most common (O'Hara & Swain, 1996). These mental health issues may lead to adverse pregnancy and neonatal outcomes such as small for gestational age baby (Dejin-Karlsson *et al.*, 2000), developmental delays (Bernard-Bonnin, 2004), poor mother-infant interaction (Cohn & Tronick, 1989), negative affect in the infant (Tronick & Reck, 2009), problems with cognitive development (Singer & Fagen, 1992) and affective disorders / attention-deficit/hyperactivity disorder (ADHD) in children (Lesesne *et al.*, 2003). This makes the antenatal period a critical time point for intervention on maternal mental health.

Despite having a strong, community-based public health care system, addressing maternal mental health has yet to gain the spotlight in Sri Lanka. The prevalence estimate for antenatal depression in Anuradhapura District, Sri Lanka is 16% (Agampodi & Agampodi, 2013) and the national estimate for postpartum depression is as high as 27% (Agampodi *et al.*, 2011). Furthermore, a secondary analysis of maternal death investigation reports from 2005–2011 in the North Central Province revealed that suicide was the leading cause of maternal mortality in the province, with 17.8% of maternal deaths attributed to suicide (Agampodi *et al.*, 2014). These findings show the urgency of addressing the mental health of pregnant women in Sri Lanka.

Mindfulness is an emerging concept in mental health promotion. It was the original healing method used for centuries in ancient cultures in Asia. Clinical use of mindfulness was widely discussed in late 20th century (Baer, 2003; Deatherage, 1975). Several recent studies have shown that mindfulness-based interventions for pregnant women have been effective in increasing positive affect, decreasing negative affect (Duncan & Bardacke, 2010; Vieten & Astin, 2008a), decreasing anxiety (Duncan & Bardacke, 2010; Dunn *et al.*, 2012; Vieten & Astin, 2008a), decreasing depression (Duncan & Bardacke, 2010; Dunn *et al.*, 2012), and decreasing stress (Dunn *et al.*, 2012) during pregnancy. These benefits were seen to extend into the postpartum period in some cases (Dunn *et al.*, 2012; Tomfohr-Madsen *et al.*, 2016; Vieten & Astin, 2008a). Nevertheless, two recent systematic reviews (Dhillon *et al.*, 2017; Roy Malis *et al.*, 2017) shows mixed results on mindfulness based interventions in pregnancy. One of these reviews found that “anxiety, depression and perceived stress indicated no differences between the mindfulness intervention group and the control group” in pooled results of RCTs. However, pooled results of non-RCTs showed a “significant benefit for the mindfulness group”. Other review also showed similar results with probable effect on maternal anxiety. Both reviews suggested that more work is needed on this area of research

to produce evidence-based guidelines. However, studies on integrating the mindfulness-based activities to routine program to improve maternal and newborn health outcomes are scarce. We report here a feasibility study on developing and incorporating a mental health promotion program based on mindfulness for pregnant women of Sri Lanka.

Methods

Study location

This preliminary study was carried out in Anuradhapura, Sri Lanka. The study was planned as a part of field healthcare delivery in the field practice area of the Department of Community Medicine, Faculty of Medicine and Allied Sciences, Rajarata University of Sri Lanka. In the first phase of this study, we translated and culturally adapted the Mindful Motherhood training curriculum and in the second phase, feasibility testing was done to see whether it is possible to incorporate this training to the public health system. Both phases of the study were carried out from May 2015 to April 2016.

Phase I (May–July 2016)

Phase I of this study was to develop a culturally acceptable mindful motherhood training curriculum. We used the “Mindful Motherhood” training curriculum (Vieten, 2009), published by the Institute of Noetic Sciences (IONS) and made available for professional use for our training programme. The general structure and content of the IONS training was left the same; however, some of the reading materials and activities were modified to be more relevant and acceptable to a Sri Lankan population. In addition, supplemental readings were added from other sources. In the Sri Lankan community, “loving kind meditation” is a well-established practice. We included supplementary reading and practice component of loving kind meditation in to all eight sessions. The sources used were simple Buddhist teaching, without addition religious components. The changes were done after extensive discussions among three community physicians, a public health physician, maternal health service providers at divisional and grassroots level and medical officers. All supplemental readings were translated from English to Sinhalese, the main language spoken in Sri Lanka. Translations were performed by two medical graduates, who were fluent in both languages as well as mindfulness teaching. Tamil translations were not made at this time, as the majority population of the Nuwera Palath Central (NPC) region, where the study was conducted, speaks Sinhalese. Back translation was not done, as the procedure was rather a cultural adaptation than tool translation. However, three investigators individually looked at the translations and discussed and finalized the materials.

The aim of our programme was to introduce the concept of mindfulness and develop the skills of the pregnant mothers to be mindful in day-to-day life, with the ultimate goal of promoting mental well-being. During the programme development, four authors (T.A., S.K., T.P. and N.R.) who did the training, had undergone mindful training. Three of the authors involved in the training (T.A., S.K. and N.R.) were regular practitioners of Buddhist mindfulness in daily life. Our curriculum consisted of eight, 2- to 3-hour weekly training sessions. Each session consisted

of several components; introduction of a short mindfulness concept; practice of mindfulness sitting/reclining meditation (i.e. breathing meditation, meditation with focus on thoughts/feelings/sensations, and several types of body scans); performing mindful movement (i.e. prenatal yoga, mindful walking, simple mindful movement series); and sometimes a mindfulness in daily life activity or discussion (i.e. mindful eating, how to complete daily tasks mindfully, mindfulness in relationships). In addition to the weekly sessions, each week the women were provided with readings summarizing or related to the week's mindfulness concepts. The initial 8-week curriculum, supplementary materials, and all training related documents are publically available on OSF (Agampodi *et al.*, 2018a).

Phase II (August 2015–April 2016)

In the second phase, we field-tested the feasibility of incorporating a mindfulness-training program into routine antenatal care (ANC) in Sri Lanka. For the intervention, we recruited pregnant women at less than 32 weeks gestational age, and who were able to read and write Sinhalese, from the NPC region. Those who were not permanent residents and planning to leave the area for delivery (in Sri Lanka, some females go to their parents' place for delivery) were excluded from the study. Participants were recruited at the antenatal clinics. We also included the Public Health Midwife (PHM), the grassroots level healthcare provider for pregnant women in the program. The PHM was included because she has extensive experience working with pregnant women in rural areas and has a unique understanding of their needs and perspectives on a range of pregnancy-related issues, providing valuable information regarding the feasibility of our training. In addition, if any intervention to be included into the routine public health system in Sri Lanka, it needs to be carried out by the PHM. The PHM selected the participants for the study purposefully, based on interest and availability to participate. Pregnant women with a history of mental disorder with a psychotic component were excluded from the study. The sample size was predetermined (not calculated) as 12–15 participants, to ensure an adequate level of interaction between the participants and instructors and based on feasibility to visit the mindfulness-training center at Faculty of Medicine and Allied Sciences.

In addition to attending the weekly training sessions developed in phase 1, as a homework assignment, women were asked to practice each of the mindfulness practices learned in the session at home. The women recorded their practice on a weekly record sheet and handed it in the following week. They were also asked to keep a diary and write any thoughts or feelings about the sessions and how they felt during the week while practicing. After seven weeks of the program, participants were given an anonymous, detailed, self-administered, semi-structured questionnaire to obtain their feedback. The questionnaire sought to gain the participants' detailed insights into how the program affected their lives (if at all), the most useful aspects of the training, their suggestions for improvement, and whether and how a similar training should be incorporated into the current ANC program. We ask about acceptability of the programme and also feasibility of practicing it at home. No maternal mental

health outcome variables were evaluated in this pilot, as it was only a feasibility study.

To evaluate the feasibility and to triangulate data for future system incorporation, facilitators of the training also recorded their thoughts on the program content, structure, and the logistics of conducting. The self-reported descriptions were collected separately to analyses and thematic analysis was done for these feedbacks as well.

Ethical approval

Ethical clearance was obtained from the Ethics Review Committee, Faculty of Medicine and Allied Sciences, Rajarata University of Sri Lanka (ERC/2015/15). Written informed consent was obtained from all the participants included in the study.

Results

In total, 12 women were recruited and participated in the training. The women ranged in age from 18 to 30 years, and were between 10 and 27 weeks gestational age. After completion of the program, participants described many benefits they received through the training. A total of eight participants completed the diaries and the feedback.

Perceived psychological benefits

Daily life stress

Most of the women (7/8) reported that the training reduced daily life stressors and brought a sense of calmness to their mind. All the women stated that participating in the training had also improved the way they responded to stressful situations.

"I was bored and stressed with the daily routine. With the training I was able to face day-to-day problems in a calm mindful manner." 22-year-old pregnant female.

Improvement in personal communication

The women reported improvements in interpersonal communication. The training had helped them to improve the way they interacted with their spouse and/or current child(ren), or that their view of how best to interact with their yet unborn child had changed based on what they had learned.

"When my elder child was stubborn, I was able to modestly tackle it. Even when other people came to argue, I was able to face them with patience. It was successful." 30-year-old pregnant female.

Controlling emotions

Some women also reported that the training helped them to control their emotions, with several (3 of 8) stating that they were better able to control their anger.

"My elder child had a fall and was injured. All the family members were panicked. I was able to remain calm and do the needful. I was not like this before. This was a new experience for me. I acquired this skill from the training." 28-year-old pregnant female.

Improving quality of work and life

The women told that they were able to do their duties at home and work place in a more efficient manner than in their past.

“Now I’m able to do each and every task with more understanding. This gives me clarity and calmness in mind. I feel less exhausted. I’m able to sleep well.” 18-year-old pregnant female.

Perceived physical benefits

In addition to the mental and emotional benefits, all the women reported physical benefits, including feeling a sense of comfort and relaxation for the body when practicing the mindful movement at home. One of the women reported successfully using several of the meditation and body scan techniques help her cope with an experience of abdominal pain, stating,

“when I fell ill and I started practicing the things we learned, putting them into action one by one, the body which was full of pain became normal in an unbelievable way.” 25-year-old pregnant female.

Every participant reported that they intended to continue the daily practice of various techniques to cultivate mindfulness, learned during the training.

Evaluation of content

Participants evaluated the different components or strategies used to cultivate mindfulness during this training. The mindful movement series was the most popular strategy among women. Of the mindfulness concepts taught, the women found the topics of “introduction to mindfulness,” “staying in the present moment,” “acceptance,” and “avoiding controlling and judging” to be the most interesting and helpful in their daily lives. Of the sitting/lying down meditations, women enjoyed and perceived the most usefulness from the body scan, loving kindness meditation, and mindful breathing practice. Everyone reported that they found the “Mindful Mother Check-in” cards (cards given to them at the beginning of the training with simple steps to bring the mind to the present moment applying the practices they had learned), which they posted around their home, to be helpful to calm their mind, particularly during stressful situations. Of the mindful movements taught, the majority of the women (6 of 8) enjoyed the short mindful movement series the most.

In addition to participant feedback, the facilitators of the training also recorded their thoughts on the program content, structure, and the logistics of conducting it. We experienced that the depth of the material being allocated for each session was advance, so that it cannot be grasped with just a simple explanation for a participant who missed a session. For this same reason, we found it a challenge to complete all the material in the allotted 2-hour time, often running over. Additionally, the women seldom asked questions about the material (which may be a cultural factor) or for clarification during the sessions.

Feasibility and timing

Each participant felt time allocated for each of the training segments was appropriate, except for one who felt it would

be better if more time was given for the mindful movement section. Although we took into consideration the demands on the average pregnant woman in our area when deciding a meeting time, we still found that it was very difficult to get the women to come consistently to every session. This is likely due to several cultural factors. In Sri Lanka, many women in more rural areas are stay-at-home wives/mothers, taking care of the household and the children. So unless there is someone at home, usually a relative, who can watch the children or pick them up from school while the mother is out, she may not be able to come, may come late or need to leave early, which we experienced on multiple occasions. All of the women also had to take public transportation to get to the trainings, which added to the challenge.

Feasibility of incorporation into ANC system

Women were asked their thoughts on the potential benefit of incorporating a mindfulness training into routine ANC in Sri Lanka. All the participants felt that this training would be very useful and important for other pregnant women. While agreeing with this, one woman raised the concern of whether one two-hour session every three months would provide sufficient time and guidance for women to fully understand the meaning and feel the value of the concepts taught. It was also suggested that it would be beneficial to have the husbands of the pregnant women involved in the program as well. Overall, all the participants, including the PHM, felt that it would be beneficial and feasible to incorporate a mindfulness program into the national ANC system if the current curriculum was shortened and included in the once per trimester antenatal sessions that are already part of the ANC system in Sri Lanka.

Based on Phase I and Phase II of this feasibility assessment, we prepared the final version of intervention. The final structure of the mindfulness based intervention to improve mental health of pregnant mothers is illustrated in [Table 1](#).

Discussion

This preliminary study shows that a incorporating a mindfulness based mental health program for pregnant mothers to the pregnancy care program in Sri Lanka is feasible and culturally acceptable. It also suggests that learning mindfulness concepts and the techniques to cultivate mindfulness may improve a woman’s ability to cope with stress during her pregnancy. Important future steps should include efforts to capture the latter outcome quantitatively, as has been done in prior pilot studies ([Duncan & Bardacke, 2010](#); [Dunn et al., 2012](#); [Vieten & Astin, 2008a](#)).

The program developed in this pilot study was based on the standard 8-week mindfulness intervention used in many other previous studies ([Astin et al., 2003](#))([Byrne et al., 2014](#))([Dunn et al., 2012](#))([Vieten & Astin, 2008b](#)). However, a brief mindfulness interventions even for a 3-week duration has proven the potential usefulness of mindfulness interventions ([Beattie et al., 2017](#); [Matvienko-Sikar & Dockray, 2017](#)). Although there are number of studies on mindfulness interventions, there is not enough evidence found about the most effective duration for a

Table 1. Proposed 8-weeks mindfulness-based training program to be tested in a larger study in Sri Lanka for pregnant women.

Session	Theme	Activities	Duration
Day 1	Introduction to mindful motherhood training	Introduction to mindfulness and mindful motherhood training	30 minutes
		Mindful daily awareness using mindful eating of a simple food	15 minutes
		Introduce mindful movement	15 minutes
		Introduce mindful sitting meditation and awareness in breathing	15 minutes
		Loving kindness meditation	15 minutes
Day 2	Control vs mindfulness, what is mindfulness? Acceptance/Willingness	Mindful movements	20 minutes
		Recap the purpose of mindful awareness Explain further mindfulness	20 minutes
		Concept: acceptance	15 minutes
		Mindful awareness of breathing	15 minutes
		Introduce mindful mothers' check-in	10 minutes
Day 3	The observing self	Loving kindness meditation	15 minutes
		Mindful movement practice	20 minutes
		Mindful mothers' check-in	10 minutes
		Concept: observing self	30 minutes
		Mindful awareness practice (sitting) -body scan	20 minutes
Day 4	Train of thoughts Metacognition	Loving kindness meditation	15 minutes
		Mindful walking exercise	20 minutes
		Mindful mothers' check-in	10 minutes
		Concept: Train of thoughts	30 minutes
		Observing the activities of mind	10 minutes
Day 5	Present movement focus	Observing thoughts	20 minutes
		Loving kindness meditation	15 minutes
		Mindful movement and body scan	30 minutes
		Mindful mothers' check-in	10 minutes
		Concept: present movement focus	30 minutes
Day 6	Additional qualities of mindfulness	Mindful awareness of present movement sitting meditation	20 minutes
		Loving kindness meditation	15 minutes
		Exercise to identify mindful/unmindful moments	20 minutes
		Concept: qualities of mindfulness (3 new qualities) nonstriving, beginners mind, being vs. doing	30 minutes
		Mindful movement – walking around the place	20 minutes
Day 7	Self-compassion	Mindful awareness of thoughts, feelings and body sensation	20 minutes
		Loving kindness meditation	15 minutes
		Mindful movement and body scan	20 minutes
		Mindful mothers' check-in	10 minutes
		Concept: Self-compassion	30 minutes
Day 8	Mindful awareness in everyday life Mindfulness in childbirth	Self-compassion – practical exercises and discussion	30 minutes
		Refresh concepts learnt so far	15 minutes
		Loving kindness meditation	15 minutes
		Mindful movement	15 minutes
		Concept: wise action and intention	30 minutes
		Quick awareness and action exercise	15 minutes
		Mindful decision making	20 minutes
		Mindful awareness as a refuge and source of strength	15 minutes
		Mindful mom's promise	10 minutes
		Mindful motherhood as a whole	15 minutes

mindfulness program. Thus, further studies has to be conducted in the future to assess the effective time duration for mindfulness interventions. Taking into account the challenges the facilitators experienced during the intervention, trying to cover all the material in a comprehensible manner within the allotted 16-hour time period of the training may not be feasible. Careful consideration will be needed to select some of these concepts that could practically be covered during the three ANC sessions, and would likely be of greatest benefit and/or interest to the pregnant women. The feedback from our pilot study participants will be valuable when making these decisions.

Given the positive response from our small feasibility pilot, the next steps will include using the participants' feedback to create a modified, shortened mindfulness training curriculum for pregnant women, and running a larger field study using this revised curriculum. Maternal mental health outcomes should also be evaluated using pre- and post-intervention questionnaires that have been validated for use in Sri Lankan populations. This will be an important step in establishing any effect that the mindfulness training may have on maternal mental health-related outcomes, including distress, depression, and coping skills. The possibility of any benefits of the training extending into the postpartum period, as suggested by pilot studies (Roy Malis *et al.*, 2017; Vieten & Astin, 2008a) should also be evaluated. The degree of success of launching such a program nationally, or even locally, will depend heavily on the willingness and enthusiasm of the PHMs, as eventually, if the program is to be sustainable, it will be trained PHMs who will be leading each of the mindfulness sessions. To date, the support from the local PHM community and Medical Officer of Health (MOH) has been strong for this program. When presented an overview of the program and the evidence behind the teachings, many of the PHMs were visibly enthusiastic about the benefits such a program could have for their patients and were eager to offer their support.

Limitations of our study include the small sample size, the inconsistent attendance of some of the participants, and the sample of

only Sinhalese-speaking women. To serve both the Sinhalese and Tamil populations in Sri Lanka, the curriculum will also need to be translated and tested in a Tamil-speaking population.

Conclusion

This pilot study suggests that a mindfulness-training program is a culturally appropriate and feasible way to incorporate promotion of maternal mental health into routine ANC in Sri Lanka. Further research must investigate the effectiveness of a shortened, modified program on impacting maternal mental health-related outcomes during pregnancy and beyond.

Data availability

Underlying data

Raw data from the participants' answers to the questionnaires are available on OSF. Please note that data are provided in Sinhalese. DOI: <https://doi.org/10.17605/OSF.IO/QFA6X> (Agampodi *et al.*, 2018b).

Extended data

All documents related to the project, including the initial proposal, ERC certificate, questionnaire (with translation) and training material are available on OSF. DOI: <https://doi.org/10.17605/OSF.IO/45HNR> (Agampodi *et al.*, 2018a).

All data are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

Grant information

This study was funded by the University grants of Rajarata University of Sri Lanka under the grant number RJT/RP and HDC/2015/FMAS/R/06.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- Agampodi SB, Agampodi TC: **Antenatal depression in Anuradhapura, Sri Lanka and the factor structure of the Sinhalese version of Edinburgh post partum depression scale among pregnant women.** *PLoS One*. 2013; 8(7): e69708. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Agampodi SB, Agampodi TC, Katumuluwa S: **Mindful Motherhood Training Materials [WWW Document]**. 2018a. <http://www.doi.org/10.17605/OSF.IO/45HNR>
- Agampodi SB, Agampodi TC, Katumuluwa S: **Mindful Motherhood Data [WWW Document]**. 2018b. <http://www.doi.org/10.17605/OSF.IO/QFA6X>
- Agampodi TC, Agampodi SB, Wickramasinghe WA, *et al.*: **Post partum depression - a problem that needs urgent attention.** *Ceylon Med J*. 2011; 56(4): 183–4. [PubMed Abstract](#) | [Publisher Full Text](#)
- Agampodi S, Wickramage K, Agampodi T, *et al.*: **Maternal mortality revisited: the application of the new ICD-MM classification system in reference to maternal deaths in Sri Lanka.** *Reprod Health*. 2014; 11(1): 17. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Astin JA, Berman BM, Bausell B, *et al.*: **The efficacy of mindfulness meditation plus Qigong movement therapy in the treatment of fibromyalgia: a randomized controlled trial.** *J Rheumatol*. 2003; 30(10): 2257–2262. [PubMed Abstract](#)
- Baer RA: **Mindfulness training as a clinical intervention: A conceptual and empirical review.** *Clin Psychol-Sci Pr*. 2003; 10: 125–143. [Publisher Full Text](#)
- Beattie J, Hall H, Biro MA, *et al.*: **Effects of mindfulness on maternal stress, depressive symptoms and awareness of present moment experience: A pilot randomised trial.** *Midwifery*. 2017; 50: 174–183. [PubMed Abstract](#) | [Publisher Full Text](#)
- Bernard-Bonnin AC: **Maternal depression and child development.** *Paediatr Child Health*. 2004; 9(8): 575–598. [PubMed Abstract](#) | [Free Full Text](#)
- Byrne J, Hauck Y, Fisher C, *et al.*: **Effectiveness of a Mindfulness-Based Childbirth Education pilot study on maternal self-efficacy and fear of childbirth.** *J Midwifery Womens Health*. 2014; 59(2): 192–197. [PubMed Abstract](#) | [Publisher Full Text](#)
- Cohn JF, Tronick E: **Specificity of infants' response to mothers' affective**

behavior. *J Am Acad Child Adolesc Psychiatry*. 1989; **28**(2): 242–8.

[PubMed Abstract](#) | [Publisher Full Text](#)

Deatherage G: **The Clinical Use of “Mindfulness” Meditation Techniques in Short-Term Psychotherapy**. *Journal of Transpersonal Psychology*. 1975; **7**: 133–143.

[Reference Source](#)

Dejin-Karlsson E, Hanson BS, Ostergren PO, *et al.*: **Association of a lack of psychosocial resources and the risk of giving birth to small for gestational age infants: a stress hypothesis**. *BJOG*. 2000; **107**(1): 89–100.

[PubMed Abstract](#) | [Publisher Full Text](#)

Dhillon A, Sparkes E, Duarte RV: **Mindfulness-Based Interventions During Pregnancy: a Systematic Review and Meta-analysis**. *Mindfulness (N Y)*. 2017; **8**(6): 1421–1437.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Duncan LG, Bardacke N: **Mindfulness-Based Childbirth and Parenting Education: Promoting Family Mindfulness During the Perinatal Period**. *J Child Fam Stud*. 2010; **19**(2): 190–202.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Dunn C, Hanieh E, Roberts R, *et al.*: **Mindful pregnancy and childbirth: effects of a mindfulness-based intervention on women’s psychological distress and well-being in the perinatal period**. *Arch Womens Ment Health*. 2012; **15**(2): 139–143.

[PubMed Abstract](#) | [Publisher Full Text](#)

Fisher J, Cabral de Mello M, Patel V, *et al.*: **WHO | Prevalence and determinants of common perinatal mental disorders in women in low- and lower-middle-income countries: a systematic review**. *Bull World Health Organ*. 2012; **90**(2): 139–149H.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Hendrick V, Altshuler L, Cohen L, *et al.*: **Evaluation of mental health and depression during pregnancy: position paper**. *Psychopharmacol Bull*. 1998; **34**(3): 297–9.

[PubMed Abstract](#)

Lesesne CA, Visser SN, White CP: **Attention-deficit/hyperactivity disorder in**

school-aged children: association with maternal mental health and use of health care resources. *Pediatrics*. 2003; **111**(5 Pt 2): 1232–1237.

[PubMed Abstract](#)

Matvienko-Sikar K, Dockray S: **Effects of a novel positive psychological intervention on prenatal stress and well-being: A pilot randomised controlled trial**. *Women Birth*. 2017; **30**(2): e111–e118.

[PubMed Abstract](#) | [Publisher Full Text](#)

O’hara MW, Swain AM: **Rates and risk of postpartum depression—a meta-analysis**. *Int Rev Psychiatry*. 1996; **8**: 37–54.

[Publisher Full Text](#)

Roy Malis F, Meyer T, Gross MM: **Effects of an antenatal mindfulness-based childbirth and parenting programme on the postpartum experiences of mothers: a qualitative interview study**. *BMC Pregnancy Childbirth*. 2017; **17**(1): 57.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Singer J, Fagen J: **Negative affect, emotional expression, and forgetting in young infants**. *Dev Psychol*. 1992; **28**: 48–57.

[Publisher Full Text](#)

Tomfohr-Madsen LM, Campbell TS, Giesbrecht GF, *et al.*: **Mindfulness-based cognitive therapy for psychological distress in pregnancy: study protocol for a randomized controlled trial**. *Trials*. 2016; **17**(1): 498.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Tronick E, Reck C: **Infants of depressed mothers**. *Harv Rev Psychiatry*. 2009; **17**(2): 147–56.

[PubMed Abstract](#) | [Publisher Full Text](#)

Vieten C: **Mindful Motherhood Training: Guidelines for Facilitators**. 2009.

Vieten C, Astin J: **Effects of a mindfulness-based intervention during pregnancy on prenatal stress and mood: results of a pilot study**. *Arch Womens Ment Health*. 2008a; **11**(1): 67–74.

[PubMed Abstract](#) | [Publisher Full Text](#)

Vieten C, Astin J: **Effects of a mindfulness-based intervention during pregnancy on prenatal stress and mood: results of a pilot study**. *Arch Womens Ment Health*. 2008b; **11**(1): 67–74.

[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status: ?

Version 1

Reviewer Report 17 December 2018

<https://doi.org/10.5256/f1000research.18639.r41129>

© 2018 Gunathunga M et al. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Moraendage Wasantha Gunathunga

Department of Community Medicine, Faculty of Medicine, University of Colombo, Colombo, Sri Lanka

Aruni Gallage

Department of Community Medicine, Faculty of Medicine, University of Colombo, Colombo, Sri Lanka

Maternal health is an important concern in Sri Lanka as in many other countries. Though maternal well being is addressed with vigor in Sri Lanka mental well being is not equally addressed. Hence, exploring possibilities of incorporating mindfulness into maternal care services is of much importance and has the potential to contribute significantly to scientific literature. It is commendable that authors embarked on this much needed research.

In phase one, what the authors mean by "Buddhist teaching, without addition religious components" is unclear. If any use can be obtained, for health benefits from any religious material we do not see any reason of not using it and acknowledging it.

Among those who took part in extensive discussions no mention is made about experience in actually practicing mindfulness. The fact that three researchers have been trained in mindfulness and are regular practitioners is noteworthy. This is essential as mindfulness involves a phenomenological component. Without substantial direct experience researchers will not understand certain components of behaviours related to mindfulness. It is also required to mention whether the researchers are capable of dealing with advanced practitioners' queries or such services are made available when needed. This is important as all mothers who start mindfulness practice are not at the same baseline and progress is not uniform. They can be expected to behave very close to a normal distribution where some progress very fast and needing advanced help.

In phase two "Pregnant women with a history of mental disorder with a psychotic component were excluded from the study". It is unclear why only psychotic component was excluded. It would have been more appropriate to exclude those who have a known psychiatric illness in this phase.

It has been mentioned that mental health variables were not assessed. However, perceived psychological benefits were assessed and positive. It is too early to count on these as the benefits cannot be directly attributed to the mindfulness practice in this design. This association, though, is noteworthy.

The sample size of 12 is inadequate to have a sufficient idea on the feasibility of the intervention due to the wide variation in the variable related to feasibility.

As mentioned in "Feasibility of incorporating into Antenatal Care(ANC) system" Participants opinion regarding incorporating this intervention to national antenatal care in Sri Lanka has to be taken with reservation as they do not see the operational aspects of such a program when it is scaled up. In the responses there is also an element of social desirability bias.

"The degree of success of launching such a program nationally, or even locally, will depend heavily on the willingness and enthusiasm of the Public Health Midwife(PHM)s, as eventually, if the program is to be sustainable, it will be trained PHMs who will be leading each of the mindfulness session". This is asking too much from already over burdened PHM in Sri Lanka. Our experience is that, out of the numbers who take up mindfulness training, only a small proportion will qualify to be trainers. Hence we consider planning to train PHMs as trainers is a weakness in the design.

As mentioned in the discussion inconsistent attendance can be a difficult challenge to meet as take-up of this kind of interventions by different individuals vary widely.

It is still premature to conclude "...a feasible way to incorporate promotion of maternal mental health into routine ANC in Sri Lanka". Though it appears feasible to implement the intervention whether it improves mental well being as an outcome has not been assessed in this work. Hence this claim is not justified.

In the abstract, there is no conclusion in relation to the set objective of evaluating feasibility of intervention.

Overall, considering the dearth of literature in this area of research and the importance in relation to promotion of maternal mental well being this work warrants indexing with modifications.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Partly

Are sufficient details of methods and analysis provided to allow replication by others?

Partly

If applicable, is the statistical analysis and its interpretation appropriate?

Not applicable

Are all the source data underlying the results available to ensure full reproducibility?

No source data required

Are the conclusions drawn adequately supported by the results?

No

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Prof. Wasantha Gunathunga: Body mind and consciousness, mental well being, happiness, mindfulness

We confirm that we have read this submission and believe that we have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however we have significant reservations, as outlined above.

Author Response 29 Dec 2018

Suneth Agampodi, Rajarata University of Sri Lanka, Saliyapura, Sri Lanka

Reply to reviewer's comments

We would like to appreciate the comments by the reviewer and it definitely helped us to enhance the quality of this manuscript. Based on the points raised by the reviewer, we have revised the manuscript. The point by point explanations for reviewers comments are below.

In phase one, what the authors mean by “Buddhist teaching, without addition religious components” is unclear. If any use can be obtained, for health benefits from any religious material we do not see any reason of not using it and acknowledging it.

In the phase one we had several meetings with stakeholders, before starting the project. Service providers had a concern that this intervention looks as a Buddhist intervention. To overcome this barrier, we mentioned the term “mindfulness” though the teaching and training were obtained from Buddhist practices. We acknowledge the materials in the article, but not during the training.

Among those who took part in extensive discussions no mention is made about experience in actually practicing mindfulness. The fact that three researchers have been trained in mindfulness and are regular practitioners is noteworthy. This is essential as mindfulness involves a phenomenological component. Without substantial direct experience researchers will not understand certain components of behaviours related to mindfulness. It is also required to mention whether the researchers are capable of dealing with advanced practitioners' queries or such services are made available when needed. This is important as all mothers who start mindfulness practice are not at the same baseline and progress is not uniform. They can be expected to behave very close to a normal distribution where some progress very fast and needing advanced help.

Of the six authors, three were practicing mindfulness for a long period. However, it is difficult to reply to this comment about needing advanced help. The purpose of this training

was to improve day to day mindful practices and to help the pregnant mothers to practice it up to a certain extent. It was neither our intention to go up to the advance mindful practices nor in assisting pregnant mothers who are already having such practices, because it was beyond any routine intervention.

In phase two “Pregnant women with a history of mental disorder with a psychotic component were excluded from the study”. It is unclear why only psychotic component was excluded. It would have been more appropriate to exclude those who have a known psychiatric illness in this phase.

In the research protocol, we had this as an exclusion criteria, but none of the selected pregnant mothers had psychiatric illnesses. We revised the manuscript to explain this.

It has been mentioned that mental health variables were not assessed. However, perceived psychological benefits were assessed and positive. It is too early to count on these as the benefits cannot be directly attributed to the mindfulness practice in this design. This association, though, is noteworthy.

We fully agree with this comment. We got a qualitative feedback and a proper trial is required to comment on the benefits. We have included this suggestion in the discussion.

The sample size of 12 is inadequate to have a sufficient idea on the feasibility of the intervention due to the wide variation in the variable related to feasibility.

This also we agree fully. This was a “pilot study” to assess feasibility as mentioned in the title. The study is not even a full feasibility study. We emphasize this in the revised limitation component of the manuscript.

As mentioned in “Feasibility of incorporating into Antenatal Care (ANC) system” Participants opinion regarding incorporating this intervention to national antenatal care in Sri Lanka has to be taken with reservation as they do not see the operational aspects of such a program when it is scaled up. In the responses there is also an element of social desirability bias.

Again we agree with this comment fully and included in the discussion.

“The degree of success of launching such a program nationally, or even locally, will depend heavily on the willingness and enthusiasm of the Public Health Midwife (PHM)s, as eventually, if the program is to be sustainable, it will be trained PHMs who will be leading each of the mindfulness session”. This is asking too much from already over burdened PHM in Sri Lanka. Our experience is that, out of the numbers who take up mindfulness training, only a small proportion will qualify to be trainers. Hence, we consider planning to train PHMs as trainers is a weakness in the design.

First, we also share the same concern about overburden PHM duties. At the same time, in Sri Lanka, national maternal health intervention through other means has never had sustainability. The second point is well taken and only a small proportion will be qualify

to be mindfulness trainers; not only among PHMs but also among other groups. This limitation and challenge was added to the revised discussion.

As mentioned in the discussion inconsistent attendance can be a difficult challenge to meet as take-up of this kind of interventions by different individuals vary widely.

Agree and it's already in the discussion.

It is still premature to conclude "...a feasible way to incorporate promotion of maternal mental health into routine ANC in Sri Lanka". Though it appears feasible to implement the intervention whether it improves mental well being as an outcome has not been assessed in this work. Hence this claim is not justified.

Agree and this conclusion was revised.

In the abstract, there is no conclusion in relation to the set objective of evaluating feasibility of intervention.

Conclusion of the abstract is revised.

Competing Interests: None

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research