REVIEW

Recent advances in understanding the physiology of hypoxic sensing by the carotid body [version 1; peer review: 3 approved]

Nanduri R. Prabhakar , Ying-Jie Peng, Jayasri Nanduri

Institute for Integrative Physiology and Center for Systems Biology of O₂ Sensing, The University of Chicago, Chicago, IL, 60637, USA

V1 First published: 06 Dec 2018, 7(F1000 Faculty Rev):1900
<https://doi.org/10.12688/f1000research.16247.1>

Latest published: 06 Dec 2018, 7(F1000 Faculty Rev):1900
<https://doi.org/10.12688/f1000research.16247.1>

Abstract

Hypoxia resulting from reduced oxygen (O₂) levels in the arterial blood is sensed by the carotid body (CB) and triggers reflex stimulation of breathing and blood pressure to maintain homeostasis. Studies in the past five years provided novel insights into the roles of heme oxygenase-2 (HO-2), a carbon monoxide (CO)-producing enzyme, and NADH dehydrogenase Fe-S protein 2, a subunit of the mitochondrial complex I, in hypoxic sensing by the CB. HO-2 is expressed in type I cells, the primary O₂-sensing cells of the CB, and binds to O₂ with low affinity. O₂-dependent CO production from HO-2 mediates hypoxic response of the CB by regulating H₂S generation. Mice lacking NDUFS2 show that complex I-generated reactive oxygen species acting on K⁺ channels confer type I cell response to hypoxia. Whether these signaling pathways operate synergistically or independently remains to be studied.

Keywords

Carotid body, Heme-oxygenase, Gasotransmitter, NADH dehydrogenase Fe-S protein 2

Open Peer Review

Approval Status

	1	2	3
version 1
06 Dec 2018			

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Norbert Weissmann**, Justus Liebig University Giessen, Giessen, Germany
2. **Prem Kumar**, University of Birmingham, Birmingham, UK
3. **Plamena R Angelova**, UCL Queen Square Institute of Neurology, London, UK

Any comments on the article can be found at the end of the article.

Corresponding author: Nanduri R. Prabhakar (nprabhak@medicine.bsd.uchicago.edu)

Author roles: Prabhakar NR: Writing – Original Draft Preparation, Writing – Review & Editing; Peng YJ: Writing – Review & Editing; Nanduri J: Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: Work in the authors' laboratory is supported by National Institutes of Health grant P01-HL-90554. *The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

Copyright: © 2018 Prabhakar NR *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Prabhakar NR, Peng YJ and Nanduri J. **Recent advances in understanding the physiology of hypoxic sensing by the carotid body [version 1; peer review: 3 approved]** F1000Research 2018, 7(F1000 Faculty Rev):1900 <https://doi.org/10.12688/f1000research.16247.1>

First published: 06 Dec 2018, 7(F1000 Faculty Rev):1900 <https://doi.org/10.12688/f1000research.16247.1>

Introduction

Systemic hypoxia, which arises from decreased oxygen (O_2) levels in the arterial blood, is a fundamental physiological stimulus. The duration of hypoxia can be acute, ranging from seconds to minutes, or chronic, lasting hours to days. Acute hypoxia evokes rapid changes in the cardiorespiratory systems to ensure optimal O_2 delivery to tissues. Cardiorespiratory responses to acute hypoxia are primarily reflexive in nature, initiated by sensory organs located in the carotid artery and aorta. Carotid bodies (CBs), which reside at the bifurcation of the common carotid arteries, are the major sensory organs for monitoring arterial blood O_2 levels¹. Although structures similar to CBs are seen at the aortic arch and in the abdominal arteries, much of the information on the mechanisms of hypoxic sensing has come from studies on the CB¹. Here, we present studies reported in the past five years on the roles for heme oxygenase-2 (HO-2), a carbon monoxide (CO)-synthesizing enzyme, and NDUFS2, a mitochondrial complex I subunit, in hypoxic sensing by the CB.

Physiology of hypoxic sensing by the carotid body

The CB receives sensory innervation from the carotid sinus nerve, whose cell bodies reside in the petrosal ganglion. Under basal conditions (arterial blood pO_2 of about 100 mmHg), sensory nerve discharge (that is, frequency of action potentials) is low. In response to even a modest decrease in arterial blood pO_2 from 100 to 80 mmHg, the sensory discharge increases and the response is fast, occurring within a few seconds after the onset of hypoxia¹. The increased sensory discharge is non-adapting and is maintained during the entire duration of hypoxia² or may progressively increase during sustained hypoxia, lasting several hours³. The exquisite sensitivity and the speed of the response with little or no adaptation are the unique features of hypoxic sensing by the CB. The increased CB sensory nerve activity is relayed to brainstem neurons, leading to reflex stimulation of breathing and blood pressure (CB chemo reflex)¹.

The CB tissue is made of two major cell types: type I cells (also called glomus cells), which are of neuronal origin, and type II cells, which resemble glial cells of the nervous system. Type I cells along with the nearby sensory nerve ending function as a “sensory unit”¹. Stimulus response of breathing to graded hypoxia parallels the CB sensory nerve activity¹. Consequently, carotid sinus nerve activity is measured as an index of CB hypoxic sensing¹. Type I cell responses to acute hypoxia are measured by monitoring exocytosis and changes in $[Ca^{2+}]_i$ and K^+ channel conductance¹. Although type I cells respond to hypoxia with elevated $[Ca^{2+}]_i$ or K^+ channel inhibition (or both), they are not always reflected in the sensory nerve activity^{4,5}, which is essential for evoking the physiologically important CB chemo reflex. Therefore, it is necessary to corroborate the cellular responses to hypoxia with the sensory nerve discharge for assessing the physiological relevance of CB hypoxic sensing.

Transduction mechanisms

The consensus is that hypoxia inhibits certain K^+ channels in type I cells and the resulting depolarization leads to Ca^{2+} -dependent release of neurotransmitter or neurotransmitters, which stimulate the nearby sensory nerve ending, leading to increased sensory discharge¹. The roles for K^+ channels and AMP kinase (AMPK) in CB hypoxic sensing have been discussed in detail

elsewhere^{1,6,7} and will not be presented in this commentary. The following section presents studies conducted in the past five years that have provided novel insights into the roles for HO-2 and mitochondrial complex subunit NDUFS2 in hypoxic sensing by the CB.

Heme oxygenase-2

Type I cells express HO-2-like immunoreactivity⁸. HO-2 is remarkably sensitive to O_2 availability, and graded hypoxia progressively decreases CO production in the CB⁹. Reduced CO production by hypoxia is also seen in HEK-293 cells with heterologous expression of HO-2⁹, suggesting that HO-2 is inherently sensitive to O_2 . HO-2 binds to O_2 with low affinity with an apparent K_m of 65 ± 5 mmHg (about 80 μM). The O_2 sensitivity of HO-2 is due to Cys²⁶⁵ and Cys²⁸² residues in the heme regulatory motif⁹. Intact Cys²⁶⁵ and Cys²⁸² residues lower the affinity of HO-2 for O_2 and thereby enable the enzyme to transduce changes in O_2 into changes in CO production. Substituting Cys²⁶⁵ and Cys²⁸² with alanine allows the HO-2 to bind to O_2 with high affinity⁹.

It was proposed, based on the findings that CO inhibits CB sensory nerve excitation by hypoxia^{8–10} and that hypoxia reduces CO production in a stimulus-dependent way⁹, that low sensory nerve activity during normoxia is due to high CO levels inhibiting CB sensory nerve activity but that hypoxia, by reducing CO production, relieves the inhibition and thereby increases the sensory nerve discharge⁸. Recent studies determined how CO inhibits CB sensory nerve activity under normoxia^{9,10}. Type I cells also express cystathionine gamma-lyase (CSE), an enzyme catalyzing hydrogen sulfide (H_2S) production¹¹. H_2S is a potent stimulator of CB sensory nerve activity in rats, mice, rabbits, and cats^{11–13}. During normoxia, H_2S levels are low and hypoxia increases H_2S levels in a stimulus-dependent manner¹¹. The increased H_2S production is not due to inherent O_2 sensitivity of CSE; rather, it is due to changes in CO production⁹. High CO levels during normoxia inhibit CSE-derived H_2S generation through protein kinase G (PKG)-dependent phosphorylation of Ser³⁷⁷ of CSE, and reduced CO generation during hypoxia relieves the inhibition of CSE, leading to increased H_2S generation in the CB⁹. CSE inhibitors and CSE knockout mice exhibit impaired type I cell and sensory nerve and breathing response to hypoxia^{11,14}. These findings suggest that low sensory discharge during normoxia is due to inhibition of H_2S generation by high levels of HO-2-derived CO but that the increased sensory nerve activity by hypoxia is due to relieving inhibition of H_2S synthesis by CO (Figure 1).

Genetic disruption of HO-2 increases baseline CB sensory nerve activity and elevates H_2S levels under normoxia, a phenotype similar to hypoxia⁹. However, hypoxia further increased CB sensory activity and elevated H_2S levels in HO-2 null CBs, indicating the existence of a redundant hypoxic sensing mechanism (or mechanisms). This redundant hypoxic sensing was due to compensatory upregulation of neuronal nitric oxide synthase (nNOS) in type I cells of HO-2 null mice⁹. nNOS, like HO-2, also binds to O_2 with low affinity, and nitric oxide (a product of nNOS), like CO, also inhibits CSE-derived H_2S through PKG signaling⁹. Blockade of nNOS in HO-2 null mice renders CBs completely insensitive to hypoxia⁹. These findings suggest

Figure 1. Heme oxygenase-2 (HO-2) signaling in hypoxic sensing by the carotid body (CB). Ca^{2+} , calcium channel; CO, carbon monoxide; CSE, cystathionine gamma-lyase; H_2S , hydrogen sulfide; K^+ , potassium channel; mito, mitochondria.

that, in the absence of HO-2, nNOS contributes to CB hypoxic sensing by regulating CSE-derived H_2S production⁹.

A recent study examined the role for HO-2–CO signaling in CB hypoxic sensing of three genetically distinct rat strains that are commonly used in experimental research¹⁰. As compared with Sprague-Dawley rat CB, Brown-Norway (BN) rat CB showed markedly attenuated sensory nerve and type I cell responses to hypoxia, and this phenotype was associated with higher CO and lower H_2S levels in the glomus tissue. The elevated CO levels in the BN rat CB were due to high affinity of HO-2 to its substrate heme¹⁰. The attenuated CB response to hypoxia is associated with a blunted chemo reflex. The CB chemo reflex is essential for ventilatory adaptations to high-altitude hypoxia^{15,16}. BN rats exposed to hypobaric hypoxia showed severe pulmonary edema¹⁰, which is a sign of chronic mountain sickness. Treating BN rats with a HO inhibitor restored the hypoxic response of the CB and prevented pulmonary edema caused by hypobaric hypoxia¹⁰.

In contrast to BN rat CBs, CBs of spontaneous hypertensive (SH) rats showed heightened sensitivity to hypoxia, and this phenotype is associated with low CO levels and high H_2S levels in the CB¹⁰. The low CO levels in SH rat CBs were due to low heme affinity of HO-2¹⁰. Current evidence suggests that a hyperactive CB chemo reflex is a major contributor of hypertension in SH rats^{17,18}. Systemic administration of a CSE inhibitor normalized CB sensory nerve and type I cell responses to hypoxia and reduced hypertension¹⁰. Although chronic ablation of CB also reduced hypertension to the same level as seen with a CSE inhibitor, combined CB ablation and CSE inhibitor treatment had no further effect on blood pressure¹⁰. These findings suggest that CO-regulated H_2S contributes to a hyperactive CB

in SH rats. It has long been known that the CB chemo reflex exhibits substantial inter-individual variations in humans and experimental animals^{19–21}. Studies on BN and SH rats indicate that inter-individual variations in chemo reflex are due in part to variations in HO-2–CO signaling in the CB.

A recent study suggests that HO-2–CO signaling in the CB also plays an important role in the pathology of sleep apnea (SA), which is a highly prevalent respiratory disease²². SA is characterized by episodic cessation of breathing leading to chronic intermittent hypoxia (CIH). Patients with SA and CIH-exposed rodents exhibit a heightened CB chemo reflex, leading to chronic elevation of sympathetic nerve activity and hypertension^{23–26}. Rodents exposed to CIH exhibit elevated reactive oxygen species (ROS) levels in the CB^{27,28}. Peng *et al.* showed that ROS inhibit CO generation by HO-2 by acting on the Cys²⁶⁵ residue in the heme regulatory motif, thereby increasing H_2S levels in the CB²². Pharmacological or genetic blockade of CSE-derived H_2S prevents CIH-induced CB hyperactivity, sympathetic nerve excitation, and hypertension²². Collectively, these studies suggest that disrupted HO-2–CO signaling in the CB leads to dire physiological consequences.

Although the studies described above suggest that CO-regulated H_2S is an important mediator of CB hypoxic sensing, a recent study questioned this possibility. Kim *et al.*²⁹ reported that inhibitors of H_2S synthesis had no effect on $[\text{Ca}^{2+}]_i$ and TASK K^+ channel responses of type I cells to anoxia (pO_2 of about 5 mmHg). Peng *et al.*⁵ re-examined the role for CSE-derived H_2S in the CB sensory nerve and type I cell $[\text{Ca}^{2+}]_i$ responses to hypoxia (pO_2 of about 37 mmHg) and anoxia (pO_2 of about 5 mmHg). The authors found that hypoxia increased H_2S levels in the CB, stimulated sensory nerve activity, and elevated $[\text{Ca}^{2+}]_i$ in

type I cells and all of these responses were blocked by a CSE inhibitor and in CSE knockout mice. In striking contrast, anoxia, though producing very low pO_2 , had no effect on H_2S levels in the CB and produced only a weak CB sensory nerve excitation as compared with hypoxia. CB sensory and type I cell responses to anoxia were unaffected by CSE inhibitors and in CSE knockout mice. Moreover, anoxia (100% N_2) depressed breathing whereas hypoxia (12% O_2) stimulated breathing⁵. CSE knockout mice showed an absence of breathing stimulation by hypoxia, whereas the depressed breathing by anoxia was unaffected in these mice⁵. These findings suggest that hypoxia and anoxia are not the same stimuli for studying CB physiology and that HO-2-CO-regulated H_2S mediates CB response to “physiologically relevant” hypoxia but not anoxia.

NADH dehydrogenase Fe-S protein 2 (NDUFS2) mitochondrial complex I subunit

Mitochondrial electron transport chain (ETC) inhibitors mimic the effects of hypoxia on CB sensory nerve activity¹ and type I cells^{30–33}. Mills and Jöbsis³⁴ reported that CBs express a putative cytochrome aa3 with two O_2 affinities: one with high and another with low affinity for O_2 . Based on spectral analysis, subsequent studies suggested that CBs express a cytochrome, which is half-reduced at a pO_2 of 60 to 80 mmHg, and this cytochrome is not expressed in either the superior cervical or the nodose ganglion^{35,36}. Acute hypoxia increased the NADH/NAD ratio and decreased mitochondrial membrane potential in type I cells, and these effects were not seen in other non-chemoreceptor tissues such as dorsal root ganglion³⁷. These studies led to the suggestion that mitochondrial ETC participates in CB hypoxic ensing.

Rotenone, an inhibitor of the mitochondrial complex I, selectively blocks the type I cell response to hypoxia³². NDUFS2 is a

component of the complex I, which binds to ubiquinone^{38–40}. Recent studies examined the role for complex I in CB hypoxic sensing in mice with targeted deletion of *Ndufs2* in tyrosine hydroxylase-positive (TH^+) cells such as type I cells⁴¹. Mice lacking NDUFS2 in TH^+ cells showed an absence of breathing stimulation by hypoxia and of hypoxia-evoked exocytosis and K^+ channel inhibition in type I cells. However, type I cell responses to severe hypercapnia (20% CO_2) were intact⁴¹. Given that these mice have a deletion of NDUFS2 since birth, the lack of breathing response to hypoxia might be secondary to metabolic adaptations during development. Additional studies were performed on adult (2-month-old) mice with conditional knockout of *Ndufs2* (ESR-NDUFS2 mice). Like the TH -NDUFS2 mice, mice with conditional knockdown of *Ndufs2* showed an absence of stimulation of breathing as well as type I cell responses to hypoxia⁴¹. The lack of cellular responses to hypoxia was associated with decreased complex I activity, complex I formation, and complex I-dependent O_2 consumption, whereas the functions of other mitochondrial complexes were intact. These studies suggest that NDUFS2 of the mitochondrial complex I contributes to hypoxic sensing by the CB.

How might NDUFS2 confer hypoxic sensitivity on type I cells? Mitochondrial ETC-generated ROS have been implicated in pulmonary artery myocyte responses to acute hypoxia^{42–44}. Acute hypoxia increased ROS in wild-type type I cell cytosol and mitochondrial intermembrane space, and these responses were attenuated in NDUFS2 null type I cells⁴¹. Intracellular application of H_2O_2 , like hypoxia, inhibited background K^+ currents in type I cells⁴¹. These findings led to the suggestion that inhibition of NDUFS2 leads to an increase in ROS production, which in turn (by inhibiting K^+ currents) leads to depolarization of type I cells by hypoxia (Figure 2).

Figure 2. NADH dehydrogenase Fe-S protein 2 (NDUFS2), a mitochondrial complex I subunit, signaling in hypoxic sensing by the carotid body (CB). Ca^{2+} , calcium channel; K^+ , potassium channel; ROS, reactive oxygen species.

Summary and future directions

It has long been thought that an O₂ sensor or sensors in type I cells initiate hypoxic sensing in the CB^{1,45,46}. To be considered an O₂ sensor, a molecule should satisfy certain criteria, namely (a) its presence in type I cells, (b) low-affinity binding to O₂, (c) altered function by hypoxia should initiate signaling events leading to increased CB sensory nerve activity, and (d) loss of CB hypoxic sensing by disrupting its function. HO-2 satisfies the proposed criteria for an O₂ sensor in the CB and contributes to CB sensory excitation by regulating H₂S production through O₂-dependent CO production. However, further studies are needed to delineate the cellular mechanism(s) underlying CB activation by H₂S. H₂S donors, like hypoxia, depolarize⁴⁷ and inhibit K⁺ channel conductance in type I cells^{12,48} and increase NADH auto-fluorescence in type I cells, an effect attributed to the inhibition of mitochondrial ETC⁴⁷. It is likely that H₂S mediates sensory nerve excitation by hypoxia by inhibiting mitochondrial ETC, thereby affecting K⁺ conductance of type I cells (Figure 1).

Studies with genetically engineered mice suggest that the inactivation of NDUFS2 is an important step for the type I cell response to hypoxia. However, it remains to be determined whether graded hypoxia inhibits NDUFS2 and establishes the affinity of O₂ for this enzyme. NDUFS2 is a ubiquitously expressed enzyme in the body. However, unlike many other tissues, CBs are highly

sensitive to changes in O₂ levels. Consequently, the uniqueness of NDUFS2 signaling in the CB remains to be established.

Finally, whether HO-2 and NDUFS2 signaling operates independently or works in concert is not clear. The CB responds to a wide range of pO₂ values (about 80–20 mmHg). It was proposed that interactions between multiple signaling pathways working in concert as a “chemosome” enable the CB to sense a wide range of pO₂ values^{45,46}. Given that high concentrations of H₂S can inhibit mitochondrial ETC⁴⁷, HO-2 and NDUFS2 signaling might work in concert as a “chemosome” for the full expression of the CB to a wide range of hypoxia, a possibility that remains to be investigated.

Author contributions

NRP contributed to the preparation of the original draft. Y-JP contributed to the preparation of the figures and editing. JN contributed to the review and editing.

Grant information

Work in the authors' laboratory is supported by National Institutes of Health grant P01-HL-90554.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- Kumar P, Prabhakar NR: **Peripheral chemoreceptors: function and plasticity of the carotid body.** *Compr Physiol.* 2012; 2(1): 141–219.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kou YR, Ernsberger P, Cragg PA, et al.: **Role of alpha 2-adrenergic receptors in the carotid body response to isocapnic hypoxia.** *Respir Physiol.* 1991; 83(3): 353–364.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Nielsen AM, Bisgard GE, Vidruk EH: **Carotid chemoreceptor activity during acute and sustained hypoxia in goats.** *J Appl Physiol (1985).* 1988; 65(4): 1796–1802.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Donnelly DF: **Chemoreceptor nerve excitation may not be proportional to catecholamine secretion.** *J Appl Physiol (1985).* 1996; 81(2): 657–664.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Peng YJ, Makarenko VV, Gridina A, et al.: **H₂S mediates carotid body response to hypoxia but not anoxia.** *Respir Physiol Neurobiol.* 2019; 259: 75–85.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- F** Peers C, Wyatt CN, Evans AM: **Mechanisms for acute oxygen sensing in the carotid body.** *Respir Physiol Neurobiol.* 2010; 174(3): 292–298.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- F** Rakoczy RJ, Wyatt CN: **Acute oxygen sensing by the carotid body: a rattlebag of molecular mechanisms.** *J Physiol.* 2018; 596(15): 2969–2976.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Prabhakar NR, Dinerman JL, Agani FH, et al.: **Carbon monoxide: a role in carotid body chemoreception.** *Proc Natl Acad Sci U S A.* 1995; 92(6): 1994–1997.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Yuan G, Vasavda C, Peng YJ, et al.: **Protein kinase G-regulated production of H₂S governs oxygen sensing.** *Sci Signal.* 2015; 8(373): ra37.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- F** Peng YJ, Makarenko VV, Nanduri J, et al.: **Inherent variations in CO-H₂S-mediated carotid body O₂ sensing mediate hypertension and pulmonary edema.** *Proc Natl Acad Sci U S A.* 2014; 111(3): 1174–1179.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- F** Peng YJ, Nanduri J, Raghuraman G, et al.: **H₂S mediates O₂ sensing in the carotid body.** *Proc Natl Acad Sci U S A.* 2010; 107(23): 10719–10724.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- F** Li Q, Sun B, Wang X, et al.: **A crucial role for hydrogen sulfide in oxygen sensing via modulating large conductance calcium-activated potassium channels.** *Antioxid Redox Signal.* 2010; 12(10): 1179–1189.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- F** Jiao Y, Li Q, Sun B, et al.: **Hydrogen sulfide activates the carotid body chemoreceptors in cat, rabbit and rat ex vivo preparations.** *Respir Physiol Neurobiol.* 2015; 208: 15–20.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Makarenko VV, Nanduri J, Raghuraman G, et al.: **Endogenous H₂S is required for hypoxic sensing by carotid body glomus cells.** *Am J Physiol Cell Physiol.* 2012; 303(9): C916–923.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- F** Dempsey JA, Powell FL, Bisgard GE, et al.: **Role of chemoreception in cardiorespiratory acclimatization to, and deacclimatization from, hypoxia.** *J Appl Physiol (1985).* 2014; 116(7): 858–866.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Dempsey JA, Forster HV: **Mediation of Ventilatory Adaptations.** *Physiol Rev.* 1982; 62(1): 262–346.
[PubMed Abstract](#) | [Publisher Full Text](#)
- F** Paton JF, Sobotka PA, Fudim M, et al.: **The carotid body as a therapeutic target for the treatment of sympathetically mediated diseases.** *Hypertension.* 2013; 61(1): 5–13.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Przybylski J: **Do arterial chemoreceptors play a role in the pathogenesis of hypertension?** *Med Hypotheses.* 1981; 7(2): 127–131.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Hodges MR, Forster HV, Papanek PE, et al.: **Ventilatory phenotypes among four strains of adult rats.** *J Appl Physiol (1985).* 2002; 93(3): 974–983.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Strohl KP, Thomas AJ, St Jean P, et al.: **Ventilation and metabolism among rat strains.** *J Appl Physiol (1985).* 1997; 82(1): 317–323.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Weil JV: **Variation in human ventilatory control-genetic influence on the hypoxic ventilatory response.** *Respir Physiol Neurobiol.* 2003; 135(2–3): 239–246.
[PubMed Abstract](#) | [Publisher Full Text](#)

22. Yuan G, Peng YJ, Khan SA, *et al.*: **H₂S production by reactive oxygen species in the carotid body triggers hypertension in a rodent model of sleep apnea.** *Sci Signal.* 2016; **9**(441): ra80.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
23. Peng YJ, Yuan G, Khan S, *et al.*: **Regulation of hypoxia-inducible factor- α isoforms and redox state by carotid body neural activity in rats.** *J Physiol.* 2014; **592**(17): 3841–3858.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
24. Kara T, Narkiewicz K, Somers VK: **Chemoreflexes—physiology and clinical implications.** *Acta Physiol Scand.* 2003; **177**(3): 377–384.
[PubMed Abstract](#) | [Publisher Full Text](#)
25. Narkiewicz K, van de Borne PJ, Pesek CA, *et al.*: **Selective potentiation of peripheral chemoreflex sensitivity in obstructive sleep apnea.** *Circulation.* 1999; **99**(9): 1183–1189.
[PubMed Abstract](#) | [Publisher Full Text](#)
26. Lesske J, Fletcher EC, Bao G, *et al.*: **Hypertension caused by chronic intermittent hypoxia—Influence of chemoreceptors and sympathetic nervous system.** *J Hypertens.* 1997; **15**(12 Pt 2): 1593–1603.
[PubMed Abstract](#) | [Publisher Full Text](#)
27. Peng YJ, Overholt JL, Kline D, *et al.*: **Induction of sensory long-term facilitation in the carotid body by intermittent hypoxia: implications for recurrent apneas.** *Proc Natl Acad Sci U S A.* 2003; **100**(17): 10073–10078.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
28. Pawar A, Nanduri J, Yuan G, *et al.*: **Reactive oxygen species-dependent endothelin signaling is required for augmented hypoxic sensory response of the neonatal carotid body by intermittent hypoxia.** *Am J Physiol Regul Integr Comp Physiol.* 2009; **296**(3): R735–742.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
29. Kim D, Kim I, Wang J, *et al.*: **Hydrogen sulfide and hypoxia-induced changes in TASK (K_v3/9) activity and intracellular Ca²⁺ concentration in rat carotid body glomus cells.** *Respir Physiol Neurobiol.* 2015; **215**: 30–38.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
30. Duchon MR, Biscoe TJ: **Mitochondrial function in type I cells isolated from rabbit arterial chemoreceptors.** *J Physiol.* 1992; **450**: 13–31.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
31. Buckler KJ, Vaughan-Jones RD: **Effects of mitochondrial uncouplers on intracellular calcium, pH and membrane potential in rat carotid body type I cells.** *J Physiol.* 1998; **513**(Pt 3): 819–833.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
32. Ortega-Sáenz P, Pardo R, García-Fernández M, *et al.*: **Rotenone selectively occludes sensitivity to hypoxia in rat carotid body glomus cells.** *J Physiol.* 2003; **548**(Pt 3): 789–800.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
33. Wyatt CN, Buckler KJ: **The effect of mitochondrial inhibitors on membrane currents in isolated neonatal rat carotid body type I cells.** *J Physiol.* 2004; **556**(Pt 1): 175–191.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
34. Mills E, Jöbsis FF: **Simultaneous measurement of cytochrome a_3 reduction and chemoreceptor afferent activity in the carotid body.** *Nature.* 1970; **225**(5238): 1147–1149.
[PubMed Abstract](#) | [Publisher Full Text](#)
35. Lahiri S, Ehleben W, Acker H: **Chemoreceptor discharges and cytochrome redox changes of the rat carotid body: role of heme ligands.** *Proc Natl Acad Sci U S A.* 1999; **96**(16): 9427–9432.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
36. Streller T, Huckstorf C, Pfeiffer C, *et al.*: **Unusual cytochrome a_{355} with low PO₂ affinity correlates as putative oxygen sensor with rat carotid body chemoreceptor discharge.** *FASEB J.* 2002; **16**(10): 1277–1279.
[PubMed Abstract](#) | [Publisher Full Text](#)
37. Duchon MR, Biscoe TJ: **Relative mitochondrial membrane potential and [Ca²⁺]_i in type I cells isolated from the rabbit carotid body.** *J Physiol.* 1992; **450**: 33–61.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
38. Kashani-Poor N, Zwicker K, Kerschner S, *et al.*: **A central functional role for the 49-kDa subunit within the catalytic core of mitochondrial complex I.** *J Biol Chem.* 2001; **276**(26): 24082–24087.
[PubMed Abstract](#) | [Publisher Full Text](#)
39. Baradaran R, Berrisford JM, Minhas GS, *et al.*: **Crystal structure of the entire respiratory complex I.** *Nature.* 2013; **494**(7438): 443–448.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
40. Rhein VF, Carroll J, Ding S, *et al.*: **NDUFAF7 methylates arginine 85 in the NDUFS2 subunit of human complex I.** *J Biol Chem.* 2013; **288**(46): 33016–33026.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
41. Fernández-Agüera MC, Gao L, González-Rodríguez P, *et al.*: **Oxygen Sensing by Arterial Chemoreceptors Depends on Mitochondrial Complex I Signaling.** *Cell Metab.* 2015; **22**(5): 825–837.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
42. Archer SL, Huang J, Henry T, *et al.*: **A redox-based O₂ sensor in rat pulmonary vasculature.** *Circ Res.* 1993; **73**(6): 1100–1112.
[PubMed Abstract](#) | [Publisher Full Text](#)
43. Leach RM, Hill HM, Snetkov VA, *et al.*: **Divergent roles of glycolysis and the mitochondrial electron transport chain in hypoxic pulmonary vasoconstriction of the rat: identity of the hypoxic sensor.** *J Physiol.* 2001; **536**(Pt 1): 211–224.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
44. Waypa GB, Chandel NS, Schumacker PT: **Model for hypoxic pulmonary vasoconstriction involving mitochondrial oxygen sensing.** *Circ Res.* 2001; **88**(12): 1259–1266.
[PubMed Abstract](#) | [Publisher Full Text](#)
45. Prabhakar NR: **Sensing hypoxia: physiology, genetics and epigenetics.** *J Physiol.* 2013; **591**(9): 2245–2257.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
46. Prabhakar NR: **O₂ sensing at the mammalian carotid body: why multiple O₂ sensors and multiple transmitters?** *Exp Physiol.* 2006; **91**(1): 17–23.
[PubMed Abstract](#) | [Publisher Full Text](#)
47. Buckler KJ: **Effects of exogenous hydrogen sulphide on calcium signalling, background (TASK) K channel activity and mitochondrial function in chemoreceptor cells.** *Pflügers Arch.* 2012; **463**(5): 743–754.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
48. Telezhkin V, Brazier SP, Cayzac SH, *et al.*: **Mechanism of inhibition by hydrogen sulfide of native and recombinant BK_{Ca} channels.** *Respir Physiol Neurobiol.* 2010; **172**(3): 169–178.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. **Plamena R Angelova**

Department of Clinical and Movement Neurosciences, UCL Queen Square Institute of Neurology, London, UK

Competing Interests: No competing interests were disclosed.

2. **Prem Kumar**

Institute of Clinical Sciences, College of Medical and Dental Sciences, University of Birmingham, Birmingham, UK

Competing Interests: No competing interests were disclosed.

3. **Norbert Weissmann**

German Center for Lung Research (DZL), ECCPS, Justus Liebig University Giessen, Giessen, Germany

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research