RESEARCH ARTICLE

REVISED Considerations related to vaping as a possible gateway into cigarette smoking: an analytical review [version 2; peer review: 1 approved, 1 approved with reservations]

Peter N. Lee ¹, Katharine J. Coombs¹, Esther F. Afolalu ²

¹P.N.Lee Statistics and Computing Ltd, Sutton, Surrey, SM2 5DA, UK

²PMI R&D, Philip Morris Products S.A., Quai JeanRenaud 5, Neuchâtel, CH-2000, Switzerland

v2 First published: 10 Dec 2018, 7:1915
<https://doi.org/10.12688/f1000research.16928.1>
 Second version: 21 May 2019, 7:1915
<https://doi.org/10.12688/f1000research.16928.2>
 Latest published: 22 Jul 2019, 7:1915
<https://doi.org/10.12688/f1000research.16928.3>

Abstract

Background: Toxicant levels are much lower in e-cigarettes than cigarettes. Therefore, introducing e-cigarettes into the market seems likely to reduce smoking-related diseases (SRD). However, vaping might provide a gateway into cigarette smoking for those who otherwise would never have smoked, a concern fuelled by cohort studies showing vaping predicts subsequent smoking initiation in young people.

Methods: In this discussion paper, we consider various aspects of the gateway issue in youths. We provide a descriptive critical review of results from prospective studies relating to the gateway effect and the extent to which the studies considered other potential confounding variables associated with smoking initiation. We then estimate the effects of omitting a confounding variable, or misclassifying it, on the association between vaping and subsequent smoking initiation, and determine how the prevalence of smoking might be affected by any true gateway effects of vaping. Finally, we examine trends in e-cigarette and smoking prevalence in youths based on national surveys.

Results: First, we demonstrate that although studies report that vaping significantly predicts smoking initiation following adjustment for various other predictors, the sets of predictors considered are quite incomplete. Furthermore, no study considered residual confounding arising from inaccurate measurement of predictors. More precise adjustment may substantially reduce the association. Second, we show any true gateway effect would likely affect smoking prevalence only modestly. Third, we show smoking prevalence in U.S. and U.K. youths in 2014–2016 declined somewhat faster than predicted by the preceding trend; a substantial gateway effect suggests the opposite. Finally, we argue that even if some gateway

Open Peer Review

Approval Status

	1	2
version 3		
(revision)	view	
22 Jul 2019		
version 2		
(revision)	view	view
21 May 2019		
version 1		
10 Dec 2018	view	view

1. **Lion Shahab** , University College London (UCL), London, UK

2. **Sandra I. Sulsky** , Ramboll US Corporation, Amherst, USA

Any reports and responses or comments on the article can be found at the end of the article.

effect exists, introducing e-cigarettes still likely reduces SRDs.

Conclusions: Given that the existence of any true gateway effect in youth is not yet clearly demonstrated the population health impact of introducing e-cigarettes is still likely to be beneficial.

Keywords

Cigarettes e-cigarettes, gateway effects

Corresponding author: Peter N. Lee (PeterLee@pnlee.co.uk)

Author roles: **Lee PN:** Conceptualization, Data Curation, Formal Analysis, Funding Acquisition, Investigation, Methodology, Project Administration, Resources, Software, Supervision, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Coombs KJ:** Data Curation, Formal Analysis, Investigation, Validation, Writing – Review & Editing; **Afolalu EF:** Data Curation, Formal Analysis, Investigation, Methodology, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: PNL and KJC consult for various tobacco companies. EFA is a current employee of Philip Morris International.

Grant information: This work was supported by PMI R&D, Philip Morris Products S.A., Quai Jeanrenaud 5, CH-2000 Neuchâtel, Switzerland.

Copyright: © 2019 Lee PN *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Lee PN, Coombs KJ and Afolalu EF. **Considerations related to vaping as a possible gateway into cigarette smoking: an analytical review [version 2; peer review: 1 approved, 1 approved with reservations]** F1000Research 2019, 7:1915 <https://doi.org/10.12688/f1000research.16928.2>

First published: 10 Dec 2018, 7:1915 <https://doi.org/10.12688/f1000research.16928.1>

REVISED Amendments from Version 1

Following the comments made by the two reviewers the following changes have been made to the paper:

- The terms “gateway-in” and “gateway-out” have been abandoned. The term “gateway-in” is now referred to as the gateway effect, and “gateway-out” is explained more generally.
- The term concordance is now no longer used, reference being made to the specific odds ratio.
- A new Additional File 4 has been added to OSF, giving fuller details of the analyses summarized in the “Has introducing e-cigarettes affected smoking trends?” section.
- A paragraph has been added at the end of the “Association of smoking with other risk factors” section which notes that though our search terms were limited, our conclusion still applies.
- A paragraph has been added at the end of the “Additional evidence on the likelihood of any marked gateway effect on smoking prevalence” section to point out some limitations of this additional evidence.
- A new theoretical beneficial effect ‘B4’ has been added to Table 9 and the text related to it has been extended.
- In the Discussion we now mention our recent publication based on data from the PATH study.
- Two paragraphs starting “While various authors” and “These analyses have” have been moved from the Results to the Discussion section.
- In discussing the limitations of our work, we now include failing to consider persistence of any true gateway effects.
- Numerous amendments have been made to clarify some of our assumptions and to make the text easier to follow.
- Some minor changes have been made to the English following a detailed reread.

See referee reports

Introduction

Recent publications make it clear that, in youths, vaping (i.e., use of e-cigarettes) and cigarette smoking (subsequently referred to as “smoking”) are strongly associated. In the U.S., for example, a survey of ninth and tenth grade students in Hawaii in 2014 (Wills *et al.*, 2017) revealed 195 ever-users of both products, 250 ever-vapers only, 37 ever-smokers only, and 820 never-users of either. From these data, the odds ratio (OR) relating ever-vaping to ever-smoking can be estimated as 17.3 (95% confidence interval (CI) 11.8–25.3). A strong association can also be seen for sixth to twelfth grade students in Texas in 2014 (Cooper *et al.*, 2016), with an OR of 28.8 (25.0–33.1), as well as nationally in 2012 (Dutra & Glantz, 2014), with an OR of 31.9 (27.6–36.8). Similarly strong relationships are reported also in Canada (Aleyan *et al.*, 2018), France (Dautzenberg *et al.*, 2016), Great Britain (Eastwood *et al.*, 2015), Poland (Goniewicz *et al.*, 2014), and Korea (Lee *et al.*, 2014). Theoretically, this association may arise if vaping encourages smoking, if smoking encourages vaping, and/or if other factors link to use of both products.

Concern that vaping may encourage subsequent smoking, the so-called “gateway” effect, was fuelled by a recent paper by

Soneji *et al.* (2017). In this manuscript, the authors combined epidemiological evidence from nine U.S. cohort studies in young people that linked smoking initiation to previous vaping. Among baseline never-smokers, baseline ever-vaping strongly predicted smoking initiation within six to 18 months (OR 3.62, 95% CI 2.42–5.41) after adjusting for various predictors of initiation. Similarly, baseline past 30-day vaping predicted subsequent 30-day cigarette use (OR 4.28, 95% CI 2.52–7.27).

Based on these results and those from one additional study (Hammond *et al.*, 2017), the National Academies Press on the Public Health Consequences of E-Cigarettes (National Academies of Sciences Engineering and Medicine, 2018) recently concluded that “there is substantial evidence that e-cigarette use increases risk of ever using combustible tobacco cigarettes among youth and young adults, noting that the relevant studies had adjusted for “a wide range of covariates” and considering that it was “unlikely that confounding entirely accounts for the association because reductions in estimates of association from unadjusted to adjusted models were not consistently observed in the literature.”

Methods

This review considers various methodological aspects of the gateway issue in youths.

Search parameters

First, based on PubMed searches carried out at intervals starting in May 2017 on “ecigarettes” or “e-cigarettes” or “e-cigs” or “electronic cigarette” or “ecigarette” or “e-cigarette”, we identified papers and reviews that presented results from prospective studies of young people relating to the gateway effect. Further publications were also sought from reference lists of selected papers and reviews. For this review, we also considered studies that provided data on trends over time in cigarette smoking by youths in relation to e-cigarette use, and information on initiation of smoking and e-cigarette smoking by youths relevant to the gateway effect.

A list of factors other than e-cigarette use that were associated with the initiation of cigarette smoking was obtained from US Surgeon General (1994) and from a separate ongoing review on determinants of smoking initiation which aims to present meta-analyses of associations between initiation of cigarette smoking and the various other factors. This identified references using an Embase search in April 2017 with the following search string:

‘smoking’/de OR ‘smoking’ AND (‘initiation’/de OR ‘initiation’ AND (‘prevalence’/de OR ‘prevalence’ OR ‘incidence’/de OR ‘incidence’ OR ‘proportion’ OR ‘age’/de OR ‘age’ OR ‘dual use’ OR ‘combined use’ OR ‘psychosocial’/de OR ‘psychosocial’ OR ‘beliefs’/de OR ‘beliefs’ OR ‘attitudes’/de OR ‘attitudes’ OR ‘perceptions’/de OR ‘perceptions’ OR ‘opinions’ OR ‘acceptance’/de OR ‘acceptance’ OR ‘predictors’/de OR ‘predictors’ OR ‘friend’/de OR ‘friend’ OR ‘family’/de OR ‘family’ OR ‘parent’/de OR ‘parent’ OR ‘propensity score’/de OR ‘propensity score’) AND [humans]/lim AND [english]/lim AND ([embase]/lim OR [medline]/lim)

For the current paper, we only use the results from this search to list those factors shown in one or more studies to strongly associate with initiation of smoking, and to consider the extent to which the identified gateway studies take these factors into account.

Analysis of search results

In addition, separate sheets of an Excel Program (Lee, 2018) were developed to:

- (a) estimate the effects of omission of a confounding variable, or misclassification of it, on the association between vaping and subsequent initiation of smoking, and to
- (b) determine how the prevalence of smoking might be affected by any true gateway effect of vaping.

Full details are given in, respectively, Additional Files 2 and 3 (Lee, 2018). Both sheets are used to provide illustrative examples of how the effects depend on the parameter values assumed.

In the first sheet, the use may determine how the observed gateway effect depends on the following eight parameters:

- The proportion of vapers in never smokers (P_1)
- The proportion with a particular smoking predictor present in never smokers (P_2)
- The odds ratio relating vaping to the smoking predictor (K)
- The probability of initiating smoking during follow-up in those who neither smoke nor have the predictor (P_A)
- The odds ratio for initiating smoking during follow-up associated with vaping - the true gateway effect (G_e)
- The odds ratio for initiating smoking during follow-up associated with the smoking predictor - the true smoking predictor effect (G_p)
- The proportion with the predictor who are misclassified as not having it (M_1), and
- The proportion without the predictor who are misclassified as having it (M_2)

P_1 , P_2 and K allow define the baseline population of N never smokers to be subdivided in a 2×2 table according to presence or absence of e-cigarettes and of the smoking predictor. P_A , G_e and G_p allow determination of the distribution of the population after a single follow-up period. M_1 and M_2 are misclassification rates of the predictor.

In the second sheet, the program considers individuals, divided into five equal strata, who initially are all never users of either cigarettes or e-cigarettes. The probability of initiating smoking or vaping increases progressively over the strata, the strata being intended to represent sets of individuals with an increasing susceptibility to tobacco. Over five time intervals, the individuals

may transfer to four other groups: current vapers only, current smokers only, current dual users and former users (those who have used one or both products but do not currently use either). Users may modify the values of seven parameters:

- The initiation rate of vaping in the first stratum (P_1)
- The relative odds of initiation for successive strata (R_1)
- The relative odds of initiation with smoking compared to vaping (R_2)
- The relative odds of quitting compared to initiation (R_3)
- The relative odds of re-initiation compared to initiation (R_4)
- The relative odds of initiating smoking for vapers compared to tobacco never-users (G_1)
- The relative odds of quitting smoking for dual users compared to smokers only (G_2)

Note that R_3 and R_4 are each taken to be applicable to either product.

Lastly, the review also includes an analysis of trends in e-cigarette and smoking prevalence in youths based on national surveys which provided annual information over a period of at least 10 years. We identified four surveys from the U.S., the Youth Risk Behavior Survey (YRBS: available from <https://www.cdc.gov/healthyyouth/data/yrbs/index.htm>), the National Youth Tobacco Survey (NYTS: available from https://www.cdc.gov/tobacco/data_statistics/surveys/nyts/index.htm), the Monitoring the Future study (MTF: available from <https://www.icpsr.umich.edu/icpsrweb/NAHDAP/series/35>), and the National Survey on Drug Use and Health (NSDUH: available from <https://www.samhsa.gov/data/data-we-collect/nsduh-national-survey-drug-use-and-health>). We also identified two surveys from the UK, the Smokefree Youth Survey in Great Britain (SYSGB: available from <http://ash.org.uk/download/use-of-electronic-cigarettes-among-children-in-great-britain/>) and the Office for National Statistics (ONS: available from <https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/drugusealcoholandsmoking/datasets/ecigaretteuseingreatbritain>). All six databases were accessed on 16 Oct 2017.

Based on linear regression of $\log_e(p/(1-p))$ on year, where p is the prevalence of smoking, observed prevalences in the years 2014–2016 (a period where the advent of e-cigarettes might have had a measurable effect on smoking prevalence if an important gateway effect existed) was compared with those expected based on the trend in those previous years for which results were available.

Results

Evidence relating vaping in youths to subsequent initiation of cigarette smoking

From the literature search, we identified a key systematic review and meta-analysis by Soneji *et al.* (2017) on the association between initial use of e-cigarettes and subsequent cigarette smoking among adolescents and young adult. Table 1 provides

Table 1. Association between vaping at baseline and subsequent smoking among youths.

Source ^a	Location	Sample size ^b	Age/grade	Follow-up period (months)	Odds ratio ^c	Unadjusted OR (95% CI) ^d	Adjusted OR (95% CI) ^d
Studies considered by Soneji <i>et al.</i> (2017)							
Leventhal <i>et al.</i> (2015)	USA, Los Angeles	2558	14	12	A	7.78 (6.15–9.84)	1.75 (1.10–2.78)
Primack <i>et al.</i> (2015)	USA, National	694	16–26	12	A	5.66 (1.99–16.07)	8.30 (1.19–58.00)
Barrington-Trimis <i>et al.</i> (2016)	USA, Southern California	298	16–18	16	A	5.76 (3.12–10.66)	6.17 (3.29–11.57)
Primack <i>et al.</i> (2016)	USA, National	1506	18–30	18	A	6.06 (2.15–17.10)	8.80 (2.37–32.69)
Miech <i>et al.</i> (2017)	USA, National	246	17–20	13	A	6.23 (1.57–24.63)	4.78 (1.91–11.96)
Spindle <i>et al.</i> (2017)	USA, National	2316	18–25	12	A	3.50 (2.41–5.09)	3.37 (1.91–5.54)
Wills <i>et al.</i> (2017)	USA, Hawaii	1141	14–16	12	A	4.25 (2.74–6.61)	2.87 (2.03–4.05)
Hornik <i>et al.</i> (2016)	USA, National	1028	13–25	6	B	11.18 (5.41–23.13)	5.43 (2.59–11.58)
Unger <i>et al.</i> (2016)	USA, Los Angeles	1056	22–24	12	B	4.71 (2.27–9.77)	3.32 (1.55–7.11)
Other studies							
Best <i>et al.</i> (2018)	UK, Scotland	2125	11–18	12	A	4.62 (3.34–6.38)	2.42 (1.63–3.60)
Conner <i>et al.</i> (2018)	UK, England	1726	13–14	12	A	5.38 (4.02–7.22)	4.06 (2.94–5.60)
Loukas <i>et al.</i> (2018)	USA, Texas	2558	18–25	18	A	2.73 (2.11–3.54)	1.36 (1.01–1.83)
Watkins <i>et al.</i> (2018)	USA, National	10384	12–17	12	A	3.50 (2.48–4.94)	2.53 (1.80–3.56)
Hammond <i>et al.</i> (2017)	Canada, 2 states	19310	14–18	12	A	4.81 (3.90–5.94)	2.12 (1.68–2.66)
Lozano <i>et al.</i> (2017)	Mexico, 3 cities	4695	12–13	20	A	1.82 (1.54–2.14)	1.40 (1.22–1.60)

^aTwo studies (Hornik *et al.*, 2016; Primack *et al.*, 2016) were reported only as abstracts, but fuller details were supplied to Soneji *et al.* for their meta-analyses. ^bOf never- (or noncurrent) smokers at baseline. ^cA = Odds of smoking initiation, among never-smokers at baseline, for ever- compared with never-vapers at baseline. B = Odds of smoking at baseline, among noncurrent smokers at baseline, for current compared with noncurrent vapers at baseline. ^dORs for U.S. studies as given by Soneji *et al.* ORs for U.K. studies come from the source, except that the unadjusted estimates for Best *et al.* and Loukas *et al.* were estimated from data given.

details of the nine U.S. studies analyzed by Soneji *et al.* (2017). A total of five studies, conducted with participants with a maximum age of 24–30 years old, collected information using internet-based surveys. The four other studies, involving participants aged less than 20 years old were based on self-completed questionnaires. The follow-up period was 6 months in one study (Hornik *et al.*, 2016), 1 year in five studies (Leventhal *et al.*, 2015; Primack *et al.*, 2015; Spindle *et al.*, 2017; Unger *et al.*, 2016; Wills *et al.*, 2017), and 13 to 18 months in the other three studies (Barrington-Trimis *et al.*, 2016; Miech *et al.*, 2017; Primack *et al.*, 2016). While most of the studies (marked with an A in Table 1) concerned ever- and never-use, two studies (marked with a B) considered current and noncurrent use. While the ORs marked B are less informative as they relate partly to relapse rates, they have been included for completeness.

Each study found that baseline vaping significantly predicted subsequent smoking, regardless of covariate adjustment. Adjusted ORs were lower than unadjusted ORs in six studies, particularly in two (Hornik *et al.*, 2016; Leventhal *et al.*, 2015). However, two studies (Primack *et al.*, 2015; Primack *et al.*, 2016) showed a moderately increased OR after adjustment.

Table 1 also includes results from six later studies cited in a recent review (Glasser *et al.*, 2018): two from the U.S. (Loukas *et al.*, 2018; Watkins *et al.*, 2018), two from the U.K.

(Best *et al.*, 2018; Conner *et al.*, 2018), and one each from Canada (Hammond *et al.*, 2017) and Mexico (Loukas *et al.*, 2018). These results showed an association that reduced but remained significant after adjustment. Two further studies could not be included in Table 1, as they did not provide comparable results. One was a study on young adults in the Chicago area (Selya *et al.*, 2018) that used path analysis and concluded that “E-cigarette use was not significantly associated with later conventional smoking...” The other was a study in the Netherlands (Treur *et al.*, 2018) that reported a strong association after covariate adjustment but did not present unadjusted results for comparison.

There were an additional 15 publications (Amato *et al.*, 2016; Ambrose, 2017; Barrington-Trimis *et al.*, 2015; Bold *et al.*, 2016; de Lacy *et al.*, 2017; Doran *et al.*, 2017; Etter & Bullen, 2014; Hanewinkel & Isensee, 2015; Huh & Leventhal, 2016; Kaufman *et al.*, 2015; Leventhal *et al.*, 2016; Loukas *et al.*, 2015; Sutfin *et al.*, 2015; Westling *et al.*, 2017; Zhong *et al.*, 2016) identified in our searches as of possible relevance based on the abstract. However, none of these provided useful data for various reasons. These included studies which: were conducted in adults; predicted e-cigarette use rather than cigarette smoking; did not present results in a form allowing the gateway effect to be estimated; were superseded by later publications; concerned intent to smoke cigarettes and not actual smoking; or even did not consider e-cigarettes at all.

Association of smoking with other risk factors

Initiation of smoking is long-established to be associated with various sociodemographic, environmental, and behavioral factors. Those identified by an authoritative source over 20 years ago (US Surgeon General, 1994) include low education level (Gritz *et al.*, 1998; Sargent *et al.*, 1997), internalizing/externalizing disorders (de Leon *et al.*, 2002; Ernst *et al.*, 2010; Rohde *et al.*, 2003), outcome expectancies (Barrington-Trimis *et al.*, 2015; Simons-Morton *et al.*, 1999), susceptibility to smoking (Huang *et al.*, 2005; Jackson, 1998), conduct problems (Dalton *et al.*, 2003; Scal *et al.*, 2003), substance use (Reed *et al.*, 2007; Scal *et al.*, 2003), risk-taking behaviour (Coogan *et al.*, 1998; Dalton *et al.*, 2003), poor school performance (O'Connor *et al.*, 2003; Sargent *et al.*, 1997), anxiety (Coogan *et al.*, 1998; Scal *et al.*, 2003), household smoking (Picotte *et al.*, 2006; Sargent *et al.*, 1997), peer smoking (Barrington-Trimis *et al.*, 2015; Picotte *et al.*, 2006), peer attitudes to smoking (Barrington-Trimis *et al.*, 2015; Daly *et al.*, 1993), low self-esteem (Dalton *et al.*, 2003; Weiss *et al.*, 2006), and other tobacco product use (Barrington-Trimis *et al.*, 2016; Jordan *et al.*, 2014). Currently available evidence reveals that many of these predictors consistently show a strong association with initiation, with reported ORs often exceeding 10. While these predictors are clearly not all independent, their number illustrates the difficulty in ensuring that gateway studies consider an adequate list. To avoid unfeasibly long questionnaires or huge studies, further work is needed to define an agreed minimum list of factors.

For the 15 studies included in Table 1, Additional File 1 (Lee, 2018) describes in detail how these risk factors were taken into account. While standard demographics were generally considered, and smoking susceptibility, substance use, risk-taking behaviour, other tobacco use, parental education, family smoking, and peer smoking were considered in at least five studies, many other relevant factors were rarely considered. These include internalizing/externalizing disorders, outcome expectancies, school performance, anxiety, parental smoking, and peer attitudes to smoking. The studies vary in the number of factors accounted for: some (Conner *et al.*, 2018; Leventhal *et al.*, 2015; Watkins *et al.*, 2018; Wills *et al.*, 2017) considered more than 10 factors, others (Miech *et al.*, 2017; Unger *et al.*, 2016) only five. As shown in Additional File 1 (Lee, 2018), the questions used to assess these factors also varied considerably between studies.

Unfortunately, no study reported which factors contributed most to their adjustment. Thus, for example, Leventhal *et al.* (2015) adjusted for the most factors and found that the unadjusted OR of 7.78 (6.15–9.84) reduced dramatically after adjustment to 1.75 (1.10–2.78); however, the authors did not clarify which factors mainly contributed to this reduction.

While the search terms we considered may not have been fully comprehensive, it is clear from the material considered that a wide range of risk factors are related to initiation of smoking and that many of them were considered rarely, if at all, in the gateway studies.

Bias in the estimated gateway effect due to failure to adjust for a relevant covariate – simple confounding

What effect might failure to adjust for a relevant predictor of smoking have on the estimated gateway effect observed between vaping and smoking? To answer this question, we consider (using the first sheet of the Excel program we developed) a hypothetical baseline population of N never-smokers, of which P_1 vape, and P_2 have the smoking predictor. P_1 and P_2 are correlated, with an odds ratio of K . We assume that, during follow up, those who neither have smoked nor have the predictor have a probability of initiating smoking of P_A and that the odds of smoking initiation are independently increased by G_E for vapers and by G_p in those with the smoking predictor.

The illustrative results in Table 2, supported by mathematical detail in Additional File 2 (Lee, 2018), demonstrate the problem. In the basic Situation 1, we set $P_1 = 0.2$, $P_2 = 0.2$, $K = 5$, $P_A = 0.05$, $G_E = 1$, and $G_p = 4$. Instead of observing the true gateway effect (G_E) of 1, we observe a spurious gateway effect of 1.633 due to the uncontrolled confounding. In Situations 2–6, G_E remains at 1, but other parameters are varied. There is a clear tendency for the confounding effect to increase with increasing K (Situation 2) and G_p (Situation 3), but varying P_A (Situation 4), P_1 (Situation 5), or P_2 (Situation 6) has less effect. Increasing G_E , with the other parameters fixed (Situation 7), increases the observed gateway effect, but the ratio of the observed to the true gateway effect is somewhat reduced, from 1.633 for $G_E = 1$ to 1.352 for $G_E = 10$.

Bias in the estimated gateway effect due to adjustment for an inaccurately measured covariate – residual confounding

While confounding is well understood by epidemiologists, and is the reason why the authors of the gateway papers made some adjustments, “residual confounding,” arising from inaccurately determining confounding variables, is rarely considered. Many statisticians have highlighted this problem of residual confounding. Almost 40 years ago, Greenland (1980) noted that “misclassification of a confounder” leads to “partial loss of ability to control confounding,” while Tzonou *et al.* (1986) later noted that “even misclassification rates as low as 10% can prevent adequate control of confounding.” Other publications (Ahlbom & Steineck, 1992; Fewell *et al.*, 2007; Greenland & Robins, 1985; Phillips & Smith, 1994; Savitz & Barón, 1989) show that if X is an inaccurately measured true cause of disease, and Y , precisely measured, is not a cause but is correlated with X , one may conclude incorrectly that Y , not X , is the cause. None of the gateway papers cited in Table 1, nor Soneji *et al.* (2017), made attempts to adjust for bias from residual confounding, although all except two (Primack *et al.*, 2015; Primack *et al.*, 2016) mention possible bias from incomplete adjustment.

Table 3 (see also Additional File 2 (Lee, 2018)) gives illustrative examples of the effect of residual confounding. Situations 1 to 4 concern misclassification occurring in each direction, and correspond to situations 1, 2(a), 3(a), and 7(a) in Table 2, where no misclassification is assumed. Whereas in Table 2, the

Table 2. Bias in the gateway effect due to failure to adjust for the smoking predictor.

Situation	Proportion of vapers in never-smokers (P_1)	Proportion with predictor in never-smokers (P_2)	Odds ratio relating vaping and predictor (K)	Probability of smoking in non-vapers where the predictor is absent (P_A)	True OR of smoking for predictor (G_P)	True gateway effect (G_E)	Observed gateway effect
1 – basic	0.2	0.2	5	0.05	4	1	1.633
2 (a)	0.2	0.2	2	0.05	4	1	1.233
(b)			3		4	1	1.399
(c)			10		4	1	1.980
3 (a)	0.2	0.2	5	0.05	2	1	1.266
(b)					7	1	1.983
(c)					10	1	2.218
4 (a)	0.2	0.2	5	0.02	4	1	1.644
(b)				0.1	4	1	1.614
(c)				0.2	4	1	1.573
5 (a)	0.1	0.2	5	0.05	4	1	1.658
(b)	0.3				4	1	1.599
(c)	0.5				4	1	1.520
6 (a)	0.2	0.1	5	0.05	4	1	1.438
(b)		0.3			4	1	1.666
(c)		0.5			4	1	1.510
7 (a)	0.2	0.2	5	0.05	4	2	3.149
(b)					4	4	5.956
(c)					4	10	13.522

Table 3. Residual confounding due to misclassification of the smoking predictor.

Situation	Odds ratio relating vaping and predictor (K)	True OR of smoking for predictor (G _P)	True RR gateway Effect (G _E)	Observed gateway effect						
				Unadjusted	Adjusted for smoking predictor, with misclassification rate of					
					1%	2%	5%	10%	20%	50%
				Misclassification is in both directions						
1	5	4	1	1.633	1.034	1.067	1.157	1.281	1.455	1.633
2	2	4	1	1.233	1.013	1.026	1.060	1.106	1.169	1.233
3	5	2	1	1.266	1.015	1.029	1.068	1.120	1.193	1.266
4	5	4	2	3.149	2.058	2.115	2.270	2.486	2.802	3.149
				Only false positives						
5	5	4	1	1.633	1.030	1.058	1.136	1.246	1.408	1.625
				Only false negatives						
6	5	4	1	1.633	1.005	1.010	1.028	1.062	1.144	1.414

Notes: A misclassification rate of x% in both directions (considered in situations 1 to 4) implies that x% of those who have the factor that predicts smoking (e.g. risk takers) are wrongly classified as not having it, and x% of those who do not have the factor are wrongly classified as having it. Situations 5 and 6 consider misclassification in one direction. All models have $P_1 = P_2 = 0.2$ and $P_A = 0.5$.

smoking predictor is assumed to be measured accurately, and adjustment for it would correct the observed gateway effect back to its true value (G_E); this is not true when misclassification is present. Thus, in Situation 1, where G_E is set at 1, adjustment for the misclassified predictor does not fully correct for the confounding. Adjustment is useless where the misclassification rate is 50% with the observed gateway effect staying at its unadjusted value of 1.633. With a 10% misclassification rate, the value reduces to 1.281 so that almost half ($281/633 = 44.4\%$) of the spurious increase in the OR remains. As misclassification rates reduce, the adjusted rate approaches its true value.

The pattern is similar in Situations 2 and 3, where G_E remains at 1, but K and R_p are varied. Again, all of the bias remains with 50% misclassification, and almost half remains with 10% misclassification. This is also observed in Situation 4, where $G_E = 2$. Situations 5 and 6 are similar to Situation 1, except that misclassification is assumed to be unidirectional. The bias is similar to that observed in Situation 1, where only false positives occur (Situation 5), but is less where only false negatives are present (Situation 6). Thus poor specificity is more important than poor sensitivity as a source of bias.

How might smoking prevalence be affected by any true effects of vaping?

It is important to understand how vaping might affect the prevalence of cigarette smoking. Not only might there be gateway effects,

with vaping increasing the probability of subsequent smoking, but vaping by smokers might also increase their probability of smoking cessation. In an attempt to gain some understanding, we used the second sheet of the Excel program we developed (see also the mathematical details in Additional File 3 (Lee, 2018)). The program considers 10,000 individuals, 2,000 in each of five strata that represent groups with varying smoking susceptibility. All individuals start as never users of either product and over time may switch to become current users of either product only, current dual users, or former users of either or both products (who currently use neither).

Where there is no effect of vaping on initiation or cessation, the user may vary the values of P_1 , R_1 , R_2 , R_3 and R_4 as defined in the methods section. These parameters are then used to update tobacco use status over time. Where there is an effect of vaping, the user may define G_1 , the gateway effect, and G_2 , the effect on the probability of smoking cessation.

Table 4 presents illustrative examples where G_1 and G_2 are set at 1.0. The five parameters P_1 , R_1 , R_2 , R_3 and R_4 are varied in turn, with the others held constant. Increasing e-cigarette initiation rates by increasing P_1 (Block 1) reduces never-users, with a compensating increase in dual users and in the odds ratio relating current vaping to current smoking. In Block 1, it is assumed that there is no quitting or re-initiation ($R_3 = R_4 = 0$), the relative odds of initiation for successive

Table 4. Effect of varying parameters on tobacco use and on the odds ratio relating vaping and smoking.
All models assume G_1 and G_2 are 1.

Block	Parameters					Percentage distribution of tobacco use					Odds ratio relating current vaping and smoking
	P_1	R_1	R_2	R_3	R_4	Never use	E-cigarettes only	Cigarettes only	Dual use	Former use	
1	0.0001	5	1	0	0	88.0	5.3	5.3	1.4	0.0	4.52
	0.0002	5	1	0	0	80.6	7.6	7.6	4.3	0.0	5.93
	0.0005	5	1	0	0	69.3	9.1	9.1	12.5	0.0	10.44
	0.001	5	1	0	0	60.8	9.2	9.2	20.9	0.0	15.05
	0.002	5	1	0	0	52.1	9.3	9.3	29.4	0.0	17.85
2	0.0002	3	1	0	0	95.5	2.2	2.2	0.1	0.0	2.67
	0.0002	10	1	0	0	59.2	6.8	6.8	27.2	0.0	34.65
3	0.0002	5	0.25	0	0	85.6	10.5	2.3	1.3	0.0	4.83
	0.0002	5	0.5	0	0	83.9	9.4	4.3	2.5	0.0	5.19
	0.0002	5	2	0	0	75.7	5.3	12.5	6.5	0.0	7.43
	0.0002	5	4	0	0	69.5	3.2	18.7	8.7	0.0	10.15
4	0.0002	5	1	0.1	0	80.6	7.5	7.5	4.0	0.4	5.77
	0.0002	5	1	0.2	0	80.6	7.4	7.4	3.9	0.7	5.62
	0.0002	5	1	0.5	0	80.6	6.8	6.8	2.7	1.7	5.25
	0.0002	5	1	1.0	0	80.6	6.8	6.8	2.7	3.2	4.79
5	0.0002	5	1	0.2	0.1	80.6	7.4	7.4	3.9	0.7	5.61
	0.0002	5	1	0.2	0.2	80.6	7.4	7.4	3.9	0.7	5.60
	0.0002	5	1	0.2	0.5	80.6	7.4	7.5	3.9	0.6	5.58

P_1 , initiation rate of vaping in stratum 1; R_1 , relative odds of initiation for successive strata; R_2 , relative odds of initiation for smoking compared to vaping; R_3 , relative odds of quitting compared to initiation; R_4 , relative odds of re-initiation compared to initiation.

strata (R_1) are set at 5, and the initiation rates are assumed the same for both tobacco products ($R_2 = 1$). Increasing R_1 (Block 2), which increases the between stratum variation in smoking susceptibility, markedly increases the odds ratio relating current vaping to current smoking. The other Blocks keep P_1 at 0.0002 and R_1 at 5. Increasing R_2 (Block 3) increases the frequency of smoking relative to vaping, the frequency of dual use and the odds ratio relating current vaping to current smoking. Increasing

the quitting factor R_3 (Block 4) reduces smokers, but the odds ratio is less affected. Furthermore, increasing the re-initiation factor R_4 (Block 5) has little effect; in fact, the chance of someone initiating, quitting, then re-initiating during follow-up is small for the given parameter values.

Table 5, Table 6 and Table 7 show how the relative odds of smoking are affected by variations, respectively, in G_1 only,

Table 5. Effect of varying G_1 and R_2 on smoking prevalence and the relative odds of smoking.

G_1	R_2	% cigarette smokers	Relative odds of smoking
1	1	11.28	1 (base)
2	1	12.31	1.10
3	1	13.03	1.18
4	1	13.56	1.23
5	1	13.93	1.28
1	0.25	3.56	1 (base)
5	0.25	5.23	1.50
1	0.5	6.56	1 (base)
5	0.5	8.93	1.40
1	2	17.37	1 (base)
5	2	19.66	1.16
1	2	23.22	1 (base)
5	4	24.79	1.09

G_1 , is the gateway effect; R_2 , the relative odds of initiation for smoking compared to vaping. The assumed values of the other parameters are $P_1 = 0.0002$, $R_1 = 5$, $R_3 = 0.2$, $R_4 = 0$, and $G_2 = 1$. See text for the full definitions of these other parameters.

Table 6. Effect of varying G_2 , R_2 , and R_3 on % smokers and relative smoking odds.

Odds are expressed relative to the situation where G_1 and $G_2 = 1$.

G_2	R_2	R_3	% smokers	Relative odds of smoking
1	1	0.2	11.28	1 (base)
2	1	0.2	11.16	0.99
3	1	0.2	11.06	0.98
4	1	0.2	10.96	0.97
5	1	0.2	10.87	0.96
1	1	0.5	11.28	1 (base)
5	1	0.5	9.82	0.93
1	0.25	0.2	3.56	1 (base)
5	0.25	0.2	3.52	0.99
1	0.5	0.2	6.56	1 (base)
5	0.5	0.2	6.42	0.98
1	2	0.2	17.37	1 (base)
5	2	0.2	16.40	0.93
1	4	0.2	23.22	1 (base)
5	4	0.2	21.55	0.91

G_2 is the effect on the probability of smoking cessation. R_2 is the relative odds of initiation for smoking compared to vaping. R_3 is the relative odds of quitting compared to initiation. The assumed values of the other parameters are $P_1 = 0.0002$, $R_1 = 5$, $R_4 = 0$, and $G_1 = 1$. See text for the definitions of these other parameters.

Table 7. Odds of smoking according to variation in G_1 , G_2 , R_2 , and R_3 .

R_2	R_3	G_1	1	1	1	2	2	2	5	5	5
R_2	R_3	G_2	1	2	5	1	2	5	1	2	5
0.25	0.2		1	0.997	0.989	1.143	1.140	1.129	1.497	1.491	1.473
0.25	0.5		1	0.993	0.975	1.136	1.128	1.104	1.473	1.459	1.421
0.5	0.2		1	0.994	0.978	1.129	1.121	1.101	1.398	1.387	1.358
0.5	0.5		1	0.987	0.955	1.123	1.107	1.066	1.383	1.359	1.301
1	0.2		1	0.989	0.960	1.104	1.091	1.056	1.276	1.259	1.217
1	0.5		1	0.976	0.926	1.102	1.073	1.012	1.272	1.236	1.160
2	0.2		1	0.979	0.933	1.072	1.050	0.999	1.164	1.140	1.086
2	0.5		1	0.961	0.897	1.073	1.031	0.959	1.169	1.124	1.046
4	0.2		1	0.967	0.908	1.043	1.011	0.952	1.090	1.059	1.002
4	0.5		1	0.952	0.890	1.048	0.999	0.936	1.103	1.055	0.990

G_1 is the gateway effect. G_2 is the effect on the probability of smoking cessation. R_2 is the relative odds of initiation for smoking compared to vaping. R_3 is the relative odds of quitting compared to initiation. The assumed values of the other parameters are $P_1 = 0.0002$, $R_1 = 5$, and $R_4 = 0$. See text for the definitions of these other parameters.

G_2 only, or both. All these results set $P_1 = 0.0002$, $R_1 = 5$, and $R_4 = 0$. With G_1 and G_2 set at 1.0, there are 11.28% cigarette smokers at the end of follow up: 7.43% smoking cigarettes only, and 3.85% dual users.

As G_1 increases to 5 (Table 5), the percentage of cigarette smokers rises to 13.96%, giving an OR for smoking of 1.28 compared with $G_1 = 1$. The increase in the percentage of smokers is proportionally much less than the increase in G_1 . Table 5 also shows that the gateway effect becomes more important as the relative initiation rate of cigarettes compared with that of e-cigarettes (R_2) decreases.

As G_2 increases from 1 to 5 (Table 6), the percentage of smokers declines. The effect is less than for increasing G_1 , and it is in the opposite direction. This effect is increased as R_3 , the relative frequency of quitting to initiation, is increased. Table 6 also shows that G_2 becomes more important as R_2 increases.

In Table 7, both G_1 and G_2 are varied, with the relative odds of smoking shown for combinations of G_1 and $G_2 = 1, 2$, or 5 and for varying values of R_2 and R_3 . The highest relative odds of smoking (1.497) are seen for the highest values of G_1 and R_2 and the lowest values of G_2 and R_3 , while the lowest odds of smoking (0.890) are seen in the reverse situation.

The effects sometimes approximately cancel out. Given $R_3 < 1$ (and it seems unlikely that quit rates would exceed initiation rates), effects of varying G_1 tend to exceed effects of varying G_2 where initiation rates are similar for vaping and smoking. However, where initiation rates for smoking are substantially higher ($R_2 = 4$), effects of similar G_1 and G_2 values also approximately cancel out. Here, with relatively more smokers to quit, effects of varying G_2 are more relevant.

While the above results are illustrative, one can derive five main general conclusions from them:

1. Large increases in G_1 , the gateway effect, result in proportionately much smaller increases in the smoking prevalence.
2. Gateway effects increase if initiation with vaping is more frequent than initiation with smoking.
3. Increasing G_2 decreases smoking prevalence, but less than that from a similar increase in G_1 .
4. Effects of varying G_2 increase as R_2 and R_3 increase, where there are more smokers who can quit.
5. With both G_1 and $G_2 > 1$, the overall effect on smoking prevalence may be in either direction.

Has introducing e-cigarettes affected smoking trends?

Smoking prevalence in U.S. youths had declined substantially even before vaping became popular. For example, the YRBS reported a decline in past 30-day smoking from 34.8% in 1995 to 15.7% in 2013. Has introducing e-cigarettes halted or even reversed this decline?

The NYTS is the only U.S. youth survey providing trend data on e-cigarette use over a reasonably long time period. In that survey, the prevalence of past 30-day vaping in each year from 2011–2015 was 1.0%, 2.0%, 2.9%, 9.2%, and 11.1%, respectively, for students in grades 6–12. In 2011–2013, vaping seems too rare to allow for assessment of any effect on smoking prevalence. To detect any effect, it seems more appropriate to compare the prevalence in 2014–2016 with that predicted from pre-existing trends.

Table 8 presents results from this comparison based on the U.S. NYTS, YRBS, MTF, and NSDUH surveys. All estimates are for past 30-day smoking. In all these surveys, smoking prevalence in 2014–2016 was less than predicted from the underlying trend, contrary to expectations of a definitive gateway effect being present. The tendency for the decline in prevalence to accelerate over 2014–2016 is evident regardless of sex or age. Additional File 4 (Lee, 2019) provides further detail, including the observed data for the years before 2014, the fitted linear regression of $\log(p/(1-p))$ on year for years up to 2013, and graphical illustration that this linear regression model fits the data well.

The Smokefree Youth Survey in Great Britain also reported increasing vaping, with percentages of 4%, 6%, 11%, and 10% each year from 2013–2016, respectively, for 11–18-year-olds. As shown in Table 8, ONS data (for an older age group) also show no tendency for a rise in cigarette smoking given increasing vaping. Here, very small annual declines (about 0.3% in males and 0.7% in females) over 2006–2015 were followed by a much larger decline of 7% between 2015 and 2016.

These analyses, though clearly limited by the fact that we do not know what smoking prevalence would have been in the absence of e-cigarettes, do not suggest that introducing e-cigarettes has had any material adverse effect on smoking prevalence trends. If there were any gateway effect, it would be clearly outweighed by other issues, such as vaping providing an alternative to cigarettes for tobacco users, or changes in attitudes to smoking resulting from anti-tobacco prevention measures.

Additional evidence on the likelihood of any marked gateway effect on smoking prevalence

Some additional evidence highlights the improbability of any substantial gateway effect of vaping on smoking prevalence.

Based on the U.S. PATH study, Pearson *et al.* (2018) recently reported that only a small percentage of nonsmokers would be interested in using a hypothetical modified risk tobacco product. While these analyses were based on adults, it was possible to confirm this finding from the publicly available data files for 18–24-year-olds. Thus, the proportion “very or somewhat likely” to use such a product was much lower in those who had never used tobacco ($57/1745 = 3.3\%$) than in those who had ever done so ($2375/7282 = 32.7\%$). This difference was similar in both sexes. Those aged 12–17 years old were also asked in the PATH study if they had seen a tobacco product that claims to be safer than other tobacco products in the past 12 months

and their likelihood of using such a product in the next 30 days. Again, the proportion answering “very or somewhat likely” was much lower in tobacco never-users ($174/4673 = 3.72\%$) than in ever-users ($264/1565 = 16.87\%$).

Based on the Canadian COMPASS study, [Aleyan et al. \(2018\)](#) compared rates of smoking initiation over a two-year follow-up period among a sample of 9,501 students between grades 9 and 11 (cigarette never-smokers), classified at baseline into four groups by current e-cigarette use and susceptibility to smoke and assessed using a three-item validated measure. Though the data in Figure 1 of that paper showed that in both unsusceptible and susceptible never-smokers, rates of smoking initiation were higher in current than noncurrent e-cigarette users, only an estimated $33/2646 = 1.2\%$ of those who had tried smoking during the follow-up period were unsusceptible current e-cigarette users.

While evidence on the reported likelihood of future use of a new product or on susceptibility to smoking is less robust than that based on actual initiation of smoking, the results from

these studies are consistent with the conclusion that even when there is some gateway effect, any resulting increase in smoking prevalence would be quite small.

How might any true gateway effect modify the population health impact of introducing e-cigarettes?

It is possible for users of alternative tobacco products to have similar nicotine exposure to that of cigarette smokers but potentially much lower risk of smoking-related diseases (SRD). This is illustrated by the experience and prevalence of Swedish snuff (“snus”) use ([Agewall et al., 2002](#); [Bolinder et al., 1997a](#); [Bolinder et al., 1997b](#); [Lee, 2011](#); [Lee, 2013](#)). As vaping presents significantly reduced exposure to toxicants and harmful and potentially harmful constituents compared with smoking ([National Academies of Sciences Engineering and Medicine, 2018](#)), it would be expected that any harmful effects would be much lower. This hypothesis was subsequently endorsed by an expert group in 2014 ([Nutt et al., 2014](#)).

This hypothesis is also further supported by various theoretical beneficial and adverse effects of vaping (see [Table 9](#)). The first

Table 8. Observed smoking percentages in five studies compared to predictions from previous trends^a.

Study ^b	Age/grade	Observed/predicted	2014	2015	2016
NYTS	6th–12th grade	Observed	6.2	6.1	-
		Predicted	8.22	7.77	-
MTF	8th grade	Observed	4.0	3.6	2.6
		Predicted	4.42	4.07	3.75
MTF	10th grade	Observed	7.2	6.3	4.9
		Predicted	9.56	9.03	8.53
MTF	12th grade	Observed	13.6	11.4	10.5
		Predicted	16.00	15.33	14.68
YRBS	9th–12th grade	Observed	-	11.8	-
		Predicted	-	14.58	-
YRBS	9th–12th grade	Observed	-	9.7	-
		Predicted	-	12.86	-
NSDUH	12–17 years	Observed	5.1	4.6	3.8
		Predicted	5.29	4.68	4.14
NSDUH	12–17 years	Observed	4.6	3.8	3.1
		Predicted	4.99	4.43	3.94
ONS	16–24 years	Observed	25.2	24.1	17.2
		Predicted	23.0	22.7	22.4
ONS	16–24 years	Observed	20.9	22.9	16.0
		Predicted	21.3	20.6	20.0

^aPrevious years used for estimation of trend: NYTS: 2004, 2006, 2009, 2011–2013; MTF, ONS: 2006, 2007, 2008, 2009, 2010, 2011, 2012, 2013; YRBS: 1995, 1997, 1999, 2001, 2003, 2005, 2007, 2009, 2011, 2013; NSDUH: 2009, 2010, 2011, 2012, 2013. ^bMTF, Monitoring the Future; NSDUH, National Survey on Drug Use and Health; NYTS, National Youth Tobacco Survey; ONS, Office for National Statistics; YRBS, Youth Risk Behavior Survey.

Table 9. Theoretical adverse and beneficial effects of e-cigarettes.

	Effect	Adverse or beneficial
A1	Vaping encourages initiation of smoking (gateway)	Adverse
A2	Smokers intending to quit switch to vaping instead	Adverse
A3	Smokers vape in addition to their normal cigarette consumption	Adverse
B1	Individuals who would otherwise have smoked vape instead	Beneficial
B2	Smokers who would have continued to smoke switch instead to vaping	Beneficial
B3	Vaping helps established smokers to quit	Beneficial
B4	Vaping helps established smokers to materially reduce cigarette consumption	Beneficial

major benefit of vaping (B1) concerns individuals who, in the absence of e-cigarettes, would have initiated smoking but instead take up vaping. They should have a much lower risk of SRDs than if they had smoked, given, for example, that the expert panel (Nutt *et al.*, 2014) estimated that relative to cigarettes, e-cigarettes likely pose a reduction in harm of up to 95%. This likely reduction in risk and harm also applies to benefit B2, where smokers who would otherwise have continued to smoke instead switch to vaping. While the benefits would take longer to emerge, as the adverse effects of past smoking need time to disappear, the reduction in risk associated with a switch to vaping should be a considerable proportion of that associated with quitting smoking. The health benefit would even be greater for established smokers, where vaping helps them quit (B3).

Effects A3 and B4 in Table 9 both relate to those who switch from cigarettes to dual use of cigarettes and e-cigarettes. In theory, adverse effects might arise if smokers vape in addition to their usual cigarette consumption (A3), but this is probably implausible, because most dual users are likely to control their nicotine intake and actually reduce their cigarette consumption (B4), partially replacing cigarettes with e-cigarettes (Berry *et al.*, 2018; McNeill *et al.*, 2014; McRobbie *et al.*, 2014). Note that we use the word “materially” in the definition of effect B4 as theoretically, an adverse health impact might occur if a very small reduction in cigarettes is counterbalanced by a large increase in use of e-cigarettes.

While smokers intending to quit will do worse if they switch to e-cigarettes than if they had quit (A2), they will still do better than if they had continued to smoke.

Clearly, the greatest adverse effect arises for individuals who otherwise would not have smoked but are encouraged by vaping to do so (A1). However, considering the overall population health impact of this gateway effect, two points require emphasis. First, this is only likely to affect young people. Older people who decided not to start tobacco product use seem unlikely to start vaping, and even if they did so, it would probably be because they believe vaping to be much safer than the smoking they have already rejected (Majeed *et al.*, 2017; Popova *et al.*, 2018; Xu *et al.*, 2016; Zhu *et al.*, 2013). Second, the arguments made earlier suggest that the number of new smokers originating from any true gateway-in effect should be quite low. Any adverse

effects resulting from a gateway effect in young people (adverse effect A1) are likely to be outweighed in the general population by beneficial effect B1, as evidenced by the substantial numbers who have vaped but not smoked. The reduction in risk of SRDs for such individuals is likely to be almost as great as any increase resulting from a gateway effect.

Discussion

Our analyses yield four major conclusions:

1. While the consistently increased adjusted gateway effect estimates appear compelling (Table 1), they may be severely biased by ignoring established predictors of smoking initiation and residual confounding – a true causal effect remains to be demonstrated.
2. If a true gateway effect were to exist, it would probably have little effect on smoking prevalence.
3. No available evidence exists that increasing e-cigarette use has slowed the decline in smoking prevalence; indeed, the decline appears to have accelerated.
4. Finally, introducing e-cigarettes should lead to a reduced risk of SRD.

Though some publications have used the evidence for a gateway effect in youths to advocate for regulations to curb vaping in the general population (Miech *et al.*, 2017; Primack *et al.*, 2015; Soneji *et al.*, 2017), others suggest that this over-interprets the evidence. Kozlowski & Warner (2017) point out the need “to better understand and assess confounding variables” and consider that the prospective studies merely “support that a minority of the relatively small number of e-cigarette triers—who haven’t also been experimenting with other tobacco products already—will go on to some experimentation with cigarettes.” Also, Levy *et al.* (2018) present calculations showing that “a strategy of replacing cigarette smoking with vaping would yield substantial life year gains, even under pessimistic assumptions [. . .].” Our argument that the gateway estimates are subject to relevant uncontrolled confounding is also supported by publications showing that vaping is associated with many factors associated with smoking (Barrington-Trimis *et al.*, 2015; Hanewinkel & Isensee, 2015; Temple *et al.*, 2017).

Unlike the implicit conclusion of the National Academies report (National Academies of Sciences Engineering and Medicine, 2018), which found that control of confounding in the gateway-in studies has been adequate, our conclusions would suggest otherwise. Our conclusion is consistent with those of a recent publication (Lee & Fry, 2019) which, based on data from the PATH study, found that making extensive adjustment for other risk factors, as well as limited adjustment for inaccuracy in some of these, explained as much as 87% of the observed gateway effect.

Our calculations indicating a lack of effect of e-cigarettes on smoking trends are further supported by Dutra & Glantz (2017), who concluded that “the introduction of e-cigarettes was not associated with a change in the linear decline in cigarette smoking among youth.” While studies (McNeill *et al.*, 2014) have argued against the likely population benefit of e-cigarettes due to their pervasive use among youths and young adults, as we have discussed, these analyses often fail to fully consider the limitations and weaknesses of the current evidence of the associated probabilities of e-cigarette use and cigarette smoking initiation in youths.

While various authors (Hill & Camacho, 2017; Levy *et al.*, 2018; Nutt *et al.*, 2014) estimate that the introduction of e-cigarettes will have a beneficial population health impact, a recent publication Soneji *et al.* (2018) argues the opposite, presenting analyses estimating that “e-cigarette use in 2014 would lead to 1,510,000 years of life lost in the U.S. population.” The study reported that the large adverse effects from an estimated 168,000 cigarette never-smokers between 12 and 29 years old initiating cigarette smoking in 2015 and going on to become daily smokers would heavily outweigh the very small beneficial effects from an additional 2,070 current cigarette smoking adults aged 25–69 quitting smoking in 2015 and remaining abstinent for seven or more years.

These analyses have various weaknesses. First, they assume that there is no reduction in risk of SRDs for cigarette smokers who become dual users but do not quit cigarettes. Dual users are likely to reduce cigarette consumption (Brose *et al.*, 2015; Etter & Bullen, 2014; Farsalinos *et al.*, 2016; Farsalinos *et al.*, 2017; McRobbie *et al.*, 2014) and hence their risk of SRD. Second, their analysis showing large adverse effects in youths relies heavily on the estimate of the gateway effect given by Soneji *et al.* (2017), which may largely result from confounding. Third, their analysis showing very small benefits from increased smoking quit rates in adults depended on an estimate derived from a prior review and meta-analysis by Kalkhoran & Glantz (2016) that estimated the OR of quitting among smokers with an interest in quitting. The estimate, and indeed the whole meta-analysis, has since drawn strong criticism from experts in the field for being inaccurate and misleading (West *et al.*, 2016).

Some limitations of our work and of the available evidence must be noted. First, many risk factors for smoking

initiation are correlated, and had even more been adjusted for, this might not have significantly changed the adjusted effect estimates. Second, while we suggest that residual confounding may be important, we have not considered data on how inaccurately specific risk factors are measured. However, nor has anyone else in this context, as far as we are aware. Third, we have not considered the likely persistence of any true gateway effects. Some youths who initiate tobacco use with vaping may try cigarettes but prefer vaping in the long run. Fourth, we did not consider evidence (Doran *et al.*, 2017; Leventhal *et al.*, 2016) that vaping may be associated with heavier smoking, although this factor may also be subject to confounding problems. Finally, our evidence on trends is based on data where the prevalence of vaping is little more than 10%. Where vaping is infrequent, any true gateway effect will only slightly increase smoking prevalence, as never smokers who have not vaped will far outweigh those who have. Any gateway effect will be greater if vaping is more common, so it is important to continue to monitor smoking and e-cigarette use trends to provide more conclusive insights.

Conclusions

The existence of any true gateway effect has not been clearly demonstrated. Even if it does exist, and subsequently, some individuals who otherwise would not have done so start to smoke cigarettes, any effect on smoking prevalence will be quite small, and any overall population health impact of introducing e-cigarettes still likely be beneficial.

Data availability

Additional File 1. Confounding factors considered by studies of vaping as a possible gateway to smoking. This .docx file lists those factors with published evidence of a relationship with smoking and gives those factors not considered in any of the 15 studies considered in Table 1.

Additional File 2. Estimating the effects of omission of a confounding variable, or misclassification of it, on the association between vaping and subsequent initiation of smoking. This .docx file presents details of the methodology used.

Additional File 3. How might the prevalence of smoking be affected by any true gateway effects of vaping? This .docx file presents details of the methodology used.

Additional File 4. Comparing smoking prevalence in 2014–2016 with that predicted from preceding years. For each of the 10 datasets considered in Table 8, this .xlsx file gives the model fitted to the prevalence in the years preceding 2014, as well as the observed and predicted prevalence, with graphs illustrating the appropriateness of the model and the tendency for the observed prevalence in 2014–2016 to be lower than predicted.

Gateway calcs2. This .xlsx file is the Excel program used to produce the results presented in Additional Files 2 and 3.

All data are available on OSF, DOI: <https://doi.org/10.17605/OSF.IO/Z3ST5> (Lee, 2018).

Data are available under the terms of the [Creative Commons Zero “No rights reserved” data waiver](#) (CC0 1.0 Public domain dedication).

Author contributions

PNL conceived the paper, developed the analyses reported in Additional Files 2 and 3, and drafted the manuscript. KJC discussed the paper with PNL, searched for relevant literature, checked tables, and commented on drafts. EFA was responsible for the work described in Additional Files 1 and 4, collation of data

used in [Table 8](#), commented on drafts, and contributed to the writing of the manuscript.

Grant information

This work was supported by PMI R&D, Philip Morris Products S.A., Quai Jeanrenaud 5, CH-2000 Neuchâtel, Switzerland.

Acknowledgements

We thank Mrs. Y. Cooper and Mrs. D. Morris for typing various drafts of this paper and assembling relevant literature. We also thank Ms. J. Hamling for assistance in preparing the figures in Additional File 4, as well as reviewers and colleagues for helpful comments and Philip Morris International for financial support.

References

- Agewall S, Persson B, Lindstedt G, *et al.*: **Smoking and use of smokeless tobacco in treated hypertensive men at high coronary risk: utility of urinary cotinine determination.** *Br J Biomed Sci.* 2002; **59**(3): 145–149.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Ahlbom A, Steineck G: **Aspects of misclassification of confounding factors.** *Am J Ind Med.* 1992; **21**(1): 107–112.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Aleyan S, Cole A, Qian W, *et al.*: **Risky business: a longitudinal study examining cigarette smoking initiation among susceptible and non-susceptible e-cigarette users in Canada.** *BMJ Open.* 2018; **8**(5): e021080.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Amato MS, Boyle RG, Levy D: **E-cigarette use 1 year later in a population-based prospective cohort.** *Tob Control.* 2016; **26**(e2): e92–e96.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Ambrose B: **E-cigarette use transitions: a case study from Waves 1 & 2 of the PATH study.** In: *Proceedings from 2017 SRNT Annual Meeting. Pre-Conference Workshop 6 - FDA's Population Health Standard: Balancing the Risks and Benefits in Regulatory Decision-making*, Florence, Italy, 2017; 29.
[Reference Source](#)
- Barrington-Trimis JL, Berhane K, Unger JB, *et al.*: **Psychosocial Factors Associated With Adolescent Electronic Cigarette and Cigarette Use.** *Pediatrics.* 2015; **136**(2): 308–17.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Barrington-Trimis JL, Uman R, Berhane K, *et al.*: **E-Cigarettes and Future Cigarette Use.** *Pediatrics.* 2016; **138**(1): pii: e20160379.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Berry KM, Reynolds LM, Collins JM, *et al.*: **E-cigarette initiation and associated changes in smoking cessation and reduction: the Population Assessment of Tobacco and Health Study, 2013-2015.** *Tob Control.* 2018; pii: tobaccocontrol-2017-054108.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Best C, Haseen F, Currie D, *et al.*: **Relationship between trying an electronic cigarette and subsequent cigarette experimentation in Scottish adolescents: a cohort study.** *Tob Control.* 2018; **27**(4): 373–378, pii: tobaccocontrol-2017-053691.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Bold KW, Kong G, Cavallo DA, *et al.*: **Reasons for Trying E-cigarettes and Risk of Continued Use.** *Pediatrics.* 2016; **138**(3): pii: e20160895.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Bolinder G, Norén A, de Faire U, *et al.*: **Smokeless tobacco use and atherosclerosis: an ultrasonographic investigation of carotid intima media thickness in healthy middle-aged men.** *Atherosclerosis.* 1997a; **132**(1): 95–103.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Bolinder G, Norén A, Wahren J, *et al.*: **Long-term use of smokeless tobacco and physical performance in middle-aged men.** *Eur J Clin Invest.* 1997b; **27**(5): 427–433.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Brose LS, Hitchman SC, Brown J, *et al.*: **Is the use of electronic cigarettes while smoking associated with smoking cessation attempts, cessation and reduced cigarette consumption? A survey with a 1-year follow-up.** *Addiction.* 2015; **110**(7): 1160–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Conner M, Grogan S, Simms-Ellis R, *et al.*: **Do electronic cigarettes increase cigarette smoking in UK adolescents? Evidence from a 12-month prospective study.** *Tob Control.* 2018; **27**(4): 365–372, pii: tobaccocontrol-2016-053539.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Coogan PF, Adams M, Geller AC, *et al.*: **Factors associated with smoking among children and adolescents in Connecticut.** *Am J Prev Med.* 1998; **15**(1): 17–24.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Cooper M, Case KR, Loukas A, *et al.*: **E-cigarette Dual Users, Exclusive Users and Perceptions of Tobacco Products.** *Am J Health Behav.* 2016; **40**(1): 108–16.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Dalton MA, Sargent JD, Beach ML, *et al.*: **Effect of viewing smoking in movies on adolescent smoking initiation: a cohort study.** *Lancet.* 2003; **362**(9380): 281–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Daly KA, Lund EM, Harty KC, *et al.*: **Factors associated with late smoking initiation in Minnesota women.** *Am J Public Health.* 1993; **83**(9): 1333–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Dautzenberg B, de Souza Moura MA, Rieu N, *et al.*: **[The e-cigarette disrupts other consumptions in Parisian teenagers (2012-2014)].** *Rev Mal Respir.* 2016; **33**(3): 225–34.
[PubMed Abstract](#) | [Publisher Full Text](#)
- de Lacy E, Fletcher A, Hewitt G, *et al.*: **Cross-sectional study examining the prevalence, correlates and sequencing of electronic cigarette and tobacco use among 11-16-year olds in schools in Wales.** *BMJ Open.* 2017; **7**(2): e012784.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- de Leon J, Diaz FJ, Rogers T, *et al.*: **Initiation of daily smoking and nicotine dependence in schizophrenia and mood disorders.** *Schizophr Res.* 2002; **56**(1–2): 47–54.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Doran N, Brikmanis K, Petersen A, *et al.*: **Does e-cigarette use predict cigarette escalation? A longitudinal study of young adult non-daily smokers.** *Prev Med.* 2017; **100**: 279–284.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Dutra LM, Glantz SA: **Electronic cigarettes and conventional cigarette use among U.S. adolescents: a cross-sectional study.** Erratum appears in JAMA 2014; **168** (7): 684. *JAMA Pediatr.* 2014; **168**(7): 610–617.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Dutra LM, Glantz SA: **E-cigarettes and National Adolescent Cigarette Use: 2004-2014.** *Pediatrics.* 2017; **139**(2): pii: e20162450.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Eastwood B, Dockrell MJ, Arnott D, *et al.*: **Electronic cigarette use in young people in Great Britain 2013-2014.** *Public Health.* 2015; **129**(9): 1150–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Ernst M, Luckenbaugh DA, Moolchan ET, *et al.*: **Decision-making and facial emotion recognition as predictors of substance-use initiation among adolescents.** *Addict Behav.* 2010; **35**(3): 286–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Etter JF, Bullen C: **A longitudinal study of electronic cigarette users.** *Addict Behav.* 2014; **39**(2): 491–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Farsalinos KE, Poulas K, Voudris V, *et al.*: **Electronic cigarette use in the European Union: analysis of a representative sample of 27 460 Europeans from 28 countries.** *Addiction.* 2016; **111**(11): 2032–2040.
[PubMed Abstract](#) | [Publisher Full Text](#)

- Farsalinos KE, Poulas K, Voudris V, *et al.*: **Prevalence and correlates of current daily use of electronic cigarettes in the European Union: analysis of the 2014 Eurobarometer survey.** *Intern Emerg Med.* 2017; 12(6): 757–763.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Fewell Z, Davey Smith G, Sterne JA: **The impact of residual and unmeasured confounding in epidemiologic studies: a simulation study.** *Am J Epidemiol.* 2007; 166(6): 646–55.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Glasser A, Abudayyeh H, Cantrell J, *et al.*: **Patterns of E-Cigarette Use Among Youth and Young Adults: Review of the Impact of E-Cigarettes on Cigarette Smoking.** *Nicotine Tob Res.* 2018.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Goniewicz ML, Gawron M, Nadolska J, *et al.*: **Rise in electronic cigarette use among adolescents in Poland.** *J Adolesc Health.* 2014; 55(5): 713–715.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Greenland S: **The effect of misclassification in the presence of covariates.** *Am J Epidemiol.* 1980; 112(4): 564–569.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Greenland S, Robins JM: **Confounding and misclassification.** *Am J Epidemiol.* 1985; 122(3): 495–506.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Gritz ER, Prokhorov AV, Hudmon KS, *et al.*: **Cigarette smoking in a multiethnic population of youth: methods and baseline findings.** *Prev Med.* 1998; 27(3): 365–84.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Hammond D, Reid JL, Cole AG, *et al.*: **Electronic cigarette use and smoking initiation among youth: a longitudinal cohort study.** *CMAJ.* 2017; 189(43): E1328–E1336.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Hanewinkel R, Isensee B: **Risk factors for e-cigarette, conventional cigarette, and dual use in German adolescents: a cohort study.** *Prev Med.* 2015; 74: 59–62.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Hill A, Camacho OM: **A system dynamics modelling approach to assess the impact of launching a new nicotine product on population health outcomes.** *Regul Toxicol Pharmacol.* 2017; 86: 265–278.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Hornik RC, Gibson L, Lerman C: **Prediction of cigarette use from six month prior electronic and combustible cigarette use for a U.S. national sample of 13-25 year olds [abstract POS5-30].** *Society for Research on Nicotine & Tobacco.* 2016 Rapid Response Posters. 2016.
[Reference Source](#)
- Huang M, Hollis J, Polen M, *et al.*: **Stages of smoking acquisition versus susceptibility as predictors of smoking initiation in adolescents in primary care.** *Addict Behav.* 2005; 30(6): 1183–94.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Huh J, Leventhal AM: **Progression of Poly-tobacco Product Use Patterns in Adolescents.** *Am J Prev Med.* 2016; 51(4): 513–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Jackson C: **Cognitive susceptibility to smoking and initiation of smoking during childhood: a longitudinal study.** *Prev Med.* 1998; 27(1): 129–34.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Jordan JN, McElroy JA, Everett KD: **Smoking initiation, tobacco product use, and secondhand smoke exposure among general population and sexual minority youth, Missouri, 2011-2012.** *Prev Chronic Dis.* 2014; 11: E113.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kalkhoran S, Glantz SA: **E-cigarettes and smoking cessation in real-world and clinical settings: a systematic review and meta-analysis.** *Lancet Respir Med.* 2016; 4(2): 116–28.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kaufman AR, Land S, Parascandola M, *et al.*: **Tobacco use transitions in the United States: The National Longitudinal Study of Adolescent Health.** *Prev Med.* 2015; 81: 251–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kozlowski LT, Warner KE: **Adolescents and e-cigarettes: Objects of concern may appear larger than they are.** *Drug Alcohol Depend.* 2017; 174: 209–214.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lee PN: **Summary of the epidemiological evidence relating snus to health.** *Regul Toxicol Pharmacol.* 2011; 59(2): 197–214.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lee PN: **Epidemiological evidence relating snus to health—an updated review based on recent publications.** *Harm Reduct J.* 2013; 10(1): 36.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Lee P: **Considerations Related to Vaping as a Possible Gateway into Cigarette Smoking.** *OSF* 2019.
<http://www.doi.org/10.17605/OSF.IO/Z3ST5>
- Lee P, Fry J: **Investigating gateway effects using the PATH study [version 1; peer review: awaiting peer review].** *F1000Res.* 2019; 8: 264.
[Publisher Full Text](#)
- Lee S, Grana RA, Glantz SA: **Electronic cigarette use among Korean adolescents: a cross-sectional study of market penetration, dual use, and relationship to quit attempts and former smoking** *J Adolesc Health.* 2014; 54(6): 684–90.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Leventhal AM, Stone MD, Andrabi N, *et al.*: **Association of e-Cigarette Vaping and Progression to Heavier Patterns of Cigarette Smoking.** *JAMA.* 2016; 316(18): 1918–1920.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Leventhal AM, Strong DR, Kirkpatrick MG, *et al.*: **Association of Electronic Cigarette Use With Initiation of Combustible Tobacco Product Smoking in Early Adolescence.** *JAMA.* 2015; 314(7): 700–707.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Levy DT, Borland R, Lindblom EN, *et al.*: **Potential deaths averted in USA by replacing cigarettes with e-cigarettes.** *Tob Control.* 2018; 27(1): 18–25.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Loukas A, Batanova M, Fernandez A, *et al.*: **Changes in use of cigarettes and non-cigarette alternative products among college students.** *Addict Behav.* 2015; 49: 46–51.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Loukas A, Marti CN, Cooper M, *et al.*: **Exclusive e-cigarette use predicts cigarette initiation among college students.** *Addict Behav.* 2018; 76: 343–347.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Lozano P, Barrientos-Gutierrez I, Arillo-Santillan E, *et al.*: **A longitudinal study of electronic cigarette use and onset of conventional cigarette smoking and marijuana use among Mexican adolescents.** *Drug Alcohol Depend.* 2017; 180: 427–430.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Majeed BA, Weaver SR, Gregory KR, *et al.*: **Changing Perceptions of Harm of E-Cigarettes Among U.S. Adults, 2012-2015.** *Am J Prev Med.* 2017; 52(3): 331–338.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- McNeill A, Etter JF, Farsalinos K, *et al.*: **A critique of a World Health Organization-commissioned report and associated paper on electronic cigarettes.** *Addiction.* 2014; 109(12): 2128–34.
[PubMed Abstract](#) | [Publisher Full Text](#)
- McRobbie H, Bullen C, Hartmann-Boyce J, *et al.*: **Electronic cigarettes for smoking cessation and reduction.** *Cochrane Database Syst Rev.* 2014; (12): CD010216.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Miech R, Patrick ME, O'Malley PM, *et al.*: **E-cigarette use as a predictor of cigarette smoking: results from a 1-year follow-up of a national sample of 12th grade students.** *Tob Control.* 2017; 26(e2): e106–e111.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- National Academies of Sciences Engineering and Medicine: **Public Health Consequences of E-Cigarettes.** The National Academies Press, Washington DC. 2018.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Nutt DJ, Phillips LD, Balfour D, *et al.*: **Estimating the harms of nicotine-containing products using the MCDA approach.** *Eur Addict Res.* 2014; 20(5): 218–225.
[PubMed Abstract](#) | [Publisher Full Text](#)
- O'Connor RJ, Flaherty BP, Edwards BQ, *et al.*: **Regular smokeless tobacco use is not a reliable predictor of smoking onset when psychosocial predictors are included in the model.** *Nicotine Tob Res.* 2003; 5(4): 535–543.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Pearson JL, Johnson AL, Johnson SE, *et al.*: **Adult interest in using a hypothetical modified risk tobacco product: findings from wave 1 of the Population Assessment of Tobacco and Health Study (2013-14).** *Addiction.* 2018; 113(1): 113–124.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Phillips AN, Smith GD: **Cigarette smoking as a potential cause of cervical cancer: has confounding been controlled?** *Int J Epidemiol.* 1994; 23(1): 42–49.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Picotte DM, Strong DR, Abrantes AM, *et al.*: **Family and peer influences on tobacco use among adolescents with psychiatric disorders.** *J Nerv Ment Dis.* 2006; 194(7): 518–23.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Popova L, Owusu D, Weaver SR, *et al.*: **Affect, risk perception, and the use of cigarettes and e-cigarettes: a population study of U.S. adults.** *BMC Public Health.* 2018; 18(1): 395.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Primack BA, Shensa A, Sidani JE, *et al.*: **Initiation of cigarette smoking after e-cigarette use: a nationally representative study.** *Ann Behav Med.* 2016; 50(Suppl 1): S68.
- Primack BA, Soneji S, Stoolmiller M, *et al.*: **Progression to Traditional Cigarette Smoking After Electronic Cigarette Use Among US Adolescents and Young Adults.** *JAMA Pediatr.* 2015; 169(11): 1018–23.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Reed MB, Wang R, Shillington AM, *et al.*: **The relationship between alcohol use and cigarette smoking in a sample of undergraduate college students.** *Addict Behav.* 2007; 32(3): 449–64.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Rohde P, Lewinsohn PM, Brown RA, *et al.*: **Psychiatric disorders, familial factors and cigarette smoking: I. Associations with smoking initiation.** *Nicotine Tob Res.* 2003; 5(1): 85–98.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Sargent JD, Dalton MA, Beach M, *et al.*: **Cigarette promotional items in public schools.** *Arch Pediatr Adolesc Med.* 1997; 151(12): 1189–96.
[PubMed Abstract](#) | [Publisher Full Text](#)

- Savitz DA, Barón AE: **Estimating and correcting for confounder misclassification.** Erratum appears in American Journal of Epidemiology 1989; 130: 1260. *Am J Epidemiol.* 1989; **129**(5): 1062–1071.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Scal P, Ireland M, Borowsky IW: **Smoking among American adolescents: a risk and protective factor analysis.** *J Community Health.* 2003; **28**(2): 79–97.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Selya AS, Rose JS, Dierker L, *et al.*: **Evaluating the mutual pathways among electronic cigarette use, conventional smoking and nicotine dependence.** *Addiction.* 2018; **113**(2): 325–333.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Simons-Morton B, Crump AD, Haynie DL, *et al.*: **Psychosocial, school, and parent factors associated with recent smoking among early-adolescent boys and girls.** *Prev Med.* 1999; **28**(2): 138–48.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Soneji S, Barrington-Trimis JL, Wills TA, *et al.*: **Association Between Initial Use of e-Cigarettes and Subsequent Cigarette Smoking Among Adolescents and Young Adults: A Systematic Review and Meta-analysis.** *JAMA Pediatr.* 2017; **171**(8): 788–797.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Soneji SS, Sung HY, Primack BA, *et al.*: **Quantifying population-level health benefits and harms of e-cigarette use in the United States.** *PLoS One.* 2018; **13**(3): e0193328.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Spindle TR, Hiler MM, Cooke ME, *et al.*: **Electronic cigarette use and uptake of cigarette smoking: A longitudinal examination of U.S. college students.** *Addict Behav.* 2017; **67**: 66–72.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Sutfin EL, Reboussin BA, Debinski B, *et al.*: **The Impact of Trying Electronic Cigarettes on Cigarette Smoking by College Students: A Prospective Analysis.** *Am J Public Health.* 2015; **105**(8): e83–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Temple JR, Shorey RC, Lu Y, *et al.*: **E-cigarette use of young adults motivations and associations with combustible cigarette alcohol, marijuana, and other illicit drugs.** *Am J Addict.* 2017; **26**(4): 343–348.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Treur JL, Rozema AD, Mathijssen JJP, *et al.*: **E-cigarette and waterpipe use in two adolescent cohorts: cross-sectional and longitudinal associations with conventional cigarette smoking.** *Eur J Epidemiol.* 2018; **33**(3): 323–334.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Tzonou A, Kaldor J, Smith PG, *et al.*: **Misclassification in case-control studies with two dichotomous risk factors.** *Rev Epidemiol Sante Publique.* 1986; **34**(1): 10–17.
[PubMed Abstract](#)
- Unger JB, Soto DW, Leventhal A: **E-cigarette use and subsequent cigarette and marijuana use among Hispanic young adults.** *Drug Alcohol Depend.* 2016; **163**: 261–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
- US Surgeon General: **Preventing tobacco use among young people. A report of the Surgeon General.** US Department of Health and Human Services; Public Health Service, Atlanta, Georgia. 1994.
[Reference Source](#)
- Watkins SL, Glantz SA, Chaffee BW: **Association of Noncigarette Tobacco Product Use With Future Cigarette Smoking Among Youth in the Population Assessment of Tobacco and Health (PATH) Study, 2013-2015.** *JAMA Pediatr.* 2018; **172**(2): 181–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Weiss JW, Garbanati JA, Tanjasiri SP, *et al.*: **Effects of family functioning and self-image on adolescent smoking initiation among Asian-American subgroups.** *J Adolesc Health.* 2006; **39**(2): 221–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
- West R, Bauld L, O'Connor R, *et al.*: **Expert reaction to meta-analysis looking at e-cigarette use and smoking cessation.** Accessed: 2016 March.
[Reference Source](#)
- Westling E, Rusby JC, Crowley R, *et al.*: **Electronic Cigarette Use by Youth: Prevalence, Correlates, and Use Trajectories From Middle to High School.** *J Adolesc Health.* 2017; **60**(6): 660–666.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Wills TA, Knight R, Sargent JD, *et al.*: **Longitudinal study of e-cigarette use and onset of cigarette smoking among high school students in Hawaii.** *Tob Control.* 2017; **26**(1): 34–39.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Xu Y, Guo Y, Liu K, *et al.*: **E-Cigarette Awareness, Use, and Harm Perception among Adults: A Meta-Analysis of Observational Studies.** *PLoS One.* 2016; **11**(11): e0165938.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Zhong J, Cao S, Gong W, *et al.*: **Electronic Cigarettes Use and Intention to Cigarette Smoking among Never-Smoking Adolescents and Young Adults: A Meta-Analysis.** *Int J Environ Res Public Health.* 2016; **13**(5): pii: E465.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Zhu SH, Gamst A, Lee M, *et al.*: **The use and perception of electronic cigarettes and snus among the U.S. population.** *PLoS One.* 2013; **8**(10): e79332.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 2

Reviewer Report 02 July 2019

<https://doi.org/10.5256/f1000research.21197.r48811>

© 2019 Shahab L. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Lion Shahab

Department of Behavioural Science and Health, University College London (UCL), London, UK

The authors have addressed most of the comments. I would still suggest to include some illustrative results in the abstract. Further, the fourth conclusion ("Finally, introducing e-cigarettes should lead to a reduced risk of SRD."), as previously mentioned, is not based on primary data and has to be acknowledged to be speculative at this stage, without using more rigorous modelling (e.g. see papers by Levy *et al.* or the models included in the NASEM report) to quantify this statement.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Epidemiology, Health Psychology, Tobacco Control, Psychopharmacology, Neuroscience

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 17 Jul 2019

Peter Lee, P.N.Lee Statistics and Computing Ltd, Sutton, UK

We thank Dr Shahab for his further comments. As a result we have extensively rewritten the abstract to include some of the illustrative results he required, though the abstract limit of 300 words precluded us giving very much detail.

We have also made a number of changes to the paper to make the limitations of our work clearer, and to improve the English in one or two places. We have removed the conclusion 'Finally, introducing e-cigarettes should lead to a reduced risk of SRD' from the four main conclusions listed at the beginning of the discussion, and made it clearer that this

conclusion is based on general considerations rather than rigorous modelling.

Competing Interests: No competing interests were disclosed.

Reviewer Report 13 June 2019

<https://doi.org/10.5256/f1000research.21197.r48812>

© 2019 Sulsky S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Sandra I. Sulsky

Ramboll US Corporation, Amherst, MA, USA

The revised paper addresses prior comments. Overall, this work demonstrates the importance of considering the totality of evidence when a public health relevant question is the topic of debate. Its quantitative evaluation of the potential effects of uncontrolled confounding on risk estimation are useful for any research setting (i.e., not just tobacco-related or harm reduction research).

Competing Interests: Through my employment at Ramboll US Corporation, I participate in research to support the development of evidence-based US tobacco policies. This work is funded through contracts between tobacco companies and Ramboll.

Reviewer Expertise: Epidemiology; population health modeling; tobacco harm reduction.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 12 April 2019

<https://doi.org/10.5256/f1000research.18509.r45925>

© 2019 Sulsky S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Sandra I. Sulsky

Ramboll US Corporation, Amherst, MA, USA

This ambitious paper by Lee and colleagues seems to have several aims, and I fear that the paper

suffers from including them all, rather than separating them out into two or more focused manuscripts.

The authors provide a review of the literature discussing what is currently known about the association between use of combusted and electronic tobacco products, they evaluate the likelihood of electronic products providing a gateway in to smoking based on seemingly informal, qualitative trend analyses, and they provide an important discussion about the potential for residual or otherwise uncontrolled confounding to explain or partially explain associations that have been noted in the literature. They also provide data from a series of simulations. The simulations demonstrate the interdependence among various tobacco use patterns in predicting population harm or benefit resulting from electronic tobacco products.

The most important points from the literature-based sections of the paper are a) the difference between predicted and observed smoking prevalence over time, which should reduce the fear that a large gateway to smoking effect is currently operating; b) the likelihood that misclassification and uncontrolled confounding play a role in the results reported by others, and probably lead to an overestimate of gateway-in effect.

The key findings from the simulations are a bit harder to uncover, due to a lack of detail in the methods section and some glossing over of assumptions. The authors found that a gateway-in effect is unlikely to be large enough to substantially affect population health. This is based on the assumptions that a) the current situation, in which smoking is far more prevalent than electronic cigarette use, will remain stable in at least the short run; b) the current higher likelihood of initiating tobacco use with cigarettes vs electronic products will remain stable in at least the short run; and c) the assumptions that the risks associated with electronic products are understood and are definitely lower than risks associated with smoking. If any of these input assumptions were varied, the simulation results would probably be different, and possibly much different. Some of the authors' strongest statements are due to differences between relative and absolute effects: Small percentages of a large underlying population (smokers) affect a large number of people, and large percentages of a small underlying population (electronic product users or initiators) affect a small number of people.

Apart from these conceptual concerns, it would strengthen the paper if the input into the simulations were better defined and justified.

The analyses presented in Table 8 seem to be the most informative of the simulations. Estimating the rate of the combination of smoking OR vaping OR dual use and comparing the prevalence in each year to the prevalence predicted by trends in smoking would give clues about the population level effects of e-products. Has overall use increased or stayed the same?

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Partly

If applicable, is the statistical analysis and its interpretation appropriate?

Partly

Are all the source data underlying the results available to ensure full reproducibility?

Partly

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: Through my employment at Ramboll US Corporation, I participate in research to support the development of evidence-based US tobacco policies. This work is funded through contracts between tobacco companies and Ramboll.

Reviewer Expertise: Epidemiology; population health modeling; tobacco harm reduction.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 13 May 2019

Peter Lee, P.N.Lee Statistics and Computing Ltd, Sutton, UK

Dr Sulsky makes three main points:

The paper may be better as multiple focussed manuscripts. We did not attempt this, our main purpose being to cover a range of issues relevant to vaping. Note that two of us also recently published in F1000Research on "Investigating gateway effects using the PATH study".

Some key findings from the simulations are hard to uncover and some assumptions not clearly stated. We made numerous amendments to answer these points.

It would also be useful to study trends in combined prevalence of vaping and/or smoking. As the paper's interest concerns effects of vaping on smoking prevalence, and as interpretation of upward trends in combined use is problematic (they may well reduce risk) we did not change the paper here.

Competing Interests: None

Reviewer Report 28 March 2019

<https://doi.org/10.5256/f1000research.18509.r45926>

© 2019 Shahab L. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Lion Shahab

Department of Behavioural Science and Health, University College London (UCL), London, UK

This is an interesting and valuable contribution to the field on the hotly debated topic of whether or not e-cigarettes function as a gateway into subsequent combustible cigarette use. Of particular value here is the analysis of the impact of residual confounding on creating spurious associations in observational studies. However, there are a number of issues in the other analyses I would recommend addressing.

1. Abstract: This is currently rather data free. I would suggest including some illustrative results here.
2. Results: Particularly in relation to Tables 5-7, I wonder if it would be possible to condense results somewhat (or to move some of these results into supplementary files). It seems that it is not strictly necessary to include all modelled effects, especially since not all are discussed. This makes this section at times difficult to follow. A better way to structure the Tables might be to link these explicitly to the 5 conclusions presented on pages 9 and 10.
3. Terminology: I am not sure that the term "gateway-out" is helpful. At the very least, it would not represent the only way to construe this concept insofar as it could be argued that in a hypothetical scenario the opposite to the "gateway-in" might be the avoidance of uptake of smoking in the first place (another "gateway-out" not related to smoking cessation where adolescents who would have smoked try an e-cigarette and never start smoking). It might therefore be helpful to clarify the difference between those gateway effects away from smoking
4. Change in smoking trends: I am not entirely convinced by this analysis for a number of reasons. First, very little information is provided about the actual analysis undertaken. I would expect to see fitting parameters for various functions to model trends in smoking prevalence in the period prior to 2014. Are these assumed to be linear, logarithmic, polynomial etc.? More detail is required here. A more appropriate way to analyse the data would be to conduct segmented regression or timeseries analysis. Second, based on the current analysis, the problem (somewhat acknowledged on top of page 11) is that it might still be the case that e-cigarettes functioned as a "gateway in" since we do not know what smoking prevalence would have been in the absence of e-cigarettes (and may have been even lower). I don't think this is likely but this issue should be acknowledged more explicitly, perhaps making it clear that such arguments would require "special pleading", i.e. one would need to postulate additional environmental changes which could explain a continued decrease in smoking prevalence even in the presence of a gateway effect.
5. Second paragraph on page 11 does not seem to add anything (it's not clear how these cognitive measures in PATH relate to actual behaviour and so it's quite weak evidence). Suggest dropping this.
6. Table 9: The counterpart to A3 is not presented, i.e. dual use which results in reduction of combustible cigarettes. If there is a commensurate reduction in cigarette consumption associated with an increase in e-cigarette use (if dual users reduce cigarette consumption sufficiently) there may be beneficial health effects (so perhaps this effect should be added).

In fact, perhaps a more nuanced discussion of dual use is needed on page 12, given that it may have both positive and negative effects and encompasses such a wide range of behavioural patterns. Please note that the statement on page 11 "The adverse effect for smokers switching to vaping instead of quitting smoking (A2) is not ideal but would still confer some benefits, given what is known regarding the reduced harm and toxicity of e-cigarettes" is logically inconsistent. If a smoker intending to quit starts using e-cigarettes instead of quitting, there is no benefit (as e-cigarette use even if less harmful than smoking does not confer any benefit over stopping completely).

7. The discussion on end of page 11 and top of page 12 feels rather qualitative. Either provide more quantitative analysis of modelled impact on health effects or move this to the discussion. You don't really provide any primary data to support your conclusion "Finally, introducing e-cigarettes should lead to a reduced risk of SRD." This is rather speculative at this stage without more rigorous modelling as was done e.g. in the NASEM report.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Partly

Are sufficient details of methods and analysis provided to allow replication by others?

Partly

If applicable, is the statistical analysis and its interpretation appropriate?

Partly

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Epidemiology, Health Psychology, Tobacco Control, Psychopharmacology, Neuroscience

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 13 May 2019

Peter Lee, P.N.Lee Statistics and Computing Ltd, Sutton, UK

Taking Dr Shahab's numbered points:

1. As the paper covers numerous issues and the conclusions are intended to be general,

and as abstract length is limited, we left the abstract unaltered.

2. We prefer not to change Tables 5 and 7. They are short and reducing them makes it harder to see effects of variation in the different parameters.
3. We have abandoned the terms gateway-in and gateway-out. Gateway-in is now called the gateway effect, and gateway-out is explained more generally.
4. A new additional file (4) gives fuller detail. The model used – linear in $\log(p/(1-p))$ – was actually described originally, the additional information suggesting this is a reasonable approach. The discussion has been increased here.
5. Rather than dropping some material in the section “Additional evidence on ...”, we prefer adding a paragraph clarifying the data limitations.
6. Table 9 and the related discussion have been amended along the lines suggested.
7. We moved the last two paragraphs of results to the discussion.

Competing Interests: None

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research