CASE REPORT

Case Report: Enlarging symmetrical masses of the palate of idiopathic etiology [version 1; peer review: 1 approved, 1 approved with reservations]

Michelangelo Vestita , Eleonora Nacchiero, Michele Maruccia, Giuseppe Giudice

Section of Plastic and Reconstructive Surgery, Department of Emergency and Organ Transplantation, University of Bari, Bari, 70124, Italy

V1 First published: 23 Feb 2018, 7:223
<https://doi.org/10.12688/f1000research.14033.1>

Latest published: 23 Feb 2018, 7:223
<https://doi.org/10.12688/f1000research.14033.1>

Abstract

We report the case of a 33 year-old woman who came to our attention with slowly enlarging exophytic masses of the palate, histologically characterized by sub-epithelial fibrous proliferation with packed collagen bundles and increased fibroblasts number. We describe the condition of idiopathic fibrous hyperplasia, its diagnosis and its surgical treatment, which in our case was carried out with the aid of a custom made thermal printed plaque used as a scaffold.

Keywords

fibrous hyperplasia, palate, idiopathic, gingival hyperplasia

Open Peer Review

Approval Status

	1	2
version 1 23 Feb 2018	 view	 view
1. Ilaria Mataro , Antonio Cardarelli Hospital, Naples, Italy		
2. André Salval , IEI - Istituto Estetico Italiano, Milan, Italy		
Francesco Ciano , Università degli Studi di Bari, Bari, Italy		

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Michelangelo Vestita (michelangelovestita@gmail.com)

Author roles: **Vestita M:** Conceptualization, Data Curation, Formal Analysis, Investigation, Supervision, Validation, Writing – Original Draft Preparation, Writing – Review & Editing; **Nacchiero E:** Investigation, Methodology, Resources, Validation, Visualization, Writing – Original Draft Preparation; **Maruccia M:** Conceptualization, Investigation, Validation, Writing – Original Draft Preparation, Writing – Review & Editing; **Giudice G:** Methodology, Project Administration, Supervision, Validation, Visualization

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2018 Vestita M *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Vestita M, Nacchiero E, Maruccia M and Giudice G. **Case Report: Enlarging symmetrical masses of the palate of idiopathic etiology [version 1; peer review: 1 approved, 1 approved with reservations]** F1000Research 2018, 7:223 <https://doi.org/10.12688/f1000research.14033.1>

First published: 23 Feb 2018, 7:223 <https://doi.org/10.12688/f1000research.14033.1>

Introduction

Idiopathic fibrous hyperplasia is a rare benign condition, characterized by a slow and progressive increase in gingival volume^{1,2}. It manifests as a rosy swelling of hard consistency while, at histological examination, it is characterized by a proliferation of fibroblasts in a myxomatous stroma. We describe and discuss a case of idiopathic fibrous hyperplasia of the palate.

Case

A 33 year-old woman came to our attention with slowly enlarging exophytic masses of the palate, which had begun to grow 2 years before and caused her disturbances of phonation as well as in swallowing solids and liquids. The patient did not take any drugs; however she had been a frequent user of nonsteroidal anti-inflammatory drugs for the last 3 years because of chronic back pain. Remote personal and family histories were negative, except for recurrent gastric nuisance and back pain.

Clinical and rhinoscopy examination demonstrated bilateral and symmetrical exuberant hypertrophic tissue, of hard consistency and rosy color, at the posterior-lateral area of the palate, with a tendency to coalesce medially. This tissue was contiguous to the adjacent gingiva (Figure 1).

Computerized tomography scan showed such lesions to be limited to the mucosal palate, with no underlying bone involvement (Figure 2).

An incisional biopsy demonstrated a sub-epithelial fibrous proliferation with packed collagen bundles and increased fibroblasts number (Figure 3). We concluded for a diagnosis of localized idiopathic fibrous hyperplasia. We treated the patient with a personalized approach using surgical resection and inseting of a thermal-printed palate plaque (Figure 4). We obtained good functional results at 20 days post-op (Figure 5), and no sign of recurrence at the 12 months follow up (Figure 6).

Figure 1. Clinical aspect: Enlarging symmetrical masses involving the palate.

Figure 2. Computerized tomography scan: Soft tissue masses, with absence of underlying bone involvement.

Figure 3. Histologic aspect: Hematoxylin & eosin (magnification, x10). Consistent with idiopathic fibrous hyperplasia.

Figure 4. Custom made, thermal printed plaque.

Figure 5. Clinical aspect, plaque in place after hyperplasia removal.

Figure 6. Clinical aspect, 12 months post-op: Satisfactory functional and cosmetic outcome and absence of recurrence.

Discussion

Two main subtypes of fibrous hyperplasia are known: generalized and localized. Generalized form has a genetic predisposition, appears at decimal or definitive dental eruption, and usually demonstrates a tendency to recur after surgery¹⁻⁴.

Secondary forms associated to pregnancy, scurvy, leukemia and drugs are also known¹⁻³. Among the latter, various chemotherapy agents can elicit secondary forms⁵, including ipilimumab and vemurafenib⁶. However, no reports link gingival fibrous hyperplasia to the drugs administered in our patient. The localized form has its onset from the second decade, does not generally recur after surgery, and normally is not associated to genetic predisposition^{7,8} although investigations to exclude syndromes commonly associated to gingival fibromatosis should always be carried out in our experience; these include Laband, Rutherford, Cross and Ramon syndromes⁵. In both the localized and the generalized forms, local factors such as dental plaques, caries, and the action of chemical substances and their metabolites might contribute to the onset in susceptible patients⁶. The precise pathogenic mechanism of idiopathic forms is unknown, but it appears to confine to the gingival and mucosal fibroblasts with no involvement of the periodontal ligament or the palate underlying bones^{1,2}.

Regardless of the etiology, excess tissue removal is the treatment of choice in localized fibrous hyperplasia, either by scalpel or by CO₂ laser^{7,8}. Our personal approach includes the use of a thermal-printed palate plaque (Figure 4), to be left in place for 20 days after surgery (Figure 5), which in our experience yields excellent hemostasis by exerting compression and, at the same time, functions as a scaffold that promotes and guides second intention healing, preventing recurrence of exuberant tissue growth. The long term (12 months) results reported in our case testify to the efficacy of such an approach.

Consent

Written informed consent for publication of their clinical details and clinical images was obtained from the patient.

Competing interests

No competing interests were disclosed.

Grant information

The author(s) declared that no grants were involved in supporting this work.

References

- Cekmez F, Pirgon O, Tanju IA: **Idiopathic gingival hyperplasia**. *Int J Biomed Sci*. 2009; 5(2): 198–200.
[PubMed Abstract](#) | [Free Full Text](#)
- Sapp JP, Eversole LR, Wysocki GP: **Connectivetissue lesions**. *Contemporary Oral and Maxillofacial Pathology*. 2004; 294–297, Mosby, London, UK 2nd edition.
[Reference Source](#)
- Pappachan B, Narayan JV, Nayak A: **Idiopathic gingival fibromatosis: A neglected case**. *Indian J Radiol Imaging*. 2002; 12: 335–338.
- Hart TC, Pallos D, Bozzo L, et al.: **Evidence of genetic heterogeneity for hereditary gingival fibromatosis**. *J Dent Res*. 2000; 79(10): 1758–1764.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Guida M, Cramarossa A, Fistola E, et al.: **High activity of sequential low dose chemo-modulating Temozolomide in combination with Fotemustine in metastatic melanoma. A feasibility study**. *J Transl Med*. 2010; 8: 115.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Mangold AR, Bryce A, Sekulic A: **Vemurafenib-associated gingival hyperplasia in patient with metastatic melanoma**. *J Am Acad Dermatol*. 2014; 71(5): e205–206.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Shetty AK, Shah HJ, Patil MA, et al.: **Idiopathic gingival enlargement and its management**. *J Indian Soc Periodontol*. 2010; 14(4): 263–265.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Palaia G: **Idiopathic fibrous hyperplasia of the palate**. *Ann Stomatol (Roma)*. 2013; 4(Suppl 2): 35.
[PubMed Abstract](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 08 March 2018

<https://doi.org/10.5256/f1000research.15255.r31155>

© 2018 Salval A et al. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

André Salval

IEI - Istituto Estetico Italiano, Milan, Italy

Francesco Ciancio

Università degli Studi di Bari, Bari, Italy

This Case Report is of scientific interest and well written. A more detailed explanation of therapy would be appreciated. About the thermal printed palate plaque for instance: what material is it made of? Is it a mold or 3D printed? Is it placed immediately after surgery and left in place 24 hours a day? Is thermal plaque therapy a novel approach or are there any references about this kind of therapy?

Is the background of the case's history and progression described in sufficient detail?

Yes

Are enough details provided of any physical examination and diagnostic tests, treatment given and outcomes?

Yes

Is sufficient discussion included of the importance of the findings and their relevance to future understanding of disease processes, diagnosis or treatment?

Yes

Is the case presented with sufficient detail to be useful for other practitioners?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Plastic surgery, burns, aesthetic surgery

We confirm that we have read this submission and believe that we have an appropriate level

of expertise to confirm that it is of an acceptable scientific standard, however we have significant reservations, as outlined above.

Reviewer Report 26 February 2018

<https://doi.org/10.5256/f1000research.15255.r31157>

© 2018 Mataro I. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Ilaria Mataro

Department of Plastic and Reconstructive Surgery and Burn Unit, Antonio Cardarelli Hospital, Naples, Italy

I believe the article is well structured and describes an interesting presentation of a rare benign condition.

I must congratulate the authors on the quality and quantity of the photographic material. The case is really well documented with clinical pictures, histologic images and radiologic findings.

I have personally treated a couple of these cases but often experienced short to medium term recurrence. The alternative treatment described by the authors (the use of a guiding plaque after removal) is interesting and seems to warrant better long term outcome.

Is the background of the case's history and progression described in sufficient detail?

Yes

Are enough details provided of any physical examination and diagnostic tests, treatment given and outcomes?

Yes

Is sufficient discussion included of the importance of the findings and their relevance to future understanding of disease processes, diagnosis or treatment?

Yes

Is the case presented with sufficient detail to be useful for other practitioners?

Yes

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research