REVIEW

Unite to divide: Oligomerization of tubulin and actin homologs regulates initiation of bacterial cell division

[version 1; peer review: 3 approved]

Marcin Krupka, William Margolin

Department of Microbiology and Molecular Genetics, McGovern Medical School, Houston, USA

V1 First published: 28 Feb 2018, 7(F1000 Faculty Rev):235
<https://doi.org/10.12688/f1000research.13504.1>
 Latest published: 28 Feb 2018, 7(F1000 Faculty Rev):235
<https://doi.org/10.12688/f1000research.13504.1>

Abstract

To generate two cells from one, bacteria such as *Escherichia coli* use a complex of membrane-embedded proteins called the divisome that synthesize the division septum. The initial stage of cytokinesis requires a tubulin homolog, FtsZ, which forms polymers that treadmill around the cell circumference. The attachment of these polymers to the cytoplasmic membrane requires an actin homolog, FtsA, which also forms dynamic polymers that directly bind to FtsZ. Recent evidence indicates that FtsA and FtsZ regulate each other's oligomeric state in *E. coli* to control the progression of cytokinesis, including the recruitment of septum synthesis proteins. In this review, we focus on recent advances in our understanding of protein-protein association between FtsZ and FtsA in the initial stages of divisome function, mainly in the well-characterized *E. coli* system.

Keywords

cytokinesis, bacteria, *Escherichia coli*, FtsZ, FtsA, protofilaments

Open Peer Review

Approval Status

	1	2	3
version 1			
28 Feb 2018			

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Jie Xiao**, Johns Hopkins University School of Medicine, Baltimore, USA
2. **Elizabeth J Harry**, The ithree Institute, University of Technology Sydney, Sydney, Australia
3. **Anuradha Janakiraman**, City College of New York, New York, USA

Any comments on the article can be found at the end of the article.

Corresponding author: William Margolin (William.Margolin@uth.tmc.edu)

Author roles: **Krupka M:** Writing – Review & Editing; **Margolin W:** Conceptualization, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The authors declare they received funding from National Institutes of Health grant GM61074 to support this work. *The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

Copyright: © 2018 Krupka M and Margolin W. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Krupka M and Margolin W. **Unite to divide: Oligomerization of tubulin and actin homologs regulates initiation of bacterial cell division [version 1; peer review: 3 approved]** F1000Research 2018, 7(F1000 Faculty Rev):235 <https://doi.org/10.12688/f1000research.13504.1>

First published: 28 Feb 2018, 7(F1000 Faculty Rev):235 <https://doi.org/10.12688/f1000research.13504.1>

Introduction

For a cell even as simple as *Escherichia coli*, splitting in half is a complex maneuver that needs to be finely regulated in time and space. For this purpose, *E. coli* has evolved a protein complex (the divisome) that spans the cytoplasmic membrane and forms a circumferential ring around midcell¹. Seen in cross-section, the innermost part of this complex, termed the proto-ring, consists of membrane-anchored cytoskeletal proteins that form polymers on the cytoplasmic side². These polymers interact with other proteins that span the membrane and guide the inward synthesis of cell wall material, called the division septum, which eventually will close to form the poles of the newborn cells. Within the proto-ring, FtsZ forms a dynamic scaffold for the recruitment of other septum synthesis proteins. FtsA and another protein, called ZipA, serve as dual membrane tethers for FtsZ; without either one, FtsZ forms a ring but cannot divide, but if both FtsA and ZipA are missing, FtsZ remains diffuse in the cytoplasm³. As FtsZ and FtsA are the most conserved across diverse bacterial species while ZipA is not, this review will focus on these two proteins.

FtsZ, the tubulin homolog

The most highly conserved divisome protein is FtsZ, a tubulin homolog. FtsZ assembles into a polymeric ring-like structure at the inner surface of the cytoplasmic membrane, marking the future division site and organizing the assembly of the divisome complex⁴. The ring then constricts in front of the invaginating division septum, reminiscent of the function of the actin ring in animal cells. Although early fluorescence microscopy studies suggested that FtsZ forms a continuous ring⁵, more recent super-resolution microscopy demonstrated that the ring in *E. coli* actually consists of multiple patches several hundred nanometers apart in a circumferential path^{6,7}. Other species as diverse as *Caulobacter crescentus*, *Bacillus subtilis*, *Streptococcus pneumoniae*, and *Prochlorococcus* show similar patchy localization of FtsZ⁶⁻¹⁰, suggesting that a symmetrical, continuous ring of FtsZ is not required for cell division. Indeed, mutants of *E. coli* that constrain active FtsZ to a partial ring structure can still divide, although the cells display abnormal morphologies^{11,12}. Recent studies have found that many bacterial species naturally initiate constriction on one side of the division site prior to becoming more symmetrical and that these asymmetric constrictions are likely promoted by relatively short FtsZ filaments that do not form a continuous ring structure^{13,14}.

Longitudinal interactions within FtsZ protofilaments

Like tubulin, FtsZ binds and hydrolyzes GTP^{15,16} but forms long single-stranded protofilaments in the presence of GTP instead of hollow microtubules¹⁷ (Figure 1A and Figure 2A). Two recent reports demonstrated that in both *E. coli* and *B. subtilis* cells, FtsZ protofilaments exhibit polarity and undergo treadmilling^{18,19}, during which subunits are selectively added to one end and removed from the other end (Figure 1A). The result is that one or more FtsZ protofilaments migrate as a unit around the cell circumference, roughly following the path of the ring. Although the movement of each treadmilling filament is processive, other FtsZ protofilaments in the same cell can move

in the opposing direction, so it seems that each protofilament unit has its own independent directionality.

FtsZ subunit turnover can occur within protofilaments or by treadmilling at protofilament ends. This turnover due to treadmilling is proportional to FtsZ's GTPase activity, indicating that GTP hydrolysis regulates circumferential protofilament migration. Consistent with the role of GTP hydrolysis in protofilament turnover and disassembly²⁰, the thermosensitive mutant *ftsZ84* exhibits much lower GTP binding and hydrolysis activity than wild-type (WT) FtsZ and is unable to support cell division at 42°C^{15,21}. Nevertheless, normal GTPase activity is clearly not essential for FtsZ function, as *ftsZ84* mutants form normal FtsZ rings and divide normally at the permissive temperature.

Through their interactions with other divisome proteins, FtsZ treadmilling protofilaments guide the key septal synthesis enzyme, FtsI (PBP3), which harbors the transpeptidase activity required for gradual inward cell wall growth eventually leading to cell splitting^{22,23}. It is important to note that in *E. coli*, the activity of cell wall synthesis enzymes drives septum synthesis independently of FtsZ treadmilling but that *B. subtilis* septal wall synthesis is dependent on FtsZ treadmilling speed^{18,19}. Although FtsZ subunits exchange rapidly within protofilaments as well as at the ends^{24,25}, it remains to be seen exactly how internal subunits of FtsZ within a processively treadmilling filament help to guide a moving septum synthesis machine²⁶. One possible mechanism is that the septum synthesis machine continuously shuttles between different FtsZ subunits at the growing end, although it is difficult to imagine how this could occur processively. Recent evidence suggests that any physical links between FtsZ and the septum synthesis proteins are likely indirect²⁷.

It is also not clear how single-stranded FtsZ protofilaments treadmill (that is, although they have inherent polarity, why one end adds net subunits while the other loses them). A recent study suggests a possible mechanism. Examining crystal structures of FtsZ, Wagstaff *et al.* found a correlation between different conformational states of FtsZ and the ability to form polymers²⁸. The closed form of FtsZ has an incomplete subunit interface that correlates with the monomeric state, whereas the open form has a more complete subunit interface and can be modeled to fit into long straight protofilaments. This difference in interfaces between closed and open conformations of the subunits may result in distinct "on-off" rates from either end; if one end has a stronger "on" rate than the other, it would become the growing end of the filament. This interesting model awaits further experimental confirmation.

As FtsZ polymerization and treadmilling are essential for normal septal morphology, the inhibition of those activities should inhibit cell division. Indeed, there are numerous endogenous regulators that target FtsZ polymerization by various mechanisms, although inactivating any one of these regulators has only small effects on cell division. For example, the SOS-inducible Sula protein sequesters FtsZ subunits to prevent

Figure 1. Model for how changing oligomeric structures of FtsA (red) and FtsZ (green) might regulate *Escherichia coli* cell division. A scheme is shown for how FtsZ and FtsA oligomeric states influence and potentially reinforce each other by positive feedback. Depicted are their structures on membranes and subunit exchange by treadmilling (curved arrows) (A) and how several treadmilling complexes containing these structures at the inner surface of the cytoplasmic membrane might function in the cell for inward synthesis of the division septum (B). The large green arrow denotes circumferential movement of a patch of FtsZ-FtsA; blue arrows show the direction of division septum synthesis that pushes the cytoplasmic membrane inward. FtsA, filamentous thermosensitive protein A; FtsZ, filamentous thermosensitive protein Z.

polymer formation or growth or both^{29–31}. SlmA, a DNA-bound dimer that inhibits FtsZ polymerization over the nucleoid^{32,33}, seems to function by binding to the conserved C-terminal peptide of FtsZ and subsequently severing the FtsZ filament^{34,35} similar to how another FtsZ inhibitor, MinC, depolymerizes FtsZ to prevent inappropriate division at cell poles³⁶. Interestingly, inactivating MinC in the GTPase-defective *ftsZ84* mutant significantly exacerbates its thermosensitivity³⁷. In addition, several peptides and small molecules, either natural or synthetic, target FtsZ polymerization and hold promise as potential antimicrobials^{38,39}.

Lateral interactions between protofilaments

In addition to longitudinal interactions between FtsZ subunits within a protofilament, lateral interactions between protofilaments play an important role. Such interactions, which result

in crosslinking or bundling of FtsZ protofilaments, may be facilitated either by intrinsic attraction between FtsZ subunits or through accessory proteins. The general role of Zaps (Z-associated proteins), including ZapA, ZapC, and ZapD in *E. coli*, in promoting lateral interactions between FtsZ protofilaments is well described in recent reviews^{40,41} and will not be discussed in detail here. Although the loss of any one of these proteins has little effect on cell division, inactivation of more than one significantly reduces the division efficiency, suggesting that they have overlapping roles⁴². However, it is not yet clear whether they work together or whether their mechanisms of action are distinct.

The physiological role of intrinsic lateral interactions between FtsZ protofilaments is less clear, although several interesting mutants of *E. coli* FtsZ provide clues. FtsZ_{R85Q}, for instance,

Figure 2. Oligomeric structures of FtsZ and FtsA. Shown are negatively stained transmission electron micrographs of FtsZ (A) and FtsZ* (B) from *Escherichia coli* and lipid monolayers containing minirings of *E. coli* FtsA (C), FtsA* arcs (D), FtsA + FtsZ (E), and FtsA* + FtsZ* (F). Scale bar = 100 nm. Arrows highlight double FtsZ protofilaments (magenta), single FtsZ protofilaments (yellow), or FtsA arcs (green). FtsA, filamentous thermosensitive protein A; FtsZ, filamentous thermosensitive protein Z.

is partially defective for cell division and displays lower lateral interactions *in vitro*^{43,44}. Another lateral mutant, FtsZ_{R174D}, is non-functional for *E. coli* cell division and is unable to form bundled protofilaments *in vitro* when assembled in millimolar Ca⁺⁺, a condition that stimulates strong bundling^{45,46}. Although this deficiency of FtsZ_{R174D} in bundling *in vitro* has been challenged recently⁴⁷, a genetic suppressor screen showed that this mutant protein can be rescued for function by amino acid changes at the nearby residue L169. Interestingly, protofilaments of FtsZ_{L169R}, also called FtsZ*, make larger bundles than WT FtsZ *in vitro* (Figure 2A, B) and exhibit lower GTPase activity⁴⁸. Consistently, FtsZ* was able to suppress the ability of Kil, a peptide from coliphage lambda that inhibits FtsZ function, to depolymerize FtsZ^{49,50}. In addition, amino acid substitutions at residues E93 and D86 result in increased FtsZ filament bundling^{34,51–53}. Like the substitutions at L169, they are

still able to function in cell division, although cells with these variants have defects, forming spiral-shaped FtsZ structures that cause abnormally shaped division septa and deformed cell poles. Hyper-bundled protofilaments have lower GTPase activities that should reduce treadmilling speed, thus perturbing the timing of septum synthesis relative to cell growth, leading to the observed shape abnormalities.

The accumulated data suggest a model in which lateral interactions between FtsZ protofilaments are regulated for optimal function. This is supported by the important role that intrinsic lateral interactions play in FtsZs from other species. For example, *B. subtilis* FtsZ protofilaments intrinsically bundle more than *E. coli* FtsZ, and these lateral interactions are mediated by the conserved C-terminal peptide of FtsZ⁵⁴. *C. crescentus* FtsZ also exhibits intrinsic lateral interactions mediated by its

C-terminal domain⁵⁵. Moreover, *S. pneumoniae* FtsZ forms mostly double-stranded filaments, reflecting robust lateral interactions⁵⁶.

In addition to the Zap proteins, recent evidence indicates that other proteins specifically regulate lateral FtsZ interactions in diverse species. For example, ARC3, a protein in plant chloroplasts, antagonizes the filament bundling of FtsZ2, one of three FtsZ homologs used for chloroplast fission⁵⁷. FzlA and FzlC, both FtsZ-binding proteins in *C. crescentus*, regulate lateral interactions between FtsZ protofilaments⁵⁵, but the ZapA homolog in *C. crescentus* does not affect FtsZ lateral interactions *in vitro*, despite its ability to condense the FtsZ ring *in vivo*⁵⁸.

FtsA, the actin homolog for cell division

FtsZ–FtsZ interactions, whether lateral or longitudinal, are clearly important for cell division. Nevertheless, several other proteins that interact directly with FtsZ also regulate FtsZ–FtsZ interactions. One is the aforementioned ZipA, which is narrowly conserved in the enterobacteriaceae and forms membrane-tethered homodimers⁵⁹. Another is SepF, which is present in a diverse group of Gram-positive bacteria. In *B. subtilis*, SepF assembles into sharply curved protofilaments at the membrane that regulate FtsZ assembly^{60,61} and has been proposed to act like a bacterial homolog of spectrin⁶².

For the purposes of this brief review, we will focus on the most well characterized and highly conserved FtsZ interactor, FtsA, an actin homolog that binds ATP and forms polymers^{63–66}. FtsA differs from actin in that it uses its C-terminal amphipathic helix for binding to the cytoplasmic membrane⁶⁷ and binds directly to the conserved C-terminal peptide of FtsZ⁶⁸. As a result, FtsA molecules provide important membrane tethers and potential guides for treadmilling FtsZ polymers. As FtsA binds to a region of FtsZ that also interacts with other key divisome proteins such as ZipA and is implicated in lateral interactions between protofilaments, it is important to understand how FtsA and FtsZ interact *in vitro*.

Until very recently, understanding this interaction was complicated by the difficulties in purifying active *E. coli* FtsA (EcFtsA) containing the amphipathic helix. However, earlier studies of FtsA from *S. pneumoniae* (SpFtsA), which was more amenable to purification, provided some important clues about FtsA function. Purified SpFtsA polymerizes bidirectionally in solution and its polymerization is facilitated upon membrane binding or when its C-terminal end is removed. Both polymerization and membrane attachment depend on ATP binding, suggesting that ATP induces a conformational switch to liberate FtsA's C-terminal amphipathic helix for membrane binding and to induce polymerization^{69–71}. Complementary physiological studies of SpFtsA are now catching up with its biochemical characterization^{72,73} which should greatly facilitate our understanding of FtsA function in this species and others.

Recent studies using purified EcFtsA (hereafter referred to as FtsA) indicate that it, too, can polymerize, but only on membranes

and not as linear polymers⁷⁴. When added to lipid monolayers, FtsA assembles mainly into closed minirings containing 12 subunits each. These minirings, 20 nm in diameter, can be single or grouped into arrays (Figure 2C). FtsZ, in the presence of GTP and near physiological concentrations, forms long single-stranded protofilaments that are strikingly aligned by these FtsA minirings but spaced apart by more than 15 nm (Figure 2E). Notably, using fluorescence microscopy of purified FtsZ and FtsA on supported lipid bilayers, Loose and Mitchison⁷⁵ also observed the assembly of FtsZ into aligned polymer rafts on top of membrane-bound FtsA, although the ultrastructure of FtsA polymers was not examined. Consistent with the tendency of FtsZ protofilaments to curve under some conditions^{76–78}, these aligned FtsZ polymers form large circular vortices that swirl unidirectionally by treadmilling. This behavior probably reflects the directional treadmilling of individual FtsZ complexes observed in cells^{18,19}.

Just as mutants of FtsZ that decrease polymerization or GTPase activity are defective in cell division, mutants of FtsA that compromise the ATP-binding site or FtsA–FtsA interactions also have cell division phenotypes. Amino acid substitutions that inactivate EcFtsA function mostly map to the ATP-binding site and decrease ATPase activity, implicating nucleotide binding and hydrolysis in normal FtsA function⁷⁹. A very recent study showed that ATP binding and hydrolysis by *E. coli* FtsA promote membrane remodeling⁸⁰, analogous to the ability of ATP to promote membrane binding of *S. pneumoniae* FtsA⁷¹. Such an activity might stimulate membrane invagination during septation. Likewise, residue substitutions at the subunit interface of *B. subtilis* FtsA that are predicted to affect FtsA oligomerization inhibit its cell division activity⁶⁸. Surprisingly, however, many substitutions at or near the subunit interface that decrease FtsA self-interaction actually enhance FtsA activity. For example, the R286W substitution in *E. coli* FtsA (FtsA*) results in a gain of function, bypassing the requirement for several other cell division proteins, including ZipA, and permits cells to divide at abnormally short cell lengths^{81–83}. Other FtsA*-like substitutions have similar gain-of-function characteristics^{79,84,85}.

One model arising from these data proposes that, at least in *E. coli*, when an FtsA subunit is not engaged with another FtsA subunit in an oligomer, it is free to interact with later-division proteins, helping to drive cytokinesis forward⁸⁴. In support of this idea, purified FtsA* does not form closed minirings on lipid monolayers but instead mostly short arcs with the same curvature (Figure 1A and Figure 2D)⁷⁴. These arcs, with potentially free protein domains available at subunit ends unlike in minirings, might be more likely to interact with late-division proteins⁸⁴. In addition, the dynamics of treadmilling might be significantly influenced by the oligomerization state of FtsA on the membrane and its ability to tether FtsZ to the membrane (see “A role for FtsA in regulating lateral interactions between FtsZ protofilaments” section below). Future high-resolution electron microscopic studies with purified

FtsA bound to other cell division proteins on membranes should help to test this model and to elucidate factors that regulate treadmilling dynamics.

A role for FtsA in regulating lateral interactions between FtsZ protofilaments

As mentioned above, minirings of *E. coli* FtsA inhibit lateral interactions between FtsZ protofilaments. If FtsZ bundling is important for cell division to proceed, the prediction is that FtsA minirings switch to a form that promotes FtsZ bundling. FtsA* may be an example of this form because when FtsZ is assembled on lipid monolayers seeded with FtsA* arcs, FtsZ protofilaments are no longer separated by more than 15 nm and instead become highly bundled, with lateral distances of about 7 nm⁷⁴. This stimulation of FtsZ bundling by FtsA* has also been observed in solution⁸⁶, indicating that some behaviors in solution may mimic effects seen on the membrane. Strikingly, when the FtsZ* hyper-bundled variant is assembled on lipid monolayers with FtsA*, large sheets of FtsZ* with protofilaments spaced about 5 nm apart are formed (Figure 2F). This over-bundling suggests that FtsA*-mediated bundling of FtsZ is synergistic with the intrinsically increased bundling of FtsZ*. The physiological consequences are clear, as cells with both FtsA* and FtsZ* do not divide normally, often exhibiting twisted cell division septa⁴⁸. Interestingly, when FtsZ* is added to lipid monolayers containing FtsA minirings, the minirings become disrupted⁷⁴, hinting at a positive feedback loop (see further below in this section).

It is not yet known which molecular mechanisms control either intrinsic or FtsA-mediated FtsZ lateral interactions or which residues are involved in direct contacts. Nonetheless, such interactions likely involve not only the core polymerizing domain and C-terminal peptides of FtsZ but also the conserved, intrinsically disordered linker that connects them^{87–90}. Recent results indicate that changing the length of the linker in *E. coli* FtsZ can also alter the distance between bundled protofilaments, although the effects on lipid membranes with natural FtsZ tethers have not yet been explored⁹⁰.

An attractive model to explain these results is that FtsA minirings antagonize bundling of FtsZ protofilaments by holding them apart for some period of time and that this constraint is a key checkpoint in the progression of cell division in *E. coli* (Figure 1A). In this model, gain-of-function variants of FtsA such as FtsA* ignore this checkpoint because they do not form minirings on the membrane and instead would transition directly to FtsA*-like non-miniring oligomers. One consequence of this transition would be premature bundling of FtsZ protofilaments. Although the overall morphology of the FtsZ ring does not change much during constriction^{8,9,91}, a more condensed state of FtsZ^{91,92}, potentially caused by increased protofilament bundling, is thought to be important for promoting septum formation (Figure 1B). Indeed, in FtsA*-like mutants that may prematurely bundle FtsZ, cell division is accelerated, potentially by bypassing a checkpoint, resulting in cells that divide at shorter cell lengths and overcome perturbations to the normal cell division pathway such as inactivation of ZipA. Considering the ability of bundled FtsZ to disrupt FtsA

minirings *in vitro*⁷⁴, we propose that factors that bundle FtsZ and factors that disrupt FtsA minirings act in the same pathway to promote septation. As FtsZ becomes more bundled, FtsA minirings are disrupted, which promotes more FtsZ bundles, which in turn disrupts more FtsA minirings. Therefore, according to this model, the increased bundling of FtsZ and decreased oligomerization of FtsA reinforce each other in a positive feedback loop that helps to maintain constriction of the inner membrane in the forward direction in concert with septum synthesis. Genetic evidence suggests that FtsEX, an ATPase complex that senses septum synthesis but also contacts FtsZ and FtsA, may be involved in such a feedback loop, as the FtsX protein likely regulates FtsA's oligomerization state^{93,94}. As more mechanistic details in the FtsA–FtsZ interaction are revealed, interesting parallels with actin and tubulin in eukaryotic cells should emerge. For example, FtsA control of FtsZ bundling is reminiscent of F-actin bundling by septins⁹⁵.

Although the above model is based on *in vitro* data and FtsA minirings have not yet been observed in *E. coli* cells, other genetic data support the idea that FtsA antagonizes FtsZ bundling. For example, a moderate excess of FtsA in *E. coli* cells lacking the FtsZ bundling proteins ZapA or ZapC significantly exacerbates the cell division defects caused by ZapA/C deficiency alone; conversely, excess FtsA can rescue cell division deficiencies in cells with over-bundled FtsZ caused by excess ZapA⁷⁴. FtsA minirings are reminiscent of the structures formed by *B. subtilis* SepF but differ from the straight filaments formed by *T. maritima* FtsA reported by Szwedziak *et al.*⁹⁶. We speculate that FtsA proteins from distinct bacterial species assemble into a variety of forms and that the miniring form is potentially exclusive to *E. coli* and related gamma-proteobacteria that have multiple FtsZ membrane tethers but need to impose a checkpoint delay on septum formation. In bacteria with only one membrane anchor, perhaps FtsA polymerization into straight actin-like filaments serves to tether more FtsZ to the membrane and consequently reduce its treadmilling, keeping septation in check by a different mechanism.

Interestingly, gain-of-function mutants in other essential *E. coli* divisome genes (*ftsL* and *ftsB*) have similar properties as *ftsA**, hinting that other parts of the cell division apparatus, including multiple periplasmic proteins with no known enzymatic activities, are also involved in regulating the progression of septum formation^{85,97}. It is notable that the FtsZ bundling-stimulator FtsA* can rescue the toxicity caused by excess ZapC, which is the opposite of what would be predicted if ZapC also induces FtsZ over-bundling⁹⁸. These effects may reflect the increased ability of FtsA* to recruit downstream divisome proteins. Clearly, the effects of gain-of-function mutants such as FtsA* cannot be limited to FtsZ bundling alone.

Conclusions and perspectives

We have discussed important recent breakthroughs in our understanding of how FtsZ and FtsA interact to organize bacterial cell division, mainly in *E. coli* but also increasingly in other model systems. FtsZ protofilaments treadmill around the circumference of the division ring and are connected—directly or indirectly—to the membrane and periplasmic peptidoglycan synthesis

machinery by FtsA and other proteins to construct the septum. The next key challenge is to elucidate the mechanism of treadmilling and how it guides septum synthesis. In addition, important new evidence for the role of lateral interactions in FtsZ function and the ability of FtsA's oligomeric state to regulate these interactions, including the roles of ATP and GTP in these processes, needs to be explored more in detail. These experiments should be greatly facilitated by improved lipid membrane systems, such as lipid monolayers, supported lipid bilayers, and lipid discs, as well as rapidly improving direct visualization methods such as super-resolution microscopy and cryo-electron tomography. Finally, it is not yet settled whether the primary purpose of dynamic FtsZ filaments is to generate a physiologically relevant inward constriction force, supported by experiments with liposomes⁹⁹, or to interface with the septal synthesis machinery and respond flexibly to numerous inputs^{100,101} or to do both.

Abbreviations

ATP, adenosine 5' triphosphate; EcFtsA, *Escherichia coli* FtsA; FtsA, filamentous thermosensitive protein A; FtsZ, filamentous

thermosensitive protein Z; FzlA, FtsZ-localized protein A; FzlC, FtsZ-localized protein C; GTP, guanosine 5' triphosphate; SepF, septation protein F; SlmA, synthetic lethal with Min protein A; SpFtsA, *Streptococcus pneumoniae* FtsA; WT, wild-type; ZapA, ZapC, ZapD, FtsZ-associated proteins A, C, D; ZipA, FtsZ-interacting protein A.

Competing interests

The authors declare that they have no competing interests.

Grant information

The authors declare they received funding from National Institutes of Health grant GM61074 to support this work.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgments

We apologize to colleagues whose work could not be cited due to space limitations.

References

- Haeusser DP, Margolin W: **Splitsville: structural and functional insights into the dynamic bacterial Z ring.** *Nat Rev Microbiol.* 2016; 14(5): 305–19.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Li Z, Trimble MJ, Brun YV, *et al.*: **The structure of FtsZ filaments in vivo suggests a force-generating role in cell division.** *EMBO J.* 2007; 26(22): 4694–708.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
- Pichoff S, Lutkenhaus J: **Unique and overlapping roles for ZipA and FtsA in septal ring assembly in *Escherichia coli*.** *EMBO J.* 2002; 21(4): 685–93.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Bi EF, Lutkenhaus J: **FtsZ ring structure associated with division in *Escherichia coli*.** *Nature.* 1991; 354(6349): 161–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Ma X, Ehrhardt DW, Margolin W: **Colocalization of cell division proteins FtsZ and FtsA to cytoskeletal structures in living *Escherichia coli* cells by using green fluorescent protein.** *Proc Natl Acad Sci U S A.* 1996; 93(23): 12998–3003.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Fu G, Huang T, Buss J, *et al.*: **In vivo structure of the *E. coli* FtsZ-ring revealed by photoactivated localization microscopy (PALM).** *PLoS One.* 2010; 5(9): e12682.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
- Rowlett VW, Margolin W: **3D-SIM super-resolution of FtsZ and its membrane tethers in *Escherichia coli* cells.** *Biophys J.* 2014; 107(8): L17–20.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
- Strauss MP, Liew AT, Turnbull L, *et al.*: **3D-SIM super resolution microscopy reveals a bead-like arrangement for FtsZ and the division machinery: implications for triggering cytokinesis.** *PLoS Biol.* 2012; 10(9): e1001389.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Holden SJ, Pengo T, Meibom KL, *et al.*: **High throughput 3D super-resolution microscopy reveals *Caulobacter crescentus* in vivo Z-ring organization.** *Proc Natl Acad Sci U S A.* 2014; 111(12): 4566–71.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
- Liu R, Liu Y, Liu S, *et al.*: **Three-dimensional superresolution imaging of the FtsZ ring during cell division of the cyanobacterium *Prochlorococcus*.** *mBio.* 2017; 8(6): pii: e00657–17.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
- Addinall SG, Lutkenhaus J: **FtsZ-spirals and -arcs determine the shape of the invaginating septa in some mutants of *Escherichia coli*.** *Mol Microbiol.* 1996; 22(2): 231–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Rowlett VW, Margolin W: **Asymmetric constriction of dividing *Escherichia coli* cells induced by expression of a fusion between two min proteins.** *J Bacteriol.* 2014; 196(11): 2089–100.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Leisch N, Pende N, Weber PM, *et al.*: **Asynchronous division by non-ring FtsZ in the gammaproteobacterial symbiont of *Robbea hypermnestra*.** *Nat Microbiol.* 2016; 2: 16182.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
- Yao Q, Jewett AI, Chang YW, *et al.*: **Short FtsZ filaments can drive asymmetric cell envelope constriction at the onset of bacterial cytokinesis.** *EMBO J.* 2017; 36(11): 1577–89.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
- de Boer P, Crossley R, Rothfield L: **The essential bacterial cell-division protein FtsZ is a GTPase.** *Nature.* 1992; 359(6392): 254–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Wang X, Lutkenhaus J: **The FtsZ protein of *Bacillus subtilis* is localized at the division site and has GTPase activity that is dependent upon FtsZ concentration.** *Mol Microbiol.* 1993; 9(3): 435–42.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Erickson HP, Taylor DW, Taylor KA, *et al.*: **Bacterial cell division protein FtsZ assembles into protofilament sheets and minirings, structural homologs of tubulin polymers.** *Proc Natl Acad Sci U S A.* 1996; 93(1): 519–23.
[PubMed Abstract](#) | [Free Full Text](#)
- Bisson-Filho AW, Hsu YP, Squyres GR, *et al.*: **Treadmilling by FtsZ filaments drives peptidoglycan synthesis and bacterial cell division.** *Science.* 2017; 355(6326): 739–43.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
- Yang X, Lyu Z, Miguel A, *et al.*: **GTPase activity-coupled treadmilling of the bacterial tubulin FtsZ organizes septal cell wall synthesis.** *Science.* 2017; 355(6326): 744–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
- Mukherjee A, Lutkenhaus J: **Dynamic assembly of FtsZ regulated by GTP hydrolysis.** *EMBO J.* 1998; 17(2): 462–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Raychaudhuri D, Park JT: ***Escherichia coli* cell-division gene *ftsZ* encodes a novel GTP-binding protein.** *Nature.* 1992; 359(6392): 251–4.
[PubMed Abstract](#) | [Publisher Full Text](#)

22. Weiss DS, Pogliano K, Carson M, *et al.*: Localization of the *Escherichia coli* cell division protein FtsI (PBP3) to the division site and cell pole. *Mol Microbiol.* 1997; 25(4): 671–81.
[PubMed Abstract](#) | [Publisher Full Text](#)
23. Nguyen-Distèche M, Fraipont C, Buddelmeijer N, *et al.*: The structure and function of *Escherichia coli* penicillin-binding protein 3. *Cell Mol Life Sci.* 1998; 54(4): 309–16.
[PubMed Abstract](#) | [Publisher Full Text](#)
24. **F** Stricker J, Maddox P, Salmon ED, *et al.*: Rapid assembly dynamics of the *Escherichia coli* FtsZ-ring demonstrated by fluorescence recovery after photobleaching. *Proc Natl Acad Sci U S A.* 2002; 99(5): 3171–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
25. Niu L, Yu J: Investigating intracellular dynamics of FtsZ cytoskeleton with photoactivation single-molecule tracking. *Biophys J.* 2008; 95(4): 2009–16.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
26. Schoenemann KM, Margolin W: Bacterial Division: FtsZ treadmills to build a beautiful wall. *Curr Biol.* 2017; 27(8): R301–R303.
[PubMed Abstract](#) | [Publisher Full Text](#)
27. **F** Söderström B, Chan H, Shilling PJ, *et al.*: Spatial separation of FtsZ and FtsN during cell division. *Mol Microbiol.* 2018; 107(3): 387–401.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
28. **F** Wagstaff JM, Tsim M, Oliva MA, *et al.*: A polymerization-associated structural switch in FtsZ that enables treadmilling of model filaments. *mBio.* 2017; 8(3): pii: e00254-17.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
29. Mukherjee A, Cao C, Lutkenhaus J: Inhibition of FtsZ polymerization by SulA, an inhibitor of septation in *Escherichia coli*. *Proc Natl Acad Sci U S A.* 1998; 95(6): 2885–90.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
30. Trusca D, Scott S, Thompson C, *et al.*: Bacterial SOS checkpoint protein SulA inhibits polymerization of purified FtsZ cell division protein. *J Bacteriol.* 1998; 180(15): 3946–53.
[PubMed Abstract](#) | [Free Full Text](#)
31. Chen Y, Milam SL, Erickson HP: SulA inhibits assembly of FtsZ by a simple sequestration mechanism. *Biochemistry.* 2012; 51(14): 3100–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
32. **F** Bernhardt TG, de Boer PA: SlmA, a nucleoid-associated, FtsZ binding protein required for blocking septal ring assembly over Chromosomes in *E. coli*. *Mol Cell.* 2005; 18(5): 555–64.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
33. **F** Schumacher MA, Zeng W: Structures of the nucleoid occlusion protein SlmA bound to DNA and the C-terminal domain of the cytoskeletal protein FtsZ. *Proc Natl Acad Sci U S A.* 2016; 113(18): 4988–93.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
34. Du S, Lutkenhaus J: SlmA antagonism of FtsZ assembly employs a two-pronged mechanism like MinCD. *PLoS Genet.* 2014; 10(7): e1004460.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
35. Cabrè EJ, Monterroso B, Alfonso C, *et al.*: The nucleoid occlusion SlmA protein accelerates the disassembly of the FtsZ protein polymers without affecting their GTPase activity. *PLoS One.* 2015; 10(5): e0126434.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
36. **F** Shen B, Lutkenhaus J: Examination of the interaction between FtsZ and MinC⁺ in *E. coli* suggests how MinC disrupts Z rings. *Mol Microbiol.* 2010; 75(5): 1285–98.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
37. Yu XC, Margolin W: Deletion of the *min* operon results in increased thermosensitivity of an *ftsZ84* mutant and abnormal FtsZ ring assembly, placement, and disassembly. *J Bacteriol.* 2000; 182(21): 6203–13.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
38. **F** Haydon DJ, Stokes NR, Ure R, *et al.*: An inhibitor of FtsZ with potent and selective anti-staphylococcal activity. *Science.* 2008; 321(5896): 1673–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
39. den Blaauwen T, Andreu JM, Monasterio O: Bacterial cell division proteins as antibiotic targets. *Bioorg Chem.* 2014; 55: 27–38.
[PubMed Abstract](#) | [Publisher Full Text](#)
40. Monahan LG, Robinson A, Harry EJ: Lateral FtsZ association and the assembly of the cytokinetic Z ring in bacteria. *Mol Microbiol.* 2009; 74(4): 1004–17.
[PubMed Abstract](#) | [Publisher Full Text](#)
41. Huang KH, Durand-Heredia J, Janakiraman A: FtsZ ring stability: of bundles, tubules, crosslinks, and curves. *J Bacteriol.* 2013; 195(5): 1859–68.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
42. Durand-Heredia J, Rivkin E, Fan G, *et al.*: Identification of ZapD as a cell division factor that promotes the assembly of FtsZ in *Escherichia coli*. *J Bacteriol.* 2012; 194(12): 3189–98.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
43. Shin JY, Vollmer W, Lagos R, *et al.*: Glutamate 83 and arginine 85 of helix H3 bend are key residues for FtsZ polymerization, GTPase activity and cellular viability of *Escherichia coli*: lateral mutations affect FtsZ polymerization and *E. coli* viability. *BMC Microbiol.* 2013; 13: 26.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
44. Márquez IF, Mateos-Gil P, Shin JY, *et al.*: Mutations on FtsZ lateral helix H3 that disrupt cell viability hamper reorganization of polymers on lipid surfaces. *Biochim Biophys Acta.* 2017; 1859(10): 1815–27.
[PubMed Abstract](#) | [Publisher Full Text](#)
45. Yu XC, Margolin W: Ca²⁺-mediated GTP-dependent dynamic assembly of bacterial cell division protein FtsZ into asters and polymer networks *in vitro*. *EMBO J.* 1997; 16(17): 5455–63.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
46. Koppelman C, Aarsman ME, Postmus J, *et al.*: R174 of *Escherichia coli* FtsZ is involved in membrane interaction and protofilament bundling, and is essential for cell division. *Mol Microbiol.* 2004; 51(3): 645–57.
[PubMed Abstract](#) | [Publisher Full Text](#)
47. Gardner KAJA, Osawa M, Erickson HP: Whole genome re-sequencing to identify suppressor mutations of mutant and foreign *Escherichia coli* FtsZ. *PLoS One.* 2017; 12(4): e0176643.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
48. Haeusser DP, Rowlett VW, Margolin W, *et al.*: A mutation in *Escherichia coli* ftsZ bypasses the requirement for the essential division gene zipA and confers resistance to FtsZ assembly inhibitors by stabilizing protofilament bundling. *Mol Microbiol.* 2015; 97(5): 988–1005.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
49. **F** Haeusser DP, Hoashi M, Weaver A, *et al.*: The Kil peptide of bacteriophage λ blocks *Escherichia coli* cytokinesis via ZipA-dependent inhibition of FtsZ assembly. *PLoS Genet.* 2014; 10(3): e1004217.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
50. Hernández-Rocamora VM, Alfonso C, Margolin W, *et al.*: Evidence that bacteriophage λ Kil peptide inhibits bacterial cell division by disrupting FtsZ protofilaments and sequestering protein subunits. *J Biol Chem.* 2015; 290(33): 20325–35.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
51. Stricker J, Erickson HP: *In vivo* characterization of *Escherichia coli* ftsZ mutants: effects on Z-ring structure and function. *J Bacteriol.* 2003; 185(16): 4796–805.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
52. Jaiswal R, Patel RY, Asthana J, *et al.*: E93R substitution of *Escherichia coli* FtsZ induces bundling of protofilaments, reduces GTPase activity, and impairs bacterial cytokinesis. *J Biol Chem.* 2010; 285(41): 31796–805.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
53. Encinar M, Kralicek AV, Martos A, *et al.*: Polymorphism of FtsZ filaments on lipid surfaces: role of monomer orientation. *Langmuir.* 2013; 29(30): 9436–46.
[PubMed Abstract](#) | [Publisher Full Text](#)
54. Buske PJ, Levin PA: Extreme C terminus of bacterial cytoskeletal protein FtsZ plays fundamental role in assembly independent of modulatory proteins. *J Biol Chem.* 2012; 287(14): 10945–57.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
55. **F** Sundararajan K, Goley ED: The intrinsically disordered C-terminal linker of FtsZ regulates protofilament dynamics and superstructure *in vitro*. *J Biol Chem.* 2017; 292(50): 20509–27.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
56. Salvarelli E, Krupka M, Rivas G, *et al.*: The Cell Division Protein FtsZ from *Streptococcus pneumoniae* Exhibits a GTPase Activity Delay. *J Biol Chem.* 2015; 290(41): 25081–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
57. **F** Shaik RS, Sung MW, Vitha S, *et al.*: Chloroplast division protein ARC3 acts on FtsZ2 by preventing filament bundling and enhancing GTPase activity. *Biochem J.* 2018; 475(1): 99–115.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
58. **F** Woldemeskel SA, McQuillen R, Hessel AM, *et al.*: A conserved coiled-coil protein pair focuses the cytokinetic Z-ring in *Caulobacter crescentus*. *Mol Microbiol.* 2017; 105(5): 721–40.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
59. Skoog K, Daley DO: The *Escherichia coli* cell division protein ZipA forms homodimers prior to association with FtsZ. *Biochemistry.* 2012; 51(7): 1407–15.
[PubMed Abstract](#) | [Publisher Full Text](#)
60. Gündoğdu ME, Kawai Y, Pavlendova N, *et al.*: Large ring polymers align FtsZ polymers for normal septum formation. *EMBO J.* 2011; 30(3): 617–26.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
61. Duman R, Ishikawa S, Celik I, *et al.*: Structural and genetic analyses reveal the protein SepF as a new membrane anchor for the Z ring. *Proc Natl Acad Sci U S A.* 2013; 110(48): E4601–10.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
62. **F** Cleverley RM, Barrett JR, Baslé A, *et al.*: Structure and function of a spectrin-like regulator of bacterial cytokinesis. *Nat Commun.* 2014; 5: 5421.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
63. Sánchez M, Valencia A, Ferrándiz MJ, *et al.*: Correlation between the structure and biochemical activities of FtsA, an essential cell division protein of the actin family. *EMBO J.* 1994; 13(20): 4919–25.
[PubMed Abstract](#) | [Free Full Text](#)
64. van den Ent F, Löwe J: Crystal structure of the cell division protein FtsA from *Thermotoga maritima*. *EMBO J.* 2000; 19(20): 5300–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
65. **F** van den Ent F, Amos LA, Löwe J: Prokaryotic origin of the actin

- cytoskeleton. *Nature*. 2001; **413**(6851): 39–44.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
66. Fujita J, Maeda Y, Nagao C, *et al.*: Crystal structure of FtsA from *Staphylococcus aureus*. *FEBS Lett*. 2014; **588**(10): 1879–85.
[PubMed Abstract](#) | [Publisher Full Text](#)
 67. Pichoff S, Lutkenhaus J: Tethering the Z ring to the membrane through a conserved membrane targeting sequence in FtsA. *Mol Microbiol*. 2005; **55**(6): 1722–34.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 68. Szwedziak P, Wang Q, Freund SM, *et al.*: FtsA forms actin-like protofilaments. *EMBO J*. 2012; **31**(10): 2249–60.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 69. Lara B, Rico AI, Petruzzelli S, *et al.*: Cell division in cocci: localization and properties of the *Streptococcus pneumoniae* FtsA protein. *Mol Microbiol*. 2005; **55**(3): 699–711.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 70. Krupka M, Rivas G, Rico AI, *et al.*: Key role of two terminal domains in the bidirectional polymerization of FtsA protein. *J Biol Chem*. 2012; **287**(10): 7756–65.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 71. Krupka M, Cabré EJ, Jiménez M, *et al.*: Role of the FtsA C terminus as a switch for polymerization and membrane association. *MBio*. 2014; **5**(6): e02221.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 72. Mura A, Fadda D, Perez AJ, *et al.*: Roles of the essential protein FtsA in cell growth and division in *Streptococcus pneumoniae*. *J Bacteriol*. 2016; pii: JB.00608-16.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 73. Zheng JJ, Perez AJ, Tsui HT, *et al.*: Absence of the KhpA and KhpB (JAG/EloR) RNA-binding proteins suppresses the requirement for PBP2b by overproduction of FtsA in *Streptococcus pneumoniae* D39. *Mol Microbiol*. 2017; **106**(5): 793–814.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 74. Krupka M, Rowlett VW, Morado D, *et al.*: *Escherichia coli* FtsA forms lipid-bound minirings that antagonize lateral interactions between FtsZ protofilaments. *Nat Commun*. 2017; **8**: 15957.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 75. Loose M, Mitchison TJ: The bacterial cell division proteins FtsA and FtsZ self-organize into dynamic cytoskeletal patterns. *Nat Cell Biol*. 2014; **16**(1): 38–46.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 76. Lu C, Reedy M, Erickson HP: Straight and curved conformations of FtsZ are regulated by GTP hydrolysis. *J Bacteriol*. 2000; **182**(1): 164–70.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 77. Mingerance J, Tadros M, Vicente M, *et al.*: Visualization of single *Escherichia coli* FtsZ filament dynamics with atomic force microscopy. *J Biol Chem*. 2005; **280**(21): 20909–14.
[PubMed Abstract](#) | [Publisher Full Text](#)
 78. Li Y, Hsin J, Zhao L, *et al.*: FtsZ protofilaments use a hinge-opening mechanism for constrictive force generation. *Science*. 2013; **341**(6144): 392–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 79. Herricks JR, Nguyen D, Margolin W: A thermosensitive defect in the ATP binding pocket of FtsA can be suppressed by allosteric changes in the dimer interface. *Mol Microbiol*. 2014; **94**(3): 713–27.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 80. Conti J, Viola MG, Camberg JL: FtsA reshapes membrane architecture and remodels the Z-ring in *Escherichia coli*. *Mol Microbiol*. 2018; **107**(4): 558–76.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 81. Geissler B, Elraheb D, Margolin W: A gain-of-function mutation in *ftsA* bypasses the requirement for the essential cell division gene *zipA* in *Escherichia coli*. *Proc Natl Acad Sci U S A*. 2003; **100**(7): 4197–202.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 82. Geissler B, Margolin W: Evidence for functional overlap among multiple bacterial cell division proteins: compensating for the loss of FtsK. *Mol Microbiol*. 2005; **58**(2): 596–612.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 83. Geissler B, Shiomi D, Margolin W: The *ftsA** gain-of-function allele of *Escherichia coli* and its effects on the stability and dynamics of the Z ring. *Microbiology*. 2007; **153**(Pt 3): 814–25.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 84. Pichoff S, Shen B, Sullivan B, *et al.*: FtsA mutants impaired for self-interaction bypass ZipA suggesting a model in which FtsA's self-interaction competes with its ability to recruit downstream division proteins. *Mol Microbiol*. 2012; **83**(1): 151–67.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 85. Liu B, Persons L, Lee L, *et al.*: Roles for both FtsA and the FtsBLQ subcomplex in FtsN-stimulated cell constriction in *Escherichia coli*. *Mol Microbiol*. 2015; **95**(6): 945–70.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 86. Chen Y, Huang H, Osawa M, *et al.*: ZipA and FtsA* stabilize FtsZ-GDP minoring structures. *Sci Rep*. 2017; **7**(1): 3650.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 87. Buske PJ, Levin PA: A flexible C-terminal linker is required for proper FtsZ assembly *in vitro* and cytokinetic ring formation *in vivo*. *Mol Microbiol*. 2013; **89**(2): 249–63.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 88. Gardner KA, JMoore DA, Erickson HP: The C-terminal linker of *Escherichia coli* FtsZ functions as an intrinsically disordered peptide. *Mol Microbiol*. 2013; **89**(2): 264–75.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 89. Sundararajan K, Miguel A, Desmarais SM, *et al.*: The bacterial tubulin FtsZ requires its intrinsically disordered linker to direct robust cell wall construction. *Nat Commun*. 2015; **6**: 7281.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 90. Huecas S, Ramírez-Aportela E, Vergoñós A, *et al.*: Self-Organization of FtsZ Polymers in Solution Reveals Spacer Role of the Disordered C-Terminal Tail. *Biophys J*. 2017; **113**(8): 1831–44.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 91. Coltharp C, Buss J, Plumer TM, *et al.*: Defining the rate-limiting processes of bacterial cytokinesis. *Proc Natl Acad Sci U S A*. 2016; **113**(8): E1044–53.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 92. Lan G, Daniels BR, Dobrowsky TM, *et al.*: Condensation of FtsZ filaments can drive bacterial cell division. *Proc Natl Acad Sci U S A*. 2009; **106**(1): 121–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 93. Yang DC, Peters NT, Parzych KR, *et al.*: An ATP-binding cassette transporter-like complex governs cell-wall hydrolysis at the bacterial cytokinetic ring. *Proc Natl Acad Sci U S A*. 2011; **108**(45): E1052–60.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 94. Du S, Pichoff S, Lutkenhaus J: FtsEX acts on FtsA to regulate divisome assembly and activity. *Proc Natl Acad Sci U S A*. 2016; **113**(34): E5052–61.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 95. Mavrakīs M, Azou-Gros Y, Tsai F, *et al.*: Septins promote F-actin ring formation by crosslinking actin filaments into curved bundles. *Nat Cell Biol*. 2014; **16**(4): 322–34.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 96. Szwedziak P, Wang Q, Bharat TA, *et al.*: Architecture of the ring formed by the tubulin homologue FtsZ in bacterial cell division. *eLife*. 2014; **3**: e04601.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 97. Tsang MJ, Bernhardt TG: A role for the FtsQLB complex in cytokinetic ring activation revealed by an *ftsL* allele that accelerates division. *Mol Microbiol*. 2015; **95**(6): 925–44.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 98. Ortiz C, Casanova M, Palacios P, *et al.*: The hypermorph FtsA* protein has an *in vivo* role in relieving the *Escherichia coli* proto-ring block caused by excess ZapC. *PLoS One*. 2017; **12**(9): e0184184.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 99. Erickson HP, Anderson DE, Osawa M: FtsZ in bacterial cytokinesis: cytoskeleton and force generator all in one. *Microbiol Mol Biol Rev*. 2010; **74**(4): 504–28.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 100. Egan AJ, Vollmer W: The physiology of bacterial cell division. *Ann N Y Acad Sci*. 2013; **1277**: 8–28.
[PubMed Abstract](#) | [Publisher Full Text](#)
 101. Coltharp C, Xiao J: Beyond force generation: Why is a dynamic ring of FtsZ polymers essential for bacterial cytokinesis? *Bioessays*. 2017; **39**(1): 1–11.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. **Anuradha Janakiraman**

Department of Biology, City College of New York, New York, USA

Competing Interests: No competing interests were disclosed.

2. **Elizabeth J Harry**

The ithree Institute, University of Technology Sydney, Sydney, Australia

Competing Interests: No competing interests were disclosed.

3. **Jie Xiao**

Department of Biophysics and Biophysical Chemistry, Johns Hopkins University School of Medicine, Baltimore, USA

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research