REVIEW

Recent advances in understanding how rod-like bacteria stably maintain their cell shapes [version 1; peer review: 3 approved]

Sven van Teeffelen¹, Lars D. Renner¹ ²

¹Department of Microbiology, Institut Pasteur, 75724 Paris Cedex 15, France

²Leibniz Institute of Polymer Research and the Max Bergmann Center of Biomaterials, 01069 Dresden, Germany

V1 First published: 28 Feb 2018, 7(F1000 Faculty Rev):241
<https://doi.org/10.12688/f1000research.12663.1>
Latest published: 28 Feb 2018, 7(F1000 Faculty Rev):241
<https://doi.org/10.12688/f1000research.12663.1>

Abstract

Cell shape and cell volume are important for many bacterial functions. In recent years, we have seen a range of experimental and theoretical work that led to a better understanding of the determinants of cell shape and size. The roles of different molecular machineries for cell-wall expansion have been detailed and partially redefined, mechanical forces have been shown to influence cell shape, and new connections between metabolism and cell shape have been proposed. Yet the fundamental determinants of the different cellular dimensions remain to be identified. Here, we highlight some of the recent developments and focus on the determinants of rod-like cell shape and size in the well-studied model organisms *Escherichia coli* and *Bacillus subtilis*.

Keywords

bacterial functions, cell shape, cell size, bacterial metabolism, rod-like, *E.coli*, *B.subtilis*, bacterial physiology

Open Peer Review

Approval Status

	1	2	3
version 1
28 Feb 2018			

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Tobias Doerr**, Cornell University, Ithaca, USA
2. **Rut Carballido-Lopez**, Université Paris-Saclay, Jouy-en-Josas, France
3. **Kevin D Young**, University of Arkansas for Medical Sciences, Little Rock, USA

Any comments on the article can be found at the end of the article.

Corresponding author: Sven van Teeffelen (sven.van-teeffelen@pasteur.fr)

Author roles: van Teeffelen S: Writing – Original Draft Preparation, Writing – Review & Editing; Renner LD: Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: S.v.T. and L.D.R. were supported by the Volkswagen Foundation. S.v.T. was supported by funds from the European Research Council (ERC-2015-STG RCSB 679980), the Mairie de Paris 'Emergence(s)' program, and the ANR 'Investissement d'Avenir Programme' (10-LABX-62-IBEID).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2018 van Teeffelen S and Renner LD. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: van Teeffelen S and Renner LD. **Recent advances in understanding how rod-like bacteria stably maintain their cell shapes [version 1; peer review: 3 approved]** F1000Research 2018, 7(F1000 Faculty Rev):241 <https://doi.org/10.12688/f1000research.12663.1>

First published: 28 Feb 2018, 7(F1000 Faculty Rev):241 <https://doi.org/10.12688/f1000research.12663.1>

Introduction

Shape and size of bacteria are important for many aspects of bacterial physiology, such as proper growth, motility, multicellularity, and host invasion¹. The wide range of diverse cell shapes and sizes in bacteria—from spherical to rod to even more bizarre star shapes covering 4 orders of magnitude in size—is astounding^{1,2}. What is even more impressive is the precision of how bacterial cells control and maintain their cell shapes. Accordingly, cells have evolved the capacity to tightly regulate their own morphologies. Physically, cell shape is determined by the geometry of the peptidoglycan (PG) cell wall. The PG cell wall is a covalently bound meshwork of glycan strands that are interconnected by peptide bonds^{3,4}. The glycan strands are likely oriented around the cell circumference in both Gram-negative and Gram-positive bacteria, which is supported by both cryo-electron tomography imaging^{5,6} and live-cell imaging of circumferential cell-wall insertion^{7–9}. The PG meshwork mechanically resists the high turgor pressure and gives the cell its specific cell shape. As a consequence, without cell wall, bacteria lose their well-defined cell shapes (spheroplasts or L-forms^{10–13}). At the same time, carefully isolated cell-wall sacculi retain the shape of the cell^{5,14,15}. While the cell wall is an elastic material that can undergo small changes of its surface area upon changes of turgor pressure (surface area can be stretched or compressed by about 5–20% in *Escherichia coli*¹⁶), larger changes of morphology during growth require enzymatic remodeling of the cell wall. Specifically, existing bonds in the cell wall must be cleaved and new PG material must be inserted to avoid the formation of large holes that otherwise would lead to lysis¹⁷. Regulation of cell-shape and cell-envelope integrity therefore requires the tight organization of cell-wall remodeling in space and time.

Bacteria organize the expansion of their cell walls in different ways. Some species add new cell wall exclusively at their poles, whereas others insert material in particular zones and yet others insert material at multiple (dispersed) locations along the long axis of the cell^{18,19}. In this review, we will focus on the two best-studied model rods—*E. coli* and *Bacillus subtilis*—that display dispersed lateral cell-wall synthesis.

Although many of the molecular players involved in cell-shape regulation are known today²⁰, neither the physical mechanisms nor the regulatory cascades underlying different variables of shape (for example, cell diameter, cell length, and cell volume) are fully understood and they will be the subject of this review with a specific focus on rod shape.

Technological advances have made it possible to observe cell-wall synthesis in real time, either indirectly by following the dynamics of individual proteins involved in cell-wall expansion (for example, 7–9,21,22) or directly by imaging the incorporation of fluorescently labeled D-amino acids into the PG wall^{23,24}. These tools paired with novel techniques to perturb cell shape^{25,26} and the cell envelope-synthesizing machinery have revealed potential physical determinants of when and where cell-wall expansion happens. For example, experiments of physically bending cells have led to insights into the role of physical forces in shape regulation^{27–29}. At the same time, advances have been

made to connect cell shape with cell physiology at the level of both side-wall elongation^{30,31} and cell-division timing^{32–37}. Here, we will highlight recent advances and open questions. We will first focus on the molecular and physical determinants of cell shape and then summarize newly identified and hypothesized connections between cell shape and cell physiology (in particular, metabolism and the chromosome replication cycle).

The molecular and physical determinants of cell-wall remodeling

Changes of cell shape during growth require enzymatic remodeling of the cell wall^{3,4,20}. Proper cell-wall expansion, in turn, requires the coordination of multiple cytoplasmic and extra-cellular or periplasmic steps: (i) precursor synthesis and flipping from the cytoplasm to the extra-cellular/periplasmic side, (ii) cell-wall cleavage by DD-endopeptidases and possibly other hydrolases for cell-wall expansion along the long axis of the cell, (iii) glycan-strand polymerization (transglycosylation), and (iv) cross-linking to neighboring strands (transpeptidation). The PG cell wall is modified through other processes. For example, lytic transglycosylases cleave glycan strands, and PG material is taken out of the cell wall and recycled. The role of these processes might be to aid in the relaxation of mechanical stresses, but their mechanistic roles are still not understood. Most of the enzymes responsible for the different steps are known and described in great genetic and biochemical detail in previous reviews^{4,20}. Here, we will discuss recent developments toward a better understanding of how the different processes of remodeling the cell wall are implicated in cell-shape regulation in rod-like bacteria, and we will focus on the model systems *E. coli* and *B. subtilis*.

The role of synthesis and hydrolysis for cell-wall expansion

Physically, cleavage of the cell wall is strictly required for cell-wall expansion during growth. In agreement with this idea, endopeptidases have been found to be essential for Gram-negative *E. coli*³⁸ and *Vibrio cholerae*³⁹ and for Gram-positive *B. subtilis*^{40,41}. Endopeptidases are thought to be particularly important, as they cut the bonds between neighboring glycan strands to allow the expansion of the cell wall along the long axis of the cell. Nevertheless, cell-wall cleavage has long been thought to be enslaved to cell-wall insertion in Gram-negative bacteria, where the cell wall is thought to be a thin layer^{5,42,43}. According to the “make-before-break” hypothesis, peptide bonds are cleaved only once a nascent glycan strand is ready for insertion. Höltje suggested that one glycan strand is cleaved out of the wall only after three new glycan strands have been connected to the neighboring strands—all performed by the same multi-enzyme rod complex (the 3-for-1 model)⁴. Similarly, a recent study by Billaudeau *et al.*⁴⁴ suggests that three new glycan strands are added to two existing strands at the closest PG layer in *B. subtilis*⁸. So far, co-localization of cell-wall synthases and endopeptidases has been observed only in *B. subtilis*⁴⁵ but not in Gram-negative bacteria. It is therefore also conceivable that cell-wall hydrolysis proceeds somewhat independently and possibly even upstream of cell-wall insertion³⁸. In Gram-positive *B. subtilis*, the cell wall is about 30 nm thick^{46,47}. As new cell wall is added underneath the existing cell wall, old material must be cleaved at least partially at locations that are spatially

disconnected from regions of synthesis. According to the “inside-out” model proposed by Koch and Doyle, new layers of PG are added underneath the cell wall while old outer layers are taken away by hydrolases⁴⁸. Thus, the coupling between hydrolysis and synthesis might be largely indirect⁴⁶, such that the turgor pressure drives PG cleavage, and PG synthesis is required merely to prevent the cell wall from thinning over time.

Recently, a long-standing paradigm for a single multi-enzyme machinery that inserts PG into the cylindrical part of rod-shaped *E. coli* and *B. subtilis* has been challenged. Instead, evidence suggests that two machineries contribute to side-wall synthesis somewhat independently²¹ (see Figure 1B, C): on the one hand, the rod complex consisting of the mono-functional transpeptidase PBP2 (PBP2A and PBPH in *B. subtilis*) and

the transglycosylase RodA in concert with the cytoskeletal MreB (the rod complex); on the other hand, the bi-functional class A penicillin-binding proteins (aPBPs) (importantly, PBP1a and PBP1b in *E. coli*²¹ and PBP1, PBP2c, and PBP4 in *B. subtilis*^{49,50}). The two types of machineries behave qualitatively differently and might have distinct functions. The rod complex moves around the cell in a persistent manner corresponding to processive insertion of long glycan strands^{7–9,21}, while the aPBPs move seemingly diffusively with intermittent pauses potentially corresponding to localized cell-wall insertion events^{21,22}. Yet the machineries seem to largely depend on each other’s functionalities, as cell-wall synthesis is reduced by about 80% upon inhibition of either complex²¹. It is thus possible that they work together and even form transient joint complexes. Elucidating the strengths and durations of these interactions will be

Figure 1. Schematic representation of the prevailing mechanisms of cell-wall synthesis in rod-shaped bacteria. (A) Illustration of a rod-shaped cell indicating the rod complex, hydrolases, and class A penicillin-binding proteins (aPBPs) during cell growth (peptidoglycan [PG] elongation) and of the divisome during cell division (PG constriction). Despite decades of research on bacterial cell-wall growth, there are still many open questions about the factors that can actively influence PG assembly and cell growth, such as metabolism, mechanical stresses, and cell shape. (B) Inset showing the major cell-wall synthesis machineries recently suggested to work partially independently²¹: (i) the processively moving rod complex containing the transglycosylase RodA, the transpeptidase PBP2, and circumferentially oriented MreB filaments and (ii) the bi-functional aPBPs. Hydrolases may be actively engaged in cell-wall cleavage during cell-wall synthesis. Questions remain of how hydrolases interact with the rod complex or aPBPs or both. (C) aPBPs are activated through outer membrane (OM) lipoproteins LpoA and LpoB. This interaction could provide a mechanism to sense pore sizes in the cell wall to direct PG synthesis. (D) Division in rod-shaped bacteria is guided by treadmilling FtsZ filaments that are oriented along the circumference of the constricting cell. IM, inner membrane.

important to fully understand the importance of the different components for cell-wall integrity and rod shape.

The Bernhardt lab recently found evidence for cell-wall synthesis through aPBPs partially being activated downstream of cell-wall cleavage in *E. coli*⁵¹. Upon overexpression of the DD-endopeptidase MepS, cell-wall synthesis was found to increase and synthesis was ascribed largely to aPBP activity. Thus, aPBPs might sense the holes generated by DD-endopeptidases (see Figure 1B, C). We expand on the possibility of sensing pore size in the next section. Whether DD-endopeptidases can act fully independently of cell-wall synthesis will have to be seen in future experiments (for example, by reducing or inhibiting cell-wall synthesis and monitoring cell-wall hydrolysis).

The role of mechanical forces for cell-envelope expansion

Prior to many molecular observations of cell-wall remodeling, Koch had suggested, in a pioneering discussion paper, that the rate of cell elongation is controlled by “smart autolysins” that cleave the PG cell wall in response to the buildup of mechanical stress in the cell wall in Gram-negative and Gram-positive bacteria⁴². Stress in the cell wall, in turn, is due to the high intracellular turgor pressure. The pressures of about 0.3–3 atm in *E. coli*^{52,53} and 10 atm in *B. subtilis*⁵⁴ are caused by high concentrations of osmolites in the cytoplasm⁵⁵. Pressure-dependent cell-wall expansion has since been confirmed for different walled organisms, in particular for fission yeast⁵⁶ and plant cells⁵⁷ (see a recent review by Rojas and Huang⁵⁸). Very recent experiments by Rojas *et al.* in *B. subtilis* suggest that pressure also has an important role in regulating the elongation rate of Gram-positive bacteria⁵⁹. The authors found that a rapid reduction or a rapid increase of pressure through osmotic shocks reduces the rate of cell elongation and also the rate of cell-wall synthesis. A decrease of cell-wall synthesis upon hypoosmotic shock is likely caused by an increase in membrane tension and possibly by a tension-induced depolarization of the membrane. Rojas *et al.* suggest that the interplay among pressure, cell-wall stress, and membrane polarization guarantees that membrane surface area and cell-wall area remain coupled at all times. However, whether pressure is also responsible for coupling overall cellular biomass growth to the rate of cell-envelope expansion remains to be studied by assessing biomass growth and cell elongation independently. In the same realm, it will be interesting to elucidate how the reduction of pressure leads to a reduction of cell elongation mechanistically—through smart autolysins, through tension-dependent synthases as proposed by Rojas *et al.*, or completely indirectly through changes of the overall growth rate of the bacteria. For the Gram-negative *E. coli*, the behavior is very different: the same authors showed, in a previous paper, that osmotic pressure was neither required nor responsible for cell-wall expansion on short time scales⁶⁰.

Despite the robustness of PG expansion in *E. coli* with respect to pressure, differential mechanical stresses in the cell wall have an impact on cell-wall expansion also in *E. coli*. Previous work has demonstrated that cells grow into curved or pancake-like shapes if physically bent or squeezed^{28,61–65}. In one of the first studies, Takeuchi *et al.* showed that rod-shaped *E. coli* can be

subjected, confined, and adapted to various shapes upon antibiotic-induced filamentous growth²⁸. Recent studies by Amir *et al.* have demonstrated that cells fully recover from bent shapes: they straighten out at a rate almost twice as fast as their rate of elongation^{27,65}. In quantitative agreement with a theoretical study of the mechanical stresses in the cell envelope during bending and straightening, the study suggests that both bending and straightening are due to asymmetric mechanical stress distributions in the cell wall²⁷. As a subtlety, Wong *et al.*²⁷ found that differential stresses along the long axis of the cell were too small to explain straightening. This is in agreement with previous work from Emonet *et al.*⁶⁶. However, they found relevant differences in stresses along the circumference of the cell, which could be responsible for differences in the pore size of the PG meshwork. An attractive hypothesis for the sensing of pore sizes in Gram-negative bacteria is that the aPBPs PBP1a and PBP1b together with their cognate outer-membrane lipoprotein activators LpoA and LpoB^{67,68} (Figure 1C), respectively, sense the pore size in the bacterial cell wall²⁰ and thus insert more material at sites where pore sizes are increased because of mechanical stress. However, whether cell-wall synthases or hydrolases are responsible for shape changes upon mechanical forces and for the recovery to normal rod-like shape remains to be studied.

The role of the MreB cytoskeleton for rod-like cell shape

The model of stress-induced bending and straightening is in contrast to a previous geometry-based model by Ursell *et al.*⁶⁹. The bacterial cytoskeleton MreB localizes to the inner regions of bent cells^{64,69}. Huang and colleagues⁶⁹ argued that the intrinsic curvature and twist of MreB filaments might be responsible for localizing MreB and thus the cell-wall synthesis machinery differentially to these regions. However, according to the more recent study by Wong *et al.*²⁷, differences in MreB localization were found to be too weak to account for the observed straightening dynamics. Instead, the MreB asymmetric localization observed could also be due to a combination of curved cell geometry and persistent circumferential motion²⁷, as previously discussed⁶⁶.

In the same study by Ursell *et al.*, the authors found that MreB filaments are attracted to regions of negative Gaussian curvature and excluded from regions of positive Gaussian curvature, such as the cell poles⁶⁹, even in cells that are not bent. However, it remains to be studied how strong and important this effect is for the cylindrical part of the cell. In *E. coli*, the exclusion of MreB from the cell poles might be due to a curvature-independent, pole-specific effect: Kawazura *et al.* recently showed that the pole-specific enrichment for anionic phospholipids is required to exclude MreB filaments from cell poles⁷⁰.

The filamentous nature of the MreB cytoskeleton was long under debate^{71,72}. Most recent measurements of fluorescent-protein fusions to MreB indicate that MreB forms independent filaments of a few hundred nanometers to a few micrometers in length in *E. coli*^{73,74} and *B. subtilis*⁷⁵ that are oriented circumferentially in both *E. coli* and *B. subtilis*^{73,76}, with a slight chiral twist in *E. coli*⁷³. Interestingly, the orientation of MreB filaments seems to

be directly coupled to the orientation of MreB tracks around the cell circumference and thus to the orientation of newly inserted PG material^{7,73,76}, in agreement with the macroscopic chiral organization of the cell wall²⁹. As MreB forms intrinsically curved filaments *in vitro*⁷⁷, curvature and potentially twist thus make MreB a strong candidate for independently mediating its own orientation inside the cell^{73,78}. This picture is supported by a recent publication from the Garner and Löwe labs, which suggests that the intrinsic orientation of MreB is responsible for the circumferential orientation of MreB filaments and tracks in *B. subtilis*⁷⁹. They demonstrate that MreB filaments assume the direction of largest principal curvature if cell shape is imposed in microchambers, and they conclude that MreB's track orientation is independent of any major local cell-wall structure. They therefore conclude that MreB's intrinsic orientation solely determines the orientation of newly inserted glycan strands⁷⁹. The orientation preference of MreB might also be responsible for the exclusion of MreB from the poles: MreB filaments that move into the poles rapidly leave the poles again, while they are stabilized in the cylindrical part of the cell.

In *E. coli*, the picture could be more complicated. The correlations between MreB and cell-wall orientations could also result from a more subtle feedback loop between MreB's filament properties and the properties of the cell-wall synthesis process. While MreB likely biases cell-wall insertion toward a near-circumferential orientation, the rod complex might also follow the existing glycan strands as a template for the insertion of new strands. The glycan-strand orientation, in turn, could come about due to a combination of MreB's intrinsic orientation and chiral biases during cell-wall insertion. This hypothesis is in agreement with Huang *et al.*, who showed that the chirality of MreB tracks can be affected by perturbations different from direct perturbations of MreB itself⁸⁰.

Ultimately, it seems reasonable to assume that the local properties of the cell wall and cell shape jointly determine where new cell-wall remodeling takes place. Jensen *et al.* recently suggested a numerical simulation of the cell wall on the basis of this principle that maintains cell shape over many generations of growth⁸¹. The model showed that careful local organization of cell-wall synthesis based on the microscopic cell-wall geometry together with a bias of cell-wall insertion toward a circumferential direction is sufficient to maintain rod-like cell shape for many generations⁸¹. Therefore, local interactions of enzymes with the cell wall together with the circumferential orientation sensed by MreB could be responsible for cell-wall integrity and cell-wall orientation. However, neither this nor previous models have established a physical determinant of cell diameter, and we simply do not yet understand how cell diameter can be highly adaptive to different growth environments but precisely maintained in cells growing in identical environmental conditions. We will come back to the determinants of cell diameter in one of the following sections.

Cytokinesis and the septal ring

During cell division, both Gram-positive and Gram-negative rods ultimately form two hemispherical poles at midcell⁸². The coordination of cell constriction and cell-wall synthesis is

complex and requires the interplay of dozens of proteins^{82,83}. A major determinant of the localization and orientation of the cell-division machinery is the tubulin homolog, FtsZ, together with accessory Zap proteins⁸⁴. FtsZ filaments are oriented circumferentially at midcell and thus are responsible for the circumferential orientation of glycan-strand insertion⁸⁵. FtsZ filaments were recently found to effectively move by treadmilling in *B. subtilis* and *E. coli*^{86,87} (Figure 1D). Unlike the way in which MreB is driven around the cell by the rod complex, FtsZ treadmilling is responsible and rate-limiting for cell-wall insertion and septation in *B. subtilis*⁸⁷. In *E. coli*, treadmilling is required for proper cell septation, but the rate of septation is surprisingly reported not to be limited by FtsZ treadmilling⁸⁸.

FtsZ was previously argued to have a role in exerting a rate-limiting constrictive force during septation, based on structural data⁸⁹ and *in vitro* work⁹⁰. However, more recent work has questioned this function⁸⁸, as changes in the GTPase activity of the FtsZ ring did not significantly change the rate of constriction. On the contrary, perturbing the activity of major cell wall-modifying enzymes (notably, FtsI) led to changes of the rate of constriction. Therefore, it remains to be discovered how the dynamics of FtsZ and the cell-wall synthesis machinery jointly lead to the formation of two hemispherical poles at midcell.

Cell shape, metabolism, and the cell cycle

One, if not the foremost, function of cell shape is to provide a well-defined volume for all intracellular processes required for the completion of every cell cycle. Empirically, average cell size is an exponentially increasing function of growth rate if growth rate changes because of nutrient limitation^{91–93}. This fundamental “growth law” was first identified by Schaechter *et al.*⁹¹. In rod-like bacteria, both cell diameter and average cell length contribute to the change in average cell volume. Yet the physical processes responsible for cell diameter and average cell volume are quite different: cell diameter is physically constrained by the cell wall. Accordingly, perturbations of cell-wall synthesis at the level of precursor synthesis, the rod complex, or the MreB cytoskeleton increase the cell diameter^{73,80,94}. During growth, the cell wall constantly expands to create space for the accumulation of intracellular biomass; hence, the rate of expansion must be tied to the rate of biomass growth. Average cell volume, on the contrary, is determined largely by the timing of cell division. Timing of cell division has often been ascribed to DNA replication^{33,95,96}, but recent studies suggest that the cell envelope could also play an important role in cell-size regulation^{30,31}. Here, we discuss recent findings and remaining questions regarding both aspects of rod elongation and division and put particular emphasis on the potential positive regulatory roles of PG precursor production, PG hydrolysis, and membrane synthesis. For a more detailed account of upstream components that might be responsible for changes in precursor levels, see an excellent review by Sperber and Herman⁹⁷.

The plasticity of cell diameter upon metabolic limitations of cell-envelope expansion

It was recently demonstrated by different labs that reducing cell-envelope capacity affects cell diameter in different ways^{30,31,98}. Most recently, Levin *et al.* showed that limitations of

phospholipid synthesis jointly reduce cell diameter and cell length according to Schaechter's growth law³⁰. Upon chemical perturbations of fatty-acid metabolism, cells produced fewer phospholipids, which led to reductions of both growth rate and cell size. Importantly, they also found that synthetic upregulation of lipid synthesis increased cell diameter, even if the growth rate was reduced. They thus proposed that membrane capacity was responsible for cell dimensions also under normal growth conditions. However, the mechanism responsible for the proposed coupling between lipid metabolism and the cell wall, which is ultimately responsible for cell diameter, remains to be identified. Levin *et al.* suggested that such coupling could happen indirectly through PG metabolism or through a common pathway component, undecaprenyl pyrophosphate. Following a similar logic, Harris and Theriot had previously hypothesized that both cell diameter and the timing of cell division could be governed largely by cell-wall metabolism³¹.

Similarly to Levin *et al.*, Harris and Theriot titrated the capacity of bacteria to build their cell envelope, here by using a chemical inhibitor of PG-precursor synthesis (fosfomycin)³¹. Unlike in the perturbations of lipid metabolism, cells increased rather than decreased their cell diameter upon mild perturbations of PG availability while maintaining a constant growth rate. A similar observation was made by Dörr *et al.*⁹⁹ in *V. cholerae*: cells became wider or thinner upon a decrease or increase of cell-wall synthesis, respectively⁹⁹. Harris and Theriot argued that cells adjust their cell diameter to attain a desired ratio of surface area to volume (SA/V ratio), which is compatible with the limited resources of PG production. The SA/V ratio is governed predominantly by cell diameter and the natural quantity of interest under two interesting assumptions we will discuss further: first, the rate of PG synthesis per total cytoplasmic proteome is given by the physiological state of the cell and is limiting for the rate of cell-wall expansion. Second, the density of PG per cell-wall area is constant and is based on the conservation of this quantity both under normal growth conditions and upon perturbations of the MreB cytoskeleton¹⁰⁰. Under these two assumptions, the SA/V ratio is fully determined by the rate of PG synthesis per cytoplasmic volume and the density of PG per cell-wall area. But as a complication and challenge to the simple model, the SA/V ratio changes during the cell cycle. The ratio decreases after birth and then increases toward division because of the formation of a septum. To rescue the model, Harris and Theriot hypothesized a pool of PG precursor material that buffers changes of PG synthesis during the cell cycle. Although the model is generally attractive in that it postulates diameter control on the basis of precursor metabolism rather than mechanical details of cell-wall insertion and the cytoskeleton¹⁰¹, the mechanistic implementation of the different proposed roles of PG for the regulation of cell diameter and cell division remains to be discovered.

As an alternative to the hypothesis for cell-shape regulation through precursor production, cell-wall expansion could be controlled at the level of cell-wall cleavage, specifically through the hydrolysis of PG bonds, as already indicated above. Redundantly essential sets of DD-endopeptidases have been identified in different model bacteria^{38,41}. Thus, the regulation of cell-wall

hydrolases is highly important for the maintenance of rod-like shape. Accordingly, variations in hydrolase activity through substrate modification¹⁰² or through changes of hydrolase abundance lead to changes in cell shape¹⁰³. Thus, it is conceivable that the rate of cell-surface expansion is regulated through hydrolase activity while cell-wall synthesis would be responsible merely for maintaining cell-wall integrity.

The studies by Levin *et al.* and Theriot *et al.* provide valuable insights into the plasticity of cell diameter upon perturbations of different aspects of cell-envelope expansion. While each of the studies proposes a single pathway of cell-envelope biogenesis to be responsible for cell diameter under normal growth conditions, it is also conceivable that fluxes or pool sizes of PG precursors and phospholipids, together with other regulatory components, jointly determine cell diameter during regular growth. Indeed, the surface areas of cell wall and cell membranes are almost equal during normal growth⁴³. Their synthesis is therefore likely coordinated. Coordination could happen both on a metabolic and on a physical level. As already indicated above, changes of membrane metabolism could feed back on PG metabolism (for example, through the regulatory molecule ppGpp). Physically, cell wall and membranes are coupled in multiple ways: as already discussed above, mechanical stress in the cell membrane could serve to couple cell-wall expansion to membrane synthesis⁵⁹. Previous work by Ehlert and Hölte¹⁰⁴ demonstrated that membrane synthesis is required for the flipping of PG precursor material, providing yet another way of coupling the two envelope components. Furthermore, many cell-wall remodeling enzymes are embedded in the cytoplasmic membrane and thus could serve as sensors of the local distance between lipid membrane and cell wall.

Together, cell diameter and the rates of membrane and PG synthesis could be limited by multiple pathways and could feed back on each other. This cycle would be controlled analogously to other processes such as overall growth rate, which is determined and limited simultaneously by different processes, in particular through ribosome levels, amino acid concentration, and nutrient availability^{105,106}. Determining causality and identifying the role of each component for diameter control under normal growth conditions will likely require a more mechanistic understanding of their respective contributions.

Average cell size is governed by the timing of cell division

Contrary to Schaechter's growth law, if the growth rate is reduced because of the excess production of unnecessary proteins, cell size increases with decreasing growth rate⁹³. Both Schaechter's growth law and the inverse behavior are intuitive. A cell requires a larger number of ribosomal and metabolic proteins at fast growth compared with slow growth to complete a larger set of tasks in a shorter time. The production of unnecessary proteins requires space for the protein itself and for additional proteins and ribosomes devoted to their production. Although these relationships are intuitive, we still have not understood the regulatory mechanisms underlying cell size control in different physiological conditions, and their discovery is a very active field of research³⁷. Following the establishment of Schaechter's growth law⁹¹, Donachie¹⁰⁷ and Helmstetter *et al.*¹⁰⁸ found a striking

relationship between chromosome replication and cell size that since has served as a major motivation to study DNA replication as a candidate for the regulatory process upstream of size control: Donachie found that cells initiate new rounds of DNA replication on average at a well-defined cell size that is independent of growth rate¹⁰⁷. This study was recently extended and confirmed for many alternative ways of changing growth rate, cell size, and the replication cycle¹⁰⁹.

From a physiological point of view, it makes sense that DNA replication serves as a checkpoint for cell division (that is, that cells divide only once every daughter cell has inherited one complete chromosome). Accordingly, cells with a reduced rate of DNA replication show a larger average cell size. However, alternatively, it is also possible that there are other checkpoints for cell division that might be limiting depending on physiological conditions. Harris and Theriot suggested that cells accumulate the PG material required for cell septation during the course of a cell cycle and that this “excess material” is responsible for the timing of cell division³¹. Levin *et al.* suggested that membrane capacity and carbon catabolism limit the time of cell septation^{30,110,111}. Similarly to the question of what determines cell diameter, it remains to be discovered which of the processes is limiting under which conditions.

A powerful method to identify causal relationships between different cell cycle-related quantities is to study correlations at the single-cell level. With this approach, bacteria and eukaryotes were recently found to divide after adding a well-defined volume to their cell size at birth, as opposed to dividing at a predetermined cell size (see, e.g., 33,93). Whether the adder principle is a result of coupling cell division to replication or whether there is an independent mechanism coupling cell division to cell size or added cell size will have to be studied in the future.

Conclusions

Recent experiments and theoretical models have led to interesting new findings and hypotheses about the determinants of cell shape in rod-like bacteria, of which we could present only a subset in this review because of space constraints. We can now observe the behavior of the microscopic machinery responsible for cell-wall remodeling at an unprecedented resolution at the

single-cell level. At the same time, genetics, biochemistry, and structural biology provide insights into candidate regulators and modulators that can be tested for their physical role in cell-shape regulation. Thus, coarse-grained physical models of cell shape can now be related to the action of specific proteins. At the same time, we can observe aspects of cellular physiology, such as growth rate, metabolism, and energetic state, with increasing resolution and accuracy. We can thus relate the physics of cell shape to other physiological observables by using coarse-grained descriptions of regulatory interactions and identifying specific molecular pathways responsible for the interactions.

Whereas the last 30 years have focused largely on the role of individual genes and proteins for cellular physiology in isolation, the coming years will likely provide a plethora of insights into their actual physical role in the cell by taking the response of the cell into account. We will thus likely obtain answers to questions that were often raised decades ago on the basis of empirical relationships such as Schaechter’s growth law of cell size or the Donachie relationship between DNA replication and division. Yet many challenges remain, as causative relationships have to be deduced from correlations and as the potential interactions between different modules of the cell are almost countless. An important role will thus be played by mathematically and physically modeling the cell at different levels of resolution. Very successful examples of such models can be found in the study of cellular growth^{112–114} and the cell cycle¹¹⁵.

Competing interests

The authors declare that they have no competing interests.

Grant information

S.v.T. and L.D.R. were supported by the Volkswagen Foundation. S.v.T. was supported by funds from the European Research Council (ERC-2015-STG RCSI 679980), the Mairie de Paris ‘Emergence(s)’ program, and the ANR ‘Investissement d’Avenir Programme’ (10-LABX-62-IBEID).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- Young KD: **The selective value of bacterial shape.** *Microbiol Mol Biol Rev.* 2006; **70**(3): 660–703.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Jiang C, Caccamo PD, Brun YV: **Mechanisms of bacterial morphogenesis: evolutionary cell biology approaches provide new insights.** *Bioessays.* 2015; **37**(4): 413–25.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Vollmer W, Blanot D, de Pedro MA: **Peptidoglycan structure and architecture.** *FEMS Microbiol Rev.* 2008; **32**(2): 149–67.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Höltje JV: **Growth of the stress-bearing and shape-maintaining murein sacculus of *Escherichia coli*.** *Microbiol Mol Biol Rev.* 1998; **62**(1): 181–203.
[PubMed Abstract](#) | [Free Full Text](#)
- Gan L, Chen S, Jensen GJ: **Molecular organization of Gram-negative peptidoglycan.** *Proc Natl Acad Sci U S A.* 2008; **105**(48): 18953–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Beeby M, Gumbart JC, Roux B, *et al.*: **Architecture and assembly of the Gram-positive cell wall.** *Mol Microbiol.* 2013; **88**(4): 664–72.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- van Teeffelen S, Wang S, Furchtgott L, *et al.*: **The bacterial actin MreB rotates, and rotation depends on cell-wall assembly.** *Proc Natl Acad Sci U S A.*

- 2011; **108**(38): 15822–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
8. **F** Domínguez-Escobar J, Chastanet A, Crevenna AH, *et al.*: **Processive movement of MreB-associated cell wall biosynthetic complexes in bacteria.** *Science*. 2011; **333**(6039): 225–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 9. **F** Garner EC, Bernard R, Wang W, *et al.*: **Coupled, circumferential motions of the cell wall synthesis machinery and MreB filaments in *B. subtilis*.** *Science*. 2011; **333**(6039): 222–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 10. Lederberg J, St Clair J: **Protoplasts and L-type growth of *Escherichia coli*.** *J Bacteriol*. 1958; **75**(2): 143–60.
[PubMed Abstract](#) | [Free Full Text](#)
 11. Errington J: **L-form bacteria, cell walls and the origins of life.** *Open Biol*. 2013; **3**(1): 120143.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 12. Dörr T, Davis BM, Waldor MK: **Endopeptidase-mediated beta lactam tolerance.** *PLoS Pathog*. 2015; **11**(4): e1004850.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 13. Monahan LG, Turnbull L, Osvath SR, *et al.*: **Rapid conversion of *Pseudomonas aeruginosa* to a spherical cell morphology facilitates tolerance to carbapenems and penicillins but increases susceptibility to antimicrobial peptides.** *Antimicrob Agents Chemother*. 2014; **58**(4): 1956–62.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 14. Poindexter JS, Hagenzieker JG: **Constriction and septation during cell division in caulobacters.** *Can J Microbiol*. 1981; **27**(7): 704–19.
[PubMed Abstract](#) | [Publisher Full Text](#)
 15. Koch AL, Woeste S: **Elasticity of the sacculus of *Escherichia coli*.** *J Bacteriol*. 1992; **174**(14): 4811–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 16. Koch AL: **Shrinkage of growing *Escherichia coli* cells by osmotic challenge.** *J Bacteriol*. 1984; **159**(3): 919–24.
[PubMed Abstract](#) | [Free Full Text](#)
 17. Huang KC, Mukhopadhyay R, Wen B, *et al.*: **Cell shape and cell-wall organization in Gram-negative bacteria.** *Proc Natl Acad Sci U S A*. 2008; **105**(49): 19282–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 18. Kysela DT, Brown PJ, Huang KC, *et al.*: **Biological consequences and advantages of asymmetric bacterial growth.** *Annu Rev Microbiol*. 2013; **67**: 417–35.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 19. Randich AM, Brun YV: **Molecular mechanisms for the evolution of bacterial morphologies and growth modes.** *Front Microbiol*. 2015; **6**: 580.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 20. Typas A, Banzhaf M, Gross CA, *et al.*: **From the regulation of peptidoglycan synthesis to bacterial growth and morphology.** *Nat Rev Microbiol*. 2011; **10**(2): 123–36.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 21. **F** Cho H, Wivagg CN, Kapoor M, *et al.*: **Bacterial cell wall biogenesis is mediated by SEDS and PBP polymerase families functioning semi-autonomously.** *Nat Microbiol*. 2016; **1**: 16172.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 22. Lee TK, Meng K, Shi H, *et al.*: **Single-molecule imaging reveals modulation of cell wall synthesis dynamics in live bacterial cells.** *Nat Commun*. 2016; **7**: 13170.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 23. Kuru E, Hughes HV, Brown PJ, *et al.*: **In Situ probing of newly synthesized peptidoglycan in live bacteria with fluorescent D-amino acids.** *Angew Chem Int Ed Engl*. 2012; **51**(50): 12519–23.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 24. Hsu Y, Rittichier J, Kuru E, *et al.*: **Full color palette of fluorescent d-amino acids for in situ labeling of bacterial cell walls.** *Chem Sci*. 2017; **8**(9): 6313–21.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 25. Weibel DB, Diluzio WR, Whitesides GM: **Microfabrication meets microbiology.** *Nat Rev Microbiol*. 2007; **5**(3): 209–18.
[PubMed Abstract](#) | [Publisher Full Text](#)
 26. Hol FJ, Dekker C: **Zooming in to see the bigger picture: microfluidic and nanofabrication tools to study bacteria.** *Science*. 2014; **346**(6208): 1251821.
[PubMed Abstract](#) | [Publisher Full Text](#)
 27. Wong F, Renner LD, Özbaykal G, *et al.*: **Mechanical strain sensing implicated in cell shape recovery in *Escherichia coli*.** *Nat Microbiol*. 2017; **2**: 17115.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 28. **F** Takeuchi S, Diluzio WR, Weibel DB, *et al.*: **Controlling the shape of filamentous cells of *Escherichia coli*.** *Nano Lett*. 2005; **5**(9): 1819–23.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 29. Wang S, Arellano-Santoyo H, Combs PA, *et al.*: **Actin-like cytoskeleton filaments contribute to cell mechanics in bacteria.** *Proc Natl Acad Sci U S A*. 2010; **107**(20): 9182–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 30. Vadia S, Tse JL, Lucena R, *et al.*: **Fatty Acid Availability Sets Cell Envelope Capacity and Dictates Microbial Cell Size.** *Curr Biol*. 2017; **27**(12): 1757–1767.e5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 31. Harris LK, Theriot JA: **Relative Rates of Surface and Volume Synthesis Set Bacterial Cell Size.** *Cell*. 2016; **165**(6): 1479–92.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 32. Voorn W, Koppes L, Grover N: **Mathematics of cell division in *Escherichia coli*.** *Curr Top Mol Genet*. 1993; **1**(20): 187–194.
 33. Amir A: **Cell Size Regulation in Bacteria.** *Phys Rev Lett*. 2014; **112**(20): 208102.
[Publisher Full Text](#)
 34. Taheri-Araghi S, Brown SD, Sauls JT, *et al.*: **Single-Cell Physiology.** *Annu Rev Biophys*. 2015; **44**: 123–42.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 35. Campos M, Surovtsev IV, Kato S, *et al.*: **A constant size extension drives bacterial cell size homeostasis.** *Cell*. 2014; **159**(6): 1433–46.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 36. Banerjee S, Lo K, Daddysman MK, *et al.*: **Biphasic growth dynamics control cell division in *Caulobacter crescentus*.** *Nat Microbiol*. 2017; **2**: 17116.
[PubMed Abstract](#) | [Publisher Full Text](#)
 37. Willis L, Huang KC: **Sizing up the bacterial cell cycle.** *Nat Rev Microbiol*. 2017; **15**(10): 606–20.
[PubMed Abstract](#) | [Publisher Full Text](#)
 38. **F** Singh SK, SaiSree L, Amrutha RN, *et al.*: **Three redundant murein endopeptidases catalyze an essential cleavage step in peptidoglycan synthesis of *Escherichia coli* K12.** *Mol Microbiol*. 2012; **86**(5): 1036–51.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 39. Dörr T, Cava F, Lam H, *et al.*: **Substrate specificity of an elongation-specific peptidoglycan endopeptidase and its implications for cell wall architecture and growth of *Vibrio cholerae*.** *Mol Microbiol*. 2013; **89**(5): 949–62.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 40. Dubrac S, Bisicchia P, Devine KM, *et al.*: **A matter of life and death: cell wall homeostasis and the WalK (YycG) essential signal transduction pathway.** *Mol Microbiol*. 2008; **70**(6): 1307–22.
[PubMed Abstract](#) | [Publisher Full Text](#)
 41. Hashimoto M, Ooiwa S, Sekiguchi J: **Synthetic lethality of the *lytE* *cwI* genotype in *Bacillus subtilis* is caused by lack of D,L-endopeptidase activity at the lateral cell wall.** *J Bacteriol*. 2012; **194**(4): 796–803.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 42. Koch AL: **The surface stress theory for the case of *Escherichia coli*: the paradoxes of gram-negative growth.** *Res Microbiol*. 1990; **141**(1): 119–30.
[PubMed Abstract](#) | [Publisher Full Text](#)
 43. **F** Matias VR, Al-Amoudi A, Dubochet J, *et al.*: **Cryo-transmission electron microscopy of frozen-hydrated sections of *Escherichia coli* and *Pseudomonas aeruginosa*.** *J Bacteriol*. 2003; **185**(20): 6112–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 44. Billaudeau C, Chastanet A, Yao Z, *et al.*: **Contrasting mechanisms of growth in two model rod-shaped bacteria.** *Nat Commun*. 2017; **8**: 15370.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 45. Carballido-López R, Formstone A, Li Y, *et al.*: **Actin homolog MreBH governs cell morphogenesis by localization of the cell wall hydrolase LytE.** *Dev Cell*. 2006; **11**(3): 399–409.
[PubMed Abstract](#) | [Publisher Full Text](#)
 46. Misra G, Rojas ER, Gopinathan A, *et al.*: **Mechanical consequences of cell-wall turnover in the elongation of a Gram-positive bacterium.** *Biophys J*. 2013; **104**(11): 2342–52.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 47. **F** Matias VR, Beveridge TJ: **Cryo-electron microscopy reveals native polymeric cell wall structure in *Bacillus subtilis* 168 and the existence of a periplasmic space.** *Mol Microbiol*. 2005; **56**(1): 240–51.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 48. Koch AL, Doyle RJ: **Inside-to-outside growth and turnover of the wall of gram-positive rods.** *J Theor Biol*. 1985; **117**(1): 137–57.
[PubMed Abstract](#) | [Publisher Full Text](#)
 49. **F** Meeske AJ, Riley EP, Robins WP, *et al.*: **SEDS proteins are a widespread family of bacterial cell wall polymerases.** *Nature*. 2016; **537**(7622): 634–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 50. Kawai Y, Daniel RA, Errington J: **Regulation of cell wall morphogenesis in *Bacillus subtilis* by recruitment of PBP1 to the MreB helix.** *Mol Microbiol*. 2009; **71**(5): 1131–44.
[PubMed Abstract](#) | [Publisher Full Text](#)
 51. Lai GC, Cho H, Bernhardt TG: **The mecillinam resistome reveals a role for peptidoglycan endopeptidases in stimulating cell wall synthesis in *Escherichia coli*.** *PLoS Genet*. 2017; **13**(7): e1006934.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 52. **F** Deng Y, Sun M, Shaevitz JW: **Direct measurement of cell wall stress stiffening and turgor pressure in live bacterial cells.** *Phys Rev Lett*. 2011; **107**(15): 158101.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 53. Cayley S, Lewis BA, Guttman HJ, *et al.*: **Characterization of the cytoplasm of *Escherichia coli* K-12 as a function of external osmolality. Implications for protein-DNA interactions in vivo.** *J Mol Biol*. 1991; **222**(2): 281–300.
[PubMed Abstract](#) | [Publisher Full Text](#)
 54. Whatmore AM, Reed RH: **Determination of turgor pressure in *Bacillus subtilis*: a possible role for K⁺ in turgor regulation.** *J Gen Microbiol*. 1990; **136**(12): 2521–6.
[PubMed Abstract](#) | [Publisher Full Text](#)

55. Wood JM: **Bacterial osmoregulation: a paradigm for the study of cellular homeostasis.** *Annu Rev Microbiol.* 2011; 65: 215–38.
[PubMed Abstract](#) | [Publisher Full Text](#)
56. **F** Minc N, Boudaoud A, Chang F: **Mechanical forces of fission yeast growth.** *Curr Biol.* 2009; 19(13): 1096–101.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
57. Cosgrove DJ: **Growth of the plant cell wall.** *Nat Rev Mol Cell Biol.* 2005; 6(11): 850–61.
[PubMed Abstract](#) | [Publisher Full Text](#)
58. Rojas ER, Huang KC: **Regulation of microbial growth by turgor pressure.** *Curr Opin Microbiol.* 2017; 42: 62–70.
[PubMed Abstract](#) | [Publisher Full Text](#)
59. Rojas ER, Huang KC, Theriot JA: **Homeostatic Cell Growth Is Accomplished Mechanically through Membrane Tension Inhibition of Cell-Wall Synthesis.** *Cell Syst.* 2017; 5(6): 578–590.e6.
[PubMed Abstract](#) | [Publisher Full Text](#)
60. Rojas ER, Theriot JA, Huang KC: **Response of *Escherichia coli* growth rate to osmotic shock.** *Proc Natl Acad Sci U S A.* 2014; 111(21): 7807–12.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
61. Männik J, Driessen R, Galajda P, *et al.*: **Bacterial growth and motility in sub-micron constrictions.** *Proc Natl Acad Sci U S A.* 2009; 106(35): 14861–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
62. Männik J, Wu F, Hol FJ, *et al.*: **Robustness and accuracy of cell division in *Escherichia coli* in diverse cell shapes.** *Proc Natl Acad Sci U S A.* 2012; 109(18): 6957–62.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
63. Si F, Li B, Margolin W, *et al.*: **Bacterial growth and form under mechanical compression.** *Sci Rep.* 2015; 5: 11367.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
64. Renner LD, Eswaramoorthy P, Ramamurthi KS, *et al.*: **Studying biomolecule localization by engineering bacterial cell wall curvature.** *PLoS One.* 2013; 8(12): e84143.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
65. Amir A, Babaiepour F, McIntosh DB, *et al.*: **Bending forces plastically deform growing bacterial cell walls.** *Proc Natl Acad Sci U S A.* 2014; 111(16): 5778–83.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
66. Sliusarenko O, Cabeen MT, Wolgemuth CW, *et al.*: **Processivity of peptidoglycan synthesis provides a built-in mechanism for the robustness of straight-rod cell morphology.** *Proc Natl Acad Sci U S A.* 2010; 107(22): 10086–91.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
67. **F** Typas A, Banzhaf M, van den Berg van Saparoea B, *et al.*: **Regulation of peptidoglycan synthesis by outer-membrane proteins.** *Cell.* 2010; 143(7): 1097–109.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
68. **F** Paradis-Bleau C, Markovskii M, Uehara T, *et al.*: **Lipoprotein cofactors located in the outer membrane activate bacterial cell wall polymerases.** *Cell.* 2010; 143(7): 1110–20.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
69. **F** Ursell TS, Nguyen J, Monds RD, *et al.*: **Rod-like bacterial shape is maintained by feedback between cell curvature and cytoskeletal localization.** *Proc Natl Acad Sci U S A.* 2014; 111(11): E1025–34.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
70. **F** Kawazura T, Matsumoto K, Kojima K, *et al.*: **Exclusion of assembled MreB by anionic phospholipids at cell poles confers cell polarity for bidirectional growth.** *Mol Microbiol.* 2017; 104(3): 472–86.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
71. Errington J: **Bacterial morphogenesis and the enigmatic MreB helix.** *Nat Rev Microbiol.* 2015; 13(4): 241–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
72. Eun YJ, Kapoor M, Hussain S, *et al.*: **Bacterial Filament Systems: Toward Understanding Their Emergent Behavior and Cellular Functions.** *J Biol Chem.* 2015; 290(28): 17181–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
73. **F** Ouzounov N, Nguyen JP, Bratton BP, *et al.*: **MreB Orientation Correlates with Cell Diameter in *Escherichia coli*.** *Biophys J.* 2016; 111(5): 1035–43.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
74. **F** Swulius MT, Jensen GJ: **The helical MreB cytoskeleton in *Escherichia coli* MC1000/pLE7 is an artifact of the N-Terminal yellow fluorescent protein tag.** *J Bacteriol.* 2012; 194(23): 6382–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
75. **F** Reimold C, Defeu Soufo HJ, Dempwolff F, *et al.*: **Motion of variable-length MreB filaments at the bacterial cell membrane influences cell morphology.** *Mol Biol Cell.* 2013; 24(15): 2340–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
76. Olshausen PV, Defeu Soufo HJ, Wicker K, *et al.*: **Superresolution imaging of dynamic MreB filaments in *B. subtilis*—a multiple-motor-driven transport?** *Biophys J.* 2013; 105(5): 1171–81.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
77. **F** Salje J, van den Ent F, de Boer P, *et al.*: **Direct membrane binding by bacterial actin MreB.** *Mol Cell.* 2011; 43(3): 478–87.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
78. Wang S, Wingreen NS: **Cell shape can mediate the spatial organization of the bacterial cytoskeleton.** *Biophys J.* 2013; 104(3): 541–52.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
79. Hussain S, Wivagg CN, Szewdzik P, *et al.*: **MreB filaments align along greatest principal membrane curvature to orient cell wall synthesis.** *eLife.* 2018; 7: pii: e32471
[PubMed Abstract](#) | [Publisher Full Text](#)
80. Tropini C, Lee TK, Hsin J, *et al.*: **Principles of bacterial cell-size determination revealed by cell-wall synthesis perturbations.** *Cell Rep.* 2014; 9(4): 1520–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
81. Nguyen LT, Gumbart JC, Beeby M, *et al.*: **Coarse-grained simulations of bacterial cell wall growth reveal that local coordination alone can be sufficient to maintain rod shape.** *Proc Natl Acad Sci U S A.* 2015; 112(28): E3689–98.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
82. Egan AJ, Vollmer W: **The physiology of bacterial cell division.** *Ann NY Acad Sci.* 2013; 1277: 8–28.
[PubMed Abstract](#) | [Publisher Full Text](#)
83. de Boer PA: **Advances in understanding *E. coli* cell fission.** *Curr Opin Microbiol.* 2010; 13(6): 730–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
84. den Blaauwen T, Hamoen LW, Levin PA: **The divisome at 25: the road ahead.** *Curr Opin Microbiol.* 2017; 36: 85–94.
[PubMed Abstract](#) | [Publisher Full Text](#)
85. Adams DW, Errington J: **Bacterial cell division: assembly, maintenance and disassembly of the Z ring.** *Nat Rev Microbiol.* 2009; 7(9): 642–53.
[PubMed Abstract](#) | [Publisher Full Text](#)
86. **F** Yang X, Lyu Z, Miguel A, *et al.*: **GTPase activity-coupled treadmilling of the bacterial tubulin FtsZ organizes septal cell wall synthesis.** *Science.* 2017; 355(6326): 744–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
87. **F** Bisson-Filho AW, Hsu Y, Squyres GR, *et al.*: **Treadmilling by FtsZ filaments drives peptidoglycan synthesis and bacterial cell division.** *Science.* 2017; 355(6326): 739–43.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
88. **F** Coltharp C, Buss J, Plumer TM, *et al.*: **Defining the rate-limiting processes of bacterial cytokinesis.** *Proc Natl Acad Sci U S A.* 2016; 113(8): E1044–53.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
89. **F** Li Y, Hsin J, Zhao L, *et al.*: **FtsZ protofilaments use a hinge-opening mechanism for constrictive force generation.** *Science.* 2013; 341(6144): 392–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
90. **F** Osawa M, Anderson DE, Erickson HP: **Reconstitution of contractile FtsZ rings in liposomes.** *Science.* 2008; 320(5877): 792–4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
91. Schaechter M, Maaloe O, Kjeldgaard NO: **Dependency on medium and temperature of cell size and chemical composition during balanced growth of *Salmonella typhimurium*.** *J Gen Microbiol.* 1958; 19(3): 592–606.
[PubMed Abstract](#) | [Publisher Full Text](#)
92. Taheri-Araghi S, Bradde S, Sauls JT, *et al.*: **Cell-size control and homeostasis in bacteria.** *Curr Biol.* 2015; 25: 385–91.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
93. Basan M, Zhu M, Dai X, *et al.*: **Inflating bacterial cells by increased protein synthesis.** *Mol Syst Biol.* 2015; 11(10): 836.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
94. Zheng H, Ho P, Jiang M, *et al.*: **Interrogating the *Escherichia coli* cell cycle by cell dimension perturbations.** *Proc Natl Acad Sci U S A.* 2016; 113(52): 15000–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
95. Wallden M, Fange D, Lundius EG, *et al.*: **The Synchronization of Replication and Division Cycles in Individual *E. coli* Cells.** *Cell.* 2016; 166(3): 729–39.
[PubMed Abstract](#) | [Publisher Full Text](#)
96. Helmstetter CE: **DNA synthesis during the division cycle of rapidly growing *Escherichia coli* B/r.** *J Mol Biol.* 1968; 31(3): 507–18.
[PubMed Abstract](#) | [Publisher Full Text](#)
97. Sperber AM, Herman JK: **Metabolism Shapes the Cell.** *J Bacteriol.* 2017; 199(11): pii: e00039–17.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
98. **F** Yao Z, Davis RM, Kishony R, *et al.*: **Regulation of cell size in response to nutrient availability by fatty acid biosynthesis in *Escherichia coli*.** *Proc Natl Acad Sci U S A.* 2012; 109(38): E2561–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
99. Dörr T, Alvarez L, Delgado F, *et al.*: **A cell wall damage response mediated by a sensor kinase/response regulator pair enables beta-lactam tolerance.** *Proc Natl Acad Sci U S A.* 2016; 113(2): 404–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
100. Zaritsky A, Woldring CL, Mirelman D: **Constant peptidoglycan density in the**

- sacculus of *Escherichia coli* B/r growing at different rates. *FEBS Lett.* 1979; **98**(1): 29–32.
[PubMed Abstract](#) | [Publisher Full Text](#)
101. Harris LK, Dye NA, Theriot JA: **A *Caulobacter* MreB mutant with irregular cell shape exhibits compensatory widening to maintain a preferred surface area to volume ratio.** *Mol Microbiol.* 2014; **94**(5): 988–1005.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 102. Dajkovic A, Tesson B, Chauhan S, *et al.*: **Hydrolysis of peptidoglycan is modulated by amidation of meso-diaminopimelic acid and Mg²⁺ in *Bacillus subtilis*.** *Mol Microbiol.* 2017; **104**(6): 972–88.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 103. Singh SK, Parveen S, SaiSree L, *et al.*: **Regulated proteolysis of a cross-link-specific peptidoglycan hydrolase contributes to bacterial morphogenesis.** *Proc Natl Acad Sci U S A.* 2015; **112**(35): 10956–61.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 104. Ehler K, Höltje JV: **Role of precursor translocation in coordination of murein and phospholipid synthesis in *Escherichia coli*.** *J Bacteriol.* 1996; **178**(23): 6766–71.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 105. You C, Okano H, Hui S, *et al.*: **Coordination of bacterial proteome with metabolism by cyclic AMP signalling.** *Nature.* 2013; **500**(7462): 301–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 106. Hermesen R, Okano H, You C, *et al.*: **A growth-rate composition formula for the growth of *E. coli* on co-utilized carbon substrates.** *Mol Syst Biol.* 2015; **11**(4): 801.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 107. Donachie WD: **Relationship between cell size and time of initiation of DNA replication.** *Nature.* 1968; **219**(5158): 1077–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
 108. Helmstetter C, Cooper S, Pierucci O, *et al.*: **On the bacterial life sequence.** *Cold Spring Harb Symp Quant Biol.* 1968; **33**: 809–22.
[PubMed Abstract](#) | [Publisher Full Text](#)
 109. Si F, Li D, Cox SE, *et al.*: **Invariance of Initiation Mass and Predictability of Cell Size in *Escherichia coli*.** *Curr Biol.* 2017; **27**(9): 1278–87.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 110. Weart RB, Lee AH, Chien AC, *et al.*: **A metabolic sensor governing cell size in bacteria.** *Cell.* 2007; **130**(2): 335–47.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 111. Chien AC, Hill NS, Levin PA: **Cell size control in bacteria.** *Curr Biol.* 2012; **22**(9): R340–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 112. Scott M, Gunderson CW, Mateescu EM, *et al.*: **Interdependence of cell growth and gene expression: origins and consequences.** *Science.* 2010; **330**(6007): 1099–102.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 113. Erickson DW, Schink SJ, Patsalo V, *et al.*: **A global resource allocation strategy governs growth transition kinetics of *Escherichia coli*.** *Nature.* 2017; **551**(7678): 119–23.
[PubMed Abstract](#) | [Publisher Full Text](#)
 114. Walker N, Nghe P, Tans SJ: **Generation and filtering of gene expression noise by the bacterial cell cycle.** *BMC Biol.* 2016; **14**: 11.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 115. Osella M, Tans SJ, Cosentino Lagomarsino M: **Step by Step, Cell by Cell: Quantification of the Bacterial Cell Cycle.** *Trends Microbiol.* 2017; **25**(4): 250–6.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. **Kevin D Young**

Department of Microbiology and Immunology, University of Arkansas for Medical Sciences, Little Rock, AR, 72205-7199, USA

Competing Interests: No competing interests were disclosed.

2. **Rut Carballido-Lopez**

MICALIS, INRA, Agro ParisTech, Université Paris-Saclay, Jouy-en-Josas, France

Competing Interests: No competing interests were disclosed.

3. **Tobias Doerr**

Department of Microbiology, Cornell University, Ithaca, NY, USA

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research