BRIEF REPORT

Comparing organic versus conventional soil management on soil respiration [version 1; peer review: 2 approved]

Bence Mátyás ^{1,2}, Maritza Elizabeth Chiluisa Andrade ³,
Nora Carmen Yandun Chida³, Carina Maribel Taipe Velasco³,
Denisse Estefania Gavilanes Morales³, Gisella Nicole Miño Montero⁴,
Lenin Javier Ramirez Cando ², Ronnie Xavier Lizano Acevedo ²

¹Grupo de Investigación Mentoría y Gestión del Cambio, Universidad Politécnica Salesiana, Cuenca, Ecuador

²Grupo de Investigación en Ciencias Ambientales, Universidad Politécnica Salesiana, Quito, Ecuador

³Ingeniería Ambiental, Universidad Politécnica Salesiana, Quito, Ecuador

⁴Administración de Empresas, Universidad Politécnica Salesiana, Guayaquí, Ecuador

V1 First published: 02 Mar 2018, 7:258
<https://doi.org/10.12688/f1000research.13852.1>
Latest published: 02 Mar 2018, 7:258
<https://doi.org/10.12688/f1000research.13852.1>

Abstract

Soil management has great potential to affect soil respiration. In this study, we investigated the effects of organic versus conventional soil management on soil respiration. We measured the main soil physical-chemical properties from conventional and organic managed soil in Ecuador. Soil respiration was determined using alkaline absorption according to Witkamp. Soil properties such as organic matter, nitrogen, and humidity, were comparable between conventional and organic soils in the present study, and in a further analysis there was no statically significant correlation with soil respiration. Therefore, even though organic farmers tend to apply more organic material to their fields, but this did not result in a significantly higher CO₂ production in their soils in the present study.

Keywords

soil respiration, conventional soil management, organic soil management

Open Peer Review

Approval Status

	1	2
version 1		
02 Mar 2018	view	view

1. **Anita Jakab** , National Agricultural Research and Innovation Centre, Újfehértó, Hungary
2. **Ankit Singla**, Sam Higginbottom Institute of Agriculture, Technology and Sciences, Allahabad, India

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Bence Mátyás (bmatyas@ups.edu.ec)

Author roles: **Mátyás B:** Conceptualization, Investigation, Methodology, Supervision, Validation, Writing – Original Draft Preparation, Writing – Review & Editing; **Chiluisa Andrade ME:** Investigation, Methodology, Writing – Original Draft Preparation; **Yandun Chida NC:** Investigation, Methodology, Writing – Original Draft Preparation; **Taípe Velasco CM:** Investigation, Methodology, Writing – Original Draft Preparation; **Gavilanes Morales DE:** Investigation, Methodology, Writing – Original Draft Preparation; **Miño Montero GN:** Project Administration, Supervision, Writing – Original Draft Preparation; **Ramírez Cando LJ:** Data Curation, Formal Analysis, Validation; **Lizano Acevedo RX:** Conceptualization, Project Administration

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2018 Mátyás B *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. Data associated with the article are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

How to cite this article: Mátyás B, Chiluisa Andrade ME, Yandun Chida NC *et al.* **Comparing organic versus conventional soil management on soil respiration [version 1; peer review: 2 approved]** F1000Research 2018, 7:258
<https://doi.org/10.12688/f1000research.13852.1>

First published: 02 Mar 2018, 7:258 <https://doi.org/10.12688/f1000research.13852.1>

Introduction

Research related to the benefits of organic management¹ has become increasingly important in sustainable agriculture. Organic soil management can contribute to meaningful socio-economic and ecologically sustainable development. Kilcher states that “Organic agriculture reduces the risk of yield failure, stabilizes returns and improves the quality of life of small farmers’ families”². Soil management has great potential to affect soil respiration, which is an important qualitative indicator of soil microbial activity³. Soil respiration is released as a result of soil organic matter decomposition. The present study aims to investigate the effects of organic versus conventional management on CO_2 production of some Northern Ecuadorian agricultural soils. Our hypothesis was that major soil respiration will be observed in soils under organic management due to the increased amount of applied organic materials.

Methods

Sampling sites

Soil samples from 23 organic farms and conventionally managed neighbouring farms were analyzed. In total, 17 sampling sites were located in organic farms, while 6 sampling sites were located in chemical fertilizer-treated areas. The sampling sites were chosen according to proximity of organic and conventionally managed farms in which the same crops are produced. Further details about each of the sampling sites can be found in Table 1. Approximately 1000 g of soil samples of 0–20 cm depth were taken. The following crops were produced in the examined areas: broccoli, potato, tomato and carrot.

Soil properties

Soil moisture content was determined gravimetrically, drying the soil at 105°C for 24 hours according to Fernández *et al.* (2008)⁴. Soil texture was measured using sodium hexametaphosphate ($(NaPO_3)_6$) according to Bouyoucos (1962)⁵. To measure the soil chemical properties, the samples were sieved through a 2mm mesh and pre-incubated at 25° for 72 hours. Soil pH in distilled water (soil/water, 1/2.5, w/w) was determined according to Karkanis (1991)⁶. In addition, we measured the electrical conductivity (EC) using a glass electrode according to Karkanis (1991)⁶. Cylinder volume was determined according to Agostini *et al.* (2014)⁷. Soil organic matter was determined according to Walkley and Black (1934)⁸. We measured the phosphorous content according to Olsen (1954)⁹. The Sand/Silt/Clay ratio was determined by Bouyoucos’s method (1936)¹⁰, while the cation exchange capacity was determined according to ISO 11260 (1994)¹¹ protocol.

Soil respiration

The experiment was applied at 25°C. 0, 1M NaOH (10ml) was placed in laboratory bottles (250ml), a sterile gauze pad were filled with 10 g of soil sample according to Witkamp (1966)¹². After 10 days, the amount of CO_2 was subsequently measured by standardized titration against 0.1N HCl using firstly phenolphthalein and then methyl orange indicator according to Witkamp (1966)¹².

The below formula was applied to calculate soil respiration:

$$m(CO_2) = V \times N \times 22 \text{ } CO_2$$

And CO_2 production (for 10 days):

$$mg(CO_2) * 100g - 1 * 10 \text{ day} - 1 = \\ \text{methyl orange factor} * HCl - \text{phenolphthalein loss} * \\ NAOH \text{ factor} * 2, 2 * \text{Moisture multiplication factor}$$

where

$$\text{Moisture multiplication factor} = \frac{(\text{moisture content} \% + 100)}{100}$$

We determined the volume of the examined soils (counting with 0 – 20 cm depth) using topsoil calculator tool (<https://www.tillersturf.co.uk/topsoil-calculator>). The results of soil respiration was then estimated in $kg(CO_2)/ha/day$.

Statistical analysis

To evaluate the behavior within results, two types of test were performed: i) Student’s t-test for comparing means between conventional and organic crop systems in terms of soil respiration ($kg/CO_2/ha/day$), organic matter (%) and nitrogen (%). Furthermore, Person’s and Spearman’s correlation were fixed in order to test data covariation and correlation. ii) ANOVA was used to compare conventional and organic crop system and the type of crop harvested in the sampling site.

Results

The results of soil respiration from areas of organic and conventional soil management are comparable (Dataset 1).

For soil respiration, conventional soil mean was 88.50 and organic mean was 98.64, showing an increment around 10%. However, there were no statistically significant differences between group means as determined by one-way ANOVA ($p=0.15$), comparing conventional and organic systems. Pearson’s and Kendall’s tests have showed no correlation. Soil respiration correlation coefficient with organic matter was lower than 0.05 and with nitrogen content was lower than 0.12. This analysis did not consider the differences between conventional and organic systems (Figure 1).

There were statistically significant differences between group means as determined by one-way ANOVA ($p < 0.05$), comparing crop types. Furthermore, a post hoc test (Duncan) was fixed. There was only one crop (carrot) in conventional system (odds lower than 0.05) that differs drastically from the others, as pointed out in (Figure 2).

Considering soil characteristics (pH, CIC, K, and Electric conductivity), Student’s t-test was applied to identify differences between conventional and organic systems. Only the characteristics

Table 1. Characteristics of the conventional and organic farms chosen for the present study. Variables are follows: areas of examined lands (m^2), Name of crops, soil management (Organic/Conventional), Total crop production (kg), Applied fertilizer (kg), Type of fertilizers, Concentration of NPK, Amount of NPK (kg), GPS coordinates of the examined lands.

Farmer's code	Crop	Solid fertilizers	Area of land m2	Total crop production (Kg)	Fertilizer application rate on total crop production (Kg)	Concentration of NPK (%) in each fertilizer solid			Amount of NPK in kg			liquid fertilizer	Fertilizer application rate on total crop production (Kg)	Concentration of NPK (%) in each liquid fertilizer						Amount of NPK in Kg			GPS coordinates				
						N	K	P	N	K	P			N	K	P	N	K	P	N	K	P	latitude	length			
OB1	Agroecological	Compost	60.38	315	95.25	0.53	0.6345	1.322	0.504825	0.60436125	1.259205	Biol	2.63975	0.2428	0.8183	0.3061	0.00640931	0.02160107	0.00606275	0804800	0003519						
OB2	Agroecological	Bocashi	119.2	576	288.03	0.17	0.4013	0.071	0.455651	1.07560439	0.1903013	Biol	185	0.2	1.0816	0.0148	0.37	2.00096	0.02738	0809419	0006402						
OB3	Agroecological	Compost	9	79.2	20.2	0.3	0.3891	0.1221	0.0606	0.0785982	0.0246642	biol	30	0.14	0.0075	0.467	0.042	0.00225	0.1401	0809136	0003476						
OB4	Agroecological	Bocashi	144	600	300	0.5	0.8667	0.1271	1.5	2.6001	0.3813	Biol	211.18	0.24	0.4033	0.0958	0.506832	0.85168894	0.20231044	0804806	0003527						
CB1	Agroecological	Bocashi	56	326.7	101	0.43	0.4427	0.5081	0.4343	0.447127	0.513181	Biol	150	0.2202	0.2862	0.0735	0.3303	0.4293	0.11025	0811423	0003176						
		18460			82.21654	18	0	46	14.7989772	0	37.8196084																
		OO60			33.33904	0	60	0	0	20.003424	0																
	Conventional	Triple 15	2511.6	8316	44.4521	15	15	15	6.667815	6.667815	6.667815										0805608	0001169					
		UREA			2.1	46	0	0	0.966	0	0	0															
OT1	Agroecological	Gallinaza	322.766	0.0322766	1550	0.59	0.6815	0.8673	9.145	10.5625	13.44315	Biol	78.51387307	1.09	1.5659	0.5374	0.85580122	1.22944874	0.421933554	0811193	0006955						
		Compost			1550	0.89	2.5875	0.6949	13.795	40.106	1.077																
OT2	Humus 1	202.4	0.02024		2470	1.24	2.9429	1.0828	30.628	72.68963	26.74516	Biol	2086.4584	0.26	0.3443	0.2216	5.42479184	7.18367627	4.623591814								
	Humus 2				2470	0.66	0.7458	0.5232	16.302	18.42126	12.92304																
	Bocashi				2470	0.8	1.2478	0.6486	19.76	30.821	16.02																
	Bocashi negro				2470	1.29	1.0581	0.2705	31.863	26.135	6.681										0809214	0003617					
OT3	Agroecological	Compost	250.912	0.0250912	6.38	0.39	0.8731	0.2064	0.024882	0.05570378	0.001316832	Biol	11964.3692	0.220045	0.073448586	0.2859998	26.3268962	8.78766	34.218072								
CT1	Bocashi				6.38	0.43	0.5081	0.4427	0.027434	0.033	0.028									0811429	0003184						
	Nitrogen				41.32	10.7	0	0	3.79	0	0																
	Magnesium	847.132	0.0847132		41.32	13	46	0	5.3716	19.0072	0									0809021	0002732						
	Ultrasol K				99.43	8	20	20	7.95	19.89	19.89																
CT2	Conventional	MAP	4827.69	0.482769	99.43	12	0	61	11.93	0	60.65																
CT3	EC FERTILIZER				41.43	15.5	0	0	5.14	0	0									0805316	0001139						
	Fluore	1234.865	0.1234865		5.668635943	1	9.5	5	0.056686358	0.538520405	0.283431792																
	Nitrofoska biolar				40.49025602	8	24	12	3.24	9.71766145	4.8588307									0805312	0001138						
OP1	Agroecological	Compost	116	408.16	1360	0.53	0.6345	1.322	7.208	8.629	17.979	Biol	1.0559	0.2428	0.8183	0.3061	0.00256373	0.00864043	0.00323211	804851	3376						
OP2	Agroecological	Gallinaza	88.2	136	50	3.14	4.3752	6.0922	1.57	2.1876	3.0461																
		Cal Agricola			9.0719	x	x	x	x	x	x	x Biol	84.472	0.17	0.4013	0.071	0.1436024	0.33898614	0.0599712	809414	6461						
OP3	Agroecological	Bocashi	69.9	181.4	1500	0.38	0.7695	0.4772	5.7	11.5425	7.158	Biol 1	12.6706	0.22	0.3619	0.013	0.02787576	0.04585563	0.001647204								
OP4	Agroecological	Compost	13.17	45.35	4.5359	0.91	0.4283	0.865	0.04127669	0.01942726	0.03923554	Biol 2 microorga.	6.3354	0.13	0.2065	0.0065	0.00823602	0.0130826	0.000411801	808161	3438						
		Harvest waste			3000				x	x	x	x Biol	1.58385	0.18	0.224	0.0387	0.00285093	0.00354782	0.00061295	808225	3496						
		103010			750	10	10	30	75	75	225																
	Conventional	18460			250	18	46	0	45	115	0																
OP1		Stimufolk			4	11	38	5	2.2	7.6	1																
	Agriate		10 000	13610	4	19	19	19	9.5	9.5	9.5									810311	5670						

Farmer's code	Crop	Solid fertilizers	Area of land m2	Total crop production (Kg)	Fertilizer application rate on total crop production (Kg)	Concentration of NPK (%) in each fertilizer solid			Amount of NPK in kg			liquid fertilizer	Fertilizer application rate on total crop production (Kg)	Concentration of NPK (%) in each liquid fertilizer			Amount of NPK in Kg			GPS coordinates		
						N	K	P	N	K	P			N	K	P	N	K	P	latitude	length	
OC1	Agroecological	Compost	92.97	2045	146.66	0.53	0.6345	1.322	0.777298	0.9305577	1.9388452	Biol	4.06	0.2428	0.8183	0.3061	0.0085768	0.03222298	0.01242766	17°N 08°04'05"	0003544	
OC2	Agroecological		15.645	156		0	0	0	0	0	0	Biol	2	0.23	0.007	0.0181	0.0046	0.00014	0.000362	17°N 08'11.49"	0003795	
OC3	Agroecological	Bocashi	9	72	1.35	0.38	0.7695	0.4772	0.00513	0.01038825	0.0064422	Biol (1)	1.64	0.22	0.3619	0.013	0.003608	0.00593516	0.0002132	17°N 08°08'28"	0003066	
												Biol (2)	0.82	0.13	0.2065	0.0065	0.001066	0.0016933	0.0005533			
OC4	Agroecological	Bocashi	11.2	246	23.3	0.5	0.8667	0.1271	0.1165	0.2019411	0.0296143	Biol	16.425	0.24	0.4033	0.0958	0.03942	0.06624203	0.01573515	17°N 08°04'08"	0003504	
OC5	Agroecological	Compost	60	1500	134.6	0.3	0.3891	0.1221	0.4038	0.5237286	0.1643466	biol	200	0.14	0.0075	0.467	0.28	0.015	0.934	17°N 08°09'136"	0003548	
OC1	Carrot		176.56	108								Biofertilizante (lombriz)	60	0.32	0.3963	0.4595	0.192	0.00126816	0.001620999			
		Nitrato de Calcio		0.13	15	0	0	0.0195	0	0	0	0										
		Fosfato Monocamónico		0.07	11	0	52.5	0.00733333	0	0.035												
		Nitrato de Potasio		0.07	13	44	0.0066667	0	0.02933333

Figure 1. Soil respiration compared with organic matter and nitrogen in soil.

Figure 2. Boxplots showing alterations within crop systems and crop harvested in the zone.

from carrot crop systems (conventional or organic) have shown differences in terms of means ($p < 0.05$). Furthermore, the mean of conventional crop system was lower in every characteristic evaluated. Besides, these results were in congruence with Figure 2, leading us to believe that the cropping system has no influence on soil respiration, which is in contrast to the influence that soil characteristics have over soil respiration in this study.

Dataset 1. Raw data for various parameters calculated in conventional and organic managed soils

<http://dx.doi.org/10.5256/f1000research.13852.d195529>

Parameters as follows: pH, Organic material (percentage), Total Nitrogen (percentage), Match (mg/kg), Potassium (cmol/kg), Electrical conductivity (dS/m), CIC (cmol/kg), Soil moisture content (percentage), Sand (percentage), Silt-limo (percentage), Clay (percentage), Texture (class), Soil respiration (kg/CO₂/ha/day).

Conclusions

Organic farmers tend to apply more organic material to their fields, but this did not result in a significantly higher CO₂ production in their soils. The difference between organic and conventional soils (10% in mean) is not enough to conclude that the soil respiration under these two systems was different, considering the analysis of their variance.

Soil properties like organic matter, nitrogen, and humidity, were comparable between conventional and organic soils in the present study, and in a further analysis there was no statically

significant correlation with soil respiration. However, biological significance should be investigated in a posteriori research including microbial community profile of the soil and specific interactions in highlands (over 2500 m.a.s.l.).

Ethics

Oral consent was obtained from the farmers for the collection of soil samples from their land. Their only request was to inform them about the results of the soil characteristics, that we have already done personally on 9 November, 2017.

Data availability

Dataset 1: Raw data for various parameters calculated in conventional and organic managed soils. Parameters as follows: pH, Organic material (percentage), Total Nitrogen (percentage), Match (mg/kg), Potassium (cmol/kg), Electrical conductivity (dS/m), CIC (cmol/kg), Soil moisture content (percentage), Sand (percentage), Silt-limo (percentage), Clay (percentage), Texture (class), Soil respiration (kg/CO₂/ha/day). DOI, [10.5256/f1000research.13852.d195529](http://dx.doi.org/10.5256/f1000research.13852.d195529)¹³

Competing interests

No competing interests were disclosed.

Grant information

The author(s) declared that no grants were involved in supporting this work.

References

1. van Diepeningena AD, de Vosa OJ, Korthalsb GW, *et al.*: **Effects of organic versus conventional management on chemical and biological parameters in agricultural soils.** *Appl Soil Ecol.* 2006; **31**(1–2): 120–135.
[Publisher Full Text](#)
2. Kilcher L: **How organic agriculture contributes to sustainable development.** Kassel University Press, University of Kassel at Witzenhausen JARTS, 2007; (Supplement 89): 31–49.
[Reference Source](#)
3. Baldock JA, Wheeler I, McKenzie N, *et al.*: **Soils and climate change: Potential impacts on carbon stocks and greenhouse gas emissions, and future research for Australian agriculture.** *Crop Pasture Sci.* 2012; **63**(3): 269–283.
[Publisher Full Text](#)
4. Fernández L, Roldán T, Zegarra H, *et al.*: **Manual de técnicas de análisis de suelos aplicadas a la remediación de sitios contaminados.** *Can Agr Eng, Mexico.* 2008.
[Reference Source](#)
5. Bouyoucos GJ: **Hydrometer method improved for making particle size analyses of soils.** *Agron J.* 1962; **54**(5): 464–465.
[Publisher Full Text](#)
6. Karkanis PG, Au K, Schaalje GB: **Comparison of 4 Measurement Schedules for Determination of Soil Particle-Size Distribution by the Hydrometer Method.** *Can Agr Eng.* 1991; **33**(2): 211–215.
[Reference Source](#)
7. de los Ángeles Agostini M, Monterubbianesi MG, Studdert GA, *et al.*: **Un método simple y práctico para la determinación de densidad aparente.** *Ciencia Del Suelo.* 2014; **32**(2): 171–176.
[Reference Source](#)
8. Walkley A, Black IA: **An examination of the Degtjareff method for determining organic carbon in soils: Effect of variations in digestion conditions and of inorganic soil constituents.** *Soil Sci.* 1934; **63**: 251–263.
9. Olsen SR, Cole CV, Watanabe FS, *et al.*: **Estimation of available phosphorus in soils by extraction with sodium bicarbonate.** USDA, Washington, DC., 1954.
[Reference Source](#)
10. Bouyoucos GJ: **Directions for making mechanical analysis of soils by the hydrometer method.** *Soil Science.* 1936; **42**(3): 225–230.
[Publisher Full Text](#)
11. ISO 11260: **Soil quality -Determination of effective cation exchange capacity and base saturation level using barium chloride solution.** 1994.
12. Witkamp M: **Decomposition of Leaf Litter in Relation to Environment, Microflora, and Microbial Respiration.** *Ecology.* 1966; **47**: 194–201.
[Publisher Full Text](#)
13. Mátyás B, Chiluisa Andrade ME, Yandun Chida NC, *et al.*: **Dataset 1 in: Comparing organic versus conventional soil management on soil respiration.** *F1000Research.* 2018.
[Data Source](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 19 March 2018

<https://doi.org/10.5256/f1000research.15056.r31654>

© 2018 Singla A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Ankit Singla

Department of Industrial Microbiology, Sam Higginbottom Institute of Agriculture, Technology and Sciences, Allahabad, Uttar Pradesh, India

The submitted manuscript by Bence et al. is good work which is suitable for publication in F1000 research. Authors have compared the organic practices and conventional practices, and compared their effects on soil respiration which is very important aspect. Standard methodologies were followed which ensures reproducibility of the results. The findings were subjected to the statistical analysis and conclusion drawn nicely.

However, I have below suggestions for improvement which may be considered as minor revisions:

- "Physical-chemical" could be replaced by "Physico-chemical" throughout the manuscript.
- In abstract, word "statically" should be replaced by "statistically"
- In result, "showing and increment around 10%." should be "showing an increment around 10%."
- The discussion could be added more so that the findings of the study will become stronger.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 15 March 2018

<https://doi.org/10.5256/f1000research.15056.r31652>

© 2018 Jakab A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Anita Jakab

National Agricultural Research and Innovation Centre, National Agricultural Research and Innovation Centre, National Agricultural Research and Innovation Centre, Újfehértó, Hungary

This article worked at the differences between organic and conventional soil management. This research examined an important and topical issue especially the soil respiration under changing plant and soil conditions.

Introduction and methods

The research investigated 23 soil samples in Ecuador. The samples were located from organic (17 samples) and conventionally managed neighboring farms (6 samples). In the research trials broccoli, potato, tomato and carrot were applied as test plant. Soil properties were measured after 1000 g soil samples of 0-20 cm depths of soil were taken in every picked area. The soil moisture, texture, pH, electrical conductivity, cylinder volume, organic matter, phosphorus content, sand/silt/clay ratio and cation exchange capacity, and the soil respiration were analyzed in laboratory.

The values of the soil parameters are presented in a dataset, which inform about the important soil parameters especially the calculated soil respiration in kg (CO₂)/ha/day). The protocols (description of the tests) are clear and traceable, especially the formula to calculate soil respiration.

The study describes the applied type of fertilizers especially the concentration of NPK fertilizers.

Comment on the Methods

- The sampling time and vegetation status are important for the evaluation, this information is missing in the study. If it's possible, describe the followings: When the soil sampling happened? What was the state of the vegetation of test plants?
- A bit more detail of the soil properties inform us about the actual soil status. The studied soils are classified as sandy textured soil, according to the soil classification (Franco Arenoso). The most typical parameters of the samples are the following: high sandy texture, neutral pH, good/very good organic matter-nitrogen and phosphorus content, 10-20% moisture content. I suggest describing it in the Methods.

Results

The results of the study are described with sufficient statistical analysis. It also describes the statistically significant/not significant results. There were solely statistically significant differences between crop types (for soil respiration by one-way ANOVA correlation test).

- The Figure 1 contains a typographical error (Orgacin matter instead of Organic matter).
- It may be more informative, if you use a line diagram instead of dot diagrams in the first figure.
- The Figure 2 include the soil respiration values in kg CO₂/ha /day, which would be more clear with the average values.

Conclusion

The results have briefly evaluated and conclusions straightforward formulated. I quite agree with observations of the study that emphasizes the importance of further microbiological studies.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Agricultural environmental management, soil management, agricultural soil science

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research