CASE REPORT

Case Report: Kikuchi: The great mimicker [version 1; peer review: 2 approved]

Kevin Bryan Lo , Anna Papazoglou, Lorayne Chua, Nellowe Candelario

Department of Medicine, Einstein Medical Center, Philadelphia, PA, 19141, USA

V1 First published: 30 Apr 2018, 7:520
<https://doi.org/10.12688/f1000research.14758.1>
 Latest published: 30 Apr 2018, 7:520
<https://doi.org/10.12688/f1000research.14758.1>

Abstract

Kikuchi-Fujimoto disease is a form of a benign necrotizing lymphadenitis which is most commonly misdiagnosed as tuberculosis and or lymphoma, usually more common among young adults in Asia. It is a benign disease but can mimic a lot of other disease processes spanning infectious, rheumatologic and even hematologic malignancies. Our patient presented with prolonged fever and lymphadenopathy. Initial considerations were lymphoma and a nonspecific viral infection. A CT scan showed diffuse cervical lymphadenopathy with lacrimal gland involvement. An excisional lymph node biopsy was done which revealed Kikuchi disease. Patient was given steroids with immediate response with defervescence. Kikuchi is a disease with many mimics and a complete workup is needed to exclude serious disease like malignancy.

Keywords

lymphadenitis, fever, Kikuchi, autoimmune

Open Peer Review

Approval Status

	1	2
version 1		
30 Apr 2018	view	view
1. Abdou-Rajack Ndiaye , Hôpital Principal de Dakar, Dakar, Senegal		
Cheikh Lame , Hôpital Principal de Dakar, Dakar, Senegal		
Hôpital d'Instruction des Armées Sénégalaise, Dakar, Senegal		
2. Arvind P. Ganpule , Muljibhai Patel Urological Hospital (MPUH), Nadiad, India		
Tak Gopal Ramdas , Muljibhai Patel Urological Hospital (MPUH), Nadiad, India		
Any reports and responses or comments on the article can be found at the end of the article.		

Corresponding author: Kevin Bryan Lo (lokevinb@einstein.edu)

Author roles: **Lo KB:** Conceptualization, Data Curation, Writing – Original Draft Preparation, Writing – Review & Editing; **Papazoglou A:** Conceptualization, Data Curation, Formal Analysis, Supervision, Writing – Review & Editing; **Chua L:** Data Curation, Writing – Review & Editing; **Candelario N:** Data Curation, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2018 Lo KB *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Lo KB, Papazoglou A, Chua L and Candelario N. **Case Report: Kikuchi: The great mimicker [version 1; peer review: 2 approved]** F1000Research 2018, 7:520 <https://doi.org/10.12688/f1000research.14758.1>

First published: 30 Apr 2018, 7:520 <https://doi.org/10.12688/f1000research.14758.1>

Introduction

Kikuchi-Fujimoto disease (sometimes also known as Kikuchi Disease) is usually more common among young adults in Asia. It is a benign disease, but can mimic a lot of other disease processes spanning infectious, rheumatologic and even hematologic malignancies¹. It usually presents with fever and cervical lymphadenopathy but occasionally it can manifest together with other unusual symptoms further increasing the chances of misdiagnosis.

Case

A 20-year-old African American woman with no other known prior medical history, presented to our institution January 2018 with fevers of 3 weeks' duration. The fevers were predominantly in the late afternoon hours, associated with night sweats, frontal headache, tender cervical lymphadenopathy, anorexia and malaise.

Two weeks prior she saw her primary care physician who diagnosed her with viral illness and recommended supportive care. She also visited the emergency department and was diagnosed with lymphadenitis; a course of amoxicillin/clavulanic acid was prescribed of unrecalled dose and she wasn't able to finish the whole course. Symptoms however persisted, and the patient also developed bilateral periorbital swelling and non-bloody diarrhea prompting her presentation at our institution. The patient indicated they had no cough, chest pain, dysuria, abdominal pain, arthralgia, rash, recent travel or sick contacts.

The patient was not in distress, with blood pressure of 120/70 mm Hg, febrile to 39.6 C and tachycardic with heart rate of 110 bpm. Physical exam was notable for bilateral periorbital swelling with violet discoloration of the eyelids, conjunctival pallor and painless bilateral cervical lymphadenopathy. No rash or joint swelling was noted.

Complete blood count revealed leukopenia with a white cell count of 2.9×10^3 /mL (65% neutrophils, 13% lymphocytes, 13% bands), microcytic anemia with a hemoglobin of 8.5 gr/dL (mean corpuscular volume 65 fL) and 181×10^3 /mL platelets. C-reactive protein (CPR) and erythrocyte sedimentation rate (ESR) level were markedly elevated at 51 and 84 respectively. Lactate dehydrogenase (LDH), ferritin and haptoglobin were also elevated. The patient tested negative for β -human chorionic gonadotropin (hCG), HIV, hepatitis B and C, angiotensin converting enzyme (ACE), antinuclear antibodies (ANA) and rheumatoid factor (RF).

Computed Tomography of the neck revealed bilateral cervical lymphadenopathy, enhancement and mild enlargement of the parotid and lacrimal glands and diffuse swelling of the pharyngeal mucosa and marked enhancement of bilateral cervical soft tissue planes [Figure 1 and Figure 2].

She was observed off antibiotics. Blood cultures, serology for Epstein Barr virus (EBV) and cytomegalovirus (CMV), bone marrow biopsy and flow cytometry were all negative. Excisional

Figure 1. Computed tomography (CT) scan coronal view head with contrast. Bilateral lacrimal glands appear large with mild increased enhancement.

Figure 2. Computed tomography (CT) scan sagittal view neck with contrast. Bilateral left greater than right cervical jugular chain, level I, occipital and supraclavicular lymph nodes demonstrate heterogeneous enhancement and enlargement, largest demonstrating conglomeration and cystic changes along the left jugular chain measuring up to 8.5 cm in length of the conglomerate.

biopsy of the left cervical lymph node revealed characteristic findings of Kikuchi-Fujimoto disease which showed geographic necrosis with fibrinoid deposits and apoptotic cells surrounded by a mononuclear infiltrate characteristically without neutrophils and eosinophils [Figure 3]. The patient was started on prednisone 40mg per day with rapid resolution of symptoms. Steroids were tapered after one week of treatment. Upon follow up in Rheumatology clinic 4 months later, patient was noted to be completely symptom free.

Figure 3. Histopathology of cervical lymph node. Geographical necrosis with fibrinoid deposits and nuclear fragments with apoptotic cells. Surrounding this area are pale histiocytes and lymphocytes. Neutrophils and eosinophils are characteristically absent.

Discussion

Kikuchi disease was first independently described through case series in 1972 by Kikuchi and Fujimoto as a form of a benign necrotizing lymphadenitis which was most commonly misdiagnosed as tuberculosis and or lymphoma². The main etiology for Kikuchi disease is still unknown but there are various studies that implicate viruses such as EBV as a potential trigger^{2,3}. It is also closely related to systemic lupus erythematosus (SLE) and in fact, there are studies and case reports showing a strong association between the two disease processes with the diagnosis of SLE coming before, after or even simultaneously with Kikuchi disease^{4,5}. The most frequent presenting symptom was fever while the most common presenting sign was cervical lymphadenopathy⁵. It also presents together with constitutional symptoms like night sweats and weight loss which can be initially be misdiagnosed as tuberculosis or lymphoma^{1,2}. However, Kikuchi disease has also been implicated to cause a wide range of symptoms ranging from neurological, musculoskeletal, cutaneous and glandular dysfunction⁶. Eye manifestations for Kikuchi usually present as uveitis and conjunctivitis^{6,7}. Our case is unique because bilateral eyelid swelling has only been reported twice in the literature as a possible presentation of Kikuchi disease, this may be attributed to lacrimal gland involvement which was seen in the imaging findings in our patient^{6,8,9}. Definitive diagnosis is established by lymph node biopsy. Classic biopsy findings include necrosis without a

neutrophilic infiltrate with the predominance of histiocytes and T lymphocytes^{1,2}. Kikuchi is a benign self-limiting disease and symptoms usually resolves spontaneously within 4 months in majority of cases with supportive treatment^{1,6}. The use of glucocorticoids have been found to have some benefit but is usually reserved in more severe persistent cases^{5,6,10}. Kikuchi is a disease with a lot of mimics, the amount of workup alone together with the actual disease manifestations can lead to a lot of morbidity and discomfort for the patient. Nevertheless, a complete workup including an excisional biopsy is recommended to help rule out other serious diseases like malignancy. Close follow up is also needed to monitor for the development of closely associated rheumatological diseases like SLE. Strengths in the approach of the case was the exhaustive diagnostic approach used to arrive at the correct diagnosis for the patient. All possible differentials were considered especially the serious ones such as malignancy. Weakness involved were due to the extensive workup done which consisted of numerous blood tests and invasive tests such as a biopsy, this caused a significant degree of anxiety and morbidity to the patient as well.

Conclusions

Kikuchi is a great mimicker and can be confused with tuberculosis, lymphoma and other viral illnesses. A complete workup including an excisional biopsy is recommended to help rule out other serious diseases like malignancy. Close follow up is needed to monitor for the development of closely associated rheumatological diseases like SLE.

Ethics and consent

Written informed consent for publication of their clinical details and clinical images was obtained from the patient and parent.

Data availability

All data underlying the results are available as part of the article and no additional source data are required.

Competing interests

No competing interests were disclosed.

Grant information

The author(s) declared that no grants were involved in supporting this work.

References

- Hutchinson CB, Wang E: **Kikuchi-Fujimoto disease**. *Arch Pathol Lab Med*. 2010; **134**(2): 289–93.
[PubMed Abstract](#)
- Shirakusa T, Eimoto T, Kikuchi M: **Histiocytic necrotizing lymphadenitis**. *Postgrad Med J*. 1988; **64**(748): 107–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Chiu CF, Chow KC, Lin TY, et al.: **Virus infection in patients with histiocytic necrotizing lymphadenitis in Taiwan. Detection of Epstein-Barr virus, type I human T-cell lymphotropic virus, and parvovirus B19**. *Am J Clin Pathol*. 2000; **113**(6): 774–81.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Baenas DF, Diehl FA, Haye Salinas MJ, et al.: **Kikuchi-Fujimoto disease and systemic lupus erythematosus**. *Int Med Case Rep J*. 2016; **9**: 163–167.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

5. Kucukardali Y, Solmazgul E, Kunter E, *et al.*: **Kikuchi-Fujimoto Disease: analysis of 244 cases.** *Clin Rheumatol.* 2007; **26**(1): 50–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
6. Bosch X, Guilabert A, Miquel R, *et al.*: **Enigmatic Kikuchi-Fujimoto disease: a comprehensive review.** *Am J Clin Pathol.* 2004; **122**(1): 141–52.
[PubMed Abstract](#) | [Publisher Full Text](#)
7. Galor A, Georgy M, Leder HA, *et al.*: **Papillary conjunctivitis associated with Kikuchi disease.** *Cornea.* 2008; **27**(8): 944–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
8. Guerriero S, Ruggeri E, Piscitelli D, *et al.*: **Kikuchi-Fujimoto disease: a rare case of orbital involvement.** *Orbit.* 2012; **31**(6): 420–2.
[PubMed Abstract](#) | [Publisher Full Text](#)
9. Chavis PS, Fallata A, Al-Hussein H, *et al.*: **Lacrimal gland involvement in Kikuchi-Fujimoto disease.** *Orbit.* 1998; **17**(2): 113–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
10. Jang YJ, Park KH, Seok HJ: **Management of Kikuchi's disease using glucocorticoid.** *J Laryngol Otol.* 2000; **114**(9): 709–11.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 07 August 2018

<https://doi.org/10.5256/f1000research.16056.r35937>

© 2018 Ganpule A et al. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Arvind P. Ganpule

Department of Urology, Muljibhai Patel Urological Hospital (MPUH), Nadiad, India

Tak Gopal Ramdas

Muljibhai Patel Urological Hospital (MPUH), Nadiad, India

Explanation for second referee question:

Periodic acid-Schiff (PAS) stain , Ziehl Neelsen (ZN) stain, Genexpert examination for tubercle bacilli (TB), immunohistochemistry (IHC) examination for lymphoma, tests for systemic lupus erythematosus (SLE) to rule out different differentials, were not mentioned.

Key features on lymph node biopsy are fragmentation, necrosis and karyorrhexis were not mentioned by authors in the histopathology examination.

Treatment includes symptomatic care, analgesics-antipyretics, corticosteroids and not the corticosteroid alone as mentioned by authors.

Author may put "Ganpule AP, Chabra JS, Singh AG, Tak GR, Soni S, Sabnis R, Desai M. Case Report: Kikuchi-Fujimoto disease: a diagnostic and therapeutic dilemma following pretransplant nephrectomy for a 2.35 Kg kidney. F1000Research. 2016;5" as the reference for their article.

Is the background of the case's history and progression described in sufficient detail?

Yes

Are enough details provided of any physical examination and diagnostic tests, treatment given and outcomes?

Partly

Is sufficient discussion included of the importance of the findings and their relevance to future understanding of disease processes, diagnosis or treatment?

Yes

Is the case presented with sufficient detail to be useful for other practitioners?

Yes

Competing Interests: No competing interests were disclosed.

We confirm that we have read this submission and believe that we have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 25 June 2018

<https://doi.org/10.5256/f1000research.16056.r35130>

© 2018 Ndiaye A et al. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abdou-Rajack Ndiaye

Service de Médecine interne, Hôpital Principal de Dakar, Dakar, Senegal

Cheikh Lame

¹ Service d'Oto-Rhino-Laryngologie, Hôpital Principal de Dakar, Dakar, Senegal

² Hôpital d'Instruction des Armées Sénégalaise, Dakar, Senegal

The article is correctly written. It is an original one. The described pathology is uncommon and raises diagnostic problems. References are up-to-date.

However, some minor revisions must be done:

1. Serodiagnostic tests for toxoplasmosis, rubella, syphilis, and infectious mononucleosis must be performed in presence of chronic lymphadenopathy with fever, even they are often negative in such case.
2. Precise the place of immunohistochemical study (cd68, cd8) to confirm the diagnosis.
3. Deaths are rare but have been observed in systemic forms of Kikuchi disease.
4. Recent treatment with intravenous immunoglobulins has been proposed in severe and resistant forms.
5. Figure 3: Specify the used coloration and the corresponding magnification.

References

1. Lame CA, Loum B, Fall AK, Cucherousset J, et al.: Kikuchi-Fujimoto disease, a rare cause of lymphadenopathy in Africa. Description of the first case in Senegal and review of the literature. *Eur Ann Otorhinolaryngol Head Neck Dis.* 2017; **134** (5): 347-349 [PubMed Abstract](#) | [Publisher Full Text](#)

Is the background of the case's history and progression described in sufficient detail?

Yes

Are enough details provided of any physical examination and diagnostic tests, treatment given and outcomes?

Yes

Is sufficient discussion included of the importance of the findings and their relevance to future understanding of disease processes, diagnosis or treatment?

Yes

Is the case presented with sufficient detail to be useful for other practitioners?

Yes

Competing Interests: No competing interests were disclosed.

We confirm that we have read this submission and believe that we have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research