RESEARCH ARTICLE

REVISED Cigarette smoking and metabolic syndrome components: a cross-sectional study from Maracaibo City, Venezuela [version 2; peer review: 1 approved, 1 approved with reservations]

Valmore Bermudez ^{1,2}, Luis Carlos Olivar², Wheeler Torres², Carla Navarro², Robys Gonzalez², Cristobal Espinoza³, Alicia Morocho³, Andres Mindiola⁴, Maricarmen Chacin², Victor Arias¹, Roberto Añez ², Juan Salazar ², Manuel Riaño-Garzon ¹, Edgar Diaz-Camargo¹, Maria Judith Bautista¹, Joselyn Rojas^{2,5}

¹Grupo de Investigación Altos Estudios de Frontera, Universidad Simón Bolívar, Cúcuta, Colombia

²Endocrine and Metabolic Diseases Research Center, School of Medicine, University of Zulia, Maracaibo, Venezuela

³Latacunga Province General Hospital, Ministry of Public Health, Cotopaxi, Ecuador

⁴Geriatric Research Education and Clinical Center, United States Department of Veterans Affairs, Miami, Florida, USA

⁵Division of Pulmonary and Critical Care Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, Massachusetts, 02115, USA

V2 First published: 10 May 2018, 7:565
<https://doi.org/10.12688/f1000research.14571.1>
 Second version: 22 May 2018, 7:565
<https://doi.org/10.12688/f1000research.14571.2>
 Latest published: 11 Jan 2019, 7:565
<https://doi.org/10.12688/f1000research.14571.3>

Abstract

Background: A growing body of evidence suggests that cigarette smoking can cause the onset of metabolic syndrome prior to cardiovascular diseases. Therefore, the objective of this study was to evaluate the relationship between smoking habit and metabolic syndrome components in an adult population from Maracaibo city, Venezuela.

Methods: The Maracaibo City Metabolic Syndrome Prevalence Study is a descriptive, cross-sectional study with random and multi-stage sampling. In this sub-study, 2212 adults from both genders were selected. On the basis of their medical background, they were classified as smokers, non-smokers and former smokers. Metabolic syndrome was defined according to Harmonizing 2009 criteria, using population-specific abdominal circumference cut-off points. The association between risk factors was evaluated using a logistic regression model.

Results: In the studied population, 14.8% were smokers, 15.4% were former smokers. In the multivariate analysis, the presence of

Open Peer Review

Approval Status

	1	2
version 3 (revision) 11 Jan 2019	 view	
version 2 (revision) 22 May 2018		 view
version 1 10 May 2018	 view	

1. Eiji Oda, Tachikawa General Hospital, Nagaoka, Japan
2. Juan P. González-Rivas , The Andes Clinic of Cardio-Metabolic Studies, Mérida,

metabolic syndrome (smokers: OR, 1.54; 95% CI, 1.11–2.14; $p=0.010$) and its components were related to cigarette smoking, with the exception of hyperglycemia. High blood pressure was inversely associated with current smoking status (smokers: OR, 0.70 (0.51–0.95); $p=0.025$).

Conclusion: Cigarette smoking represents a related factor with metabolic syndrome, being associated with low high-density lipoprotein-cholesterol, increased abdominal circumference and elevated triacylglyceride levels. Former smokers did not present a greater risk for developing this metabolic disease when compared to non-smokers. The effect of avoiding this habit should be evaluated in future studies in our population.

Keywords

smoking habit, metabolic syndrome, smokers, hypertension, cardiovascular risk.

Venezuela

Foundation for Clinical, Public Health, and
Epidemiological Research of Venezuela
(FISPEVEN), Mérida, Venezuela

Any reports and responses or comments on the
article can be found at the end of the article.

Corresponding authors: Valmore Bermudez (valmore@gmail.com), Juan Salazar (juanjsv18@hotmail.com)

Author roles: **Bermudez V:** Conceptualization, Writing – Review & Editing; **Olivar LC:** Data Curation, Formal Analysis; **Torres W:** Data Curation, Formal Analysis; **Navarro C:** Formal Analysis, Investigation; **Gonzalez R:** Formal Analysis, Investigation; **Espinoza C:** Formal Analysis, Writing – Original Draft Preparation; **Morocho A:** Formal Analysis, Writing – Original Draft Preparation; **Mindiola A:** Formal Analysis, Writing – Original Draft Preparation; **Chacin M:** Data Curation, Investigation; **Arias V:** Data Curation, Investigation; **Añez R:** Data Curation, Formal Analysis; **Salazar J:** Conceptualization, Writing – Review & Editing; **Riaño-Garzon M:** Funding Acquisition, Writing – Original Draft Preparation; **Diaz-Camargo E:** Data Curation, Funding Acquisition; **Bautista MJ:** Formal Analysis, Funding Acquisition; **Rojas J:** Conceptualization, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This study was funded by the Council of Scientific, Humanistic and Technological Development under No. CC-0437-10-21-09-10 and Fundacite-Zulia under No. FZ-0058-2007.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2018 Bermudez V *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. Data associated with the article are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

How to cite this article: Bermudez V, Olivar LC, Torres W *et al.* **Cigarette smoking and metabolic syndrome components: a cross-sectional study from Maracaibo City, Venezuela [version 2; peer review: 1 approved, 1 approved with reservations]** F1000Research 2018, 7:565 <https://doi.org/10.12688/f1000research.14571.2>

First published: 10 May 2018, 7:565 <https://doi.org/10.12688/f1000research.14571.1>

REVISED Amendments from Version 1

We want to thank the reviewer courtesy for evaluate and comment on our manuscript.

We share your opinion about the results limitations given the study design, but the cross sectional analysis is the most viable way to evaluate risk factors in our country.

According to your suggestion, the conclusion of the manuscript has been edited, specifically substituting the assertion of smoking habit as "independent risk factor" for "related factor".

See referee reports

Introduction

Smoking is one the main causes of morbidity and mortality in the working-age population; it is responsible for approximately 7.2 million deaths per year¹. This constitutes a major public health issue. Almost one-third of the world population older than 15 years of age smokes², with a global prevalence of 21.2% in developing countries³. In the Americas, the prevalence in the general adult population is 17.1%⁴; however, this varies among different countries, with Chile having the highest (38.9%) and Panama the lowest (7.4%) rates⁴. Venezuela is a country with one of the highest prevalence (33.9%)⁵, with a frequency of 14.8% in Maracaibo City in recent studies⁶.

These percentages are important because smoking habit is a major modifiable risk factor for developing non-communicable diseases², including cardiovascular disease (CVD) and type 2 diabetes mellitus (DM2)⁷. A growing body of evidence suggests that before the onset of these two diseases, cigarette smoking favors the appearance of metabolic syndrome (MS)^{8,9} high blood pressure, dyslipidemia, obesity and high blood glucose¹⁰⁻¹⁴. Main contributors for this association include the presence of dyslipidemia and central obesity¹⁵.

The physiopathology of the relationship between cigarette smoking and MS comes from a decrease in peripheral insulin sensitivity, lipoproteins metabolism alterations and endothelial dysfunction, all present in smoking individuals¹⁶. Until now, epidemiological published results are not definitive in showing the association between cigarette smoking and MS. On the other hand, it is not certain whether this association is caused by other behavioral patterns and unhealthy habits of patients with cardiometabolic diseases¹⁷. Thus, the aim of this study was to evaluate the relationship between smoking habit and MS components in the adult population from Maracaibo City, Venezuela.

Methods

Study design and subject selection

The Maracaibo city MS prevalence study (MMSPS) was a cross-sectional, descriptive study performed in Maracaibo, Venezuela. It was designed to provide estimations about the presence of MS and associated cardiovascular risk factors in the adult population during the period between May 2007 and December 2009. The study method was reported previously¹⁸. The most important aspects of the protocol are presented here. Maracaibo city was divided into parishes, which were sampled

proportionally through a multistage random sampling, defining conglomerates in two phases: In the first phase, the conglomerates represented the sectors of the 18 parishes, selecting 4 areas per parish by means of simple random sampling; in the second phase, the conglomerates were represented by the neighborhood of each chosen area, to which a random number was assigned. To evaluate smoking habit in this sub-study, 18 subjects were excluded: 9 because of inconsistencies about when they started smoking; the other 9 subjects because they smoked cigar types that were different to cigarettes. Finally, a total of 2212 subjects were evaluated. The study was approved by the Bioethics Committee of the Endocrine and Metabolic Research Center – University of Zulia (approval number: BEC-006-0305). This ethical approval included all future studies that used the data from the MMSPS. All participants signed an informed consent form before being questioned and physically examined by a trained team.

Clinical evaluation

Every subject in the study underwent a medical examination performed by trained personnel to obtain a full medical history. During the anamnesis, past medical and family history of endocrine and metabolic disorders was collected; including age, race, marriage status, education and socioeconomic status. The latter was measured using the Graffar scale modified by Mendez-Castellano and De Mendez¹⁹.

The auscultatory method was performed to measure arterial pressure, using an adequate calibrated and validated sphygmomanometer. Korotkoff phases I and V were used to measure systolic and diastolic pressures, respectively. Subjects remained sitting still for 15 minutes before assessment, with both feet on the ground. A total of 3 measurements per day were taken in 15 minute intervals, for 2 days consecutively. The employed criteria to classify the population came from the Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC7)²⁰.

Anthropometric measures were taken using a height rod that had been previously calibrated and placed on a flat surface. Weight was measured using a digital weighing scale (Tanita, TBF-310 GS Body Composition Analyzer; Tokyo, Japan), with the patient wearing light clothes and no shoes. The body mass index (BMI) was calculated applying the Quetelet formula (weight/height²), and classified according to the WHO classification²¹, as follows: normal weight (<25 Kg/m²), overweight (25.0–29.9 Kg/m²), obese (≥30.0 Kg/m²). Abdominal circumference was measured using a plastic measuring tape, graded in centimeters and millimeters, in a spot equidistant to the lower ribcage and the anterior-superior iliac spine, according to the United States National Institute of Health protocol²².

Smoking habit evaluation

Subjects were asked about smoking habit presence and duration, being categorized as: a) current smoker, any subject who had smoked more than 100 cigarettes in his/her lifetime, is currently smoking, or less than 1 year had passed after he/she quit smoking; b) Former smoker: any subject who has quit smoking for more than 1 year; c) non-smoker, any subject who has never smoked or had smoked less than 100 cigarettes in his/her lifetime⁶. Smoking intensity was assessed posteriorly,

according to number of cigarettes per day. It was divided in the following tertiles: T1 <3 cigarettes/day; T2 = 3–9 cigarettes/day; and T3 ≥10 cigarettes/day.

Physical activity and alcohol consumption evaluations

Physical activity was evaluated using the International Physical Activity Questionnaire²³. It takes into account four elements of evaluation: physical activity in transport, work, domestic and gardening, and leisure time. To quantify time investment on each element, subjects were classified in quintiles. The final scoring was reported using metabolic equivalents (METs)-min/week on each item; any subject with 0 METs was considered as physically inactive.

Subjects with ≥1 MET were classified in quintiles according to gender, resulting in six categories for physical activity: physical inactivity (MET = 0), very low (Q1), low (Q2), moderate (Q3), high (Q4), and very high (Q5) physical activity. Leisure time was classified as: a) Q1 or very low physical activity, <296.999 METs for men and <230.999 METs for women; b) Q2 or low physical activity, 297.000–791.999 METs for men and 231.000–445.499 METs for women; Q3 or moderate physical activity, 792.000–1532.399 METs for men and 445.500–742.499 METs for women; Q4 or high physical activity, 1532.400–2879.999 MET for men and 742.500–1798.499 METs for women; and e) Q5 or very high physical activity, ≥2879.000 METs for men and ≥1798.500 METs for women. For alcohol consumption, any subject that drinks ≥1 gram daily was considered as a “drinker”²⁴.

Laboratory analysis

After 8 hours of fasting, a blood sample was taken from the cubital vein, and was centrifuged to obtain the serum. Serum levels of glucose (catalog number REF-10123), total cholesterol (catalog number REF-10015) and triacylglycerides (TAG) (catalog number REF-10163) were determined using enzymatic–colorimetric kits (Human Gesellschaft für Biochemica und Diagnostica mbH) and a specialized computer system. Glycemic status was classified according to ADA 2017 criteria in normal glucose (basal glucose, <100 mg/dl), impaired fasting glucose (basal glucose, 100–125 mg/dl) and DM2 (≥126 mg/dl)²⁵. Serum hs-C reactive protein (hs-CRP) levels were quantified using immunoturbidimetric assays (Human Gesellschaft für Biochemica und Diagnostica mbH, (catalog number REF-11544), setting the cutoff point at ≥0,765 mg/l²⁶.

Basal insulin serum levels were determined using a commercial kit (catalog number EIA-2935) based on the ELISA method (DRG International, Inc.), with a detection limit of <1 mU/l. Insulin resistance (IR) was calculated using software (HOMA-Calculator v2.2.2) supplied by the Oxford Centre for Diabetes, Endocrinology and Metabolism; the cutoff-point for HOMA2-IR was 2.00²⁷.

Metabolic syndrome evaluation

MS diagnosis was made using the proposed criteria from the IDF and AHA/NHLBI in 2009²⁸. It requires three or more of the following components to achieve a diagnosis: 1) TAG ≥150 mg/dl; 2) high-density lipoprotein–cholesterol (HDL-C) <40 mg/dl for men or <50 mg/dl for women; 3) basal glucose levels

≥100 mg/dl, or a previous diagnosis of DM2 or use of an antidiabetic drug; 4) arterial pressure ≥130/85 mmHg, or a previous diagnosis of hypertension or use of an antihypertensive drug; 5) abdominal circumference with cutoff points adapted for our population, which are ≥91 for women and ≥98 cm for men²⁹.

Statistical analysis

Qualitative variables were expressed in absolute and relative frequencies. The relationship between these was examined with a χ^2 test and the difference in proportions using a Z-test. Quantitative variables were expressed in arithmetic means $\pm$ standard deviations, with prior analysis using Geary's test. Variables without a normal distribution were submitted to logarithmic transformation with posterior normality test. Multiple logistic regression models were made to estimate odds ratios (OR) and 95% confidence intervals (95% CI); they were used for the presence of MS and each of its components, adjusted for gender, age, ethnic group, marital status, education level, socioeconomic status, working status, alcohol consumption, BMI categories, insulin resistance and smoking habit. On another model, smoking intensity was assessed dividing consumption in tertiles (T1 <3 cigarettes/day; T2 = 3–9 cigarettes/day; and T3 ≥10 cigarettes/day). Data were analyzed by using SPSS v.21 for Windows (IBM SPSS), and considering statistically significant results when $p < 0.05$.

Results

General characteristics of the sample

A total of 2212 individuals were studied, of whom 52.7% (n=1166) were women. The mean age $\pm$ SD was 39.27 $\pm$ 15.38 years, and the most frequently occurring age group was 30–49 years (38.5%; n=851). For smoking habit, 14.8% were smokers (n=328), 15.4% former smokers (n=340) and 69.8% were non-smokers (n=1544). The prevalence of MS was 35.7% (n=935) in the sample. The most frequent MS components were low HDL-C (57.6%; n=1275) and abdominal obesity (48.5%; n=1072). Other general characteristics can be found in Table 1.

Smoking habit and metabolic syndrome components

Smoking habit in accordance to MS components could be seen in Table 2. It shows a statistically significant association between cigarette smoking and having MS ($\chi^2=39.285$; $p < 0.001$) with a greater percentage of individuals with MS in former smokers (47.9%) and current smokers (42.1%) than in non-smokers (31.6%), $p < 0.05$.

Each component of the MS was analyzed in relation to smoking. A higher percentage of individuals with high TAG were former (37.1%) and current (36.9%) smokers, compared with non-smokers (23.6%) ($\chi^2=41.886$; $p < 0.001$). The same happened for abdominal obesity in former smokers (62.9%) and current smokers (52.1%) ($\chi^2=40.039$, $p < 0.001$). A high percentage of former smokers presented hyperglycemia (33.5% vs 25.9%; $\chi^2=10.759$; $p < 0.005$) and high blood pressure (48.5 vs 36.9%; $\chi^2=16.88$; $p < 0.001$) in comparison to nonsmokers. No statistical association was found between low HDL-C and smoking status.

Comparing smoking habit with the number of MS criteria (Figure 1), a statistically significant association was observed ($\chi^2=49.249$, $p < 0.001$). The highest percentages were

Table 1. General characteristics of the sample population.

Variable	Female (n=1166)		Male (n=1046)		Total (n=2212)	
	n	%	n	%	n	%
Age (years)						
<30	349	29.9	410	39.2	759	34.3
30–49	464	39.8	387	37.0	851	38.5
≥50	353	30.3	249	23.8	602	27.2
Marital status						
Single	495	42.5	455	43.5	950	42.9
Married	427	36.6	446	42.6	873	39.5
Other	244	20.9	145	13.9	389	17.6
Socioeconomic status						
Stratus I: high class	17	1.5	19	1.8	36	1.6
Stratus II: middle-high class	206	17.7	204	19.5	410	18.5
Stratus III: middle class	432	37.0	440	42.1	872	39.4
Stratus IV: Working class	446	38.3	344	32.9	790	35.7
Stratus V: extreme poverty	65	5.6	39	3.7	104	4.7
Working status						
Employed	527	45.2	757	72.4	1284	58.0
Unemployed	639	54.8	289	27.6	928	42.0
Education						
Illiterate	33	2.8	19	1.8	52	2.4
Primary school	237	20.3	112	10.7	349	15.8
High school	516	44.3	519	49.6	1035	46.8
Higher education	380	32.6	396	37.9	776	35.1
Ethnic group						
Mixed	873	74.9	807	77.2	1680	75.9
White-Hispanic	189	16.2	159	15.2	348	15.7
Afro-Venezuelan	30	2.6	36	3.4	66	3.0
Indigenous American	62	5.3	43	4.1	105	4.7
Other	12	1.0	1	0.1	13	0.6
Smoking habit						
Non smoker	878	75.3	666	63.7	1544	69.8
Current smoker	134	11.5	194	18.5	328	14.8
Former smoker	154	13.2	186	17.8	340	15.4
Leisure time physical activity						
Inactive	832	71.4	529	50.9	1361	61.7
Very low	67	5.7	99	9.5	166	7.5
Low	62	5.3	102	9.8	164	7.4
Moderate	64	5.5	103	9.9	167	7.6
High	72	6.2	92	8.8	164	7.4
Very high	69	5.9	115	11.1	184	8.3
Alcohol consumption	219	18.8	505	48.3	724	32.7
Metabolic syndrome [†]	470	40.3	465	44.5	935	42.3
Metabolic syndrome [‡]	391	33.5	398	38.0	789	35.7
Low HDL [†]	747	64.1	528	50.5	1275	57.6
High TAG [†]	268	23.0	343	32.8	611	27.6
Hyperglycemia [†]	301	25.8	317	30.3	618	27.9
High blood pressure [†]	406	34.8	447	42.7	853	38.6
Abdominal obesity [†]	920	78.9	741	70.8	1661	75.1
Abdominal obesity [‡]	569	48.8	503	48.1	1072	48.5

WC, waist circumference; HDL, high-density lipoprotein; TAG, triacylglycerides.
[†]According to IDF and AHA/NHLBI-ISO-2009 Consensus²⁸. [‡]According to The Maracaibo City Metabolic Syndrome Study²⁹: WC ≥98cm for men; ≥91cm for women.

Table 2. Smoking habit in accordance with MS and its components. Maracaibo, Venezuela.

Smoking status	MS [‡]		High TAG [†]		Abdominal obesity [‡]		Hyperglycemia [†]		Low HDL-C [†]		High BP [†]	
	n	%	n	%	n	%	n	%	n	%	n	%
Nonsmoker	488	31.6	364	23.6	687	44.5	400	25.9	878	56.9	569	36.9
Current Smoker	138	42.1	121	36.9	171	52.1	104	31.7	201	61.3	119	36.3
Former Smoker	163	47.9	126	37.1	214	62.9	114	33.5	196	57.6	165	48.5

MS, Metabolic Syndrome; TAG, triacylglycerides; HDL-C, high-density lipoprotein-cholesterol; BP, blood pressure.

MS: $\chi^2 = 39.285$ ($p < 0.001$); High TAG: $\chi^2 = 41.886$ ($p < 0.001$); Abdominal Obesity: $\chi^2 = 40.039$ ($p < 0.001$); Hyperglycemia: $\chi^2 = 10.759$ ($p = 0.005$); Low HDL-C levels: $\chi^2 = 2.160$ ($p = 0.340$); High Blood Pressure: $\chi^2 = 16.883$ ($p < 0.001$). [†]According to IDF and AHA-NHLBI-ISO 2009²⁸. [‡]According to The Maracaibo City Metabolic Syndrome Study²⁹: WC ≥ 98 cm for men; ≥ 91 cm for women.

Figure 1. Smoking habit and number of metabolic syndrome criteria in subjects from Maracaibo, Venezuela.

with nonsmokers who met 0 criteria (76.31%) and 1 criterion (76.77%). However, the greatest prevalence of smokers was observed in subjects who met 4 (16.96%) and 5 (20%) criteria.

Smoking habit as a risk factor for MS and its components

In Table 3, models of multivariate logistic regression are shown for the diagnosis of MS and its components. An association between current smoking and increased risk of presenting with MS could be observed (OR, 1.54; 95% CI, 1.11–2.14; $p = 0.010$); the same was true of high TAG serum levels (OR, 1.66; 95% CI, 1.23–2.23; $p < 0.001$); abdominal obesity (OR, 1.54; 95% CI, 1.05–2.28; $p = 0.027$) and low HDL-C levels (OR, 1.32; 95% CI, 1.01–1.74; $p = 0.046$).

On the other hand, by assessing smoking intensity according to number of cigarettes per day (Table 4), an association between

the consumption tertile and high serum TAG levels was observed (T3: OR, 1.51; 95% CI, 1.03–2.22; $p = 0.036$). Also, an association was observed between smoking intensity and abdominal obesity (T3: OR, 2.05; 95% CI, 1.15–3.64; $p = 0.015$). By contrast, an inverse relationship was observed with high blood pressure (T3: OR, 0.66; 95% CI, 0.44–0.99; $p = 0.045$).

Dataset 1. Cigarette smoking and MS MMSPS dataset³⁰

<http://dx.doi.org/10.5256/f1000research.14571.d201851>

Discussion

Cigarettes are composed of more than 1000 toxic and carcinogenic elements. Nicotine is the main alkaloid in tobacco; it constitutes 1.5% of the commercial cigarette weight and 95% of the total alkaloids present³¹. Despite its effects, cigarette smoking has

Table 3. Adjusted OR for metabolic syndrome and its components according to smoking habit in Maracaibo, Venezuela.

Smoking status	MS [‡]	High TAG [†]	Abdominal Obesity [‡]	Hyperglycemia [†]	Low HDL-C [†]	High BP [†]
	OR (95% CI); p	OR (95% CI); p	OR (95%CI); p	OR (95%CI); p	OR (95%CI); p	OR (95%CI); p
Non-smokers	1.00	1.00	1.00	1.00	1.00	1.00
Current smokers	1.54 (1.11–2.14); 0.010	1.66 (1.23–2.23); <0.001	1.54 (1.05–2.28); 0.027	1.13 (0.83–1.54); 0.408	1.32 (1.01–1.74); 0.046	0.70 (0.51–0.95); 0.025
Former smokers	0.96 (0.70–1.31); 0.799	1.15 (0.86–1.53); 0.324	1.24 (0.85–1.81); 0.245	0.86 (0.64–1.16); 0.330	0.97 (0.74–1.26); 0.832	0.79 (0.58–1.06); 0.127

TAG, triacylglycerides; HDL-C, high-density lipoprotein-cholesterol; BP, blood pressure; OR, odds ratio; CI, confidence interval. [†]According to IDF and AHA/NHLBI-ISO 2009²⁸. [‡] According to The Maracaibo City Metabolic Syndrome Study²⁹; WC ≥98cm for men; ≥91cm for women.

Table 4. Adjusted odds ratios for MS and its components according to smoking habit intensity in Maracaibo, Venezuela.

Smoking status	MS [‡]	High TAG [†]	Abdominal Obesity [‡]	Hyperglycemia [†]	Low HDL-C [†]	High BP [†]
	OR (95%CI); p	OR (95%CI); p	OR (95%CI); p	OR (95%CI); p	OR (95%CI); p	OR (95%CI); p
Non-smoker	1.00	1.00	1.00	1.00	1.00	1.00
<3 cigarettes/day	1.13 (0.79–1.62); 0.487	1.25 (0.90–1.73); 0.178	1.42 (0.94–2.13); 0.092	1.02 (0.72–1.42); 0.932	1.10 (0.82–1.47); 0.539	0.75 (0.54–1.06); 0.101
3–9 cigarettes/day	1.25 (0.88–1.78); 0.208	1.41 (1.02–1.95); 0.035	1.11 (0.73–1.69); 0.620	1.01 (0.72–1.41); 0.968	1.24 (0.91–1.69); 0.166	0.81 (0.58–1.13); 0.210
≥10 cigarettes/day	1.21 (0.78–1.88); 0.381	1.51 (1.03–2.22); 0.036	2.05 (1.15–3.64); 0.015	0.90 (0.61–1.35); 0.620	1.02 (0.70–1.49); 0.912	0.66 (0.44–0.99); 0.045

MS, metabolic syndrome; TAG, triacylglycerides; HDL-C, high-density lipoprotein-cholesterol; BP, blood pressure. [†]According to IDF and AHA/NHLBI-ISO 2009²⁸. [‡]According to The Maracaibo City Metabolic Syndrome Study²⁹; WC ≥98cm for men; ≥91cm for women.

spread all over the globe, becoming a leading cause of chronic and degenerative pathologies. In Latin America, its use has markedly increased since 1950, and is now considered the second most common cardiovascular risk factor, following high blood pressure³². This has led to an increase in cancer deaths and a drop in life expectation of 2–6 years³³. In Maracaibo, high prevalence of cigarette smoking and MS has been observed, which may suggest an existing relationship between these variables^{6,34}.

In this study, MS prevalence in current smokers was 42.1% in both genders and a greater probability of having MS than in nonsmokers was observed. Kang and Song in the Korea National Health and Nutrition Examination Survey (KHANES) reported similar results with a cross-sectional study with 11559 subjects. They evaluated smoking habit by looking for nicotine in urine samples; a greater risk for developing MS was observed in those subjects³⁵. Likewise, Slagter *et al.*¹¹ conducted a study in the Netherlands which included 59,467 subjects from both sexes. In that study, a higher prevalence of MS was observed in smokers (a dose-dependent relationship), and increased the risk of MS depended neither on BMI nor gender.

On the other hand, Sun *et al.*⁸ conducted a meta-analysis from multiple cohort studies and included 13 articles. In total, 56,691

subjects and 8688 cases from Asia, Europe and North America were included. They found that cigarette smoking actively increases the risk of having MS. The effects of smoking on the cardiovascular system could be caused by increased action of nicotinic receptors. Activation of nicotinic receptor could promote the release of neurotransmitters and hormones such as vasopressin, CRH, ACTH, growth hormone, dopamine, serotonin, glutamate and GABA in the central nervous system, acetylcholine in the peripheral nervous system, and catecholamine and cortisol from the adrenal glands. All of these molecules affect metabolism and appetite regulation³⁶.

The CKB cohort study³⁷ included 487,527 adult subjects and reported that regular cigarette smoking was associated with a decrease in BMI and an increase in abdominal circumference in both men and women (they used an adjusted model for BMI). Similar results were reported from the FINRISK study³⁸, which included 5817 Finnish adults; greater abdominal circumference was observed in overweight and obese women who smoke. Clair *et al.*³⁹, in a cross-sectional study that included 6123 adult Caucasians from Switzerland, reported that both sexes had an increased risk of obesity according to the number of cigarettes they smoke per day. These results resemble those from the present study, where cigarette smoking

was associated with increased abdominal obesity. This epidemiological behavior could be explained by the recent hypothesis of the association between cigarette smoking and a decrease in body weight, using the CHRNA3 genetic variant (rs1051730); establishing that smoking does not affect body fat distribution and the increase in localized visceral fat and in abdominal obesity are due to high cortisol plasma levels and insulin resistance, respectively^{40–42}.

According to the results of our study, cigarette smoking represents a risk factor for having high TAG levels. Similar behavior was seen in the ICMR-INDIAB cross-sectional study⁴³ of 16,607 adult individuals, which showed a positive correlation between high TAG levels and smoking. Ueyama *et al.*⁴⁴ reported there was a positive association between smoking and high TAG levels in a study of 5959 Japanese individuals. This phenomenon could be explained by the fact that stimulation of the sympathetic nervous system produces the release of insulin antagonists. These antagonists, such as cortisol and growth hormone, increase lipolysis, leading to an elevation of free fatty acids in the blood^{8,42}.

Low HDL-C levels were observed more frequent in smokers than nonsmokers in our study. Sun *et al.*⁴⁵, in a cross-sectional study of 11,956 Chinese individuals, reported similar results by showing that current smokers had an increased risk of having low HDL-C levels. Takata *et al.*⁴⁶, showed that in 32 individuals who were participating in an anti-smoking program using varenicline or transdermal nicotine patches, HDL-C levels, apolipoprotein AI and HDL subfractions did not change significantly according to therapeutic strategy used. In the same study, cholesterol efflux capacity and HDL inflammatory index improved significantly with the anti-smoking program (baseline cholesterol efflux capacity: $14.15 \pm 2.46\%$ vs after smoking cessation cholesterol efflux capacity: $14.83 \pm 2.35\%$; $p=0.01$; baseline HDL inflammatory index: 1.13 ± 0.31 vs after smoking cessation HDL inflammatory index: $0.98 \pm 0.18\%$; $p=0.01$).

The inverse relationship between current cigarette smoking and high blood pressure observed in the present study is noteworthy. However, three decades ago cigarette smoking was globally reported as acutely increasing blood pressure, heart frequency and myocardial contractility⁴⁷. This was thought to be caused by increased nicotinic activity on the sympathetic nervous system. Despite this, epidemiological evidence could not confirm the role of cigarette smoking in the development of elevated blood pressure⁴⁸. On the other hand, diverse evidence suggests an inverse relationship between these factors. Kaneko *et al.*⁴⁹, in a recent study of 1297 Japanese individuals without any history of high blood pressure, showed that cigarette smoking appeared to be a protective factor against blood pressure elevation. Onat *et al.*⁵⁰ observed a similar pattern in a Turkish population. The inverse association between blood pressure and smoking habit could be related to the cigarette effect on weight loss, since obesity is associated with a high incidence of high blood pressure; explaining the rebound effect on blood pressure in obese subjects who stop smoking⁵⁰; however, this study evidenced that smokers presented with more abdominal obesity

than nonsmokers. Another theory to explain this behavior suggests that smokers show less response to psychological stress: many of them report a decrease in anxiety and stress when smoking a cigarette⁵¹. This may come from modifications to adrenal and cardiovascular responses to external stimuli caused by cigarette smoking; thus, stopping smoking would increase blood pressure⁵².

Leone⁵³ reports a two-phase effect of cigarette smoking on arterial pressure: the first phase, without a determined duration, when there is a decrease in blood pressure; and the second phase, when the smoker develops elevated blood pressure from the toxic effects of carbon monoxide⁵³. This finding shows the importance in chronologically assessing smoking habit duration. Despite this, smoking does not benefit to cardiovascular health, but increases the risk of cardiovascular disease, especially in men⁵⁰.

Similarly, with the assessment of smoking intensity according to number of cigarettes per day, a direct relationship was found between the number of cigarettes smoked and an increased risk of high serum TAG levels and abdominal obesity, and an inverse relationship with hypertension; this was seen especially in heavy smokers (≥ 10 cigarette daily). In this sense, in a study performed by Chen *et al.*¹⁰, 1146 individuals showed a significant dose-response relationship between the number of cigarettes per day and high TAG levels. Data analysis from the KHANES study revealed an increased risk of obesity and central obesity with an increase in smoking habit intensity⁵⁴. This relationship could be caused by the dose-dependent effect of nicotine on fatty acid metabolism and catecholamine release; also inducing increase in lipolysis, free fatty acids, VLDL, LDL levels, and visceral adipose tissue independent of weight gain or loss⁵⁵.

In the present study, former smokers did not exhibit an increased risk of developing MS or its components when compared with non-smokers. Similar results were reported in Korea by Oh *et al.*⁵⁶. The benefits to cardiovascular health from stopping cigarette smoking seem to depend on the following variables: first, the time since the subject stopped; and second, the length of time for which he/she was smoking and the quantity of cigarettes. A previous study showed that smoking 20 cigarettes daily increased the risk of developing MS for the next 10 years, whereas smoking 40 cigarettes daily increased the risk for the next 20 years⁵⁷. This is why in the Maracaibo population it is necessary to conduct a cohort study on subjects who stopped smoking to evaluate the long term effects on cardiometabolic health.

Regarding the limitations of this study, its cross-sectional design makes it incapable of determining causality; it is also influenced by the subjectivity of its participants regarding the intensity and duration of their smoking habit. All of this should be considered in future studies.

In conclusion, the present study showed that smoking in our population represent a related factor with MS, and is individually associated with low HDL-C levels, increased abdominal circumference and high TAG levels. Former smok-

ers did not show any increase in risk of present MS relative to non-smokers; despite this, future research studies should be conducted to evaluate how stopping cigarette smoking decreases cardiometabolic risk. Prevention measures focused on patients who smoke, especially anti-smoking counseling from medical personnel, could help to decrease any cigarette cardiometabolic consequences in the Maracaibo City population.

Data availability

Dataset 1. Cigarette smoking and MS MMSPS dataset. DOI: [10.5256/f1000research.14571.d201851](https://doi.org/10.5256/f1000research.14571.d201851)³⁰

Competing interests

No competing interests were disclosed.

Grant information

This study was funded by the Council of Scientific, Humanistic and Technological Development under No. CC-0437-10-21-09-10 and Fundacite-Zulia under No. FZ-0058-2007.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- GBD 2015 Risk Factors Collaborators: **Global, regional, and national comparative risk assessment of 79 behavioural, environmental and occupational, and metabolic risks or clusters of risks, 1990-2015: a systematic analysis for the Global Burden of Disease Study 2015.** *Lancet*. 2016; **388**(10053): 1659–1724. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- World Health Organization: **WHO global report on trends in prevalence of tobacco smoking 2015.** [Reference Source](#)
- Organización Mundial de la Salud: **Informe OMS sobre la epidemia mundial de tabaquismo: aumentar los impuestos al tabaco.** Ginebra: OMS; 2015. [Reference Source](#)
- Cayon A: **Regional Report on Tobacco Control, 2016.** Pan American Health Organization / World Health Organization. 2017 [cited 25 August 2017]. [Reference Source](#)
- Blanco A, Caixeta R, Sandoval R, *et al.*: **Informe sobre Control del Tabaco en Venezuela.** Organización Panamericana de Salud. 2011. [Reference Source](#)
- Bermúdez V, Miquilena E, Salazar J, *et al.*: **Smoking Habit in Adult Population from Maracaibo City, Venezuela.** *Int J Respir Pulm Med*. 2016; **3**: 061. [Publisher Full Text](#)
- World Health Organization: **WHO Global Status Report on Noncommunicable Diseases.** Attaining the nine global noncommunicable diseases targets: a shared responsibility. Geneva: WHO. 2014. [Reference Source](#)
- Sun K, Liu J, Ning G: **Active smoking and risk of metabolic syndrome: a meta-analysis of prospective studies.** *PLoS One*. 2012; **7**(10): e47791. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Al-khalifa I, Mohammed S, Ali Z: **Cigarette Smoking as a Relative Risk Factor for Metabolic Syndrome.** *J Endocrinol Metab*. 2016; **6**(6): 178–182. [Publisher Full Text](#)
- Chen CC, Li TC, Chang PC, *et al.*: **Association among cigarette smoking, metabolic syndrome, and its individual components: the metabolic syndrome study in Taiwan.** *Metabolism*. 2008; **57**(4): 544–548. [PubMed Abstract](#) | [Publisher Full Text](#)
- Slagter SN, van Vliet-Ostapchouk JV, Vonk JM, *et al.*: **Associations between smoking, components of metabolic syndrome and lipoprotein particle size.** *BMC Med*. 2013; **11**(1): 195. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Dedinska I, Laca L, Miklušica J, *et al.*: **Czypalenytoniu jest czynnikami ryzyka zespołometabolicznego?** *Clinical Diabetology*. 2014; **3**(4): 136–143.
- Tsai JS, Chen SC, Huang KC, *et al.*: **Plasma zinc α2-glycoprotein levels are elevated in smokers and correlated with metabolic syndrome.** *Eur J Clin Invest*. 2015; **45**(5): 452–459. [PubMed Abstract](#) | [Publisher Full Text](#)
- Yankey BN, Strasser S, Okosun IS: **A cross-sectional analysis of the association between marijuana and cigarette smoking with metabolic syndrome among adults in the United States.** *Diabetes Metab Syndr*. 2016; **10**(2 Suppl 1): S89–S95. [PubMed Abstract](#) | [Publisher Full Text](#)
- Cena H, Fonte ML, Turconi G: **Relationship between smoking and metabolic syndrome.** *Nutr Rev*. 2011; **69**(12): 745–753. [PubMed Abstract](#) | [Publisher Full Text](#)
- Jia WP: **The impact of cigarette smoking on metabolic syndrome.** *Biomed Environ Sci*. 2013; **26**(12): 947–952. [PubMed Abstract](#) | [Publisher Full Text](#)
- Rabaeus M, Salen P, de Lorgeril M: **Is it smoking or related lifestyle variables that increase metabolic syndrome risk?** *BMC Med*. 2013; **11**(1): 196. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Bermúdez V, Marcano RP, Cano C, *et al.*: **The Maracaibo city metabolic syndrome prevalence study: design and scope.** *Am J Ther*. 2010; **17**(3): 288–294. [PubMed Abstract](#) | [Publisher Full Text](#)
- Méndez-Castellano H, De Méndez M: **Estratificación social y biología humana: método de Graffar modificado.** *Arch Ven Pueric Pediatr*. 1986; **49**: 93–104.
- Chobanian AV, Bakris GL, Black HR, *et al.*: **The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: the JNC 7 report.** *JAMA*. 2003; **289**(19): 2560–72. [PubMed Abstract](#) | [Publisher Full Text](#)
- World Health Organization: **The World Health Report 2003.** [Reference Source](#)
- Health Statistics: **NHANES III reference manuals and reports (CDROM).** Hyattsville, MD: Centers for Disease Control and Prevention, 1996. [Reference Source](#)
- Guidelines for Data Processing and Analysis of the International Physical Activity Questionnaire (IPAQ).** [Reference Source](#)
- Bermúdez V, Torres Y, Apruzzese V, *et al.*: **Alcohol drinking patterns in the adult population from the Maracaibo municipality, Zulia – Venezuela.** *Revista Latinoamericana de Hipertensión*. 2014; **9**(3). [Reference Source](#)
- American Diabetes Association: **2. Classification and Diagnosis of Diabetes.** *Diabetes Care*. 2017; **40**(Suppl 1): S11–S24. [PubMed Abstract](#) | [Publisher Full Text](#)
- Bermúdez V, Cabrera M, Mendoza L, *et al.*: **High-sensitivity c-reactive protein epidemiological behavior in adult individuals from Maracaibo, Venezuela.** *Revista Latinoamericana de Hipertensión*. 2013; **8**(1): 22–29. [Reference Source](#)
- Bermúdez V, Rojas J, Martínez MS, *et al.*: **Epidemiologic Behavior and Estimation of an Optimal Cut-Off Point for Homeostasis Model Assessment-2 Insulin Resistance: A Report from a Venezuelan Population.** *Int Sch Res Notices*. 2014; **2014**: 616271. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Alberti KG, Eckel RH, Grundy SM, *et al.*: **Harmonizing the metabolic syndrome: a joint interim statement of the International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society; and International Association for the Study of Obesity.** *Circulation*. 2009; **120**(16): 1640–1645. [PubMed Abstract](#) | [Publisher Full Text](#)
- Bermúdez V, Rojas J, Salazar J, *et al.*: **Sensitivity and Specificity Improvement in Abdominal Obesity Diagnosis Using Cluster Analysis during Waist Circumference Cut-Off Point Selection.** *J Diabetes Res*. 2015; **2015**: 750265. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Bermúdez V, Olivar LC, Torres W, *et al.*: **Dataset 1 in: Cigarette smoking and metabolic syndrome components: a cross-sectional study from Maracaibo City, Venezuela.** *F1000Research*. 2018. [Data Source](#)
- Benowitz NL, Hukkanen J, Jacob P 3rd: **Nicotine chemistry, metabolism, kinetics and biomarkers.** *Handb Exp Pharmacol*. 2009; **192**: 29–60. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

32. Pereira J, Peñaranda D, Reyes A, *et al.*: **Prevalencia de factores de riesgo cardiovascular en América Latina: una revisión de la evidencia publicada de 2010 a 2015.** *Revista mexicana de cardiología*. 2015; **26**(3): 125–139.
[Reference Source](#)
33. Palloni A, Novak B, Pinto-Aguirre G: **The enduring effects of smoking in Latin America.** *Am J Public Health*. 2015; **105**(6): 1246–1253.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
34. Bermudez V, Rojas J, Salazar J, *et al.*: **The Maracaibo city metabolic syndrome prevalence study: primary results and agreement level of 3 diagnostic criteria.** *Revista Latinoamericana de Hipertensión*. 2014; **9**: 1–15.
[Reference Source](#)
35. Kang JH, Song YM: **Association between cotinine-verified smoking status and metabolic syndrome: analyses of Korean National Health and Nutrition Examination Surveys 2008-2010.** *Metab Syndr Relat Disord*. 2015; **13**(3): 140–148.
[PubMed Abstract](#) | [Publisher Full Text](#)
36. Yildiz D: **Nicotine, its metabolism and an overview of its biological effects.** *Toxicol*. 2004; **43**(6): 619–632.
[PubMed Abstract](#) | [Publisher Full Text](#)
37. Lv J, Chen W, Sun D, *et al.*: **Gender-specific association between tobacco smoking and central obesity among 0.5 million Chinese people: the China Kadoorie Biobank Study.** *PLoS One*. 2015; **10**(4): e0124586.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
38. Tuovinen E, Saarni S, Männistö S, *et al.*: **Smoking status and abdominal obesity among normal- and overweight/obese adults: Population-based FINRISK study.** *Prev Med Rep*. 2016; **4**: 324–330.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
39. Clair C, Chiolerio A, Faeh D, *et al.*: **Dose-dependent positive association between cigarette smoking, abdominal obesity and body fat: cross-sectional data from a population-based survey.** *BMC Public Health*. 2011; **11**(1): 23.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
40. Winslow UC, Rode L, Nordestgaard BG: **High tobacco consumption lowers body weight: a Mendelian randomization study of the Copenhagen General Population Study.** *Int J Epidemiol*. 2015; **44**(2): 540–550.
[PubMed Abstract](#) | [Publisher Full Text](#)
41. Fujiyoshi A, Miura K, Kadowaki S, *et al.*: **Lifetime cigarette smoking is associated with abdominal obesity in a community-based sample of Japanese men: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA).** *Prev Med Rep*. 2016; **4**: 225–232.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
42. Wilkins JN, Carlson HE, Van Vunakis H, *et al.*: **Nicotine from cigarette smoking increases circulating levels of cortisol, growth hormone, and prolactin in male chronic smokers.** *Psychopharmacology (Berl)*. 1982; **78**(4): 305–308.
[PubMed Abstract](#) | [Publisher Full Text](#)
43. Joshi SR, Anjana RM, Deepa M, *et al.*: **Prevalence of dyslipidemia in urban and rural India: the ICMR-INDIAB study.** *PLoS One*. 2014; **9**(5): e96808.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
44. Ueyama C, Horibe H, Yamase Y, *et al.*: **Association of smoking with prevalence of common diseases and metabolic abnormalities in community-dwelling Japanese individuals.** *Biomed Rep*. 2017; **7**(5): 429–438.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
45. Sun G, Li Z, Guo L, *et al.*: **High prevalence of dyslipidemia and associated risk factors among rural Chinese adults.** *Lipids Health Dis*. 2014; **13**(1): 189.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
46. Takata K, Imaizumi S, Kawachi E, *et al.*: **Impact of cigarette smoking cessation on high-density lipoprotein functionality.** *Circ J*. 2014; **78**(12): 2955–2962.
[PubMed Abstract](#) | [Publisher Full Text](#)
47. Robertson D, Tseng CJ, Appalsamy M: **Smoking and mechanisms of cardiovascular control.** *Am Heart J*. 1988; **115**(1 Pt 2): 258–63.
[PubMed Abstract](#) | [Publisher Full Text](#)
48. Green MS, Jucha E, Luz Y: **Blood pressure in smokers and nonsmokers: epidemiologic findings.** *Am Heart J*. 1986; **111**(5): 932–940.
[PubMed Abstract](#) | [Publisher Full Text](#)
49. Kaneko M, Oda E, Kayamori H, *et al.*: **Smoking was a Possible Negative Predictor of Incident Hypertension After a Five-Year Follow-up Among a General Japanese Population.** *Cardiol Res*. 2012; **3**(2): 87–93.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
50. Onat A, Uğur M, Hergenç G, *et al.*: **Lifestyle and metabolic determinants of incident hypertension, with special reference to cigarette smoking: a longitudinal population-based study.** *Am J Hypertens*. 2009; **22**(2): 156–162.
[PubMed Abstract](#) | [Publisher Full Text](#)
51. McEwen A, West R, McRobbie H: **Motives for smoking and their correlates in clients attending Stop Smoking treatment services.** *Nicotine Tob Res*. 2008; **10**(5): 843–50.
[PubMed Abstract](#) | [Publisher Full Text](#)
52. al'Absi M, Wittmers LE, Erickson J, *et al.*: **Attenuated adrenocortical and blood pressure responses to psychological stress in ad libitum and abstinent smokers.** *Pharmacol Biochem Behav*. 2003; **74**(2): 401–10.
[PubMed Abstract](#) | [Publisher Full Text](#)
53. Leone A: **Smoking and Hypertension.** *J Cardiol Curr Res*. 2015; **2**(2): 00057.
[Publisher Full Text](#)
54. Kim Y, Jeong SM, Yoo B, *et al.*: **Associations of smoking with overall obesity, and central obesity: a cross-sectional study from the Korea National Health and Nutrition Examination Survey (2010-2013).** *Epidemiol Health*. 2016; **38**: e2016020.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
55. Chelland Campbell S, Moffatt RJ, Stamford BA: **Smoking and smoking cessation -- the relationship between cardiovascular disease and lipoprotein metabolism: a review.** *Atherosclerosis*. 2008; **201**(2): 225–35.
[PubMed Abstract](#) | [Publisher Full Text](#)
56. Oh S, Yoon Y, Lee E, *et al.*: **Association between cigarette smoking and metabolic syndrome: the Korea National Health and Nutrition Examination Survey.** *Diabetes Care*. 2005; **28**(8): 2064–2066.
[PubMed Abstract](#) | [Publisher Full Text](#)
57. Wada T, Urashima M, Fukumoto T: **Risk of metabolic syndrome persists twenty years after the cessation of smoking.** *Intern Med*. 2007; **46**(14): 1079–1082.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status: ? ✓

Version 2

Reviewer Report 07 January 2019

<https://doi.org/10.5256/f1000research.16385.r42390>

© 2019 González-Rivas J. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Juan P. González-Rivas

¹ The Andes Clinic of Cardio-Metabolic Studies, Mérida, Venezuela

² Foundation for Clinical, Public Health, and Epidemiological Research of Venezuela (FISPEVEN), Mérida, Venezuela

In this cross-sectional survey, the authors assessed the relationship between metabolic syndrome and tobacco habits and found a significant association with some MS components. Here are some comments oriented to improve the manuscript:

Introduction:

1. Paragraph 2: *"these percentages are important because"* can be erased.

Methods:

1. Clinical evaluation: the classification of the JNC7 was not used in the results, this explanation about the JNC7 criteria can be avoided.

Results:

1. No comments.

Discussion:

1. The first paragraph should resume the main findings.
2. Paragraph 3: *"on the other hand"* can be erased.
Paragraph 5: change *"according to the results of our study"* for *"In this study"*.
3. The authors should be careful in presenting cigarette use as a "protective" strategy for high blood pressure.
4. Line 13, paragraph 7: the authors should instead discuss the masked hypertension effect observed in smokers - this could explain this "paradoxical result". This effect was also reported in patients with type 2 diabetes and smokeless tobacco use in Venezuela (González-Rivas *et al.*, 2017¹).

References

1. González-Rivas J, Santiago R, Mechanick J, Nieto-Martínez R: Chimó, a Smokeless Tobacco Preparation, is Associated With a Lower Frequency of Hypertension in Subjects with Type 2

Diabetes. *International Journal of Cardiovascular Sciences*. 2017. [Publisher Full Text](#)

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

No source data required

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Epidemiology, cross-sectional surveys, cardiometabolic risk factors

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 15 May 2018

<https://doi.org/10.5256/f1000research.15858.r33887>

© 2018 Oda E. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Eiji Oda

Medical Check-up Center, Tachikawa General Hospital, Nagaoka, Japan

The authors investigated cross-sectional associations between smoking and metabolic syndrome (MS) and its components in 2,212 adults from Maracaibo city, Venezuela. They reported that, in the multivariate analysis, the presence of MS, abdominal obesity, hypertriglyceridemia and hypo-HDL cholesterolemia were positively associated with current smoking while high blood pressure was inversely associated with current smoking. There was no significant association between smoking

and hyperglycemia. Their data and study methods are fair. However, these results are not novel but well known findings.

The authors stated that cigarette smoking represents an independent risk factor for the development of MS in Conclusions. However, this conclusion is incorrect because the authors stated that, regarding the limitations of this study, its cross-sectional design makes it incapable of determining causality. After all, the authors had no data regarding the association between baseline smoking status and incidence (development) of MS which can only be obtained from longitudinal studies.

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

No source data required

Are the conclusions drawn adequately supported by the results?

No

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research