Check for updates

RESEARCH ARTICLE

Reproductive performance of asian catfish (*Hemibagrus wyckii* Bagridae), a candidate species for aquaculture [version 1; peer review: 2 approved]

Netti Aryani ¹, Indra Suharman ¹, Hafrijal Syandri ²

¹Department of Aquaculture, Faculty of Fisheries and Marine Science, Universitas Riau, Pekanbaru, 28294, Indonesia

²Department of Aquaculture, Faculty of Fisheries and Marine Science, Universitas Bung Hatta, Padang, 25133, Indonesia

V1 First published: 31 May 2018, 7:683
<https://doi.org/10.12688/f1000research.14746.1>

Latest published: 03 Sep 2018, 7:683
<https://doi.org/10.12688/f1000research.14746.2>

Abstract

Background: *Hemibagrus wyckii* Bagridae is one of the most important economic fish species that lives in the rivers and reservoir in Riau Province, Indonesia. The present study aimed to determine the reproductive performance of *H. wyckii* under culture conditions.

Methods: A total of 10 female and 10 male fish were selected, and weight, length, characteristics of egg and sperm, and hatchery performance were measured. Eggs were fertilized using the dry method. Egg weight and egg diameters were measured for 50 eggs per female. Egg size (50 eggs for each fish) was measured using an Olympus microscope (CX40). Then, saline solution was added over the eggs, followed by the addition of pooled sperm from 10 males.

Results: Average relative fecundity, egg weight and egg diameter were 2060±512 eggs/kg fish, 29.86±1.21 mg and 2.67±0.26 mm, respectively. The fertilization rate and hatching rate were 60.91±4.68% and 42.91±2.92% respectively. Sperm characteristics such as volume per fish (mL), pH, concentration (per mL), motility (%) and duration of motility (second) were 0.82±0.20, 7.15±0.12, 3.68±0.15, 72.77±1.46 and 47.5±4.84, respectively.

Conclusion: The study results and scientific observations regarding reproductive performance suggest that *H. wyckii* can be considered a new candidate species for aquaculture.

Keywords

Hemibagrus wyckii, endangered species, alternative fish species, egg quality, sperm

Open Peer Review

Approval Status

	1	2
version 2		
(revision)		
03 Sep 2018	view	
version 1		
31 May 2018	view	view

1. **Zainal Abidin Muchlisin** , Syiah Kuala University, Banda Aceh, Indonesia

2. **Rudy Agung Nugroho** , Mulawarman University, Samarinda, Indonesia

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Netti Aryani (nettiaryani@yahoo.com)

Author roles: **Aryani N:** Conceptualization, Data Curation, Formal Analysis, Methodology, Writing – Original Draft Preparation; **Suharman I:** Data Curation, Formal Analysis, Supervision, Validation; **Syandri H:** Data Curation, Project Administration, Supervision, Validation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This study was funded by a study grant (Riset Dasar Unggulan Perguruan Tinggi) from the Directorate of Research and Community Service, Ministry of Research Technology and Higher Education Republic of Indonesia (No. 311/UN.19.5.1.3/PP/2018) *The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

Copyright: © 2018 Aryani N *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. Data associated with the article are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

How to cite this article: Aryani N, Suharman I and Syandri H. **Reproductive performance of asian catfish (*Hemibagrus wyckii* Bagridae), a candidate species for aquaculture [version 1; peer review: 2 approved]** F1000Research 2018, 7:683 <https://doi.org/10.12688/f1000research.14746.1>

First published: 31 May 2018, 7:683 <https://doi.org/10.12688/f1000research.14746.1>

Introduction

In Indonesia, the fisheries sector plays an important economic role through income generation, diversification of livelihoods, supply of animal proteins, and foreign exchange earnings¹. During recent decades, in the freshwater aquaculture sector, the prioritized species for culture were *Clarias*, *Pangasius*, *Tilapia*, Common carp and Giant gourami. Wild fish species in rivers, reservoirs and lakes have not been prioritized for aquaculture operations.

In the Riau Province, there are three rivers, the Kampar Kanan, Kampar Kiri, and Siak rivers, and the Koto Panjang Reservoir. The Kampar Kanan river hosts up to 34 fish species², the Kampar Kiri river hosts up to 86 fish species³, the Ukai river, a branch of Siak River hosts up to 31 fish species⁴ and Koto Panjang Reservoir hosts up to 26 fish species⁵.

Hemibagrus wyckii (Bagridae) is one of the most important economic fish species that lives in the rivers and reservoir in Riau Province. *H. wyckii* (its local name is *geso*) is a carnivorous freshwater finfish native to Indonesia^{2,6}. *H. wyckii* has been categorized as of “least concern” by the International Union for Conservation of Nature (IUCN). However, *H. wyckii* in the Kampar Kanan river was categorized as a vulnerable to endangered species^{7,8}.

Due to the endangered population of *H. wyckii*, it is necessary to domesticate this species as an aquaculture candidate in the future. Therefore, the present study aimed to determine the reproductive performance of *H. wyckii* as a potential species under culture conditions to provide preliminary scientific information and evaluation.

Methods

Ethical considerations

While *H. wyckii* is classified as vulnerable to endangered in the Kampar Kanan river, the Government of the Republic of Indonesia does not require licences to be obtained to capture and rear this species, hence no licences are applicable to this study. No animals suffered as a result of the activities of this study. *H. wyckii* was transported to the pond farm for rearing, injection, ovulation, stripping and sperm production. In the end of the experiment the *H. wyckii* still in good condition until return back to the pond.

Rearing and selection of breeders

Broodfish of *H. wyckii* were collected from upstream areas of the Kampar Kanan river in the Kouk village (0° 19' 23.44" N and 100° 56' 40.05" E), Kampar Regency, Riau Province. The broodfish kept in oxygenated polythene bag and transported by truck to Sadarlis Green Catfish Farm, Sungai Paku, Kampar Regency, Indonesia. Then, the broodstock of *H. wyckii* had been adapted and grown to maturation under the farm conditions. Prior to stocking, female and male fish were weighed using balance scale (OHAUS model CT 6000-USA), and their lengths were measured using a meter ruler with 0.01mm accuracy. During the grow-out period, fish were cultured in two ponds (4 x 4 x 3 m) separated by sex. The depth of water

in each pond was 2.0 m. The inlet of the pond water come from Sungai Paku reservoir at a rate of 2.0 m³ per sec. The broodfish fed with freshwater seashell meat (*Pilsbryconcha exilis*; Unionidae) collected from local fisherman near to Sungai Paku Reservoir. The seashell meat was kept at cold box with the temperature 5°C prior given to broodfish. According to Aryani *et al.*⁶ the proximate composition (% wet weight base) of the seashell meat was 89.37% moisture, 7.08% crude protein, 0.82% fat, 0.29% crude ash and 2.44% carbohydrate. The total of seashell meat given to broodfish everyday were 2.500g per pond (equivalent with 9% body weight of population). The feeding time at 17:00 pm due to carnivorous broodfish. The average weight and length from ten (10) of the female broodfish were 2,669.4±486.917 g and 62.84±8.20 cm, respectively. Meanwhile, the ten (10) of male broodfish were 1,769.1±401.10 g and 54.52±7.17 cm, respectively.

The fish were checked monthly for ovulation and semen production from mid November, 2017 onwards. The broodfish were captured with a gillnet formed into a net bag with the appropriate mesh size and anesthetized orally with Tricaine methanesulfonate (MS-222, ethyl 4-aminobenzoate methanesulfonate 98%, Sigma Aldrich Co, USA, MO; 50 mg L⁻¹), based on the dosage used for *Solea senegalensis*⁹. Oocyte maturation was assessed for each individual. The fish were returned to their pond after evaluation, and no mortality occurred. Fish were fasted 48 h prior to the evaluation. Oocytes sampled *in vivo* were taken from females using the method described by Nowosad *et al.*¹⁰ and were placed in Serra's solution (6:3:1 of 70% ethanol, 40% formaldehyde and 99.5% acetic acid) for clarification of the cytoplasm. After 5 min, the position of the stage IV oocyte nucleus was determined using criteria by Krejszeff *et al.*¹¹ and was classified as germinal vesicle in the periphery or germinal vesicle breakdown (GVBD).

Female reproductive performance

H. wyckii was categorized as endangered species and difficult to obtain in Kampar Kanan river. A total of 10 mature have been eligible for the experiment. 10 mature females that had oocyte stage IV were sampled from broodstock from 3rd week of February to March 2018 in the same farm, and live weights (FeW) and total lengths (FeL) were measured after anesthetization with 0.50 mg L⁻¹ MS-222⁹. For ovulation, each female broodfish received two injections of GnRH analogs with a dopamine antagonist (Ovaprim) (manufactured for Syndel Laboratories Ltd, 2595 McCullough Rd. Nanaimo, B.C.V9S 4M9 Canada) applied intraperitoneally under the left pectoral fin. The first injection was 0.2 mL kg BW⁻¹ and the second was 0.6 mL kg BW⁻¹ (total 0.8 mL kg BW⁻¹) at 12 h intervals. These dosages refer to the previous doses for ovulation of *H. wyckii*⁶. At 18 to 20 h after injection, eggs were stripped into a plastic vessel. Eggs were fertilized using the “dry method” as described by Dabrowski *et al.*¹² Egg weights of each female were by determined weighing 50 eggs to the nearest 0.01 g, and egg diameters were measured to the nearest 0.01 mm. Egg size (50 eggs for each fish) was measured using an Olympus microscope. Then, a balanced saline solution (7.5 g of NaCl, 0.2 g of KCl, 0.2 g of CaCl₂·2H₂O, and 0.02 g of NaHCO₃, in 1000 mL distilled water)

was added over the eggs¹³, followed by an addition of pooled sperm from 10 males. The eggs were then gently mixed for fertilization and left for three minutes. The fertilized eggs were rinsed several times with incubation water to remove sperm remnants as well as dead and broken eggs. The eggs were left for an additional 25 minutes to facilitate egg hardening by water absorption and disinfected with 100 ppm iodine for 10 minutes. Then, eggs were transferred to incubation trays placed in a vertical hatching system. The water flow rate to each vertical incubator was 3 L min⁻¹. Fifty eggs were randomly sampled at 15 h after fertilization to determine the fertilization rate (FR). The hatching rate (HR) was determined by counting all hatched fry.

Determination of sperm quality

Males were stimulated with a half-dose of the same hormonal preparations used to stimulate the females. Semen samples were obtained from 10 fish randomly selected from the farm. The male fish were anesthetized with 50 mg L⁻¹ of MS-222. The doses of the anaesthetic agents were prepared a few minutes before each experiment based on the methods of Weber *et al.*⁹, and then, weights (MaW) and total lengths (MaL) were measured. Special care was taken to avoid any contamination of semen with urine, feces, mucus and water. Semen samples were collected using plastic syringes in 3 mL aliquots, then placed in an insulated ice-cooled container, transported to the laboratory and analyzed within 2 h.

The sperm evaluation included gross (visual) and microscopic examination (as reviewed by Rurangwa *et al.*¹⁴, and Cabrera *et al.*¹⁵). The gross examination was based on visual and physical observation of parameters such as the semen volume by collecting the semen in a graduated cylinder and determining the level in milliliters. The microscopic examination was carried out using an Olympus model CX40, with magnification between X 10 and X 25 to determine other parameters such as: motility (duration and percentage). Motility (MO) percentage and duration

were determined by observing water activated semen placed on a glass slide under a microscope. Motile sperm were observed and expressed as a percent of non-moving sperm. Motility duration (DMO) was determined as the period between movements of the sperm to cessation of any progressive movement expressed in seconds. Sperm concentration (SC) was measured under a microscope using Neubeur's hemocytometer and calculated as the number of sperm ml⁻¹¹⁶. Semen pH was determined with a hand pH meter (HI8424 Hanna Instruments, USA).

Water quality

The water temperature of the farm was measured with a thermometer (Celsius scale), and water samples were collected to determine the dissolved oxygen (DO) concentrations. An oxygen meter (YSI model 52, Yellow Spring Instrument Co., Yellow Springs, OH, USA) was used *in situ*, and pH values were determined with a pH meter (Digital Mini-pH Meter, 0-14PH, IQ Scientific, Chemo-science Thailand Co., Ltd, Thailand). Alkalinity and hardness levels of the water were measured in each replicate according to standard procedures¹⁷. The water quality parameters were measured once per month.

Statistical analysis

Results were given as the means $\pm$ SD. Simple linear regression analyses were performed using SPSS software (version 16.0 for Windows; SPSS Inc., Chicago, IL). The standard deviation of each parameter was determined. For linear regression analysis, significant correlations were considered at $p < 0.05$.

Results

Descriptive measurements and the reproductive performance of female *H. wyckii* are presented in Table 1. Fifty percent of eggs hatched at 60 h (29–30 °C water temperature). The fertilization rate varied between 53.2 and 68.3%, whereas the hatching rate was between 39.5 and 48.3%.

Table 1. Female size, egg characteristics and hatchery performance of *Hemibagrus wyckii* (Mean $\pm$ SD).

	Variables	Range (Min–Max)
Fish weight (g)	2669.4 $\pm$ 486.917	2030 – 3296
Fish length (cm)	62.84 $\pm$ 8.20	57.5 – 78.2
Absolute fecundity (egg/fish)	5809 $\pm$ 2275	4125 – 9958
Relative fecundity (egg/kg fish)	2060 $\pm$ 512	1400 – 3000
Egg diameter (mm)	2.67 $\pm$ 0.26	2.10 – 2.86
Hardened egg diameter (mm)	2.94 $\pm$ 0.30	2.45 – 2.97
Egg diameter increase (%)	18.27 $\pm$ 10.43	4.64 – 32.36
Egg weight (mg)	29.86 $\pm$ 1.21	24.4 – 31.8
Hardened egg weight (mg)	30.08 $\pm$ 1.20	28.63 – 32.01
Egg weight increase (%)	21.59 $\pm$ 1.08	20.02 – 23.5
Fertilization rate (%)	60.91 $\pm$ 4.68	53.2 – 68.3
Hatching rate (%)	42.91 $\pm$ 2.92	39.5 – 48.3
Hatching weight (mg)	30.08 $\pm$ 1.20	28.63 – 32.01

Characteristics of male fish and sperm samples are presented in Table 2. The average live weight of the males is $1,769.1 \pm 401.1$ g. Male *H. wyckii* are found to be slightly smaller than females. In the genital maturation stage, the papilla is not prominent for all male fish as the second sexual characteristic of the other *Hemibagrus*.

According to the analysis of the linear relationship (r^2) between variables of *H. wyckii* females shown in Table 3, there was a strong linear relationship between AF and FeW, AF and RF, HEW and EW, HW and EW, HW and HEW, and HR and FR. In contrast, the analysis of the linear relationship (r^2) between variables of *H. wyckii* males shown in Table 4, show a strong linear relationship between MaW and MaL, MaW and GW, and MaL and GW.

The temperature of the pond water ranged from 28°C to 29°C, oxygen ranged from 6.5 mg L⁻¹ to 6.7 mg L⁻¹, pH ranged from 6.5 to 6.8, alkalinity ranged from 42.97 mg L⁻¹ to 57.33 mg L⁻¹ and hardness ranged from 104.83 mg L⁻¹ to 110.51 mg L⁻¹.

Dataset 1. Data of female size, egg characteristic and hatchery performance of *Hemibagrus wyckii*

<http://dx.doi.org/10.5256/f1000research.14746.d204328>

Discussion

In our study, the spawning period of *H. wyckii* started at the 3rd week of February and continued until the final examination in March. However, our monthly observations of *H. wyckii* captured by fishermen in the Kampar Kanan river found fish

Table 2. Male size and sperm characteristics of *Hemibagrus wyckii* (Mean $\pm$ SD).

	Variables	Range (Min–Max)
Fish weight (g)	1769.1 $\pm$ 401.10	1105 – 3127
Fish length (cm)	54.52 $\pm$ 7.17	43.7 – 65.2
Gonad weight (g)	29.28 $\pm$ 3.42	24.96 – 54.96
Gonadosomatic index (%)	1.95 $\pm$ 0.71	1.02 – 2.83
Semen volume (mL)	0.82 $\pm$ 0.20	0.60 – 1.20
Semen pH	7.15 $\pm$ 0.12	7.30 – 7.70
Sperm concentration (10 ⁹ /mL)	3.68 $\pm$ 0.15	3.50 – 3.90
Motility (%)	72.77 $\pm$ 1.46	70.2 – 75.50
Duration of motility (sec)	47.5 $\pm$ 4.84	40.0 – 54.0

Table 3. Correlations of variables (r^2) for *Hemibagrus wyckii* females.

	FeW	FeL	AF	RF	EW	HEW	EWI	ED	HED	HDI	FR	HR	HW
FeL	0.548												
AF	0.812	0.687											
RF	0.523	0.582	0.899										
EW	0.076	0.058	0.127	0.151									
HEW	0.074	0.059	0.126	0.152	0.999								
EWI	0.088	0.029	0.056	0.054	0.151	0.148							
ED	0.395	0.381	0.504	0.486	0.464	0.467	0.051						
HED	0.105	0.315	0.196	0.303	0.572	0.579	0.189	0.696					
HDI	0.229	0.598	0.279	0.328	0.220	0.224	0.007	0.234	0.353				
FR	0.087	0.018	0.030	0.003	0.021	0.024	0.001	0.120	0.366	0.091			
HR	0.099	0.026	0.079	0.032	0.043	0.047	0.007	0.133	0.366	0.008	0.915		
HW	0.052	0.084	0.112	0.151	0.974	0.976	0.148	0.467	0.579	0.244	0.024	0.031	
GI	0.584	0.665	0.712	0.632	0.471	0.468	0.139	0.677	0.438	0.406	0.007	0.020	0.468

Statistically important at $r^2 > 0.500$ (underlined)

Few: Female fish weight, FeL: Female fish length, AF: Absolute fecundity, RF: Relative fecundity, EW: Egg weight, HEW: Hardened egg weight, EWI: Eggs weight increase, ED: Egg diameter, HED: Hardened egg diameter, HDI: Hardened egg diameter increase, FR: Fertilization rate, HR: Hatching rate, HW: Hatching weight, GI: Gonadosomatic index.

at the gonadal I, II, III and IV stages of development (stage scale by Krejszeff *et al.*¹¹), showing a spawning type shows of a partial spawner. Spawning in the wild occurs at the start and end of the rainy season, or this species could spawn twice per year. The spawning type of *H. wyckii* is the same as that of *Hemibagrus nemurus*¹⁸. The duration from fertilization to a 50% hatching rate was 60 h with 29°C to 30°C water temperature. The fertilization rate and hatching rate of various *Hemibagrus* species are reported in Table 5. When findings are compared with other *Hemibagrus* species, the duration of hatching is longer in *H. wyckii*. In other words, *H. wyckii* has species specific characteristics the regarding hatching rate, because the egg diameter was bigger than other *Hemibagrus* eggs (Table 5).

AFs of *H. wyckii* were between 4125 and 9958 eggs/fish and RFs were between 1400 and 3000 eggs kg⁻¹. Egg production per kg fish (RF) is thought to be more informative than absolute fecundity. RF values of *H. wyckii* were lower compared with those of *H. nemurus*^{19–21}. In our study, there was no strong

linear relationship between RF and fish size. Meanwhile, there was a strong linear relationship between AF and fish size (Table 3).

EDs and EWs obtained here ranged between 2.10 and 2.86 mm and 24.4 and 31.8 mg respectively, consistent with those reported by Aryani *et al.*⁶ At the end of the hardening process, the increases in egg weight and diameter were 29.86 and 14.9% respectively. In the present study, there were strong linear relationships between EW and HEW, and between EW and HW (Table 3). Lahnsteiner and Patzner²² state that egg weight increases after the hardening process and is linearly correlated with the viability of eggs in Rainbow trout, but not in Alakir trout²³.

In this study, the FRs of *H. wyckii* were higher than those of *H. wyckii* from research conducted by Aryani *et al.*⁶. This suggests that we have improved the method for the fertilization process of sperm and eggs so that hatching rate can increase. Nevertheless, the fertilization rate of *H. wyckii* was lower than

Table 4. Correlations of variables (r^2) for *Hemibagrus wyckii* males.

	MaW	MaL	GW	GI	MV	pH	SC	MO
MaL	<u>0.810</u>							
GW	<u>0.773</u>	<u>0.727</u>						
GI	0.449	0.496	0.311					
SV	0.049	0.002	0.129	0.066				
pH	0.019	0.047	0.029	0.392	0.204			
SC	0.193	0.110	0.267	0.037	0.001	0.038		
MO	0.184	0.066	0.098	0.004	0.003	0.066	0.133	
DMO	0.121	0.077	0.035	0.015	0.016	0.191	0.139	0.499

Statistically important at $r^2 > 0.500$ (underlined)

MaW: Male fish weight, MaL: Male fish length, GW: Gonada weight, GI: Gonadosomatic index, MV: Semen volume, SC: Sperm concentration, MO: Motility, DMO: Duration of motility.

Table 5. Summary of the egg diameter, fertilization and hatching rate of *H. wyckii*, with respect to those reported for *H. nemurus* and *H. dibrugarensis*.

Species	Eggs diameter (mm)	Fertilization rate (%)	Hatching rate (%)	Reference
<i>H. wyckii</i>	2.10 – 2.86	53.2 – 68.3	39.5 – 58.3	This study
<i>H. wyckii</i>	2.79 – 2.85	45.8 – 60.2	37.33 – 55.1	6
<i>H. nemurus</i>	1.13 – 1.22	74.16 – 77.83	64.16 – 67.83	19
<i>H. nemurus</i>	1.07 – 1.21	70.50 – 81.4	63.5 – 71.4	20
<i>H. dibrugarensis</i>		34.83 – 77.54	20.61 – 74.32	24
<i>H. nemurus</i>	-	93.5 – 94.5	56.91 – 78.3	25
<i>H. nemurus</i>	1.57 – 1.75	81.57 – 98.54	81.3 – 96.72	21
<i>H. nemurus</i>		39.5 – 79.0	32.5 – 69.7	23

that of *H.nemurus*^{17,24,25} (Table 5). When HR values are compared with other *Hemibagrus* species, HR levels are lower than those of Aryani and Suharman¹⁴, Adebiyi *et al.*²⁵ and Suhenda *et al.*²¹ but higher than those of Aryani *et al.*⁶. There were strong positive correlations with FR and HR ($r^2 = 0.91$) (Table 3). Meanwhile, FR and HR were not positively correlated with ED (Table 3).

Gonadotropins (GTHs), follicle-stimulating hormone (FSH), luteinizing hormone (LH), and sex steroids are the key regulators of reproduction^{26–28}. Moreover, numerous circulating endocrine and locally acting paracrine and autocrine factors regulate the various stages of oocyte development and maturation^{29,30}. Other factors that significantly affect fish eggs are genetic, environmental and stress factors^{31–33}. However, there is no information about the effects of such kinds of factors on the embryonic development of *H.wyckii*. There are also further information requirements concerning hatchery management of *H.wyckii* such as feed levels and feed type⁶. During the adaptation period in the present instance, *H. wyckii* were fed with meat freshwater seashell (*Pilsbryoconcha exilis*; Unionidae), the local name of which is “lokan” which may not be fully suitable for this species, even though this species is carnivorous. The requirement of a balanced feed that meets the nutritional requirements of species being cultured^{34–36} and application of a proper feeding program during the ovarian development³⁷ have been emphasized. Suhenda *et al.*²¹ reported that diet should be offered to *H. nemurus* broodstock at an 8% lipid level in fish feed with 35% crude protein for 4 months to obtain high quality gametes. Meanwhile, Aryani and Suharman²⁰ suggest that a minimum of 32% crude protein should be included in the diet of female *H.nemurus* broodstock. Additionally, implantation of 17 β -estradiol has also managed to improve the reproductive performance of *H.nemurus*¹⁹. These issues in *H.wyckii* are yet to be elucidated. Therefore, such research is very important for *H.wyckii* in the future.

Average SVs determined in *H. wyckii* are higher than those in *H.nemurus* (0.10 to 0.35 mL)¹⁸ and *Clarias gariepinus*³⁸, but lower than those reported for *Ictalurus punctatus*³⁹. It appears that the semen volume in other species has a positive relationship with sperm concentration, including in *Ictalurus punctatus*³⁹. Meanwhile, in fish farms and hatcheries, the biotic and abiotic factors that affect sperm quality are diverse and dependent on complex interactions between genetic, physiological and environmental factors¹⁴. On the other hand, improvements in broodstock nutrition and feeding greatly improve gamete quality and larvae production⁴⁰.

Semen pH values of *H.wyckii* are consistent with those of other species, including *Barbus grypus*⁴¹ and *Carrasius gibelio*⁴². The sperm motility of *H.wyckii* between 70.2 and 75.50%, and the

duration of motility was between 40.0 and 54.0 sec, results that are consistent with *Ictalurus punctatus*³⁹. According to Effer *et al.*⁴³ the duration of sperm motility in fish depends on the temperature of the activation medium. Sperm of *H.wyckii* had an effective fertilizing ability according to the correlation analysis, which did not detect any significant relationship between FR and sperm parameters. However, there was a positive relationship between MO and DMO ($r^2 = 0.49$). Sperm morphology, density, volume, motility and fertilizing capacity, as well as the composition and osmolality of the seminal plasma are parameters commonly measured to assess sperm quality in fish^{14,44}. In this study, we did not investigate the ionic composition of the semen, but this phenomenon could be related to the ionic composition of semen, which has a significant influence on motility and duration of motility^{45–47}.

Conclusion

In conclusion, the reproductive performance of *H.wyckii* under culture conditions is within the range of available data for other *Hemibagrus* species. Considering that successful larvae production (and potential juvenile growth) was possible using the same methods as in Bagridae, *H.wyckii* can be an alternative species for aquaculture. However, further studies are clearly required to determine several aspects of this fish under culture conditions.

Data availability

Dataset 1: Data of female size, egg characteristic and hatchery performance of *Hemibagrus wyckii* [10.5256/f1000research.14746.d204328](https://doi.org/10.5256/f1000research.14746.d204328)⁴⁸

Competing interests

The authors have no conflicts of interest to declare.

Grant information

This study was funded by a study grant (Riset Dasar Unggulan Perguruan Tinggi) from the Directorate of Research and Community Service, Ministry of Research Technology and Higher Education Republic of Indonesia (No. 311/UN.19.5.1.3/PP/2018).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgements

The authors thank the Ministry of Research Technology and Higher Education Republic of Indonesia for supporting this study through the competitive grants schema (*Riset Dasar Unggulan Perguruan Tinggi*). Appreciation also goes to all of fisherman and students who helped the author during experiments in the farm.

References

1. Nhuong T, U-Primo R, Chin YC, *et al.*: Indonesian aquaculture futures: An analysis of fish supply and demand in Indonesia to 2030 and role of

aquaculture using the AsiaFish model. *Mar Policy*. 2017; **79**: 25–32. [Publisher Full Text](#)

2. Aryani N: **Native species in Kampar Kanan river, Riau Province Indonesia.** *Int J Fish Aquatic Stud.* 2015; 2(5): 213–217.
[Reference Source](#)
3. Simanjuntak CPH, Rahardjo MF, Sukimin S: **Ichthyofauna in floodplain of Kampar Kiri river.** *Jurnal Ikhtologi Indonesia.* 2006; 6(2): 99–109.
[Reference Source](#)
4. Kaidir PP: **The diversity of fish species in Ukai stream, branch of Siak River Riau.** *Berkala Perikanan Terubuk.* 2011; 1(39): 24–32.
5. Warsa A, Krismono ASN, Nurfiarini A, *et al.*: **The capture fishery resources in Koto Panjang Reservoir.** *Bawal.* 2009; 2(3): 93–97.
[Reference Source](#)
6. Aryani N, Suharman I, Ainul M, *et al.*: **Reproductive performance of Asian catfish (*Hemibagrus wyckii*, Bleeker, 1858)-Preliminary study.** *Pakistan J Nutr.* 2017; 16(7): 550–556.
[Publisher Full Text](#)
7. Aryani N, Suharman I, Sabrina H: **Length-weight relationship and condition factor of the critically endangered fish of Geso, *Hemibagrus wyckii* (Bleeker, 1858) Bagridae from Kampar Kanan river, Indonesia.** *J Entomol Zool Stud.* 2016; 4(2): 119–122.
[Reference Source](#)
8. Fithra RY, Siregar YI: **Fishes biodiversity of Kampar river an inventory from Kampar Kanan river.** *J Environ Sci.* 2010; 2(4): 139–147.
[Reference Source](#)
9. Weber RA, Peleteiro JB, García Martín LO, *et al.*: **The efficacy of 2-phenoxyethanol, metomidate, clove oil and MS-222 as anaesthetic agents in the Senegalese sole (*Solea senegalensis* Kaup 1858).** *Aquaculture.* 2009; 288(1–2): 147–150.
[Publisher Full Text](#)
10. Nowosad J, Targońska K, Chwaluczyk R, *et al.*: **Effect of temperature on the effectiveness of artificial reproduction of dace [Cyprinidae (*Leuciscus leuciscus* (L.))] under laboratory and field conditions.** *J Therm Biol.* 2014; 45: 62–68.
[PubMed Abstract](#) | [Publisher Full Text](#)
11. Krejszef S, Katarzyna T, Daniel Z, *et al.*: **Domestication affect spawning of the Ide (*Leuciscus idus*)—preliminary study.** *Aquaculture.* 2009; 295(1–2): 145–147.
[Publisher Full Text](#)
12. Dabrowski K, Ciereszko A, Ramseyer L, *et al.*: **Effects of hormonal treatment on induced spermiation and ovulation in the yellow perch (*Perca flavescens*).** *Aquaculture.* 1994; 120(1–2): 171–180.
[Publisher Full Text](#)
13. Kobayashi T, Fushiki S, Ueno K: **Improvement of sperm motility of sex-reversed male rainbow trout, *Oncorhynchus mykiss*, by incubation in high-pH artificial seminal plasma.** *Environ Biol Fish.* 2004; 69(1–4): 419–425.
[Publisher Full Text](#)
14. Rurangwa E, Kime DE, Ollevier F, *et al.*: **The measurement of sperm motility and factors affecting sperm quality in cultured fish.** *Aquaculture.* 2004; 234(1–4): 1–28.
[Publisher Full Text](#)
15. Cabrita E, Martínez-Páramo S, Gavaia PJ, *et al.*: **Factors enhancing fish sperm quality and emerging tools for sperm analysis.** *Aquaculture.* 2014; 432: 389–401.
[Publisher Full Text](#)
16. Mylonas CC, Duncans NJ, Asturiano JF: **Hormonal manipulations for the enhancement of sperm production in cultured fish and evaluation of sperm quality.** *Aquaculture.* 2017; 472: 21–44.
[Publisher Full Text](#)
17. Rice EW, Baird RB, Eaton AD, *et al.*: **Standard methods for the examination of water and wastewater, 22nd ed.** American Public Health Association, American Water Works Association, Water Environment Federation. 2012.
[Reference Source](#)
18. Sularto RRS, Dewi PS, Khasani I: **The influence of 17 α -methyltestosterone implantation on gonada maturation and fertility of male green catfish (*Mystus nemurus*).** *J Ris Akuakultur.* 2010; 1(5): 53–57.
[Reference Source](#)
19. Aryani N, Suharman I: **Effects of 17 β -estradiol on the reproduction of green catfish (*Hemibagrus nemurus*, Bagridae).** *J Fish Aquacult.* 2014; 5(1): 163–166.
[Reference Source](#)
20. Aryani N, Suharman I: **Effect of dietary protein level on the reproductive performance of female of green catfish (*Hemibagrus nemurus* Bagridae).** *Aquacult Res Dev.* 2015; 6: 11.
[Publisher Full Text](#)
21. Suhenda N, Samsudin R, Kristanto AH: **The role of dietary lipid level in embryo development, hatching rate and survival rate of green catfish (*Mystus nemurus*) larvae.** *J Ris Akuakultur.* 2009; 2(4): 201–211.
[Reference Source](#)
22. Lahnsteiner F, Patzner RA: **Rainbow trout egg quality determination by the relative weight increase during hardening: a practical standardization.** *J Appl Ichthyol.* 2002; 18: 24–26.
[Publisher Full Text](#)
23. Kanyilaz M: **Reproductive performance of a newly described Salmonid fish, Alakir Trout (*Salmo Kottelati*), a candidate species for aquaculture.** *Pak J Zool.* 2016; 48(1): 83–89.
[Reference Source](#)
24. Bailung B, Biswas SP: **Successful induced breeding of a Bagrid Catfish, *Mystus dibrugarensis* in captive condition.** *J Aquacult Res Dev.* 2014; 5: 7.
[Publisher Full Text](#)
25. Adebisi FA, Siraj SS, Harmin SA, *et al.*: **Induced spawning of a river catfish *Hemibagrus nemurus* (Valenciennes, 1840).** *Pertanika J Trop Agric Sci.* 2013; 36(1): 71–78.
[Reference Source](#)
26. García Ayala A, Villaplana M, García Hernández MP, *et al.*: **FSH-, LH-, and TSH-expressing cells during development of *Sparus aurata* L. (Teleostei). An immunocytochemical study.** *Gen Comp Endocrinol.* 2003; 134(1): 72–79.
[PubMed Abstract](#) | [Publisher Full Text](#)
27. Aizen J, Kobayashi M, Selicharova I, *et al.*: **Steroidogenic response of carp ovaries to piscine FSH and LH depends on the reproductive phase.** *Gen Comp Endocrinol.* 2012; 178(1): 28–36.
[PubMed Abstract](#) | [Publisher Full Text](#)
28. Nyuji M, Kazeto Y, Izumida D, *et al.*: **Greater amberjack FSH, LH, and their receptors: Plasma and mRNA profiles during ovarian development.** *Gen Comp Endocrinol.* 2016; 225: 224–234.
[PubMed Abstract](#) | [Publisher Full Text](#)
29. Lubzens E, Young G, Bobe J, *et al.*: **Oogenesis in teleosts: how eggs are formed.** *Gen Comp Endocrinol.* 2010; 165(3): 367–389.
[PubMed Abstract](#) | [Publisher Full Text](#)
30. Lubzens E, Bobe J, Young G, *et al.*: **Maternal investment in fish oocytes and eggs: The molecular cargo and its contributions to fertility and early development.** *Aquaculture.* 2017; 472: 107–143.
[Publisher Full Text](#)
31. Grande M, Andersen S: **Effect of two temperature regimes from a deep and a surface water release on early development of salmonids.** *Regulated Rivers: Research & Management.* 1990; 5(4): 355–360.
[Publisher Full Text](#)
32. Webb MA, Doroshov SI: **Importance of environmental endocrinology in fisheries management and aquaculture of sturgeons.** *Gen Comp Endocrinol.* 2011; 170(2): 313–321.
[PubMed Abstract](#) | [Publisher Full Text](#)
33. Kime DE, Nash JP: **Gamete viability as an indicator of reproductive endocrine disruption in fish.** *Sci Total Environ.* 1999; 233(1–3): 123–129.
[Publisher Full Text](#)
34. Chong ASC, Ishak SD, Osman Z, *et al.*: **Effect of dietary protein level on the reproductive performance of female swordtails *Xiphophorus helleri* (Poeciliidae).** *Aquaculture.* 2004; 234(1–4): 381–392.
[Publisher Full Text](#)
35. Coldebella IJ, Neto JR, Mallmann CA, *et al.*: **The effects of different protein levels in the diet on reproductive indexes of *Rhamdia quelen* females.** *Aquaculture.* 2011; 312(1–4): 137–144.
[Publisher Full Text](#)
36. Sink TD, Lochmann RT, Pohlenz C, *et al.*: **Effects of dietary protein source and protein–lipid source interaction on channel catfish (*Ictalurus punctatus*) egg biochemical composition, egg production and quality, and fry hatching percentage and performance.** *Aquaculture.* 2010; 298(3–4): 251–259.
[Publisher Full Text](#)
37. Bromage N, Jones J, Randall C, *et al.*: **Broodstock management, fecundity, egg quality and the timing of egg production in the rainbow trout (*Oncorhynchus mykiss*).** *Aquaculture.* 1992; 100(1–3): 141–166.
[Publisher Full Text](#)
38. Gbemisola OB, Adebayo OT: **Sperm quality and reproductive performance of male *Clarias gariepinus* induced with synthetic hormones (Ovatide and Ovaprim).** *International Journal of Fisheries and Aquaculture.* 2014; 6(1): 9–15.
[Publisher Full Text](#)
39. Jaspers EJ, Avault JE Jr, Roussel JD: **Testicular and Spermatozoal Characteristics of Channel Catfish, *Ictalurus punctatus*, outside the Spawning Season.** *Trans Am Fish Soc.* 1978; 107(2): 309–315.
[Publisher Full Text](#)
40. Effer B, Figueroa E, Augsburg A, *et al.*: **Sperm biology of *Merluccius australis*: Sperm structure, semen characteristics and effects of pH, temperature and osmolality on sperm motility.** *Aquaculture.* 2013; 408–409: 147–151.
[Publisher Full Text](#)
41. Izquierdo MS, Fernandez-Palacios H, Tacon AGJ: **Effect of broodstock nutrition on reproductive performance of fish.** *Aquaculture.* 2001; 197(1–4): 25–42.
[Publisher Full Text](#)
42. Khodadadi M, Arab A, Jaferian A: **A Preliminary study on sperm morphology, motility and composition of seminal plasma of Shirbot, *Barbus grypus*.** *Turk J Fish Aquat Sci.* 2016; 16(4): 947–951.
[Publisher Full Text](#)
43. Taati MM, Mehrad B, Shabani A, *et al.*: **Correlation between chemical composition of seminal plasma and sperm motility characteristics of Prussian carp (*Carassius gibelio*).** *AACL Bioflux.* 2010; 3(3): 233–237.
[Reference Source](#)
44. Butts IA, Litvak MK, Tripple EA: **Seasonal variations in seminal plasma and sperm characteristics of wild-caught and cultivated Atlantic cod, *Gadus morhua*.** *Theriogenology.* 2010; 73(7): 873–885.
[PubMed Abstract](#) | [Publisher Full Text](#)

45. Dadras H, Sampels S, Golpuor A, *et al.*: **Analysis of common carp *Cyprinus carpio* sperm motility and lipid composition using different *in vitro* temperatures.** *Anim Reprod Sci.* 2017; **180**: 37–43.
[PubMed Abstract](#) | [Publisher Full Text](#)
46. Zilli L, Schiavone R, Vilella S: **Role of protein phosphorylation/dephosphorylation in fish sperm motility activation: State of the art and perspectives.** *Aquaculture.* 2017; **472**: 73–80.
[Publisher Full Text](#)
47. Barman AS, Kumar P, Mariahabib, *et al.*: **Role of nitric oxide in motility and fertilizing ability of sperm of *Heteropneustes fossilis* (Bloch.).** *Anim Reprod Sci.* 2013; **137**(1–2): 119–127.
[PubMed Abstract](#) | [Publisher Full Text](#)
48. Aryani N, Suharman I, Syandri H: **Dataset 1 in: Reproductive performance of asian catfish (*Hemibagrus wyckii* Bagridae), a candidate species for aquaculture.** *F1000Research.* 2018.
[Data Source](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 22 August 2018

<https://doi.org/10.5256/f1000research.16043.r37278>

© 2018 Nugroho R. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Rudy Agung Nugroho

Department of Biology, Mulawarman University, Samarinda, Indonesia

Title

The title should be revised: delete "a candidate species for aquaculture". It needs more parameters (Growth, feed aspect, etc.) to convey that this species can be a candidate for aquaculture.

Abstract

In methods: the author stated that "Egg size (50 eggs for each fish) was measured using an Olympus microscope (CX40)". This statement should be revised, since the microscope CX40 itself cannot be used to measure the egg size. It must be a micrometer within a microscope that has been used to measure the egg size, or the author has another technique to measure the egg size. This method should be also clearly stated in the methods section.

Introduction

The introduction should be extended by providing more information about Asian catfish and their biological, physiological, and ecological aspects.

Methods

Ethical consideration: Please revise, it is seems there is a typo:

"[...]the Government of the Republic of Indonesia does not require licences to be obtain to capture and rear this species, hense no licences are applicable to this study." This sentence should be rewritten: "[...] the Government of the Republic of Indonesia does not require licenses to obtain, capture and rear this species. Hence, no licenses are applicable to this study."

"In the end of the experiment the *H. wyckii* still in good condition until return back to the pond." The word pond should be changed to river, because the fish is returned back to the original (River not pond).

Rearing and selection of breeders: "Then, the broodstock of *H. wyckii* had been adapted and

grown to maturation under the farm conditions". The word grown should be changed to reared, because this experiment is not about growth.

Water quality: "Alkalinity and hardness levels of the water were measured in each replicate according to standard procedures". It is not clear what "each replicate" is referring to. Does it mean that the measurement was done in a few replications? Why only alkalinity and hardness?

Results

Results are well presented.

Discussion

The discussion can be extended by explaining the relationship between the results of water quality and the biological reproduction aspect, since the purpose of this study is under culture condition.

Conclusion

Needs slight revision based on all aspects of the findings.

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Fish physiology, Nutrition

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 11 June 2018

<https://doi.org/10.5256/f1000research.16043.r34564>

© 2018 Muchlisin Z. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Zainal Abidin Muchlisin

Faculty of Marine and Fisheries, Syiah Kuala University, Banda Aceh, Indonesia

Title:

1. I think the title should be revised as "Reproductive biology and breeding of the Asian catfish (*Hemibagrus wyckii*) after domestication process.
2. In the title: (*Hemibagris wyckii* Bagridae) is missed understanding, the correct written should be *Hemibagrus wyckii*: Bagridae), because Bagridae is a Family level, or you can write in as (*Hemigarus wyckii* Bleeker, 1858)

Abstract:

Background: I think "most" should be deleted because sound so emotional or hyperbolic, just say ".... is one of the important economic fish species...."

Methods: Please mention where the brood fish is come from?

Conclusion: The conclusions is out of the context. You have to make conclusion just based on the data (findings), forget for candidate species for aquaculture, because, there was no detail discussion about this issue, so just focus on your present findings.

Keywords: I suggest don't use similar words which already exist in the title.

Introduction:

I think the state of the arts about the asian catfish is still shallow or unclear, so please provide more information about the previous studies on this species related to bio ecology, feeding, aquaculture etc.

Please add some information what is the advantages of the species compared to other freshwater species?

Methods:

Please cite these references to enhance your methods:

- Muchlisin *et al.*, 2010 ¹
- Muchlisin *et al.*, 2011 ²

Results:

The results have been explained in several short paragraphs, please extend the paragraph by combining paragraph 1, 2 in one paragraph and paragraph 3 and 4 in one paragraph.

Discussion:

There was no comprehensive discussion to justify why this species is potential for aquaculture. The justification is not only based on biological aspects but also on economic consideration (see Muchlisin, 2013³).

Unfortunately, there was no economic evaluation performed. Therefore, I suggest to focus only on the finding as already presented.

Conclusion:

The conclusion has to be revised. Conclusion is based on the finding, not assumption or interpretation.

References

1. Muchlisin ZA, Musman M, Azizah MN: Spawning seasons of *Rasbora tawarensis* (Pisces: Cyprinidae) in Lake Laut Tawar, Aceh province, Indonesia. *Reprod Biol Endocrinol*. 2010; **8**: 49 [PubMed Abstract](#) | [Publisher Full Text](#)
2. Muchlisin ZA, Musman M, Fadli N, Nor Siti-Azizah N: Fecundity and spawning frequency of *Rasbora tawarensis* (Pisces: Cyprinidae) an endemic species from Lake Laut Tawar, Aceh, Indonesia. *AACL Bioflux*. 2011; **4** (3).
3. Muchlisin ZA: Potency of freshwater fishes in Aceh waters as a basis for aquaculture development program. *Indonesian Journal of Ichthyology*. 2013; **13** (1).

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Partly

Are sufficient details of methods and analysis provided to allow replication by others?

Partly

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research