RESEARCH NOTE

REVISED Preliminary study on the inhibitory effect of seaweed *Gracilaria verrucosa* extract on biofilm formation of *Candida albicans* cultured from the saliva of a smoker [version 2; peer review: 1 approved, 2 approved with reservations]

Zaki Mubarak¹, Adintya Humaira¹, Basri A. Gani¹, Zainal A. Muchlisin²

¹Faculty of Dentistry, Syiah Kuala University, Banda Aceh, 23111, Indonesia

²Faculty of Marine and Fisheries, Syiah Kuala University, Banda Aceh, 23111, Indonesia

V2 First published: 31 May 2018, 7:684
<https://doi.org/10.12688/f1000research.14879.1>
 Second version: 16 Aug 2018, 7:684
<https://doi.org/10.12688/f1000research.14879.2>
 Latest published: 18 Sep 2018, 7:684
<https://doi.org/10.12688/f1000research.14879.3>

Abstract

Background: *Candida albicans* is an opportunistic fungus that might infect the oral cavity. Increased colony numbers of *C. albicans* in the mouth can be caused by multiple factors, such as smoking, weakened immune system, antibiotics use and immune-compromised condition. Smoking can increase expression of virulence factors of *C. albicans* and make it stronger. One virulence factor of *C. albicans* is biofilm formation. The ability of creating biofilm makes *C. albicans* more tolerant to commercial antifungal agents. The objective of this preliminary study was to examine the ability of the seaweed *G. verrucosa* extracts to inhibit the formation of biofilm by *C. albicans* isolated from the saliva of a smoker.

Methods: The extract of *G. verrucosa* was prepared by maceration using 96% methanol and subjected for phytochemical analysis. *C. albicans* was isolated from the saliva of a smoker who voluntarily participated in the study after providing informed consent. In triplicate, the fungus was cultured in the growth medium containing increased concentrations of *G. verrucosa* (6.25, 12.5, 25, 50, 75 and 100%). The same reaction using fluconazole 0.31 µg/ml *C. albicans* was prepared as positive control. Biofilm formation was accessed based on optical density of cell mixtures using an ELISA reader. The data obtained were subjected to Kruskal-Wallis test at a significance limit of 0.05.

Results: Methanol extract of seaweed *G. verrucosa* contained three bio-active compounds namely steroids, terpenoid, and tannins. Inhibitory activity of seaweed extracts on *C. albicans* biofilm formation increased as their concentration increased. The highest inhibitory effect was recorded at fungus culture treated with seaweed

Open Peer Review

Approval Status

	1	2	3
version 3			
(revision)			
18 Sep 2018	view		view
version 2			
(revision)			
16 Aug 2018	view	view	
version 1			
31 May 2018			
	view	view	view

1. **Heni Susilowati** , Gadjah Mada University, Yogyakarta, Indonesia
2. **Shahida Mohd-Said** , Universiti Kebangsaan Malaysia, Kuala Lumpur, Malaysia
3. **Cristiane Y. Koga-Ito** , São Paulo State University - UNESP, São José dos Campos, Brazil

Any reports and responses or comments on the article can be found at the end of the article.

concentration of 25% at 24 hours of time exposure.

Conclusions: Seaweed *G. verrucosa* extract contained steroids, terpenoids and tannins that were able to effectively inhibit the formation of biofilm by *C. albicans* at the concentration of 25% after 24 hours of time exposure.

Keywords

Candida albicans, oral candidiasis, seaweed Gracilaria verrucosa

Corresponding author: Zaki Mubarak (zakimubarak54@yahoo.com)

Author roles: **Mubarak Z:** Conceptualization, Formal Analysis, Funding Acquisition, Investigation, Methodology, Supervision, Validation; **Humaira A:** Data Curation, Investigation, Project Administration, Software, Visualization, Writing – Original Draft Preparation; **Gani BA:** Conceptualization, Formal Analysis, Investigation, Methodology, Resources, Writing – Original Draft Preparation; **Muchlisin ZA:** Software, Visualization, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2018 Mubarak Z *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. Data associated with the article are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

How to cite this article: Mubarak Z, Humaira A, Gani BA and Muchlisin ZA. **Preliminary study on the inhibitory effect of seaweed *Gracilaria verrucosa* extract on biofilm formation of *Candida albicans* cultured from the saliva of a smoker [version 2; peer review: 1 approved, 2 approved with reservations]** F1000Research 2018, 7:684 <https://doi.org/10.12688/f1000research.14879.2>

First published: 31 May 2018, 7:684 <https://doi.org/10.12688/f1000research.14879.1>

REVISED Amendments from Version 1

Some revisions and improvements have been done throughout the manuscript as responses to the good comments and suggestions from reviewers. Major revisions performed include grammar check and corrections; sentence improvements; and reorganization and adding appropriate details in each section of the Methods part. Grammar issues, a concern of all reviewers, have been addressed. The ways to present results were also changed as response to comment from Dr Heni Susilowati about the meaning of OD values in presenting the growth of *C. albicans* and how to interpret them. Restructure of Results and Discussion, therefore, was also performed. These changes became the basis for synthesizing more appropriate Conclusion sentences. One reference was also added to support discussion about the importance of extract concentration for resulting in expected medicinal effects, in this study was growth inhibition that reflected by biofilm formation. Since this study is still in preliminary step, no non-smoker participant was participated. As consequence, no comparison could be made between smoker and non-smoker groups. Yet, the study was still able to confirm potential of seaweed *Gracilaria verrucosa* extract as anti candidiasis in smokers. Potential of doing further studies was mentioned in relation to this weakness of our study.

See referee reports

Introduction

Smoking is a common problem in both developed and developing countries, including in Indonesia. Based on a survey by the Tobacco Atlas in 2015, Indonesia has the highest number of smokers in Asia, with 66% of men in Indonesia being active smokers¹. Smoking can lead to addiction owing to the nicotine contents, and harm due to the presence of toxic compounds such as carbon monoxide, ammonia and tar contents in tobacco¹. Substances in cigarettes can also contribute for the occurrence of oral candidiasis, infection of in the mouth cavity caused by the fungus *Candida albicans*². This fungus is part of the normal flora of the human mouth, but it can become pathogenic in certain conditions, for example, due to nicotine exposure³.

The *C. albicans* that infects human tissues generally form biofilm³, an extracellular matrix consisting of *C. albicans* colonies⁴. The size of the biofilm increases when the fungus was exposed to substances in cigarette smoke because cigarette contains chemicals that can initiate the growth of and nourish *C. albicans*^{5,6}.

Currently, fluconazole and nystatin are the most effective drugs for treating oral candidiasis. Unfortunately, these drugs could result in undesired side effects. Prolonged use of fluconazole, for example, can lead to resistance⁷ whereas high dosages of nystatin give gastrointestinal discomfort and increase plaque formation⁸. Therefore, plant-derived antifungals may be viable oral treatment options for candidiasis. One of these potential plants is *Gracilaria verrucosa*. This seaweed contains several bioactive compounds, including alkaloids, flavonoids, phenolics, saponins, steroids and terpenoids⁹. Aceh Province, Indonesia, has large *G. verrucosa* resources, although this aquatic plant has not been commonly used for medicinal purposes. Hence, the

objective of the present study was to examine the ability of seaweed *G. verrucosa* extract to inhibit the biofilm formation of *C. albicans* isolated from the saliva of smoker.

Methods

Time and site

The study was conducted in August 2017 at the Laboratory of Microbiology, Veterinary Faculty, Syiah Kuala University. The *G. verrucosa* seaweeds were collected from a farmer in Pulo Aceh, Aceh Province.

Ethics

All research protocols used in this study were approved by the Research Ethics Committee of the Dentistry Faculty of Syiah Kuala University No. 1741/UN11.1.21/TU/2017.

Saliva collection

The saliva was collected from an active smoker who worked as administrative staff at the Faculty of Dentistry Medicine of Syiah Kuala University and voluntarily participated in this study after providing written informed consent. Inclusion criterion of the volunteer was active smoker who smoked 20 cigarettes per day. Saliva was collected once by spitting into a glass jar (15 ml) right after the subject finished smoking, and added with 10 ml of PBS (0.01 M, pH 7.2). The jar was centrifuged at 10,000 rpm for 10 minutes. The precipitate was stored for the microbiological examination.

Candida albicans Isolation and preparation

Precipitate was cultured in ChromAgar *Candida* medium and incubated at 37 °C for 2 days. The *C. albicans* fungus grew as green colonies. One colony of *C. albicans* was mixed with 5 ml of peptone in a tube and incubated at 37°C for 24 hours. Turbidity of medium was compared to a 0.5 McFarland solution standard, which was equivalent to 1.5×10^8 CFU/ml.

Seaweed extraction

Seaweed extraction was performed by maceration using 96% methanol¹⁰. In brief, a total of 500 g of fresh seaweeds were washed with tap water then with distilled water. Seaweed samples were dried at 25°C for 24 hours, chopped into small pieces (2 mm), and soaked in 1.5 liters of 96% methanol. Macerate was filtered using Whatman filter paper No. 42. Filtrate collected was concentrated using a vacuum rotary evaporator (Laborta 4003 control, Heildolph) 60°C and at a speed of 80-90 rpm for 3 days. Concentrated extract was put in a sealed dark bottle and stored at 4 °C.

Phytochemical tests

Flavonoid test. A 0.5 cm magnesium plate was rinsed in 5 ml of seaweed extract, mixed with two drops of HCl, and heated by passing it over a Bunsen flame. Red or purple coloration formed on the heating indicated the presence of flavonoids¹¹.

Alkaloid test. Seaweed extract (5 ml) was mixed with 8 ml of HCl and filtered. Filtrate was subjected to Mayer, Wagner and Dragendroff tests for alkaloids¹¹. This was done by mixing

2 ml of filtrate with 5 g potassium mercuric iodide (Mayer test), 2 ml of Wagner reagent, or 2 ml of bismuth potassium iodide solution (Dragendroff test). The formation of white or pale precipitates (Mayer test), brown or reddish-brown precipitates (Wagner test) and red precipitates (Dragendroff test) indicated the presence of alkaloids.

Tannin/phenolic test. Two drops of 1% FeCl_3 was added to 1 ml seaweed extract. The change in the color to a blackish green indicated the presence of tannin/phenolic¹².

Saponin test. Seaweed extract, 1 ml, was diluted in 20 ml of distilled water and shaken vertically for 15 seconds. Persistent foaming indicated the presence of saponin content.

Steroid test. Two milliliters of seaweed extract was added with 2 ml of CHCl_3 , 2 drops of H_2S and 1 ml of CH_3COOH . The formation of green or blue precipitates indicated the presence of steroid¹¹.

Terpenoid test. Seaweed extracts (5 ml) was mixed with 2 ml of chloroform. Concentrated H_2SO_4 , 3 ml, were carefully added. The formation of reddish brown layer at the interface of extract and chloroform solution indicated the presence of terpenoids¹³.

Examination of Biofilm Formation

Casein-peptone lecithin polysorbate broths (Merck-1117230500), 100 μl , were poured in each well of a 96-well plate, incubated at room temperature for 5 minutes and discarded. Fifty microliter *C. albicans* culture had turbidity was equal to 0.5 McFarland standard were added to each well and incubated for 5 minutes. In triplicate, decreased concentrations of seaweed extracts (100, 75, 50, 25, 12.5 and 6.25%) were added. The same reaction using fluconazole 0.31 $\mu\text{g/ml}$ was prepared as positive control. The plate was then incubated at 37 °C for 24, 48 or 72 hours. Each well was added with 200 μl of 0.1% violet crystal, incubated at 25 °C for 15 min, and washed three times with 200 μl of 0.01 M PBS. The crystal violet in each well was then removed by adding 100 μl of 96% ethanol for 2 min. The biofilm formation was ana-

lyzed by reading optical density of mixture using an ELISA reader at 620 nm^{14,15}.

Data analysis

The data obtained were subjected to Kruskal-Wallis test using SPSS software v20.0 for windows.

Results

The results of phytochemical tests in Table 1 show that methanol extract of seaweed *G. verrucosa* contained bioactive compounds belonged to steroids, terpenoids, and tannins/polyphenols.

In general, the inhibitory effect was reduced as seaweed concentration increased (Figure 1). The highest inhibition after 24 and 48 hours time exposures was recorded from *C. albicans* culture treated with 25% seaweed extract, followed by those with 50% and 75%. The best inhibition after 75 hours time exposure, however, was found in *C. albicans* culture treated with 25% seaweed extract. In all time exposures, the inhibitory effect caused by fluconazole (control) was the worst, followed by those caused by 100% seaweed extracts, but there was no difference between these two groups. Results of Kruskal-Wallis analysis showed that seaweed extract significantly ($p < 0.05$) inhibited the formation of biofilm by *C. albicans*, indicated potential of the extract to inhibit the growth of the fungus.

Dataset 1. The raw data of the Triplo anti-Biofilm seaweed to *C. albicans* for 24, 48 and 72 h at a wavelength 620 nm

<http://dx.doi.org/10.5256/f1000research.14879.d204270>

Discussion

The study showed that 25–75% seaweed extracts are promising for inhibiting the growth of *C. albicans* as shown by much lower biofilm formation after 24, 48 and 72 hours exposure compared to those caused by positive control (fluconazole) and 100% seaweed extract. This indicated potential of seaweed extract as natural anti-fungus to treat oral candidosis caused by against *C. albicans* in smokers.

Table 1. Phytochemical contents of seaweed *Gracilaria verrucosa* extract.

Substance	Reagent	Result	Indication
Alkaloid	Mayer	-	White deposit
	Wagner	-	Brown deposit
	Dragendroff	-	Red deposit
Steroid	Uji Lieberman-Burchard	+	Green or blue colors
Terpenoid	Uji Lieberman-Burchard	+	Red or purple colors
Saponin	Shuffling method	-	Stable foams
Flavonoid	0.5 Mg and HCl	-	Red or purple colors
Tannin/Phenolic	MgCl_3	+	Dark green

Figure 1. The formation of biofilm by *Candida albicans* exposed to seaweed *Gracilaria verrucosa* extract on different time exposures and concentrations.

C. albicans is a normal micro-organism in the human mouth. This fungus, however, can be pathogenic in certain circumstances³ such as in the mouth of smokers². Smoking can stimulate synthesis of HWPI, EAP1 and SAP2 proteins in *C. albicans*, causing higher virulence of the fungus. This can lead to increased formation of biofilm and finally cause oral candidiasis⁶. Smoking also cause a decreased immune function, making individuals more susceptible to oral infections, including candidiasis^{4,6}.

The ability of 25–75% seaweed extracts to inhibit the formation of *C. albicans* biofilm significantly stronger than those of fluconazole (positive control) was probably caused by bioactive compounds presence in the extracts. These included steroids, terpenoids, and tannins/polyphenols, all of which have been known their benefits for human health. According to Sampaio *et al.*¹⁶, antifungal activity of a substance strongly depends on the composition of its bioactive compounds. Steroids can kill *C. albicans* through their lipophilic properties, interfering with the formation of fungal spores and mycelium¹⁷. This activity weakens *C. albicans*, inhibiting the formation of the biofilm. To function optimally, steroids require oligosaccharides that are also present in the seaweed content¹⁸.

Terpenoids are derivatives of saponins that may act as an antifungals by damaging organelles of the fungus and by inhibiting secretion of enzymes, leading to the inhibition of *C. albicans* cell growth. Terpenoids can also damage the morphology of *C. albicans*¹⁹. Tannins may inhibit chitin synthesis in *C. albicans*

cell walls, leading to lost of membrane cell protection and disrupted cellular metabolism. Tannins also can inhibit ergosteron activity of *C. albicans*²⁰.

The effectiveness of seaweed extracts in inhibiting fungal growth is influenced by at least three factors, namely concentration, exposure time, and contact surface media²¹. The present study showed that diluted concentration (25–75%) of extracts showed better inhibitory effect on the growth of *C. albicans* than the more concentrated 100% extract. Concentrated extract usually has lesser effectiveness *in vitro* due to solubility and import problems. In more aqueous condition plant extracts generally show better medicinal properties with increased concentration as shown by a study testing antifungal activity of ethanol extracts of *Syzygium jombolanum*, *Cassia siamea*, *Ordina wodier*, *Momodica charantia* and *Melia azedarach* as well as two algal species *Sargassum wightii* and *Saulerpa scalpelliformis* against 25 *C. albicans* isolates *in vitro*²².

This study also showed the best inhibitory effect of seaweed extracts was recorded at 24 hour of exposure. This is probably because the farnesol, a quorum-sensing molecule that has the potency to inhibit *C. albicans* growth²³, works effectively after 48–72 hours of exposure.

This study, however, was still preliminary due to limited number of subject and unavailability of non-smoker, control. Virulence factor of *C. albicans* analyzed was also only biofilm formation. Since more virulence factors are possibly synthesized

by the fungus under certain environmental condition, further studies are need to be conducted to investigate effect of individual bioactive compound contained in the seaweeds on the formation of biofilm and on the expression of other virulence factors of *C. albicans* isolated form both smoker and non-smoker individuals.

Conclusion

Extract of *Gracilaria verrucosa* seaweed could inhibit the growth of *C. albicans* isolated from the saliva of a smoker. The inhibitory effect decreased with the increase of concentration, and reached the highest at concentration of 25% and time exposure of 24 hours.

Data availability

Dataset 1. The raw data of the Triplo anti-Biofilm seaweed to *C. albicans* for 24, 48 and 72 h at a wavelength 620 nm. DOI: [10.5256/f1000research.14879.d204270](https://doi.org/10.5256/f1000research.14879.d204270)²⁴.

Competing interests

No competing interests were disclosed.

Grant information

The authors declare that no funding was involved in supporting this work.

References

- Lian TY, Dorotheo U: **The Tobacco Control Atlas**. Seutawan Co. 2016; 3. [Reference Source](#)
- Mayer FL, Wilson D, Hube B: **Candida albicans pathogenicity mechanisms**. *Virulence*. 2013; 4(2): 119–128. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Adel E, Khadijeh PS, Mohammad M: **Oral Cavity Candidiasis as a Complication of Fungal Diseases in Diabetic Patients in South-East of Iran**. *IJIAS*. 2016; 14(4): 1134–1138. [Reference Source](#)
- Little JW, Falace D, Miller C: **Dental Management of The Medically Compromised Patient**. Mosby: Elsevier; 2013.
- Keten HS, Keten D, Ucer H, *et al.*: **Prevalence of oral candida carriage and candida species among cigarette and maras powder users**. *Int J Clin Exp Med*. 2015; 8(6): 9847–54. [PubMed Abstract](#) | [Free Full Text](#)
- Semlali A, Killer K, Alanazi H, *et al.*: **Cigarette smoke condensate increases C. albicans adhesion, growth, biofilm formation, and EAP1, HWP1 and SAP2 gene expression**. *BMC Microbiol*. 2014; 14: 61. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Peron IH, Reichert-Lima F, Busso-Lopes AF, *et al.*: **Resistance Surveillance in Candida albicans: A Five-Year Antifungal Susceptibility Evaluation in a Brazilian University Hospital**. *PLoS One*. 2014; 11(7): e0158126. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Lyu X, Zhao C, Yan ZM, *et al.*: **Efficacy of nystatin for the treatment of oral candidiasis: a systematic review and meta-analysis**. *Drug Des Devel Ther*. 2016; 10: 1161–1171. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Sahat JH: **Warta Ekspor**. September ed. Jakarta: DGNED; 2013; 3.
- Prima MI: **Uji Aktivitas Antibakteri Ekstrak Metanol Ganggang Merah Gracilaria verrucosa terhadap Beberapa Bakteri Patogen Gram Positif dan Gram Negatif**. Published Only In Database. 2012. [Reference Source](#)
- Vimalakumar CS, Hosagaudar VB, Suja SR, *et al.*: **Comparative Preliminary Phytochemical Analysis Of Ethanolic Extracts of Leaves of Olea dioica Roxb., Infected with The Rust Fungus Zaghovania oleae (E.J.Butler) Cummins and Non-infected Plants**. *J Pharmacogn Phytochem*. 2014; 3(4): 69–72. [Reference Source](#)
- Hariyanto Ih, Inarah F, Suci PR, *et al.*: **Skrining Fitokimia dan Analisis Kromatografi Lapis Tipis dari Ekstrak Ethanol Herba Pacar Air (Impatiens balsamina Linn.)**. Published Only In Database. 2014. [Reference Source](#)
- Khan MA, Qureshi AR, Ullah F, *et al.*: **Phytochemical analysis of selected medicinal plants of Margalla Hills and surroundings**. *JMPR*. 2011; 5(25): 6017–6023. [Reference Source](#)
- Metzler A: **Developing a Crystal Violet Assay to Quantify Biofilm Production Capabilities of Staphylococcus aureus**. Published Only In Database. 2016. [Reference Source](#)
- Roberts SK, Wei GX, Wu CD: **Evaluating biofilm growth of two oral pathogens**. *Lett Appl Microbiol*. 2002; 35(6): 552–6. [PubMed Abstract](#) | [Publisher Full Text](#)
- Sampaio BL, Edrada-Ebel R, Da Costal FB: **Effect of the environment on the secondary metabolic profile of Tithonia diversifolia: a model for environmental metabolomics of plants**. *Sci Rep*. 2016; 6: 29265. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Subsiha S, Subramoniam A: **Antifungal Activities of Steroid From Pallavicinia lyellii, a liverwort**. *Indian J Pharmacol*. 2005; 37(5): 304–308. [Publisher Full Text](#)
- Cammarata A, Upadhyay SK, Jursic BS, *et al.*: **Antifungal activity of 2 α ,3 β -functionalized steroids stereoselectively increases with the addition of oligosaccharides**. *Bioorg Med Chem Lett*. 2011; 21(24): 7379–7386. [PubMed Abstract](#) | [Publisher Full Text](#)
- Martínez A, Rojas N, García L, *et al.*: **In vitro activity of terpenes against Candida albicans and ultrastructural alterations**. *Oral Surg Oral Med Oral Pathol Oral Radiol*. 2014; 118(5): 553–9. [PubMed Abstract](#) | [Publisher Full Text](#)
- Hastuti SU, Ummah PIY, Khasanah NH: **Antifungal Activity of Piper aduncum and Peperomia pellucida Leaf Ethanol Extract Against Candida albicans**. *AIP Conf Proc*. 2017; 1844: 020006. [Publisher Full Text](#)
- Kenakin PT: **A Pharmacology Primer: Theory, Application, and Methods**. 2nd edition. USA: Elsevier. 2006; 36–38. [Publisher Full Text](#)
- Prabhakar K, Kumar LS, Rajendran S, *et al.*: **Antifungal Activity of Plant Extracts against Candida Species from Oral Lesions**. *Indian J Pharm Sci*. 2008; 70(6): 801–803. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Alem AM, Oteef MD, Flowers TH, *et al.*: **Production of tyrosol by Candida albicans biofilms and its role in quorum sensing and biofilm development**. *Eukaryot Cell*. 2006; 5(10): 1770–1779. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Mubarak Z, Humaira A, Gani BA, *et al.*: **Dataset 1 in: Preliminary study on the inhibitory effect of seaweed Gracilaria verrucosa extract on biofilm formation of Candida albicans cultured from the saliva of a smoker**. *F1000Research*. 2018. <http://www.doi.org/10.5256/f1000research.14879.d204270>

Open Peer Review

Current Peer Review Status: ? ✓ ?

Version 2

Reviewer Report 23 August 2018

<https://doi.org/10.5256/f1000research.17357.r37284>

© 2018 Mohd-Said S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Shahida Mohd-Said

Periodontology Unit, Centre for Restorative Dentistry, Faculty of Dentistry, Universiti Kebangsaan Malaysia, Kuala Lumpur, Malaysia

Improvements made are acknowledged and approved.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Periodontology, natural product drug discovery

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 20 August 2018

<https://doi.org/10.5256/f1000research.17357.r37282>

© 2018 Susilowati H. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Heni Susilowati

Department of Oral Biology, Faculty of Dentistry, Gadjah Mada University, Yogyakarta, Indonesia

I really appreciate the efforts of the authors who have revised the report, but I would like to add suggestions to the authors to explain why the optical density value of the treatment group treated with Fluconazole and extracts of 100% was even greater than the extracts with smaller concentrations.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Molecular Biology and Oral Microbiology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Version 1

Reviewer Report 27 June 2018

<https://doi.org/10.5256/f1000research.16195.r34560>

© 2018 Koga-Ito C. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Cristiane Y. Koga-Ito

Oral Biopathology Graduate Program, Institute of Science and Technology, São Paulo State University - UNESP, São José dos Campos, Brazil

The manuscript by Mubarak et al. aimed to evaluate the effect of *G. verrucosa* extract on *Candida albicans* biofilm formation. The rationale of the study should be clearer. Also, there is lack of essential information throughout the text. The English should be revised.

Specific comments

Introduction

The period "Smoking is a common problem in most developing countries, including Indonesia.". A reference should be added. Also, smoking is not a problem only or mainly in developing countries. Please, consider revising.

The authors stated that "This fungus is part of the normal flora of the human mouth, but it can become pathogenic in certain conditions, for example, due to nicotine exposure.". The exposure to nicotine has been correlated with increasing in *C. albicans* virulence factor expression. The predisposing conditions for candidiasis are much more related to immunologic state of the host and imbalance in microflora. Please, consider revising.

Revise the period "Infection with *C. albicans* will increase the formation of a biofilm of the fungus", it is confusing.

Revise the period "The biofilm is an extracellular matrix consisting of *C. albicans* colonies", it is confusing.

The authors stated that "high dosages of nystatin give gastrointestinal discomfort and increase plaque formation". What do authors mean by "increase plaque formation"?

The rationale of the study is not clear. Why did the authors select *G. verrucosa*? Is this plant commonly used? Why did the authors decide to use saliva from a smoker individual?

Methods

Revise the period "C. albicans was extracted from the saliva", C. albicans was isolated from saliva. The authors reported that "G. verrucosa seaweed was collected from a farmer in Pulo Aceh, Aceh Province.". More information on the plant, the exact location it was collected from, the period of the year, identification procedure (how and who did the identification?), registration in herbarium, number of voucher should be included in the text.

Please, revise the period "Saliva was collected by spitting into a glass jar (15 ml), then 1 ml PBS (0.01 M, pH 7.2) was added to the jar." Why did 1 ml of PBS added to the saliva? What was the final volume of saliva collected? Was saliva stimulated?

The authors stated that "If the colour of a colony was green, this indicated that the colony was C. albicans.". However, the color of the colony in CHROMagar is only a presumptive test and phenotypic or genotypic definitive identification should be done.

The inclusion of a reference strain is highly needed.

Include the number of experiments/replicates performed.

The inclusion of more clinical isolates from non-smokers patients is needed.

The methodology of activity of extract on biofilm formation is not clear. Why and how peptone was removed from the wells? Why did the fungal suspension left in the wells for 5 min? How 100% extract concentration was obtained in the well (you already has the broth inside the well)? Why the concentration of 0.31 was chosen for fluconazole?

Figure 1 should be revised. Note that optical density at 620 nm is higher after treatment with 100% extract when compared to the other concentrations.

Discussion and Conclusion sections should be revised after the revision of the aforementioned points.

Is the work clearly and accurately presented and does it cite the current literature?

No

Is the study design appropriate and is the work technically sound?

No

Are sufficient details of methods and analysis provided to allow replication by others?

No

If applicable, is the statistical analysis and its interpretation appropriate?

Not applicable

Are all the source data underlying the results available to ensure full reproducibility?

No

Are the conclusions drawn adequately supported by the results?

No

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Microbiology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Reviewer Report 19 June 2018

<https://doi.org/10.5256/f1000research.16195.r34561>

© 2018 Mohd-Said S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Shahida Mohd-Said

Periodontology Unit, Centre for Restorative Dentistry, Faculty of Dentistry, Universiti Kebangsaan Malaysia, Kuala Lumpur, Malaysia

Overall fair manuscript with sound finding and relevant area of study. Can much be improved with grammar check and essential scientific writing reorganisation especially in Introduction and Discussion section. Inclusion of results for negative control (untreated biofilm) would critically improve Results presentation and appreciation of findings. Lacks relevant information on findings between smoker and non-smoker in the study that may not strongly support the conclusion on the anti-fungal effect of agent on smoker.

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Partly

If applicable, is the statistical analysis and its interpretation appropriate?

I cannot comment. A qualified statistician is required.

Are all the source data underlying the results available to ensure full reproducibility?

No

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Periodontology, natural product drug discovery

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 05 Aug 2018

Zaki Mubarak, Syiah Kuala University, Banda Aceh, Indonesia

Some improvement in grammar, sentences and writing reorganization have been done as a response to reviewer Dr Shahida M. Said. Some details in methods and analysis have also been provided to make better readability and potential reproducibility of the work. As a result the work in this study is much better presented. These improvements can be found in track changes.

As this study is still preliminary comparison between smoker and non-smoker has not been performed yet, but the finding clearly show potential of use of seaweed extract to treat oral candidiasis.

Competing Interests: No competing interests were disclosed.

Reviewer Report 12 June 2018

<https://doi.org/10.5256/f1000research.16195.r34557>

© 2018 Susilowati H. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Heni Susilowati

Department of Oral Biology, Faculty of Dentistry, Gadjah Mada University, Yogyakarta, Indonesia

This preliminary research is interesting enough to be developed but there are some things that need to be reconsidered:

1. ABSTRACT

- Research background that written on the abstract (lines 6 and 7) does not match the purpose of the study. The sentence can be interpreted in contrast to the antifungal potential possessed by *Garcinia verrucosa*. It is dubious to investigate the potency of biofilm inhibition effect if *Candida albicans* more tolerant to *Gracilaria* extract.

2. METHODS

Methods need to explain the following:

- Systemic conditions and state of teeth and oral soft tissue volunteers,
- The predestined determination of the *Gracilaria verrucosa* plant should be mentioned.
- Were the culture washed after the incubation period on treatment?
- How many times an experiment produces a representative result?

3. RESULTS

- Interpretation of the results is confusing; as far as I know the higher the optical density value the more biofilms are formed. The results in Fig. 1 show that 100% and Fluconazole extracts have higher optical densities rather than the lower concentration of extracts, as far as I know this shows that the mass of biofilms formed in the group is higher. Please observe the methods and results of research reported by Sebaa et al 2016.
- The statistical method used was only Kruskal-Wallis, is there a multiple difference analysis? Researchers need to discuss the effect of antibiofilm extract at lower concentrations, because of course 100% extract is not a good recommendation for subsequent experimental use.

References

1. Sebaa S, Hizette N, Boucherit-Otmani Z, Courtois P: Dose-dependent effect of lysozyme upon *Candida albicans* biofilm. *Mol Med Rep.* 2017; **15** (3): 1135-1142 [PubMed Abstract](#) | [Publisher Full Text](#)

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

No

If applicable, is the statistical analysis and its interpretation appropriate?

I cannot comment. A qualified statistician is required.

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

No

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 05 Aug 2018

Zaki Mubarak, Syiah Kuala University, Banda Aceh, Indonesia

Abstract

Comments from reviewer :

The objective has been clarified

Action done:

We think aims of study still need to be stated

Rearrangement and sentences improvement were done in some sections of abstract

Introduction

Comments from reviewer :

No comments

Action done:

Grammar check, revise and improvement were done to provide more concise and clear sentences/statements

Material and method

Comments from reviewer :

Need clarification on sentence:

"This volunteer has a bad OHIS4" (time and place)

".Then microtiter wells plate wells were washed three times with 200µl of PBS buffer and dried for 15 minutes."(biofilm examination)

"This examination was repeted three times with triplo method"(biofilm examination)

Action done:

Sentences improvement, grammar check and correction had been done throughout this section to provide better readability, clarity and possible reproducibility

Results

Comments from reviewer :

No comments

Action done:

Rewritten this part due to misinterpretations we did in the first previous manuscript about which treatment gave best inhibition effect on *C. albicans*

Discussion

Comments from reviewer :

No specific comments

Action done:

Rewritten has been done to this part in realtion to due to changes in results presentation
Sentences improvement, grammar check and reorganization had been done throughout this section to provide better readability and clarity

Conclusion

Comments from reviewer :

No specific comments

Action done:

This part had been rewritten in relation to the revision performed in result and discussion.

Competing Interests: No competing interests were disclosed.

Comments on this article

Version 1

Author Response 05 Aug 2018

Zaki Mubarak, Syiah Kuala University, Banda Aceh, Indonesia

Some revisions and improvements have been done throughout the manuscript as responses to the good comments and suggestions from reviewers. Major revisions performed include grammar check and corrections; sentence improvements; and reorganization and adding appropriate details in each section of the material and methods part. Grammar issues, were concerned by all reviewers, had been addressed. The ways to present results were also changed as response to comment from Indonesia reviewer about the meaning of OD values in presenting the growth of *C. albicans* and how to interpret them. Restructure of result and discussion, therefore, was also performed. These changes became the basis for synthesizing more appropriate conclusion sentences. One reference was also added to support discussion about the importance of extract concentration for resulting in expected medicinal effects, in this study was growth inhibition that reflected by biofilm formation. Since this study is still in preliminary step, no non-smoker participant was participated. As consequence, no comparison could be made between smoker and non-smoker groups. Yet the study was still able to confirm potential of seaweed *Gracia verrucosa* extract as anti candidiasis in smokers. Potential of doing further studies was mentioned in relation to this weakness of our study.

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research