Check for updates

RESEARCH ARTICLE

Streptococcus pneumoniae serotype epidemiology among PCV-10 vaccinated and unvaccinated children at Gertrude's Children's Hospital, Nairobi County: a cross-sectional study [version 1; peer review: 1 approved, 1 approved with reservations, 1 not approved]

Michael Walekhwa¹ , Margaret Muturi¹, Revathi Gunturu², Eucharika Kenya³, Beatrice Kabera⁴

¹Department of Pathology, Gertrude's Children's Hospital, Nairobi, Kenya

²School of Medicine, Aga Khan University Hospital, Nairobi, Kenya

³Faculty of Biological Sciences, University of Embu, Embu, Kenya

⁴School of Medicine, Kenyatta University, Nairobi, Kenya

V1 First published: 22 Jun 2018, 7:879
<https://doi.org/10.12688/f1000research.14387.1>

Latest published: 31 Jan 2019, 7:879
<https://doi.org/10.12688/f1000research.14387.2>

Abstract

Background: *Streptococcus pneumoniae* (SPn) serotype replacement and emergence of multidrug resistant SPn has exacerbated the need for continuous regional serotype surveillance. We investigated SPn serotypes circulating among children ≤5 years in Nairobi County.

Methods: *Streptococcus pneumoniae* stocks stored at −70°C in brain heart infusion medium were thawed at room temperature for 30 minutes. In total, 10 µl of the stored SPn cells were suspended in 50 µl PBS and gently vortexed. About 10 µl of the suspended cells were added on to a glass slide and mixed with 10 µl pooled antisera. The glass slide was swirled gently while observing for any reaction. The process was repeated with individual groups under various antisera pools. Those serotypes that did not belong to any pool were typed directly until a positive agglutination reaction was observed. The cells/PBS/serotype-specific antisera mixture on the glass slide were covered with a coverslip and observed under a phase contrast microscope at ×100 objective lens with oil emulsion.

Results: Out of the 206 subjects sampled, 20.39% (n=42) were found to be carriers of SPn. About 52% (n=22) of the SPn carriers had received the recommended dose of PCV-10, while 48% (n=20) of the carriers had not. Almost all (n=41; 19.90% of subjects) isolates contained non-vaccine type SPn serotypes, while n=1 of the serotypes (in 0.49% of subjects) were untypeable. Serotypes 28F, 6A, 11A, 3 and 7C were prevalent in both vaccinated and unvaccinated children,

Open Peer Review

Approval Status

	1	2	3
version 2 (revision) 31 Jan 2019	view		
version 1 22 Jun 2018	view	view	view

1. **Bartholomew N. Ondigo**, Kenya Medical Research Institute, Kisumu, Kenya
2. **Jackie K. Obey**, University of Eastern Africa, Baraton, Eldoret, Kenya
3. **Felix Dube** , University of Cape Town, Cape Town, South Africa

Any reports and responses or comments on the article can be found at the end of the article.

whereas serotypes 23A, 17F, 35F, 48, 13 and 35B, and 23B, 20, 19B, 21, untypeable, 15B and 39 were found among unvaccinated and vaccinated groups, respectively.

Conclusions: All *SPn* serotypes isolated from the subjects sampled were non PCV-10 vaccine type. Therefore Kenyan children receiving PCV-10 vaccine are not protected.

Keywords

Streptococcus pneumoniae, serotypes, Nairobi, Quellung reaction, Optochin test, Bile solubility

Corresponding authors: Margaret Muturi (muturimargaret@gmail.com), Revathi Gunturu (gunturu.revathi@aku.edu), Eucharia Kenya (eucharia.kenya2012@gmail.com), Beatrice Kabera (beats.kabera@gmail.com)

Author roles: **Walekhwa M:** Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Writing – Original Draft Preparation, Writing – Review & Editing; **Muturi M:** Conceptualization, Methodology, Project Administration, Supervision; **Gunturu R:** Methodology, Resources, Supervision; **Kenya E:** Funding Acquisition, Methodology, Supervision; **Kabera B:** Methodology, Resources, Supervision

Competing interests: No competing interests were disclosed.

Grant information: The study was in part funded by the National Commission of Science Technology & Innovation (NACOSTI) Kenya. Ref Nos. NACOSTI/RCD/ST & I/7TH CALL/PhD/148.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2018 Walekhwa M *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. Data associated with the article are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

How to cite this article: Walekhwa M, Muturi M, Gunturu R *et al.* *Streptococcus pneumoniae* serotype epidemiology among PCV-10 vaccinated and unvaccinated children at Gertrude's Children's Hospital, Nairobi County: a cross-sectional study [version 1; peer review: 1 approved, 1 approved with reservations, 1 not approved] F1000Research 2018, 7:879 <https://doi.org/10.12688/f1000research.14387.1>

First published: 22 Jun 2018, 7:879 <https://doi.org/10.12688/f1000research.14387.1>

Introduction

Streptococcus pneumoniae (SPn) is a highly invasive, Gram-positive, extracellular bacterial pathogen (Mitchell & Mitchell, 2010). It is a major cause of morbidity and mortality globally, causing more deaths than any other infectious disease (Jones *et al.*, 2010). SPn is classified into serogroups (denoted by numbers and letters, e.g. 18c, 23f) (Kellogg *et al.*, 2001). There are over 90 known serotypes and the prevalence of different serotypes varies by regions of the world (Hackel *et al.*, 2013). Different serotypes exhibit differing potentials to cause disease and may cause different syndromes in different age groups (Harboe *et al.*, 2009).

Some strains also have a greater potential to develop antibiotic resistance (Song *et al.*, 2012). The 13 most common serotypes of SPn pneumonia cause 80–93% of serious pneumococcal disease in children (Johnson *et al.*, 2010). According to the World Health Organization (WHO) and UNICEF Global Action Plan for the Prevention and Control of Pneumonia, pneumonia kills more children than any other illness in the world (WHO & UNICEF, 2009). Given the high burden of under-five mortality associated with pneumonia, control efforts are critical to achieving Sustainable Development Goal 3 (Colglazier, 2015).

WHO and UNICEF estimates indicate that over 800,000 children under 5 years of age die from pneumococcal disease each year (O'Brien *et al.*, 2009). In Kenya, an estimated one in every five children under 5 years of age dies from this disease every year (WHO, 2013).

SPn vaccines protect against several severe forms of pneumococcal disease, such as meningitis, pneumonia and bacteremia (Feldman & Anderson, 2014). These vaccines will not protect against these conditions if they are caused by agents other than SPn or from other strains of SPn that are not contained in the vaccine (Moffitt & Malley, 2011). The 10-valent pneumococcal conjugate vaccine (PCV10) was introduced into the Kenya Expanded Program on Immunization (KEPI) in February 2011 with a 2+1 schedule (at 6, 10, 14 weeks) without catch-up vaccinations (Hammit *et al.*, 2014). The vaccine covers 1, 4, 5, 6b, 7f, 9V, 14, 18c, 19f and 23f SPn serotypes.

Currently over 90 different serotypes have been identified, six of them very recently (Weinberger *et al.*, 2011). Various SPn serotypes with antigenic similarities are classified under the same groups (9A, 9L, 9N and 9V) while those lacking antigenic similarities are given numbers only (1, 2, 3, 4 and 5). The degree of interaction (cross-reactivity) between various SPn groups may vary. For instance, serotypes 6A and 6B have identical chemical composition except for one of the bonds between two sugars yet they are highly cross-reactive but serotypes 19F and 19A are less reactive.

Pneumococcal conjugate (PCVs) and polysaccharide (PPVs) vaccines are designed according to their virulence mechanisms and how they generally interact with the human immune system (Castañeda-Orjuela *et al.*, 2012). The WHO has advised that all children ≤5 years should be immunized against pneumococcal disease and continuous surveillance done to keep out the disease

especially in the developing world (Vandenbos *et al.*, 2013). The need for continuous surveillance has been exacerbated by the acute emergence of multi-drug resistant SPn strains and escalated child mortality and morbidity due to pneumococcal disease, despite the availability of PCVs and PPVs. (Väkeväinen *et al.*, 2010). This study therefore sought to establish the SPn serotypes among vaccinated and unvaccinated children ≤5 years of age in Nairobi County, Kenya.

Methods

Study location

This study was conducted among children ≤5 years attending the outpatient department of Gertrude's Children's Hospital, Muthaiga, and its satellite clinic in Githongoro, Nairobi County between May 2017 and February 2018. Subjects were clinically assessed by a physician and those who presented with pneumococcal disease symptoms recommended to the study nurse for recruitment. Gertrude's Children's Hospital is the largest standalone health care facility specializing in pediatric care in East and Central Africa. The hospital is accredited by the Joint Commission on International Accreditation (JCIA). SPn isolation and stocking was done at Gertrude's Children's Hospital Main Laboratory and capsular serotyping done at KEMRI Wellcome Trust, Kilifi, Kenya.

Study design

This was a descriptive cross-sectional study. *Streptococcus pneumoniae* serotype epidemiology among PCV-10 vaccinated and unvaccinated children between 6 months and 5 years of age was measured. Children who had no history of any chronic disease and whose parents or legal guardians consented to the study were systematically recruited. Children whose parents or legal guardians declined to give consent and those with any known immunosuppressive conditions were excluded from the study.

Sample size determination

To determine the minimum sample size, the formula developed by Chow *et al.* (2007) was used, with a prevalence rate of 16% (Agweyu *et al.*, 2014).

$$n = \frac{z^2 \hat{p}(1 - \hat{p})}{m^2}$$

Where n= desired minimal sample size; z= standard normal deviation (1.96, from the tailed normal table); $\hat{p}$ = prevalence rate; and m= the desired degree of accuracy at a 95% confidence level of 0.05. This gave a sample size of 206.

Identification of SPn

Nasopharyngeal swabs were per nasally collected using Copan flocked swabs and temporarily suspended in Armies medium for transportation to the main laboratory. Each swab was inoculated onto a selective gentamicin with 5% sheep blood agar (BA) plate. All swabs were plated within 24 h of collection. The plates were incubated at 37°C in a 5% CO₂ atmosphere and examined at 16–24 h and then again at 40–48 h for growth of SPn. Isolates were identified as SPn by colony morphology (Mucoid,

draughtsman appearance, α -haemolysis) and susceptibility to optochin (positive, ≥ 14 mm zone of inhibition; negative, < 14 mm zone of inhibition). Plates with colonies akin to SPn morphological features but with optochin clearance zones below 14 mm were further subjected to solubility in bile salts (positive, bile soluble; negative, bile insoluble).

The isolation of a single colony indicated carriage. Single colonies were picked using sterile inoculating loops and evenly plated on BA. After 24–48 h, enough inoculum was stocked in brain heart infusion (BHI) agar with 5% sheep blood (Ultralab East Africa, Ltd), gently vortexed and stored at -70°C for serotyping.

Serotyping of SPn

Capsular serotyping was done using the Quellung reaction test. Frozen vials containing SPn stocks stored at -70°C were thawed at room temperature for about 30 min. Next, 10 μl of the stored SPn cells were suspended in 50 μl PBS and gently vortexed. Subsequently, 10 μl of the suspended cells were added on to a glass slide and mixed with 10 μl pooled antisera (Statens Serum Institute, cat. No. 16744). The glass slide was swirled gently while observing for any agglutination reaction until a positive reaction was observed with various pooled antisera. The process was repeated with individual groups under various antisera pools.

After that, 10 μl of the suspended cells in PBS were added to a glass slide and mixed with various SPn serotype-specific antisera included in the antisera pools that gave a positive reaction. This was done until a positive reaction with the particular serotype specific antisera was observed. Those serotypes that did not belong to any pool were typed directly until a positive agglutination reaction was observed. The cells/PBS/serotype-specific antisera mixture on the glass slide were covered with a cover slip and observed under a phase contrast microscope with a $\times 100$ objective lens with oil emulsion.

Results

Out of $n=206$ (100%) of the subjects sampled, $n=97$ (47.1%) were male and $n=109$ (52.9%) were female. In total, 68 (33.0%) of the children studied were within the age bracket of 6–12 months, 47 (22.8%) were between the ages of 13–24 months, 46 (22.3%) were between the ages of 25–36 months, 17 (8.3%) were between

the ages of 37 and 48 months and 28 (13.6%) were between the ages of 49 and 60 months. Out of the total number of subjects ($n=206$) sampled, 20.39% ($n=42$) were found to be carriers of SPn; 52% ($n=22$) of the SPn carriers had received the recommended dose of PCV-10 immunization, while 48% ($n=20$) had not. All isolates ($n=42$; 20% of subjects) contained non-vaccine-type SPn serotypes, while $n=1$ (0.49% of the subjects) of the serotypes were untypeable (Table 1). In total, 18 different SPn serotypes were found in this population. They include: 28F (8 instances), 6A (5 instances), 3 (4 instances), 23B (3 instances), 20 (3 instances), 23A (3 instances), 19B (2 instances), 17F (2 instances), 7C (2 instances), 11A (2 instances), 35F (1 instance), 15B (1 instance), untypeable (1 instance), 48 (1 instance), 35B (1 instance), 21 (1 instance), 39 (1 instance) and 13 (1 instance).

Various serotypes were found to be prevalent in different age groups. For instance, out of the 42 serotypes found, 9 (23.53%) were prevalent among children at 6–12 months of age ($n=16$). They include: 28F (4 instances), 11A (2 instances), 23A (2 instances), 3 (2 instances), 6A (2 instances), 17F (1 instance), 35F (1 instance), 7C (1 instance) and untypeable (1 instance). There were 7 (16.67%) serotypes prevalent among children at 13–24 months ($n=8$), including: 20 (2 instances), 21 (1 instance), 39 (1 instance), 28F (1 instance), 35B (1 instance), 17F (1 instance) and 13 (1 instance). There were 8 (19%) serotypes found among children of 25–36 months of age ($n=12$), including: 23B (3 instances), 19B (2 instances), 3 (2 instances), 20 (1 instance), 28F (1 instance), 7C (1 instance), 23A (1 instance) and 48 (1 instance). There were 3 (7%) serotypes prevalent among children at 37–48 months old ($n=4$), including: 6A (2 instances), 15B (1 instance) and 28F (1 instance).

There were 2 (4.76% of the total) serotypes prevalent among children at 49–60 months ($n=2$): 6A (1 instance) and 28F (1 instance) (Table 2). Out of the 42 isolates (found in 20.39% of subjects), serotype 28F was the most prevalent (3.88% of the total), followed by 6A (2.43%), 3 (1.94%) and 20, 23A and 23B all at 1.46% ($n=3$). Each of the serotypes 7C, 11A, 17F and 19B represented 0.97% ($n=2$) of the total serotypes, while serotypes: 13, 21, 39, untypeable, 48, 15B, 35B and 35F represented 0.49% ($n=1$) each of the total serotypes found (Figure 1 and Figure 2). In total 51% ($n=106$) of the total sampled subjects were confirmed to have received a full dose of the PCV-10

Table 1. Overall *Streptococcus pneumoniae* (SPn) carriage of vaccine type and non-vaccine type serotypes. The percentage of SPn carriage status among PCV-10 vaccinated and unvaccinated children is shown.

	All children		Vaccinated children		Unvaccinated children	
	n	%	n	%	n	%
Overall SPn carriage	42	20.39	22	10.68	20	9.71
Proportion of SPn serotypes						
PCV-10	0	0.00	0	0.00	0	0.00
Non-PCV-10 serotypes	41	19.90	41	19.90	41	19.90
Non-typeable	1	0.49	1	0.49	1	0.49

Table 2. *Streptococcus pneumoniae* (SPn) Serotype distribution by age. The SPn serotypes as found among PCV-10 vaccinated and unvaccinated children of varying age groups is shown.

Variable	Subjects					
	All subjects	6–12 months	13–24 months	25–36 months	37–48 months	49–60 months
Numbers with carriage, n	42	16	8.00	12	4	2
Carriage, %	20.39	23.53	17.02	26.09	23.53	7.14
Total different serotypes, n	18	9	7	8	3	2
Serotypes seen (n)	28F (8)	28F (4)	20 (2)	23B (3)	6A (2)	6A (1)
	6A (5)	11A (2)	21 (1)	19B (2)	15B (1)	28F (1)
	3 (4)	23A (2)	39 (1)	3 (2)	28F (1)	
	23B (3)	3 (2)	28F (1)	20 (1)		
	20 (3)	6A (2)	35B (1)	28F (1)		
	23A (3)	17F (1)	17F (1)	7C (1)		
	19B (2)	35F (1)	13 (1)	23A (1)		
	17F (2)	7C (1)		48 (1)		
	7C (2)	Untypeable (1)				
	11A (2)					
	35F (1)					
	15B (1)					
	untypeable (1)					
	48 (1)					
	35B (1)					
	21 (1)					
	39 (1)					
	13 (1)

Figure 1. Serotype distribution in the *Streptococcus pneumoniae* (SPn) isolates. This figure shows the prevalence of various SPn serotypes among PCV-10-vaccinated and –unvaccinated children.

Figure 2. *Streptococcus pneumoniae* (SPn) serotype distribution by PCV-10 vaccination status.

vaccination as per the recommended schedule of immunization at 6, 10 and 14 weeks. Approximately 11% (n=12) of the immunized children were carriers of SPn in their nasopharyngeal region; 10% (n=10) of the non-immunized group were also carriers (Table 3). Serotypes 28F (5 instances), 23A (3 instances), 6A (3 instances), 17F (2 instances), 11A (1 instance), 3 (1 instance), 35F (1 instance), 48 (1 instance), 13 (1 instance), 35 (1 instance) and 7C (1 instance) were prevalent among the 9.71% (n=20) of the total sample group that had not received PCV-10 immunization. Serotypes 3 (3 instances), 28F (3 instances), 23B (3 instances), 20 (3 instances), 19B (2 instances), 6A (2 instances), 21 (1 instance), 11A (1 instance), 7C (1 instance), untypeable (1 instance), 15B (1 instance) and 39 (1 instance) were prevalent among the 10.68% (n=22) of the total sample group that received immunization (Table 4).

Dataset 1. List of basic demographic information for each subject, with the size of the optochin clearance zone and serotype of *Streptococcus pneumoniae*, if found

<http://dx.doi.org/10.5256/f1000research.14387.d207458>
(Walekhwa *et al.*, 2018).

Discussion

This study found that 20.39% of all children studied, from both the PCV-10 vaccinated and unvaccinated groups, were carriers of SPn. While this is a significant reduction from the pre-vaccine era, it is still high compared to malaria, diarrhea and HIV/AIDS (Feikin *et al.*, 2010). In total, n=41 of the serotypes found were non-vaccine type (in 19.90% of the subjects), with one additional untypeable serotype. This is a very important finding as it explains the high level of child morbidity and mortality due to pneumococcal disease despite the availability of PCV-10.

While these findings agree partially with those of Jacobs *et al.* (2008), where a significant decrease in the vaccine type SPn serotypes found in isolates was observed, a 97.6% (n=41) decrease is, at the very least, surprising. This trend may be attributed to the increased level of antimicrobial misuse by a greater percentage of the study population (Domenech de Cellès *et al.*, 2011). 10-valent pneumococcal conjugate vaccine contains 10 different serotypes, which include: 1, 4, 5, 6B, 7F, 9V, 14, 18C, 19F, 23F (Slotved *et al.*, 2016). None of these 10 serotypes was found in the study population yet this is the vaccine currently included in KEPI, targeting the same population.

Streptococcus pneumoniae carriage decreased with age as 11.65% (n=24) were obtained from children aged between 6–24 months and 8.74% (n=18) from children >24 months. The study results demonstrated a linear relationship between child age and SPn carriage. A similar study done elsewhere reported findings that partly agree with this and partly disagrees (Hill *et al.*, 2008). The former being attributable to development of SPn-specific IgG antibodies due to vaccination and during that window before most children start attending school (Corcadden *et al.*, 2013). Unlike findings from a study by de Paz *et al.* (2015), serotype 28F was the most prevalent and was present in all five age groups profiled. This is a likely scenario of serotype replacement as SPn attempts to evade the action of the immune system and eventually shares the resistant genes within the microbial community, especially in the nasopharyngeal region (Donati *et al.*, 2010).

Serotypes 28F, 6A, 11A, 3 and 7C were prevalent in both vaccinated and unvaccinated children, whereas serotypes 23A, 17F, 35F, 48, 13, 35B and 23B, 20, 19B, 21, untypeable, 15B, 39 were found among unvaccinated and vaccinated groups respectively.

Table 3. *Streptococcus pneumoniae* (SPn) serotype distribution by PCV-10 vaccination status. The percentage prevalence of SPn serotypes among PCV-10 vaccinated and unvaccinated children is shown.

Unvaccinated (n=100)			Vaccinated (n=106)		
Serotype	n	%	Serotype	n	%
28F	5	5	3	3	2.83
23A	3	3	28F	3	2.83
6A	3	3	23B	3	2.83
17F	2	2	20	3	2.83
11A	1	1	19B	2	1.89
3	1	1	6A	2	1.89
35F	1	1	21	1	0.94
48	1	1	11A	1	0.94
13	1	1	7C	1	0.94
35B	1	1	Untypeable	1	0.94
7C	1	1	15B	1	0.94
			39	1	0.94

There exists different antigenic features between and within various strains of SPn (Song *et al.*, 2012). While the majority, if not all, pneumococcal serotypes are capable of causing disease, the frequency with which they are isolated varies (Kalin, 1998). In this case, vaccination would only be partially effective and, if so, due to inter-strain antigenic characteristics.

While trying to evade the action of the immune system, SPn has a tendency to exchange resistant genes and other antigenic correlates at the nasopharyngeal region (Johnston *et al.*, 2014). Resistance to antimicrobial agents is occasioned by among other factors, misuse of antibiotics (Dinsbach, 2012). This is largely due to a lack of properly enforced antibiotic use regulations by the authorities.

Table 4. Pneumococcal carriage by vaccination status.

Child immunization status	SPn	n	%
Immunized	NGR	84	40.78
	SPn	22	10.68
Not immunized	NGR	80	38.83
	SPn	20	9.71
Total	N/A	206	N/A

NGR, no growth observed; SPn, *Streptococcus pneumoniae*; NA, not applicable.

Data availability

Dataset 1. List of basic demographic information for each subject, with the size of the optochin clearance zone and serotype of *Streptococcus pneumoniae*, if found. DOI: <http://doi.org/10.5256/f1000research.14387.d207458> (Walekhwa *et al.*, 2018).

Competing interests

No competing interests were disclosed.

Grant information

The study was in part funded by the National Commission of Science Technology & Innovation (NACOSTI) Kenya. Ref Nos. NACOSTI/RCD/ST & I/7TH CALL/PhD/148.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgements

Special thanks to Prof. Anthony Scott, Dr. Claire Gordon and Ms. Angela Karani of Kemri Wellcome Trust, Kilifi for their professional input. The laboratory and technical team: Ann Karanu, Elizabeth Mbithe, Shadrack Mutua and Alfred Too. Your technical input was amazing! The Gertrude's Hospital clinical and laboratory team (Linet Okose, Barasa Kasili, Charles Muia, Lilive Njagi, Carolyn Thumbi, and Peninah Chege), you made this possible. Thanks to Japheth Wambani Rapando for the review of my work.

References

- Agweyu A, Kibore M, Digolo L, *et al.*: **Prevalence and correlates of treatment failure among Kenyan children hospitalised with severe community-acquired pneumonia: a prospective study of the clinical effectiveness of WHO pneumonia case management guidelines.** *Trop Med Int Health*. 2014; **19**(11): 1310–20.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Castañeda-Orjuela C, Alvis-Guzmán N, Velandia-González M, *et al.*: **Cost-effectiveness of pneumococcal conjugate vaccines of 7, 10, and 13 valences in Colombian children.** *Vaccine*. 2012; **30**(11): 1936–1943.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Chow S, Wang H, Shao J: **Sample Size Calculations in Clinical Research.** New York. 2nd ed. Taylor & Francis Group. 2007.
[Reference Source](#)
- Colglazier W: **SUSTAINABILITY. Sustainable development agenda: 2030.** *Science*. 2015; **349**(6252): 1048–50.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Corscadden KJ, Kirkham LA, Thornton RB, *et al.*: **High pneumococcal serotype specific IgG, IgG1 and IgG2 levels in serum and the middle ear of children with recurrent acute otitis media receiving ventilation tubes.** *Vaccine*. 2013; **31**(10): 1393–1399.
[PubMed Abstract](#) | [Publisher Full Text](#)
- de Paz HD, Selva L, Muñoz-Almagro C: **Pneumococcal Vaccination and Consequences.** *Streptococcus Pneumoniae*. 2015; 41–57.
[Publisher Full Text](#)
- Domenech de Cellès M, Opatowski L, Salomon J, *et al.*: **Intrinsic epidemicity of *Streptococcus pneumoniae* depends on strain serotype and antibiotic susceptibility pattern.** *Antimicrob Agents Chemother*. 2011; **55**(11): 5255–5261.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Dinsbach NA: **Antibiotics in dentistry: Bacteremia, antibiotic prophylaxis, and antibiotic misuse.** *Gen Dent*. 2012; **60**(3): 200–207; quiz 208–9.
[PubMed Abstract](#)
- Donati C, Hiller NL, Tettelin H, *et al.*: **Structure and dynamics of the pan-genome**

of *Streptococcus pneumoniae* and closely related species. *Genome Biol.* 2010; 11(10): R107.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Feikin DR, Jagero G, Aura B, *et al.*: High rate of pneumococcal bacteremia in a prospective cohort of older children and adults in an area of high HIV prevalence in rural western Kenya. *BMC Infect Dis.* 2010; 10: 186.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Feldman C, Anderson R: Review: Current and new generation pneumococcal vaccines. *J Infect.* 2014; 69(4): 309–25.

[PubMed Abstract](#) | [Publisher Full Text](#)

Hackel M, Lascols C, Bouchillon S, *et al.*: Serotype prevalence and antibiotic resistance in *Streptococcus pneumoniae* clinical isolates among global populations. *Vaccine.* 2013; 31(42): 4881–4887.

[PubMed Abstract](#) | [Publisher Full Text](#)

Hammit LL, Akech DO, Morpeth SC, *et al.*: Population effect of 10-valent pneumococcal conjugate vaccine on nasopharyngeal carriage of *Streptococcus pneumoniae* and non-typeable *Haemophilus influenzae* in Kilifi, Kenya: Findings from cross-sectional carriage studies. *Lancet Glob Health.* 2014; 2(7): e397–e405.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Harboe ZB, Thomsen RW, Riis A, *et al.*: Pneumococcal serotypes and mortality following invasive pneumococcal disease: A population-based cohort study. *PLoS Med.* 2009; 6(5): e1000081.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Hill PC, Cheung YB, Akisanya A, *et al.*: Nasopharyngeal Carriage of *Streptococcus pneumoniae* in Gambian Infants: A Longitudinal Study. *Clin Infect Dis.* 2008; 46(6): 807–814.

[PubMed Abstract](#) | [Publisher Full Text](#)

Jacobs MR, Good CE, Beall B, *et al.*: Changes in serotypes and antimicrobial susceptibility of invasive *Streptococcus pneumoniae* strains in Cleveland: a quarter century of experience. *J Clin Microbiol.* 2008; 46(3): 982–990.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Johnson HL, Deloria-Knoll M, Levine OS, *et al.*: Systematic evaluation of serotypes causing invasive pneumococcal disease among children under five: The pneumococcal global serotype project. *PLoS Med.* 2010; 7(10): pii: e1000348.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Johnston C, Campo N, Bergé MJ, *et al.*: *Streptococcus pneumoniae*, le transformiste. *Trends Microbiol.* 2014; 22(3): 113–9.

[PubMed Abstract](#) | [Publisher Full Text](#)

Jones RN, Jacobs MR, Sader HS: Evolving trends in *Streptococcus pneumoniae* resistance: Implications for therapy of community-acquired bacterial pneumonia. *Int J Antimicrob Agents.* 2010; 36(3): 197–204.

[PubMed Abstract](#) | [Publisher Full Text](#)

Kalin M: Pneumococcal serotypes and their clinical relevance. *Thorax.* 1998;

53(3): 159–162.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Kellogg JA, Bankert DA, Elder CJ, *et al.*: Identification of *Streptococcus pneumoniae* revisited. *J Clin Microbiol.* 2001; 39(9): 3373–3375.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Mitchell AM, Mitchell TJ: *Streptococcus pneumoniae*: virulence factors and variation. *Clin Microbiol Infect.* 2010; 16(5): 411–8.

[PubMed Abstract](#) | [Publisher Full Text](#)

Moffitt KL, Malley R: Next generation pneumococcal vaccines. *Curr Opin Immunol.* 2011; 23(3): 407–13.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

O'Brien KL, Wolfson LJ, Watt JP, *et al.*: Burden of disease caused by *Streptococcus pneumoniae* in children younger than 5 years: global estimates. *Lancet.* 2009; 374(9693): 893–902.

[PubMed Abstract](#) | [Publisher Full Text](#)

Slotved HC, Dalby T, Hoffmann S: The effect of pneumococcal conjugate vaccines on the incidence of invasive pneumococcal disease caused by ten non-vaccine serotypes in Denmark. *Vaccine.* 2016; 34(6): 769–774.

[PubMed Abstract](#) | [Publisher Full Text](#)

Song JH, Dagan R, Klugman KP, *et al.*: The relationship between pneumococcal serotypes and antibiotic resistance. *Vaccine.* 2012; 30(17): 2728–37.

[PubMed Abstract](#) | [Publisher Full Text](#)

Väkeväinen M, Soininen A, Lucero M, *et al.*: Serotype-specific hypo-responsiveness to pneumococcal conjugate vaccine in infants carrying pneumococcus at the time of vaccination. *J Pediatr.* 2010; 157(5): 778–83.e1.

[PubMed Abstract](#) | [Publisher Full Text](#)

Vandenbos F, Gal J, Radicchi B: [Vaccination coverage against influenza and pneumococcus for patients admitted to a pulmonary care service]. *Rev Mal Respir.* 2013; 30(9): 746–751.

[PubMed Abstract](#) | [Publisher Full Text](#)

Walekhwa M, Muturi M, Gunturu R, *et al.*: Dataset 1 in: *Streptococcus pneumoniae* serotype epidemiology among PCV-10 vaccinated and unvaccinated children at Gertrude's Children's Hospital, Nairobi County: a cross-sectional study. *F1000Research.* 2018.

[Data Source](#)

Weinberger DM, Harboe ZB, Flasche S, *et al.*: Prediction of serotypes causing invasive pneumococcal disease in unvaccinated and vaccinated populations. *Epidemiology.* 2011; 22(2): 199–207.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

WHO: Pneumonia Factsheet. *WHO Media Centre.* 2013; 1–2.

WHO & UNICEF: Global Action Plan for Prevention and Control of Pneumonia (GAPP) Technical Consensus statement. *B World Health Organ.* 2009; 86(5): 1–23.

[Reference Source](#)

Open Peer Review

Current Peer Review Status: ? ✓ ✗

Version 1

Reviewer Report 15 November 2018

<https://doi.org/10.5256/f1000research.15656.r39482>

© 2018 Dube F. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Felix Dube

Division of Medical Microbiology, Department of Pathology, University of Cape Town, Cape Town, South Africa

The study reports pneumococcal carriage in a cohort of Kenyan children. This has significant implication when it comes to the evaluation of the impact of PCV10 on the population structure of pneumococcus.

Specific comments:

1. Pneumococcus is a normal commensal, not "highly invasive" as is reported in intro.
2. The authors must avoid the use of non-standard nomenclature such as SPn. Further, there is inconsistent use of *S. pneumoniae* and pneumococcus throughout the text. These cannot be used interchangeably.
3. The introduction needs to be reworked and repetitions avoided, i.e. in paragraph 1 and 5, the authors talk about 90 serotypes. The last sentence of paragraph 5 does not include references.
4. The study was conducted between 2017 and 2018, it would really have benefited from the WHO working group report on pneumococcal carriage studies¹. Most importantly, 79% (164/206) being non-viable seriously indicates problems in the experimental design. The authors don't say anything about broth enrichment in order to improve recovery of the pneumococcus. Amies media is not ideal for pneumococcus compared to STGG. A 10ul inoculum is very little, did the authors attempt to use bigger volumes especially for the samples with no growth? A 2% BA plate is used as primary culture then selective media, if possible, do this in parallel.
5. Need to be more consistent in reporting proportions.
6. What was the PCV10 vaccine coverage at each timepoint? Also authors need to report non-

pcv as non-PCV10 because some of the serotypes the report as "non-vaccine" such as serotype 3 are included in PCV13.

7. Repetition of results. The authors do not report any metadata looking at risk factors for carriage, hence the "epi" in title must fall out.
8. The authors repeatedly report "decrease" in carriage post PCV, but they really can't say this without data on Pre-PCV10.
9. Avoid sweeping statement to infer serotype replacement if they actually don't show this evidence.
10. "While this is a significant reduction from the pre-vaccine era, it is still high compared to malaria, diarrhea and HIV/AIDS" doesn't make sense, what are the authors referring to?
11. "This is a very important finding as it explains the high level of child morbidity and mortality due to pneumococcal disease despite the availability of PCV-10." How do you arrive at this if working with carriage and not invasive disease isolates.
This and other strong conclusions need to be avoided, you surely cant with 42 isolates.

References

1. Satzke C, Turner P, Virolainen-Julkunen A, Adrian PV, et al.: Standard method for detecting upper respiratory carriage of *Streptococcus pneumoniae*: updated recommendations from the World Health Organization Pneumococcal Carriage Working Group. *Vaccine*. 2013; **32** (1): 165-79 [PubMed Abstract](#) | [Publisher Full Text](#)

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Partly

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Medical microbiology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to state that I do not consider it to be of an acceptable scientific standard, for reasons outlined above.

Reviewer Report 01 November 2018

<https://doi.org/10.5256/f1000research.15656.r39125>

© 2018 Obey J. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Jackie K. Obey

School of Health Sciences, Department of Medical Laboratory Sciences, University of Eastern Africa, Baraton, Eldoret, Kenya

The study carried out by the authors on *Streptococcus pneumoniae* is extremely important for Kenya. It addresses a major health problem that has been addressed by other authors in the past and for which a lasting solution is currently being sought. The study is detailed and was able to employ modern techniques to assess the carriage state of children at The Gertrude's Children Hospital, Nairobi, Kenya. The methods used were appropriate and in line with the study's objectives. The sample size was appropriately determined and descriptive statistics have been used appropriately to describe the results obtained by the authors.

The title of the study however includes the word 'epidemiology' and this may lead the reader to think that the authors would have tried to determine factors that influenced the establishment of the research problem at the study site. The authors have not determined those factors or risks that lead to Kenyan children being vaccinated, yet not being protected by the PCV-10 vaccine. Those findings would then have given an idea of the risk of exposure of children to *Streptococcus pneumoniae* disease and given the hospital and government the strategies to employ for preventive measures against the disease. The conclusion is brief and does not give recommendations to the Government of Kenya or The Gertrude's Children Hospital on what to do after the findings of the study.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Medicinal parasitology (Malariology), immunoparasitology and medical entomology, antimicrobial activity of plant extracts

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 25 October 2018

<https://doi.org/10.5256/f1000research.15656.r39126>

© 2018 Ondigo B. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Bartholomew N. Ondigo

Centre for Global Health Research, Kenya Medical Research Institute, Kisumu, Kenya

Title: To include Kenya

Corresponding author contacts need to be indicated

Abstract:

Background:

It needs to be indicated that circulating SPn serotypes are being investigated after the introduction of 10-valent pneumococcal conjugate vaccine (PCV10) in 2011.

The methods need to be a summary of the techniques used in data collection. As it is written it appears to be copy pasted from the method section. Indicate number of children assessed. For instance:

Materials:

Two hundred and six children attending and not-attending in20-2009 were studied. Materials for study were pharyngeal swabs and sputum. Identification was performed using optochin disks, Quellung reaction,.....agglutination on the glass, viewed under a phase contrast microscope and the sent to KEMRI- Kilifi for further confirmative identification tests.

The conclusion seem alarming and need to suggest or indicate. Therefore Kenyan children

receiving PCV-10 vaccine are not protected – Revise to something like:-
This study highlights the importance of monitoring and evaluation to provide epidemiological information to determine the effectiveness of PCV10 in Kenya's Public health services.

Repeating ideas should be deleted - causing more deaths than any other infectious disease vs. kills more children than any other illness in the world.

Introduction need to be shortened, preferably to three paragraphs.

Methods:

Were the children admitted or outpatient?

All the source of equipment used and consumables need to be indicated, for instance incubator etc.

The Research Ethics Committee that approved the study need to be indicated.

Software used for calculation of %s need to be indicated?

Results:

Headings need to be introduced that are in agreement with the content. This will help the reader to when reading. Suggested possible headings include:

Demography of the Study Participants

Prevalence of SPn carriage status among PCV-10 vaccinated and unvaccinated children

Prevalence of SPn carriage status by age – You probably have several age groups on this for instance <1, 2 -4 years, 4 – 5 years.

Authors need to clarify on the following:

How many pneumococcus serotypes were identified? Which serotype was most frequent?

Can you please arrange them in a descending order?

Discussion:

Adequate in content.

Consider serotype replacement to enrich your discussion.

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

No

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Immunoparasitology

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 06 Jan 2019

Michael Walekhwa, Gertrude's Children's Hospital, Nairobi, Kenya

Dear Dr. Ondigo,

Many thanks for your review of this article.

I have read through and keenly updated the areas you highlighted during your review.
Kindly revisit.

Many thanks

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research