REVIEW

Eating disorders in children: is avoidant-restrictive food intake disorder a feeding disorder or an eating disorder and what are the implications for treatment? [version 1; peer review: 2 approved]

Grace A. Kennedy , Madeline R. Wick, Pamela K. Keel

Department of Psychology, Florida State University, 1107 West Call Street, Tallahassee, FL, 32304, USA

V1 First published: 18 Jan 2018, 7(F1000 Faculty Rev):88
<https://doi.org/10.12688/f1000research.13110.1>

Latest published: 18 Jan 2018, 7(F1000 Faculty Rev):88
<https://doi.org/10.12688/f1000research.13110.1>

Abstract

Avoidant-restrictive food intake disorder (ARFID) is a current diagnosis in the "Feeding and Eating Disorders" section of the *Diagnostic and Statistical Manual of Mental Disorders* (fifth edition) and captures a heterogeneous presentation of eating disturbances. In recent years, ARFID has been studied primarily within the context of eating disorders despite having historical roots as a feeding disorder. The following review examines ARFID's similarities with and differences from feeding disorders and eating disorders, focusing on research published within the last three years. Implications of this differentiation for treatment are discussed.

Keywords

eating disorders, Avoidant-Restrictive Food Intake Disorder, feeding disorders

Open Peer Review

Approval Status

	1	2
version 1
18 Jan 2018		

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Jennifer J Thomas**, Harvard Medical School, Boston, USA
Massachusetts General Hospital, Boston, USA
2. **Timothy D Brewerton**, Medical University of South Carolina, Charleston, USA

Any comments on the article can be found at the end of the article.

Corresponding author: Pamela K. Keel (keel@psy.fsu.edu)

Author roles: **Kennedy GA:** Conceptualization, Writing – Original Draft Preparation; **Wick MR:** Conceptualization, Resources, Writing – Review & Editing; **Keel PK:** Conceptualization, Supervision, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The authors declare that this work was funded by National Institutes of Mental Health (R01 MH 111263).
The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2018 Kennedy GA *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Kennedy GA, Wick MR and Keel PK. **Eating disorders in children: is avoidant-restrictive food intake disorder a feeding disorder or an eating disorder and what are the implications for treatment?** [version 1; peer review: 2 approved]
F1000Research 2018, 7(F1000 Faculty Rev):88 <https://doi.org/10.12688/f1000research.13110.1>

First published: 18 Jan 2018, 7(F1000 Faculty Rev):88 <https://doi.org/10.12688/f1000research.13110.1>

Introduction and context

Avoidant-restrictive food intake disorder (ARFID) is a diagnosis in the “Feeding and Eating Disorders” section of the fifth edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-5)¹. ARFID is a reformulation of the DSM-IV diagnosis of feeding disorder of infancy or early childhood (FDIEC)² expanded to capture heterogeneous eating disturbances that lead to deficits in energy or nutritional intake. Reasons for insufficient intake include lack of appetite, the sensory characteristics of food, and fears related to the act of eating, such as fear of choking¹. These disturbances can confer harmful physiological or psychological consequences, including low weight or failure to grow, severe nutritional insufficiency, reliance on nutritional supplements, and psychosocial dysfunction¹.

Following ARFID’s introduction to the DSM-5 in 2013, an emerging line of research has involved descriptive studies comparing ARFID with eating disorders, such as anorexia nervosa (AN) and bulimia nervosa (BN)^{3–5}. Furthermore, a plenary session on atypical eating disorders at the 2017 International Conference on Eating Disorders in Prague included two talks on ARFID^{6,7}. This emerging body of work suggests that ARFID may be viewed as an eating disorder by the eating disorder research community. However, examination of ARFID’s history in the DSM along with the history of eating disorders in the DSM suggests that ARFID may represent a feeding disorder. Like children with ARFID, those with FDIEC displayed persistent failure to eat, leading to weight loss or failure to grow. The FDIEC diagnosis differed from the ARFID diagnosis in that FDIEC could be diagnosed only with an age of onset by six years old and did not account for other consequences of food restriction. Thus, the diagnosis of FDIEC was rarely used and research on its presentation was stagnant⁸. Given the limited interest and utility of the diagnosis^{2,8}, the DSM-5 Eating Disorders Work Group eliminated FDIEC and replaced it with ARFID, which includes all cases previously diagnosed with FDIEC along with cases that would not have met the age or weight consequences criteria².

Further supporting ARFID’s history as a feeding disorder, FDIEC fell under the umbrella of “Disorders Primarily Occurring in Infancy, Early Childhood, and Adolescence” in the DSM-IV, alongside pica and rumination disorder, in a section that was distinct from “Eating Disorders”⁹. Interestingly, in the DSM-III-R, pica, rumination disorder, AN, and BN were all included under the umbrella of “Eating Disorders”, but it was noted that “Pica and Rumination Disorder of Infancy are [...] probably unrelated to Anorexia Nervosa and Bulimia Nervosa” (10, page 65). This may have presaged the separation of AN and BN into their own separate section as “Eating Disorders” in the DSM-IV, suggesting that FDIEC, pica, and rumination disorder were all conceptualized as feeding disorders, even though only FDIEC was explicitly labeled as such. Moreover, the decision to unite ARFID, pica, and rumination disorder with AN, BN, and binge eating disorder in the DSM-5 in a section named “Feeding and Eating Disorders” suggests that some of these conditions are feeding disorders and some are eating disorders. Yet the DSM-5 provides no guidance on which diagnoses belong to which group.

According to the National Institute of Mental Health¹¹, “Eating disorders are [...] serious and often fatal illnesses that cause severe disturbances to a person’s eating behaviors. Obsessions with food, body weight, and shape may also signal an eating disorder”. No equivalent definition for “feeding disorder” is presented; however, Sharp *et al.*¹² defined feeding disorders as “severe disruptions in nutritional and caloric intake exceeding ordinary variations in hunger, food preference, and/or interest in eating” (page 116). Understanding ARFID’s place in the nomenclature is crucial for informing future research and treatment of the disorder. Thus, the following review examines the evidence for ARFID as a feeding disorder and as an eating disorder, focusing on research published within the last three years. We then discuss implications of this differentiation for treatment.

Avoidant-restrictive food intake disorder as a feeding disorder

Within the past three years, ARFID has been studied largely within the context of eating disorders. Thus, ARFID’s similarity to posited feeding disorders, such as pica or rumination disorder, is discussed largely in terms of its dissimilarity to eating disorders. For instance, in a retrospective chart review of 7- to 17-year-olds admitted to a day program for eating disorders, Nicely *et al.*¹³ found that compared to children with AN, BN, and other eating disorders, those with ARFID were more likely to have a comorbid diagnosis of autism spectrum disorder (ASD), learning disorder, or cognitive impairment¹³. This is similar to findings primarily in pica in which common comorbidities are ASD or intellectual disability or both¹. Furthermore, individuals with ASD often demonstrate restrictive food selectivity due to sensory sensitivity to food texture and smell, which is consistent with ARFID¹⁴. For some individuals with ASD, rigid food preferences may contribute to nutritional deficits and require clinical attention, warranting a separate diagnosis of ARFID. Indeed, Lucarelli *et al.*¹⁵ presented a case study of a 4-year-old girl with ASD who presented to treatment for ARFID because of rigid food and meal rules¹⁵. In this way, ARFID resembles other feeding disorders in which neurodevelopmental comorbidity may play a role in presentation. Yet this may not be unique to ARFID given growing research suggesting that individuals with AN are higher on autistic traits than healthy controls^{16,17}. Thus, while ASD does have a high comorbidity with feeding problems related to sensory characteristics of food and rigid food preferences independent of weight concerns, the broader autistic phenotype may be over-represented in AN and ARFID, suggesting a potential link between ASD and restrictive eating and feeding disorders.

With regard to treatment, studies have repeatedly shown that children with ARFID enter care at a younger age and are less likely to be self-referred compared with children with eating disorders^{3,4,13,18,19}. In line with case reports, this suggests that ARFID onsets earlier and is more likely to be recognized as a problem by parents than are eating disorders^{15,20,21}. Indeed, numerous case reports and retrospective chart review studies have documented that individuals with ARFID enter care between the ages of 4 and 13^{3,4,22}, suggesting similarity to the classic conceptualization of feeding disorders as a childhood problem⁹.

In terms of clinical presentation, individuals with ARFID exhibit other notable differences from those with AN, BN, and their variants. By definition, ARFID may be diagnosed only in the absence of weight or shape concerns. Thus, as in pica and rumination disorder, the driving force behind feeding disturbances cannot be cognitive concerns related to the effects of food on appearance. In fact, previous research demonstrates that children presenting with ARFID often desire to gain weight rather than lose it¹⁹. These findings may explain the multiple studies documenting a greater representation of males among those with ARFID than among those with AN or BN^{3,4,13}. It may also explain the greater use and acceptance of nasogastric feeding in ARFID compared with AN. Both Peebles *et al.*²³ and Ornstein *et al.*¹⁸ have noted that children with ARFID require nasogastric feeding more often than those with AN but demonstrate less distress regarding this process. Patients with AN fear caloric content, which makes the route of caloric intake less relevant. In contrast, those with ARFID often fear choking during the process of eating but express notably less anxiety about taking in calories, making the route of intake highly relevant and the actual caloric content less relevant^{18,23}.

Taken together, these data suggest that ARFID is distinct from eating disorders on patterns of comorbidity, age of presentation, referral source, gender distribution, and response to intervention modalities in ways that make it more similar to pica and rumination disorder. This conclusion should be interpreted with some caution, however, given the paucity of research on pica or rumination disorder or research comparing these conditions with ARFID.

Avoidant-restrictive food intake disorder as an eating disorder

With regard to similarities to eating disorders, characteristics shared with AN have been identified. Like AN, ARFID often presents with low weight and malnutrition, thus requiring a multidisciplinary team, including family members, physicians, mental health professionals, and dietitians, to restore weight. Of note, a multidisciplinary team of providers and family members has been presented as the best practice for all feeding disorders¹², highlighting a broader overlap between feeding and eating disorders when they occur in children. For instance, Peebles *et al.*²³ described an inpatient refeeding protocol that could be employed for children and adolescents with either AN or ARFID²³. This protocol used a meal plan with increasingly higher caloric content and heavy involvement of parents in the refeeding process both during inpatient hospitalization and in outpatient aftercare. Furthermore, Forman *et al.*⁴ found that treatment of ARFID did not differ from that of AN or atypical AN in a large retrospective chart review of 700 adolescents attending eating disorder programs across 14 different sites⁴. Importantly, however, treatments differed considerably by site, such that site, rather than diagnosis, may have been the main determinant of what treatments were offered. Furthermore, at one-year follow-up, AN and ARFID did not differ significantly in likelihood of median body mass index being at least 90% expected; however, this may have reflected limited statistical power, as this outcome was twice as likely for AN patients compared with ARFID patients. Nonetheless, treatments of

ARFID and AN in children share components because of shared medical concerns and utility of involving parents in the care of their child's illness.

Much like children and adolescents with AN, those with ARFID present with strict food rules about what they can and cannot eat^{15,21}. Food exposure to reduce avoidance of feared foods is used in the treatment of both AN and ARFID²³. Furthermore, both retrospective chart reviews and case reports illustrate that some individuals with ARFID transition to a diagnosis of AN once weight is partially restored^{22,24}. In a chart review of 205 patients presenting to care for an eating disorder, Norris *et al.*²⁴ found that 12% of patients with ARFID transitioned to a diagnosis of AN during treatment, suggesting that a diagnosis of ARFID may serve as a risk factor for the development of AN. Case reports presented by Maertens *et al.*²² suggest that weight concerns may emerge after refeeding in some individuals with comorbid ARFID and obsessive-compulsive disorder, resulting in the later diagnosis of AN that was not evident when patients were very underweight.

Another consideration is the overlap between ARFID and non-fat phobic AN (NFP-AN). Individuals with NFP-AN restrict energy intake, resulting in low weight; however, they deny fear of fat or weight gain. In a review on NFP-AN, Becker *et al.*²⁵ described cases whereby restriction of food intake was due to religiosity and gastrointestinal discomfort as well as a lack of appetite. Given that DSM-5 AN can be diagnosed in an individual who engages in behaviors to prevent weight gain and does not recognize the seriousness of low weight but reports no other body image disturbance, some cases of ARFID bear a strong resemblance to NFP-AN. Unlike those with ARFID, however, those with NFP-AN are averse to weight gain, which is an important differentiation between the two.

In summary, despite ARFID's differentiations from eating disorders, it demonstrates key similarities to AN in regard to presenting symptoms of underweight and rigid food rules, potential to transition to endorsing concerns about shape and weight as motivating food restriction, and treatment. These similarities may explain why ARFID has been compared with eating disorders in several recent studies and why it was included in talks focusing on atypical eating disorders.

Conclusions and future directions

Research on ARFID represents an emerging field of study. ARFID resembles a feeding disorder in demographic features, comorbidity, source of presentation, and greater acceptance of invasive treatments. Conversely, ARFID resembles AN in children in terms of management and treatment of the illness and shares similar presentations with NFP-AN in some cases. Moreover, ARFID appears to precede AN in a substantial minority (12%) of patients with ARFID. Thus, rather than revealing whether ARFID is a feeding disorder or an eating disorder, this review suggests ways in which ARFID may be conceptualized as *both* a feeding and an eating disorder and proposes that the similarity and emphasis depend on clinical presentation. Indeed, those with ARFID who present with sensory sensitivities (for example, rejecting food on the basis of color or texture) may have more in common with pica

or rumination disorder, as sensory stimulation may contribute to aberrant eating behaviors in all three disorders^{26,27}. It is possible that underlying deficits in sensation processes are at play in pica and rumination disorder and, in a subset of cases, ARFID. In contrast, ARFID associated with intense fear of vomiting and gastrointestinal distress may have greater similarity to the conceptualization of NFP-AN. Migration from ARFID to AN underscores the fluidity of the distinction between ARFID and AN, which may reflect developmental shifts as well as changes in weight over the course of development and treatment.

Given ARFID's history as a feeding disorder, it is surprising that the focus of recent research has been primarily within the field of eating disorders; however, there are exceptions to this. Sharp *et al.*¹² suggested that ARFID is the psychiatric disorder that best captures feeding disorder presentations and offered a review of studies on techniques for treating ARFID-like feeding disturbances in children. Furthermore, in addition to being referred to eating disorder programs, patients with ARFID are referred to feeding disorder programs and gastroenterology clinics^{28,29}. Thus, greater communication between those who approach ARFID from a feeding disorder perspective and those who approach ARFID from an eating disorder perspective may be beneficial in both the treatment and the understanding of ARFID. Indeed, to this end, Thomas *et al.*³⁰ have drawn on the knowledge base of feeding

and eating disorders to put forth a conceptual neurobiological model of ARFID and a potential avenue for treatment enhancement.

In summary, ARFID possesses characteristics relevant to both feeding and eating disorders and may represent the missing link between these differing types of disorders. As such, research on ARFID may yield insights into underlying mechanisms contributing to disturbances in eating across the range of these disorders. More information on ARFID from both feeding and eating disorder perspectives could improve greater understanding of not only ARFID but also additional feeding or eating disorders to which ARFID has similarities.

Competing interests

The authors declare that they have no competing interests.

Grant information

The authors declare that this work was funded by National Institutes of Mental Health (R01 MH 111263).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- American Psychiatric Association: **Diagnostic and statistical manual of mental disorders (5th ed.)**. Arlington, VA: American Psychiatric Association. 2013.
[Publisher Full Text](#)
- Attia E, Becker AE, Bryant-Waugh R, *et al.*: **Feeding and eating disorders in DSM-5**. *Am J Psychiatry*. 2013; **170**(11): 1237–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Fisher MM, Rosen DS, Ornstein RM, *et al.*: **Characteristics of avoidant/restrictive food intake disorder in children and adolescents: a “new disorder” in DSM-5**. *J Adolesc Health*. 2014; **55**(1): 49–52.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Forman SF, McKenzie N, Hehn R, *et al.*: **Predictors of outcome at 1 year in adolescents with DSM-5 restrictive eating disorders: report of the national eating disorders quality improvement collaborative**. *J Adolesc Health*. 2014; **55**(6): 750–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Hay P, Mitchison D, Collado AEL, *et al.*: **Burden and health-related quality of life of eating disorders, including Avoidant/Restrictive Food Intake Disorder (ARFID), in the Australian population**. *J Eat Disord*. 2017; **5**: 21.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Kreipe RE: **Seeing Atypical Eating Disorders in Young Children: Identification and Treatment of Atypical Eating Disorders in Clinic**. Oral Presentation at International Conference for Eating Disorders Prague, Czech Republic. 2017.
- Eddy K: **Looking for signs and not symptoms: Improving precision in the measurement of eating disorders**. Oral Presentation at International Conference for Eating Disorders Prague, Czech Republic. 2017.
- Zimmerman J, Fisher M: **Avoidant/Restrictive Food Intake Disorder (ARFID)**. *Curr Probl Pediatr Adolesc Health Care*. 2017; **47**(4): 95–103.
[PubMed Abstract](#) | [F1000 Recommendation](#)
- American Psychiatric Association: **Diagnostic and statistical manual of mental disorders (4th ed.)**. Washington, DC: American Psychiatric Association. 1994.
[Reference Source](#)
- American Psychiatric Association: **Diagnostic and statistical manual of mental disorders (3rd ed., text revision)**. Washington, DC: American Psychiatric Association. 1987.
- National Institute of Mental Health: **Eating Disorders**. 2016; Retrieved December 19, 2017.
[Reference Source](#)
- Sharp WG, Volkert VM, Scallan L, *et al.*: **A Systematic Review and Meta-Analysis of Intensive Multidisciplinary Intervention for Pediatric Feeding Disorders: How Standard Is the Standard of Care?** *J Pediatr*. 2017; **181**: 116–124.e4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Nicely TA, Lane-Loney S, Masciulli E, *et al.*: **Prevalence and characteristics of avoidant/restrictive food intake disorder in a cohort of young patients in day treatment for eating disorders**. *J Eat Disord*. 2014; **2**(1): 21.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Cermak SA, Curtin C, Bandini LG: **Food selectivity and sensory sensitivity in children with autism spectrum disorders**. *J Am Diet Assoc*. 2010; **110**(2): 238–46.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Lucarelli J, Pappas D, Welchons L, *et al.*: **Autism Spectrum Disorder and Avoidant/Restrictive Food Intake Disorder**. *J Dev Behav Pediatr*. 2017; **38**(1): 79–80.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Baron-Cohen S, Jaffa T, Davies S, *et al.*: **Do girls with anorexia nervosa have elevated autistic traits?** *Mol Autism*. 2013; **4**(1): 24.
[PubMed Abstract](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Tchanturia K, Smith E, Weineck F, *et al.*: **Exploring autistic traits in anorexia: a clinical study**. *Mol Autism*. 2013; **4**(1): 44.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ornstein RM, Essayli JH, Nicely TA, *et al.*: **Treatment of avoidant/restrictive food intake disorder in a cohort of young patients in a partial hospitalization program for eating disorders**. *Int J Eat Disord*. 2017; **50**(9): 1067–74.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Strandjord SE, Sieke EH, Richmond M, *et al.*: **Avoidant/Restrictive Food Intake Disorder: Illness and Hospital Course in Patients Hospitalized for Nutritional Insufficiency**. *J Adolesc Health*. 2015; **57**(6): 673–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)

20. Bryant-Waugh R: **Avoidant restrictive food intake disorder: an illustrative case example.** *Int J Eat Disord.* 2013; **46**(5): 420–3.
[PubMed Abstract](#) | [Publisher Full Text](#)
21. Katzman DK, Stevens K, Norris M: **Redefining feeding and eating disorders: What is avoidant/restrictive food intake disorder?** *Paediatr Child Health.* 2014; **19**(8): 445–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
22. Maertens C, Couturier J, Grant C, *et al.*: **Fear of Vomiting and Low Body Weight in Two Pediatric Patients: Diagnostic Challenges.** *J Can Acad Child Adolesc Psychiatry.* 2017; **26**(1): 59–61.
[PubMed Abstract](#) | [Free Full Text](#) | [F1000 Recommendation](#)
23. Peebles R, Lesser A, Park CC, *et al.*: **Outcomes of an inpatient medical nutritional rehabilitation protocol in children and adolescents with eating disorders.** *J Eat Disord.* 2017; **5**: 7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
24. Norris ML, Robinson A, Obeid N, *et al.*: **Exploring avoidant/restrictive food intake disorder in eating disordered patients: a descriptive study.** *Int J Eat Disord.* 2014; **47**(5): 495–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
25. Becker AE, Thomas JJ, Pike KM: **Should non-fat-phobic anorexia nervosa be included in DSM-V?** *Int J Eat Disord.* 2009; **42**(7): 620–35.
[PubMed Abstract](#) | [Publisher Full Text](#)
26. Fleisher DR: **Regurgitation, Rumination and the Rumination Syndromes.** Child Health Publications, 2000.
[Reference Source](#)
27. Piazza CC, Fisher WW, Hanley GP, *et al.*: **Treatment of pica through multiple analyses of its reinforcing functions.** *J Appl Behav Anal.* 1998; **31**(2): 165–89.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
28. Williams KE, Hendy HM, Field DG, *et al.*: **Implications of Avoidant/Restrictive Food Intake Disorder (ARFID) on Children with Feeding Problems.** *Children's Health Care.* 2014; **44**(4): 307–21.
[Publisher Full Text](#)
29. Eddy KT, Thomas JJ, Hastings E, *et al.*: **Prevalence of DSM-5 avoidant/restrictive food intake disorder in a pediatric gastroenterology healthcare network.** *Int J Eat Disord.* 2015; **48**(5): 464–70.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
30. Thomas JJ, Lawson EA, Micali N, *et al.*: **Avoidant/Restrictive Food Intake Disorder: a Three-Dimensional Model of Neurobiology with Implications for Etiology and Treatment.** *Curr Psychiatry Rep.* 2017; **19**(8): 54.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. Timothy D Brewerton

Department of Psychiatry and Behavioral Sciences, Medical University of South Carolina, Charleston, SC, USA

Competing Interests: No competing interests were disclosed.

2. Jennifer J Thomas

¹ Department of Psychiatry, Harvard Medical School, Boston, MA, USA

² Eating Disorders Clinical and Research Program, Massachusetts General Hospital, Boston, MA, USA

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research