REVIEW

Understanding the CD8 T-cell response in natural HIV control

[version 1; peer review: 3 approved]

Sushma Boppana , Paul Goepfert

Department of Medicine, University of Alabama at Birmingham, Birmingham, USA

V1 First published: 02 Jul 2018, 7(F1000 Faculty Rev):985
<https://doi.org/10.12688/f1000research.15029.1>Latest published: 02 Jul 2018, 7(F1000 Faculty Rev):985
<https://doi.org/10.12688/f1000research.15029.1>

Abstract

HIV-infected individuals who maintain control of virus without antiretroviral therapy (ART) are called HIV controllers. The immune responses of these individuals suppress HIV viral replication to low levels or, in the case of elite controllers, to undetectable levels. Although some research indicates a role for inferior virulence of the infecting viral strain in natural control, perhaps by way of defective Nef protein function, we find that the majority of research in HIV controllers highlights CD8 T cells as the main suppressor of viral replication. The most convincing evidence for this argument lies in the strong correlation between certain HLA-I alleles, especially B*57, and HIV control status, a finding that has been replicated by many groups. However, natural control can also occur in individuals lacking these specific HLA alleles, and our understanding of what constitutes an effective CD8 T-cell response remains an incomplete picture. Recent research has broadened our understanding of natural HIV control by illustrating the interactions between different immune cells, including innate immune effectors and antigen-presenting cells. For many years, the immune responses of the natural HIV controllers have been studied for clues on how to achieve functional cure in the rest of the HIV-infected population. The goal of a future functional cure to HIV is one where HIV-infected individuals' immune responses are able to suppress virus long-term without requiring ART. This review highlights recent advances in our understanding of how HIV controllers' natural immune responses are able to suppress virus.

Keywords

HIV Controllers, CD8 T cells

Open Peer Review

Approval Status

	1	2	3
version 1 02 Jul 2018			

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Charles R Rinaldo**, University of Pittsburgh, Pittsburgh, USA
2. **Joel N Blankson**, Center for AIDS Research, Johns Hopkins University School of Medicine, Baltimore, USA
3. **Nicole Frahm**, Fred Hutchinson Cancer Research Center, Seattle, USA

Any comments on the article can be found at the end of the article.

Corresponding author: Paul Goepfert (paulg@uab.edu)

Author roles: **Boppana S:** Conceptualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Goepfert P:** Conceptualization, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by National Institutes of Health grants R01AI064060, R01AI112566, and T32AI007051-40. *The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

Copyright: © 2018 Boppana S and Goepfert P. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Boppana S and Goepfert P. **Understanding the CD8 T-cell response in natural HIV control [version 1; peer review: 3 approved]** F1000Research 2018, 7(F1000 Faculty Rev):985 <https://doi.org/10.12688/f1000research.15029.1>

First published: 02 Jul 2018, 7(F1000 Faculty Rev):985 <https://doi.org/10.12688/f1000research.15029.1>

Introduction

This review aims to summarize the field's current understanding of natural HIV controllers, with a focus on recent advances examining the CD8 T-cell response in these individuals and harnessing this knowledge for future therapies. A difficulty with comparing different studies on HIV controllers is the variation in definitions that different groups use, as illustrated by Gurdasani *et al.*'s systemic review of 714 definitions in 501 elite controller studies¹. For this review, we will assume the following definitions and specify any deviations: HIV controllers maintain viral loads of less than 2,000 RNA copies per milliliter (mL) of plasma off-antiretroviral therapy (ART). This group of HIV controllers can be further divided into elite controllers, individuals who maintain undetectable viremia (HIV RNA <50 copies/mL), and viremic controllers, who maintain low, but detectable, viremia (HIV RNA <2,000 copies/mL). Some studies focus on long-term non-progressors (LTNPs), who are HIV-infected individuals who remain asymptomatic for a prolonged period time off-ART with CD4 cell counts higher than 500, and do not restrict based on viral load.

A major question in the field asks what the relative contribution of viral functions and host factors are to natural HIV control. Although some cases demonstrate that infection with less-virulent HIV strains allows for better natural control, based on the evidence available, the host immune response, especially the CD8 T-cell response, appears to more strongly contribute to natural control.

Viral functions contributing to control

Several studies indicate that the function of Nef, the negative factor protein encoded by HIV, is impaired in HIV controllers. Typically, Nef is responsible for down-regulating several surface markers, including CD4, on HIV-infected cells and helps perpetuate the infection. Several groups first described the connection between infection with a Nef-deleted or -defective virus and natural viral control in 1995^{2,3}. However, more recent studies of Nef function in elite controllers show that while individuals can be infected with a Nef-defective virus, these less-virulent viruses are often a result of the controllers' CD8 T cells exerting immune pressure and causing an accumulation of HLA-I-associated polymorphisms within the Nef viral sequence. This accumulation of CD8 escape mutations then results in impaired Nef functionality^{4,5}, perhaps by increasing the sensitivity of infected cells to antibody-dependent cellular cytotoxicity (ADCC)^{6,7}.

There is also some evidence indicating that viruses isolated from elite controllers have a poorer replicative capacity than those from non-controllers. Investigations of Gag and Pol sequences indicate that HIV in controllers is less replication competent; however, like Nef impairment, this weakened replicative fitness is caused by fitness-costly escape mutations, a result of the host CD8 T-cell response^{8,9}. A recent study of the Env sequences showed that viruses isolated from elite controllers had impaired CD4 binding and signaling compared to non-controllers and asserts virologic factors play a role in HIV control in these individuals¹⁰. Other studies have described instances where the same virus caused significant pathogenicity in some

HIV-infected individuals but was effectively suppressed by the CD8 T-cell response in HIV controllers^{11–13}. Similarly, characterization of the viral fitness and early immune response in a recently infected controller demonstrated that the individual maintained viral control even with a replication-competent virus through a potent CD8 T-cell response that prevented escape mutations from occurring in the Gag protein¹⁴. Another recent study examining superinfection in a B*57-positive progressor showed that disease progression occurred in spite of significant decreases in viral fitness, indicating that decreased viral fitness is not sufficient for maintained viral control¹⁵.

CD8 response in HIV controllers

Perhaps the strongest evidence that the host CD8 T-cell response is the major player in natural control of HIV is the association of specific HLA-I alleles, like B*27 and B*57, with elite controllers; this finding is consistent across several groups^{16–19}. HLA-I alleles dictate which epitopes from HIV are presented to and recognized by the host immune response, and these protective HLA-I alleles drive potent and effective CD8 T-cell responses²⁰. In fact, this relationship is so strong that it can be seen even within a single clinic's cohort. In the University of Alabama at Birmingham's HIV clinic, 1917 clinic, B*57-positive HIV-infected patients are more likely to be controllers ($P = 0.04$, Fisher's Exact test) and have lower viral loads ($P = 0.015$ by Mann Whitney test, [Figure 1](#)). Carlson and colleagues recently demonstrated the importance of CD8 T cells in viral control by examining the relationship between the "pre-adaptation" of the infecting virus and subsequent control of that virus. They found that individuals infected by a virus already containing CD8 T-cell escape mutations had poorer clinical outcomes, including higher viral load set points and faster CD4 T cell decline. Importantly, the advantage conferred by protective HLA-I alleles, like B*57, was lost when B*57-positive individuals were infected with pre-adapted virus, indicating that viral control in elite controllers is dependent on the CD8 T-cell response's ability to prevent CD8 escape mutations from accumulating²¹.

The CD8 T-cell response of HIV controllers is more robust than that of non-controllers in several ways, displaying greater cytotoxicity^{22,23}, proliferation^{24,25}, and polyfunctionality^{26–28}. However,

Figure 1. Association of B*57 with viral control in 1917 clinic cohort. Viral loads of 1917 clinic patients with protective HLA-B*57 allele versus those of patients lacking any HLA-I alleles previously associated with delayed or accelerated disease progression.

our understanding of the CD8 T-cell response in HIV controllers is continuing to evolve. For example, a more recent study found that freshly thawed cells from controllers and progressors were similarly able to inhibit virus but that CD8 T cells from controllers exhibited better sustained polyfunctionality over time²⁹. Two recent studies transfected cells with the T-cell receptors (TCRs) from an elite controller's CD8 T-cell response and found that the superior functionality of CD8 T cells is not completely dictated by the TCR itself and that factors external to the TCR sequence contribute to antiviral functionality^{30,31}. Although many aspects of the CD8 T-cell response are superior in controllers, whether these CD8 T-cell characteristics are the cause of greater viral control or the result of a more preserved immune response because of low or absent viremia often remains uncertain.

Another beneficial aspect of the CD8 T-cell response of elite controllers is thought to be early and potent targeting of the Gag protein, a key structural and functional protein of HIV. Brockman *et al.* found that viruses from individuals with protective alleles had lower replication capacity, which inversely correlated with HLA-B-associated Gag polymorphisms in early HIV infection. This correlation suggests that the early CD8 T-cell response in these individuals created immune pressure and forced the virus to undergo fitness-costly mutations within Gag³². Additionally, the magnitude of the anti-Gag CD8 T-cell response correlates with the level of viral inhibitory activity³³. These potent CD8 T-cell responses are seen in elite controllers with and without protective HLA-I alleles³⁴.

The CD8 T-cell responses of elite controllers may also be more cross-reactive than those of HIV progressors, meaning their CD8 T cells may be better equipped to respond to not just the epitopes encoded by the infecting virus but also variant epitopes that arise as a result of HIV's rapid accumulation of mutations. One study computationally demonstrated that protective HLA-I alleles, like B*57, present fewer self-peptides in the thymus, resulting in a T-cell repertoire that relies on a lower number of residues to recognize viral peptides and, therefore, a T-cell response that is able to recognize a much broader array of variant epitopes³⁵. This increased cross-recognition was also observed *in vivo* and *in vitro* studies where higher cross-recognition scores were associated with lower viral loads³⁶ and T-cell clones derived from controllers were more cross-reactive than clones derived from HLA-matched progressors³⁷. CD8 T cells from B*57-positive elite controllers have been found to effectively suppress the replication of viruses encoding escape mutations, demonstrating an ability to suppress variant epitope-encoding viruses³⁸.

Some HIV controllers eventually lose their ability to suppress virus without therapy, and investigations into the mechanisms behind this loss of control may provide clues into which actors exert control in the first place. A recent study comparing elite controllers with and without CD4 decline found that disturbances in T-cell homeostasis, including lower levels of naïve CD8 T cells, fewer recent thymic emigrants, and greater CD4 T-cell exhaustion, appeared to lead to this loss of control and rapid CD4 T-cell decline³⁹. Most of the disturbances in homeostasis

that correlated with CD4 T-cell loss in this study involved the CD8 T-cell response, supporting the idea that CD8 T cells are the major player in the viral control existing before CD4 loss occurred.

Parallels in model systems

Some non-human primate (NHP) models are seen as comparable to natural HIV controllers. Simian immunodeficiency viruses (SIVs), which are related to HIV, replicate efficiently in their natural hosts (e.g. African green monkeys and sooty mangabeys) without causing any disease pathology, but the SIV/macaque model, which is often used for HIV studies, involves laboratory-selected SIV strains, which are pathogenic. Watkins and colleagues have linked HIV control in the presence of the protective HLA-I allele B*27 with Mamu B*08-mediated SIV control in rhesus macaques where roughly 50% of Mamu B*08-positive macaques control SIV replication similar to elite controllers⁴⁰. They have also demonstrated that an effective CD8 T-cell response in acute infection is key and that early CD8 T-cell-mediated viral escape occurs in Mamu B*08 progressors but not in Mamu B*08 controllers⁴¹. Additionally, upon depletion of CD8 T cells in controller monkeys, researchers observed a major rebound in virus, even greater than that seen in progressors^{42,43}. These studies of rhesus macaques suggest that the CD8 T-cell response is the primary effector in natural SIV control. Another recently published study lends support to this idea that viral factors are not the key influencers of disease progression in NHP controllers. Joas *et al.* examined the impact of adding or replacing pathogenic HIV-encoded Vpu and Nef, two important HIV virulence factors, to a non-pathogenic SIV strain. They found that despite high levels of viral replication, Vpu- and/or Nef-containing SIV strains remained non-pathogenic. Because the introduction of these virulence factors did not increase the pathogenicity of the tested SIV strains, the authors concluded that host factors are more likely to be responsible for protection against disease⁴⁴. Overall, these studies in NHP models of SIV infection lend strong support to the importance of CD8 T cells in natural control.

Other host functions contributing to control

Recent studies have also investigated the contribution of the innate immune system to natural HIV control. Krishnan and colleagues found that D-dimer, soluble tissue factor, and CD14⁺CD16⁺ monocyte levels were elevated in elite controllers, indicating a more activated innate immune response⁴⁵. Another recent study looked at differences in systemic cytokine responses between elite controllers and non-controllers and found that a subset of cytokines was increased in controllers and that this combination of cytokines (CCL14, CCL21, CCL27, XCL1, and SDF-1) was able to suppress viral replication *in vitro*⁴⁶. Additional research is required before we fully understand the complex roles of innate immune actors on viral control in HIV controllers.

The antigen-presenting cells (APCs) are also an important component of the immune response, and their role in HIV control is becoming clearer. Martin-Gayo and colleagues used a computational framework and single-cell RNA sequencing to describe a higher functional antiviral dendritic cell state that primes

polyfunctional T-cell responses in elite controllers⁴⁷. In addition to aiding the CD8 response, dendritic cells and other APCs may contribute to natural control through an innate resistance to spreading virus from cell to cell, or trans-infection. For dendritic cells, B lymphocytes, and macrophages, the Rinaldo group demonstrated that this decreased facilitation of trans-infection is linked to altered cholesterol metabolism. In HIV non-progressors, they found that APCs express less free cholesterol on the cell surface, leading to a decreased internalization of virus^{48,49}. To date, APCs in HIV have been understudied, and future research will likely highlight the important ways in which these cells interact with effector immune cells.

Recent research has also noted significant differences between male and female HIV controllers (reviewed in 50); women are overrepresented among HIV controllers in several studies^{51,52}. Zhang *et al.* found that female elite controllers actually had similar gene expression profiles to uninfected females but differences from male elite controllers. Because the differentially expressed genes between men and women in the study were not exclusively located on the X or Y chromosome, the authors argue that these gene expression differences are not simply due to sex-linked genes⁵³. Another study indicates that significantly fewer female elite controllers experience loss of natural HIV control as defined by CD4 T-cell decline³⁹. In fact, even within the 1917 clinic, we see that women are significantly overrepresented within the elite controller population compared to the general clinic population (Table 1, $P = 0.03$). Recent publications indicate a need for more research into sex differences in HIV controllers and a better understanding of how these differences might play into future therapeutic efforts.

The CD4 response may also play an important role in natural HIV control. A recent study of the TCRs responsible for the immunodominant CD4 response in controllers found a significant number of public clonotypes, including shared CDR3 motifs. Transfer of these Gag-specific, high-avidity TCRs to non-HIV-infected CD4 T cells resulted in high antigen sensitivity and polyfunctionality and even redirected CD8 T cells to target the HIV capsid⁵⁴. A follow-up study found that these CD4 TCRs were able to suppress HIV in the context of multiple genetic backgrounds, raising the possibility of engineering or transferring CD4 T cells as an effective therapy⁵⁵. Although CD4 T cells may play a supportive role in controlling HIV in these individuals, depletion of CD4 T cells in SIV-infected macaque elite

controllers does not impact viral load, indicating that they are not required for natural control⁵⁶.

Future directions

A current area of interest for the HIV cure field is the dynamics of the latent HIV reservoir within infected individuals because this reservoir poses a major barrier to achieving HIV cure. Boritz and colleagues studied the persistence of virus in HIV controllers and found that, despite effective antiviral immune responses in these individuals, their reservoir was maintained through viral replication, clonal expansion of latently infected cells, and a low level of circulating infected cells⁵⁷. However, it was recently demonstrated that the HIV reservoir is smaller in elite controllers, particularly in peripheral T follicular helper cells, meaning the host response within these controllers is able to limit the HIV reservoir size⁵⁸. The HIV reservoir found in lymphoid tissue is a major barrier to current CD8 T-cell-based cure efforts because without effective immune responses or daily ART, these latently infected cells reactivate and virus rebounds. Further research into how natural HIV controllers are able to suppress the emergence of virus from these pockets could be a useful step forward.

A caveat to using natural HIV control to inform functional cure strategies is recent research indicating that in spite of long-term ART-free viral control, elite controllers may still have increased levels of inflammation compared to HIV-negative individuals. Even in HIV controllers, viral evolution can be seen, indicating that these individuals have some level of ongoing viral replication^{59,60}. This low-level persistence of HIV in controllers may contribute to the described increased levels of inflammatory markers, like IFN γ , IL10, and sCD40L⁶¹. However, another publication from the same year found that elite controllers with preserved CD4 T-cell percentages do not have increased levels of T-cell activation⁶², and more recent research in *JAMA Cardiology* found that arterial and lymph node inflammation in elite controllers was comparable to that of ART-suppressed HIV-infected individuals⁶³. Additionally, analysis of non-AIDS-defining events (nADEs) found that while controllers do experience nADEs, they do so at a lower rate than do non-controllers⁶⁴. Further research into the potential long-term effects of natural HIV control is required to confirm that natural control is the ideal model for developing a functional cure for HIV.

There is still hope that our understanding of natural control can be used to inform future vaccines or therapy strategies. Although we have made significant progress in understanding the dynamics underlying the CD8 T-cell responses in these individuals, the field has successfully harnessed effective CD8 T-cell responses to counter HIV infection only in the lab and in macaque models. While antibodies are likely to play a major role in preventing HIV infection, preventative vaccines that pack a powerful CD8 arm may be able to alleviate the course of disease in vaccinated individuals who become infected. Based on our understanding of natural controllers' CD8 responses, this vaccine-induced response would likely need to effectively prevent the virus from escaping the immune response or would

Table 1. 1917 clinic demographics, female versus male elite controllers.

	Females		Males		Total
	Number	Percent	Number	Percent	Number
Active patients	839	24%	2,618	76%	3,457
Elite controllers	9	47%	10	53%	19

Active patients defined as being seen in the 1917 clinic within the past year; elite controllers defined as individuals with viral loads of <50 copies of RNA/mL of plasma.

need to force more fitness-costly mutations to occur. CD8 T cells restricted by HLA-E and possibly HLA-II could also be harnessed to prevent establishment of HIV infection, as has been elegantly demonstrated by the Picker group using cytomegalovirus recombinant SIV vectors in the NHP model system^{65,66}. CD8 T cells may also be the central effector of future functional cures, where the CD8 T-cell response of an infected individual is reinvigorated and redirected to effectively suppress virus in the absence of ART. Future research on natural HIV control can help the field move closer to achieving these goals.

Abbreviations

ADCC, antibody-dependent cellular cytotoxicity; APC, antigen-presenting cell; ART, antiretroviral therapy; LTNP, long-term

non-progressor; mL, milliliter; nADE, non-AIDS-defining event; NHP, non-human primate; SIV, simian immunodeficiency virus; TCR, T-cell receptor

Competing interests

The authors declare that they have no competing interests.

Grant information

This work was supported by National Institutes of Health grants R01AI064060, R01AI112566, and T32AI007051-40.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- Gurdasani D, Iles L, Dillon DG, *et al.*: **A systematic review of definitions of extreme phenotypes of HIV control and progression.** *AIDS*. 2014; **28**(2): 149–62. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Deacon NJ, Tsykin A, Solomon A, *et al.*: **Genomic structure of an attenuated quasi species of HIV-1 from a blood transfusion donor and recipients.** *Science*. 1995; **270**(5238): 988–91. [PubMed Abstract](#) | [Publisher Full Text](#)
- Kirchhoff F, Greenough TC, Brettler DB, *et al.*: **Brief report: absence of intact nef sequences in a long-term survivor with nonprogressive HIV-1 infection.** *N Engl J Med*. 1995; **332**(4): 228–32. [PubMed Abstract](#) | [Publisher Full Text](#)
- Kuang XT, Li X, Anmole G, *et al.*: **Impaired Nef function is associated with early control of HIV-1 viremia.** *J Virol*. 2014; **88**(17): 10200–13. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Mwimanzu P, Markle TJ, Martin E, *et al.*: **Attenuation of multiple Nef functions in HIV-1 elite controllers.** *Retrovirology*. 2013; **10**: 1. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Alsahafi N, Ding S, Richard J, *et al.*: **Nef Proteins from HIV-1 Elite Controllers Are Inefficient at Preventing Antibody-Dependent Cellular Cytotoxicity.** *J Virol*. 2015; **90**(6): 2993–3002. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Alsahafi N, Richard J, Prévost J, *et al.*: **Impaired Downregulation of NKG2D Ligands by Nef Proteins from Elite Controllers Sensitizes HIV-1-Infected Cells to Antibody-Dependent Cellular Cytotoxicity.** *J Virol*. 2017; **91**(16): pii: e00109–17. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
- Brumme ZL, Li C, Miura T, *et al.*: **Reduced replication capacity of NL4-3 recombinant viruses encoding reverse transcriptase-integrase sequences from HIV-1 elite controllers.** *J Acquir Immune Defic Syndr*. 2011; **56**(2): 100–8. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Miura T, Brumme ZL, Brockman MA, *et al.*: **Impaired replication capacity of acute/early viruses in persons who become HIV controllers.** *J Virol*. 2010; **84**(15): 7581–91. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
- Casado C, Marrero-Hernández S, Márquez-Arce D, *et al.*: **Viral Characteristics Associated with the Clinical Nonprogressor Phenotype Are Inherited by Viruses from a Cluster of HIV-1 Elite Controllers.** *MBio*. 2018; **9**(2): pii: e02338–17. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
- Bailey JR, O'Connell K, Yang HC, *et al.*: **Transmission of human immunodeficiency virus type 1 from a patient who developed AIDS to an elite suppressor.** *J Virol*. 2008; **82**(15): 7395–410. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Blankson JN, Bailey JR, Thayil S, *et al.*: **Isolation and characterization of replication-competent human immunodeficiency virus type 1 from a subset of elite suppressors.** *J Virol*. 2007; **81**(5): 2508–18. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Buckheit RW 3rd, Allen TG, Alme A, *et al.*: **Host factors dictate control of viral replication in two HIV-1 controller/chronic progressor transmission pairs.** *Nat Commun*. 2012; **3**: 716. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Walker-Sperling VE, Pohlmeier CW, Veenhuis RT, *et al.*: **Factors Associated With the Control of Viral Replication and Virologic Breakthrough in a Recently Infected HIV-1 Controller.** *EBioMedicine*. 2017; **16**: 141–9. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
- Brener J, Gall A, Hurst J, *et al.*: **Rapid HIV disease progression following superinfection in an HLA-B*27:05/B*57:01-positive transmission recipient.** *Retrovirology*. 2018; **15**(1): 7. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
- Altfield M, Kalife ET, Qi Y, *et al.*: **HLA Alleles Associated with Delayed Progression to AIDS Contribute Strongly to the Initial CD8(+) T Cell Response against HIV-1.** *PLoS Med*. 2006; **3**(10): e403. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Migueles SA, Sabbaghian MS, Shupert WL, *et al.*: **HLA B*5701 is highly associated with restriction of virus replication in a subgroup of HIV-infected long term nonprogressors.** *Proc Natl Acad Sci U S A*. 2000; **97**(6): 2709–14. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- International HIV Controllers Study, Pereyra F, Jia X, *et al.*: **The major genetic determinants of HIV-1 control affect HLA class I peptide presentation.** *Science*. 2010; **330**(6010): 1551–7. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
- Moosa Y, Tanko RF, Ramsuran V, *et al.*: **Case report: mechanisms of HIV elite control in two African women.** *BMC Infect Dis*. 2018; **18**(1): 54. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Brennan CA, Ibarondo FJ, Sugar CA, *et al.*: **Early HLA-B*57-restricted CD8+ T lymphocyte responses predict HIV-1 disease progression.** *J Virol*. 2012; **86**(19): 10505–16. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Carlson JM, Du VY, Pfeifer N, *et al.*: **Impact of pre-adapted HIV transmission.** *Nat Med*. 2016; **22**(6): 606–13. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
- Hersperger AR, Martin JN, Shin LY, *et al.*: **Increased HIV-specific CD8+ T-cell cytotoxic potential in HIV elite controllers is associated with T-bet expression.** *Blood*. 2011; **117**(14): 3799–808. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
- Migueles SA, Osborne CM, Royce C, *et al.*: **Lytic granule loading of CD8+ T cells is required for HIV-infected cell elimination associated with immune control.** *Immunity*. 2008; **29**(6): 1009–21. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
- Migueles SA, Laborico AC, Shupert WL, *et al.*: **HIV-specific CD8+ T cell proliferation is coupled to perforin expression and is maintained in nonprogressors.** *Nat Immunol*. 2002; **3**(11): 1061–8. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
- Day CL, Kiepiela P, Leslie AJ, *et al.*: **Proliferative capacity of epitope-specific CD8 T-cell responses is inversely related to viral load in chronic human immunodeficiency virus type 1 infection.** *J Virol*. 2007; **81**(1): 434–8. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ndhlovu ZM, Chibnik LB, Proudfoot J, *et al.*: **High-dimensional immunomonitoring models of HIV-1-specific CD8 T-cell responses accurately**

- identify subjects achieving spontaneous viral control. *Blood*. 2013; 121(5): 801–11.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
27. Betts MR, Nason MC, West SM, *et al.*: HIV nonprogressors preferentially maintain highly functional HIV-specific CD8⁺ T cells. *Blood*. 2006; 107(12): 4781–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 28. Peris-Pertusa A, López M, Rallón NI, *et al.*: Evolution of the functional profile of HIV-specific CD8⁺ T cells in patients with different progression of HIV infection over 4 years. *J Acquir Immune Defic Syndr*. 2010; 55(1): 29–38.
[PubMed Abstract](#) | [Publisher Full Text](#)
 29. F Shasha D, Karel D, Angiuli O, *et al.*: Elite controller CD8⁺ T cells exhibit comparable viral inhibition capacity, but better sustained effector properties compared to chronic progressors. *J Leukoc Biol*. 2016; 100(6): 1425–33.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 30. F Flieri NC, Chen H, Glover TD, *et al.*: T-Cell Receptor (TCR) Clonotype-Specific Differences in Inhibitory Activity of HIV-1 Cytotoxic T-Cell Clones Is Not Mediated by TCR Alone. *J Virol*. 2017; 91(6): pii: e02412-16.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 31. F Joglekar AV, Liu Z, Weber JK, *et al.*: T cell receptors for the HIV KK10 epitope from patients with differential immunologic control are functionally indistinguishable. *Proc Natl Acad Sci U S A*. 2018; 115(8): 1877–82.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 32. Brockman MA, Brumme ZL, Brumme CJ, *et al.*: Early selection in Gag by protective HLA alleles contributes to reduced HIV-1 replication capacity that may be largely compensated for in chronic infection. *J Virol*. 2010; 84(22): 11937–49.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 33. Turk G, Ghiglione Y, Falivene J, *et al.*: Early Gag immunodominance of the HIV-specific T-cell response during acute/early infection is associated with higher CD8⁺ T-cell antiviral activity and correlates with preservation of the CD4⁺ T-cell compartment. *J Virol*. 2013; 87(13): 7445–62.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 34. Migueles SA, Mendoza D, Zimmerman MG, *et al.*: CD8⁺ T-cell Cytotoxic Capacity Associated with Human Immunodeficiency Virus-1 Control Can Be Mediated through Various Epitopes and Human Leukocyte Antigen Types. *EBioMedicine*. 2015; 2(1): 46–58.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 35. F Kosmrlj A, Read EL, Qi Y, *et al.*: Effects of thymic selection of the T-cell repertoire on HLA class I-associated control of HIV infection. *Nature*. 2010; 465(7296): 350–4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 36. F Turnbull EL, Lopes AR, Jones NA, *et al.*: HIV-1 epitope-specific CD8⁺ T cell responses strongly associated with delayed disease progression cross-recognize epitope variants efficiently. *J Immunol*. 2006; 176(10): 6130–46.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 37. F Ladell K, Hashimoto M, Iglesias MC, *et al.*: A molecular basis for the control of preimmune escape variants by HIV-specific CD8⁺ T cells. *Immunity*. 2013; 38(3): 425–36.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 38. Pohlmeier CW, Buckheit RW 3rd, Siliciano RF, *et al.*: CD8⁺ T cells from HLA-B*57 elite suppressors effectively suppress replication of HIV-1 escape mutants. *Retrovirology*. 2013; 10: 152.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 39. F Benito JM, Ortiz MC, León A, *et al.*: Class-modeling analysis reveals T-cell homeostasis disturbances involved in loss of immune control in elite controllers. *BMC Med*. 2018; 16(1): 30.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 40. F Loffredo JT, Maxwell J, Qi Y, *et al.*: Mamu-B*08-positive macaques control simian immunodeficiency virus replication. *J Virol*. 2007; 81(16): 8827–32.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 41. Mudd PA, Ericson AJ, Burwitz BJ, *et al.*: Escape from CD8⁺ T cell responses in Mamu-B*0801⁺ macaques differentiates progressors from elite controllers. *J Immunol*. 2012; 188(7): 3364–70.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 42. Chowdhury A, Hayes TL, Bosinger SE, *et al.*: Differential Impact of In Vivo CD8⁺ T Lymphocyte Depletion in Controller versus Progressor Simian Immunodeficiency Virus-Infected Macaques. *J Virol*. 2015; 89(17): 8677–86.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 43. Friedrich TC, Valentine LE, Yant LJ, *et al.*: Subdominant CD8⁺ T-cell responses are involved in durable control of AIDS virus replication. *J Virol*. 2007; 81(7): 3465–76.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 44. F Joas S, Parrish EH, Gnanadurai CW, *et al.*: Species-specific host factors rather than virus-intrinsic virulence determine primate lentiviral pathogenicity. *Nat Commun*. 2018; 9(1): 1371.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 45. Krishnan S, Wilson EM, Sheikh V, *et al.*: Evidence for innate immune system activation in HIV type 1-infected elite controllers. *J Infect Dis*. 2014; 209(6): 931–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 46. F Jacobs ES, Keating SM, Abdel-Mohsen M, *et al.*: Cytokines Elevated in HIV Elite Controllers Reduce HIV Replication In Vitro and Modulate HIV Restriction Factor Expression. *J Virol*. 2017; 91(6): pii: e02051-16.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 47. F Martin-Gayo E, Cole MB, Kolb KE, *et al.*: A Reproducibility-Based Computational Framework Identifies an Inducible, Enhanced Antiviral State in Dendritic Cells from HIV-1 Elite Controllers. *Genome Biol*. 2018; 19(1): 10.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 48. F DeLucia DC, Rinaldo CR, Rappocciolo G: Inefficient HIV-1 trans infection of CD4⁺ T cells by macrophages from HIV-1 Nonprogressors is Associated with Altered Membrane Cholesterol and DC-SIGN. *J Virol*. 2018; 92(13): pii: e00092-18.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 49. Rappocciolo G, Jais M, Piazza P, *et al.*: Alterations in cholesterol metabolism restrict HIV-1 trans infection in nonprogressors. *MBio*. 2014; 5(3): e01031–13.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 50. Griesbeck M, Scully E, Altfeld M: Sex and gender differences in HIV-1 infection. *Clin Sci (Lond)*. 2016; 130(16): 1435–51.
[PubMed Abstract](#) | [Publisher Full Text](#)
 51. Crowell TA, Gebo KA, Blankson JN, *et al.*: Hospitalization Rates and Reasons Among HIV Elite Controllers and Persons With Medically Controlled HIV Infection. *J Infect Dis*. 2015; 211(11): 1692–702.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 52. F de Azevedo SSD, Caetano DG, Côrtes FH, *et al.*: Highly divergent patterns of genetic diversity and evolution in proviral quasiespecies from HIV controllers. *Retrovirology*. 2017; 14(1): 29.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 53. F Zhang W, Ambikan AT, Sperk M, *et al.*: Transcriptomics and Targeted Proteomics Analysis to Gain Insights Into the Immune-control Mechanisms of HIV-1 Infected Elite Controllers. *EBioMedicine*. 2018; 27: 40–50.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 54. F Benati D, Galperin M, Lambotte O, *et al.*: Public T cell receptors confer high-avidity CD4 responses to HIV controllers. *J Clin Invest*. 2016; 126(6): 2093–108.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 55. F Galperin M, Farenc C, Mukhopadhyay M, *et al.*: CD4⁺ T cell-mediated HLA class II cross-restriction in HIV controllers. *Sci Immunol*. 2018; 3(24): pii: eaat0687.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 56. Mudd PA, Ericson AJ, Price AA, *et al.*: Reduction of CD4⁺ T cells in vivo does not affect virus load in macaque elite controllers. *J Virol*. 2011; 85(14): 7454–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 57. F Boritz EA, Darko S, Swaszek L, *et al.*: Multiple Origins of Virus Persistence during Natural Control of HIV Infection. *Cell*. 2016; 166(4): 1004–15.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 58. F García M, Górgolas M, Cabello A, *et al.*: Peripheral T follicular helper Cells Make a Difference in HIV Reservoir Size between Elite Controllers and Patients on Successful cART. *Sci Rep*. 2017; 7(1): 16799.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 59. O'Connell KA, Brennan TP, Bailey JR, *et al.*: Control of HIV-1 in elite suppressors despite ongoing replication and evolution in plasma virus. *J Virol*. 2010; 84(14): 7018–28.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 60. Mens H, Kearney M, Wiegand A, *et al.*: HIV-1 continues to replicate and evolve in patients with natural control of HIV infection. *J Virol*. 2010; 84(24): 12971–81.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 61. Li JZ, Arnold KB, Lo J, *et al.*: Differential levels of soluble inflammatory markers by human immunodeficiency virus controller status and demographics. *Open Forum Infect Dis*. 2015; 2(1): ofu117.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 62. Bansal A, Sterrett S, Erdmann N, *et al.*: Normal T-cell activation in elite controllers with preserved CD4⁺ T-cell counts. *AIDS*. 2015; 29(17): 2245–54.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 63. F Tawakol A, Ishai A, Li D, *et al.*: Association of Arterial and Lymph Node Inflammation With Distinct Inflammatory Pathways in Human Immunodeficiency Virus Infection. *JAMA Cardiol*. 2017; 2(2): 163–71.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 64. F Dominguez-Molina B, Leon A, Rodriguez C, *et al.*: Analysis of Non-AIDS-Defining Events in HIV Controllers. *Clin Infect Dis*. 2016; 62(10): 1304–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
 65. F Hansen SG, Wu HL, Burwitz BJ, *et al.*: Broadly targeted CD8⁺ T cell responses restricted by major histocompatibility complex E. *Science*. 2016; 351(6274): 714–20.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
 66. F Hansen SG, Sacha JB, Hughes CM, *et al.*: Cytomegalovirus vectors violate CD8⁺ T cell epitope recognition paradigms. *Science*. 2013; 340(6135): 1237874.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. Nicole Frahm

Vaccine and Infectious Disease Division and the HIV Vaccine Trials Network, Fred Hutchinson Cancer Research Center, Seattle, USA

Competing Interests: No competing interests were disclosed.

2. Joel N Blankson

Department of Medicine, Center for AIDS Research, Johns Hopkins University School of Medicine, Baltimore, USA

Competing Interests: No competing interests were disclosed.

3. Charles R Rinaldo

Department of Infectious Diseases and Microbiology, University of Pittsburgh, Pittsburgh, USA

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research