RESEARCH ARTICLE

A fruit fly model for studying paclitaxel-induced pain [version 1; peer review: 1 approved, 1 approved with reservations]

Zina Hamoudi^{1,2}, Thang Manh Khuong^{1,2}, Tiffany Cole^{1,2}, G. Gregory Neely^{id}^{1,2}

¹University of Sydney, Charles Perkins Centre and School of Life and Environmental Sciences, Camperdown, New South Wales, Australia

²University of Sydney, Dr. John and Anne Chong Lab for Functional Genomics, Camperdown, New South Wales, Australia

V1 First published: 23 Jan 2018, 7:99
<https://doi.org/10.12688/f1000research.13581.1>
 Latest published: 16 Oct 2018, 7:99
<https://doi.org/10.12688/f1000research.13581.2>

Abstract

Background: Paclitaxel-induced peripheral neuropathy is a common and limiting side effect of an approved and effective chemotherapeutic agent. The cause of this nociception is still unknown.

Methods: To uncover the mechanism involved in paclitaxel-induced pain, we developed a *Drosophila* thermal nociceptive model to show the effects of paclitaxel exposure on third instar larvae.

Results: We found that paclitaxel increases pain perception in a dose-dependent manner, without overt morphological changes.

Conclusions: Our simple, high throughput model can be combined with genomics approaches to identify regulators of chemotherapy-induced pain to eliminate its adverse side effects.

Keywords

Drosophila, fruit fly, paclitaxel, nociception, pain, CIPN

Open Peer Review

Approval Status

	1	2	3
version 2 (revision) 16 Oct 2018			 view
version 1 23 Jan 2018	 view	 view	

1. **Adam Claridge-Chang** ^{id}, Institute of Molecular and Cell Biology (IMCB), Singapore, Singapore
2. **Salahadin Abdi**, University of Texas MD Anderson Cancer Center, Houston, USA
3. **Giorgio Gilestro** ^{id}, Imperial College London, London, UK

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: G. Gregory Neely (greg.neely@sydney.edu.au)

Author roles: **Hamoudi Z:** Conceptualization, Formal Analysis, Investigation, Methodology, Project Administration, Supervision, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Khuong TM:** Formal Analysis, Investigation, Writing – Original Draft Preparation; **Cole T:** Formal Analysis, Methodology; **Neely GG:** Conceptualization, Funding Acquisition, Methodology, Project Administration, Supervision, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This work was supported in part through NHMRC project grants APP1026310, APP1029672, APP1028887, APP1046090, APP1042416, APP1086851, and by a NHMRC career development fellowship II CDF1111940.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2018 Hamoudi Z *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. Data associated with the article are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

How to cite this article: Hamoudi Z, Khuong TM, Cole T and Neely GG. **A fruit fly model for studying paclitaxel-induced pain [version 1; peer review: 1 approved, 1 approved with reservations]** F1000Research 2018, 7:99 <https://doi.org/10.12688/f1000research.13581.1>

First published: 23 Jan 2018, 7:99 <https://doi.org/10.12688/f1000research.13581.1>

Introduction

Chemotherapy-induced peripheral neuropathy (CIPN) is a dose-limiting side effect of many effective cancer treatments (Burton *et al.*, 2007), and can have a lasting impact on the quality of life of cancer survivors (Hausheer *et al.*, 2006 and Shimozuma *et al.*, 2012). A meta-analysis of 31 studies from over 4000 chemotherapy-treated patients revealed that CIPN was prevalent in 68.1% of patients in the first month following chemotherapy, in 60% of patients at 3 months, and in 30% at 6 months or more (Seretny *et al.*, 2014).

Paclitaxel has a potent ability to cause CIPN (Addington & Freimer, 2016; Reyes-Gibby *et al.*, 2009). Derived from the bark of the western yew, *Taxus brevifolia*, it is an approved and effective treatment against breast, ovarian, lung and Kaposi sarcoma (Chang *et al.*, 1993; Gill *et al.*, 1999; Holmes *et al.*, 1991; McGuire *et al.*, 1989; Wani *et al.*, 1971). Patients treated with paclitaxel experience side effects as early as one to three days following treatment (Lipton *et al.*, 1989; Reyes-Gibby *et al.*, 2009). Common symptoms are hyperalgesia, hypoalgesia, allodynia, tingling, numbness, and shooting pain (Boland *et al.*, 2010). Paclitaxel has a direct effect on Schwann cells, promotes axonal degeneration, and can cause mitochondrial damage (André *et al.*, 2000; Cavaletti *et al.*, 1995; Sahenk *et al.*, 1994), however the molecular mechanisms causing pain are still largely unknown.

While much knowledge has been gained about the genetics of pain from vertebrate systems, high throughput dissection of pain is possible using the fruit fly *Drosophila melanogaster* (Neely *et al.*, 2010). When challenged with a noxious thermal stimulus, third instar larvae exhibit an aversive escape response that has been utilised to identify conserved genes required for nociception (Babcock *et al.*, 2009; Neely *et al.*, 2010; Tracey *et al.*, 2003). This nociceptive response is a result of activating class IV multidendritic-dendritic arborisation (md-da) sensory neurons at the site of stimulation (Hwang *et al.*, 2007). Previously in *Drosophila*, paclitaxel has been reported to be toxic in somatic cells, and causes loss of axons in peripheral nerves. (Bhattacharya *et al.*, 2012; Cunha *et al.*, 2001). However, its effects on nociception have not yet been evaluated. Here, we examined the effects of paclitaxel exposure on the fruit fly larval nociception system, and observed a robust and dose-dependent increase in pain perception. This system is amenable to high throughput screening and genetic manipulation, and may help define why chemotherapies such as paclitaxel cause pain.

Methods

Drosophila treatment

All flies were reared at 25°C and 65% humidity over a 12-hour light-dark cycle. Six female and two male *Canton S Drosophila melanogaster* were mated on food medium (5.4% sucrose, 3.6% yeast, 1% agar, 1.2% nipagin, and 0.6% propionic acid) treated with ethanol, 0 μ M, 0.1 μ M, 0.5 μ M, 2.5 μ M, 5 μ M or 10 μ M paclitaxel (Taxol®; Catalog No. A4393) purchased from ApexBio (Houston, USA). A stock of 1000 μ M paclitaxel in ethanol was prepared and diluted in food medium accordingly to create the different drug concentrated food. F0 Flies were discarded two days after mating and F1 larvae were left to grow for another three days. On the

fifth day, early third instar were collected to assess nociception or dendritic morphology.

Behavioural assay

For the thermal nociceptive assay (Tracey *et al.*, 2003), distilled water was added to experimental vials to soften the food and release the foraging third instar larvae. The softened, liquid food was then passed through mesh to catch the larvae to be transferred to a 100mm petri dish sprayed with distilled water. The larvae were touched laterally on abdominal segments four to six with a heat probe (soldering iron with narrow tip) set to 42°C or 46°C. The rolling response was measured in seconds with a cut-off of 10 seconds. For each drug concentration, five repeats were performed, with 30–40 larvae per repeat.

Larvae dissection, fixation, and mounting

Third instar larvae (*w;ppk-CD4-tdGFP*) were collected, washed, and placed onto a sylgard dissecting plate (Sylgard184 silicone elastomer kit, Dow Corning, Midland, MI). At least 10 larvae of each group were then dissected in HL-3 saline solution (70 mM NaCl, 5 mM KCl, 10 mM NaHCO₃, 5 mM HEPES, 115 mM sucrose, 5 mM threosulose, 20 mM MgCl₂, pH 7.2). Dissected larvae were fixed with 4% paraformaldehyde solution, and then washed in PBS four times. Larvae of the same genotype were then transferred to microscope slides to be mounted using Vectashield (Vector Laboratories, Burlingame, CA).

Confocal microscopy and image analysis

GFP-expressing class IV md-da sensory neurons were imaged on a Leica DMI-6000 inverted microscope and SP8 Basic Confocal system using 40x oil immersion lens (Leica, Wetzlar, Germany). Class IV neuron at abdominal segment 5 (A5) of each larva was imaged. 488 nm lasers for excitation were used, emission signal was captured at 500–550nm wave length, scanning speed of 400 Hz, zoom factor of 0.7, and gain of 600. Images of Z-stack sections were captured at 1024 × 1024 pixel resolution with section thickness of 0.4 μ m. Z-stack images were then rendered into a maximum intensity projection using ImageJ. Total dendritic surface area was quantified using ImageJ. The experiment was conducted in a blinded manner.

Statistical analysis

Data are presented as mean $\pm$ SEM and compared to vehicle control. Analysis was done using GraphPad Prism 5. Statistical analysis for response time was done using Kruskal-Wallis ANOVA, followed by Dunn's pairwise test for multiple comparisons. Statistical analysis for dendritic area percentage was done using Student's *t*-test.

Results

Our goal here was to develop a reproducible paradigm to investigate the effects of paclitaxel on nociception in the fly larvae. Based on previous studies for toxicity (Bhattacharya *et al.*, 2012; Cunha *et al.*, 2001), we selected paclitaxel doses below the lethal limit (Figure 1A), and then tested larval nociception using a heat probe set to a low intensity noxious heat (42°C; Figure 1B), which is mildly nociceptive to fly larvae (Babcock *et al.*, 2009). Our dose-response study revealed 2.5 μ M paclitaxel was sufficient

Figure 1. Paclitaxel induces heat-hyperalgesia in third instar *Drosophila* larvae. Schematic representation of the **A**) experimental design and **B**) thermal nociceptive assay in third instar *Drosophila* larvae. **C–D**) Average nociceptive latency (in seconds) in response to a 42°C or 46°C thermal stimulus, respectively. Increased paclitaxel concentration significantly induces heat-hyperalgesia in third instar larvae at 42°C. Note concentrations higher than 10 μ M paclitaxel were 100% lethal. **E**) Percentage response to each time point in seconds to 42°C thermal stimulus. All values represent mean $\pm$ SEM. p values were generated using Kruskal-Wallis ANOVA, followed by Dunn's pairwise test for multiple comparisons. Significance is relative to vehicle control. Five repeats were performed for each drug concentration with roughly 30 larvae each ($n = 130$ –180 animals).

to induce significant hyperalgesia, with a maximal hyperalgesia effect observed at 10 μM (Figure 1C). Concentrations higher than 10 μM paclitaxel were 100% lethal (not shown). Interestingly, paclitaxel did not significantly alter heat nociception latency to a 46°C heat stimulus across any of the doses (Figure 1D). Vehicle (ethanol) control and normal (no ethanol) control showed a response time of 5.71 sec (± 0.23 SEM; $n=173$) and 5.62 sec (± 0.20 SEM, $n=180$, not shown), respectively (42°C; Figure 1E). At low concentration's of 0.1 μM (5.21 sec ± 0.23 SEM; $n=150$) and 0.5 μM (5.44 sec ± 0.26 SEM; $n=131$) paclitaxel did not affect response profiles, however, concentrations of 2.5 μM paclitaxel (4.22 sec ± 0.19 SEM; $n=180$; $p<0.001$) and higher altered response distribution and significantly enhanced nociceptive latency (42°C; Figure 1E). The fastest latency response was observed at 10 μM paclitaxel (3.84 sec ± 0.24 SEM; $n=140$; $p<0.001$) with a 36.6% increase in response time relative to vehicle control (Figure 1C).

To evaluate if paclitaxel exposure caused robust morphological differences in peripheral pain sensing neurons, we fed genetically labeled (*w;ppk-CD4-tdGFP*) larvae paclitaxel and imaged the sensory neuron structure (Figures 2A–B). Surprisingly, treating larvae with 10 μM paclitaxel did not affect dendritic area

percentage compared to vehicle control (Figure C) despite significantly enhancing nociceptive sensitivity (not shown). Thus we establish that paclitaxel sensitizes larvae to heat pain via enhancing sensory neuron or higher order nociception, and not via inducing overt morphological changes.

Dataset 1. Larval response time in seconds to 42°C heat stimulus

<http://dx.doi.org/10.5256/f1000research.13581.d191022>

Paclitaxel fed larvae were touched with a 42°C heat probe and their response time was measured in seconds with a cut-off of 10 seconds. Different treatments were tested: food control, ethanol control, 0.1 μM , 0.5 μM , 2.5 μM , 5 μM , and 10 μM paclitaxel. Five repeats were performed ($n = 130 - 180$).

Dataset 2. Larval response time in seconds to 46°C heat stimulus

<http://dx.doi.org/10.5256/f1000research.13581.d191023>

Paclitaxel fed larvae were touched with a 46°C heat probe and their response time was measured in seconds with a cut-off of 10 seconds. Different treatments were tested: food control, ethanol control, 0.1 μM , 0.5 μM , 2.5 μM , 5 μM , and 10 μM paclitaxel. Five repeats were performed ($n = 130 - 180$).

Figure 2. Paclitaxel has no effect on class IV md-da sensory neuron dendritic coverage in third instar *Drosophila* larvae. A–B) Confocal images of third instar *ppk-CD4-tdGFP* larvae following (A) vehicle control or (B) 10 μM paclitaxel treatment. Images are of class IV md-da neurons at abdominal segment A5. Images are at 40x magnification and 0.7 zoom factor. Scale bar represents 40 μm . **C)** Quantification of dendritic area percentage of class IV md-da sensory neurons. Values represent mean $\pm$ SEM ($n = 10-11$ animals).

Dataset 3. Dendritic morphology of third instar ppk-CD4-tdGFP

<http://dx.doi.org/10.5256/f1000research.13581.d222138>

Confocal images of vehicle control and 0.1 μ M paclitaxel treated larvae. Images represent class IV md-da neurons at abdominal segment A5. Images are at 40x magnification and 0.7 zoom factor. Scale bar represents 40 μ m.

Conclusions

Many approved and effective anti-cancer therapeutic agents cause severe pain as a side effect. This can limit our ability to eradicate tumors, and often leaves cancer patients and survivors in intense, and untreatable pain. In this study, we describe a simple system to provide a high throughput dissection of the mechanisms involved in paclitaxel-induced pain. By combining these techniques with genomic approaches to identify regulators of chemotherapy pain, we can better understand how the pain arises, and potentially avoid these severe side effects while more effectively targeting the underlying disease.

Data availability

Dataset 1: Larval response time in seconds to 42°C heat stimulus. Paclitaxel fed larvae were touched with a 42°C heat probe and their response time was measured in seconds with a cut-off of 10 seconds. Different treatments were tested: food control, ethanol control, 0.1 μ M, 0.5 μ M, 2.5 μ M, 5 μ M, and 10 μ M paclitaxel. Five repeats were performed (n = 130 - 180). DOI, [10.5256/f1000research.13581.d191022](https://doi.org/10.5256/f1000research.13581.d191022) (Hamoudi *et al.*, 2018a).

Dataset 2: Larval response time in seconds to 46°C heat stimulus. Paclitaxel fed larvae were touched with a 46°C heat probe and their response time was measured in seconds with a cut-off of 10 seconds. Different treatments were tested: food control, ethanol control, 0.1 μ M, 0.5 μ M, 2.5 μ M, 5 μ M, and 10 μ M paclitaxel. Five repeats were performed (n = 130 - 180). DOI, [10.5256/f1000research.13581.d191023](https://doi.org/10.5256/f1000research.13581.d191023) (Hamoudi *et al.*, 2018b).

Dataset 3: Dendritic morphology of third instar ppk-CD4-tdGFP.

Confocal images of vehicle control and 0.1 μ M paclitaxel treated larvae. Images represent class IV md-da neurons at abdominal segment A5. Images are at 40x magnification and 0.7 zoom factor. Scale bar represents 40 μ m. DOI, [10.5256/f1000research.13581.d222138](https://doi.org/10.5256/f1000research.13581.d222138) (Hamoudi *et al.*, 2018c).

Competing interests

No competing interests were disclosed.

Grant information

This work was supported in part through NHMRC project grants APP1026310, APP1029672, APP1028887, APP1046090, APP1042416, APP1086851, and by a NHMRC career development fellowship II CDF1111940.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- Addington J, Freimer M: Chemotherapy-induced peripheral neuropathy: an update on the current understanding [version 1; referees: 2 approved]. *F1000Res*. 2016; 5(F1000 Faculty Rev): 1466.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- André N, Braguer D, Brasseur G, *et al.*: Paclitaxel induces release of cytochrome c from mitochondria isolated from human neuroblastoma cells. *Cancer Res*. 2000; 60(19): 5349–53.
[PubMed Abstract](#)
- Babcock DT, Landry C, Gallo MJ: Cytokine signaling mediates UV-induced nociceptive sensitization in *Drosophila* larvae. *Curr Biol*. 2009; 19(10): 799–806.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Bhattacharya MR, Gerds J, Naylor SA, *et al.*: Taxanes: the genetic toxicity of paclitaxel and docetaxel in somatic cells of *Drosophila melanogaster*. *Mutagenesis*. 2012; 16(1): 79–84.
- Boland BA, Sherry V, Polomano RC: Chemotherapy-induced peripheral neuropathy in cancer survivors. *Oncol Nurse Edn*. 2010; 24(2): 33–38, 42–43.
[Reference Source](#)
- Burton AW, Fanciullo GJ, Beasley RD, *et al.*: Chronic pain in the cancer survivor: a new frontier. *Pain Med*. 2007; 8(2): 189–198.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Cavaletti G, Tredici G, Braga M, *et al.*: Experimental peripheral neuropathy induced in adult rats by repeated intraperitoneal administration of taxol. *Exp Neurol*. 1995; 133(1): 64–72.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Chang AY, Kim K, Glick J, *et al.*: Phase II study of taxol, merbarone, and piroxanthone in stage IV non-small-cell lung cancer: The Eastern Cooperative Oncology Group Results. *J Natl Cancer Inst*. 1993; 85(5): 388–94.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Cunha KS, Reguly ML, Graf U, *et al.*: Taxanes: the genetic toxicity of paclitaxel and docetaxel in somatic cells of *Drosophila melanogaster*. *Mutagenesis*. 2001; 16(1): 79–84.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Gill PS, Tulpule A, Espina BM, *et al.*: Paclitaxel is safe and effective in the treatment of advanced AIDS-related Kaposi's sarcoma. *J Clin Oncol*. 1999; 17(6): 1876–83.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Hamoudi Z, Khuong TM, Cole T, *et al.*: Dataset 1 in: A fruit fly model for studying paclitaxel-induced pain. *F1000Research*. 2018a.
[Data Source](#)
- Hamoudi Z, Khuong TM, Cole T, *et al.*: Dataset 2 in: A fruit fly model for studying paclitaxel-induced pain. *F1000Research*. 2018b.
[Data Source](#)
- Hamoudi Z, Khuong TM, Cole T, *et al.*: Dataset 3 in: A fruit fly model for studying paclitaxel-induced pain. *F1000Research*. 2018c.
[Data Source](#)
- Hausheer FH, Schilsky RL, Bain S, *et al.*: Diagnosis, management, and evaluation of chemotherapy-induced peripheral neuropathy. *Semin Oncol*. 2006; 33(1): 15–49.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Holmes FA, Walters RS, Theriault RL, *et al.*: Phase II trial of taxol, an active drug in the treatment of metastatic breast cancer. *J Natl Cancer Inst*. 1991; 83(24): 1797–805.
[PubMed Abstract](#) | [Publisher Full Text](#)

Hwang RY, Zhong L, Xu Y, *et al.*: **Nociceptive neurons protect *Drosophila* larvae from parasitoid wasps.** *Curr Biol.* 2007; **17**(24): 2105–2116.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Lipton RB, Apfel SC, Dutcher JP, *et al.*: **Taxol produces a predominantly sensory neuropathy.** *Neurology.* 1989; **39**(3): 368–373.

[PubMed Abstract](#) | [Publisher Full Text](#)

McGuire WP, Rowinsky EK, Rosenshein NB, *et al.*: **Taxol: A unique antineoplastic agent with significant activity in advanced ovarian epithelial neoplasms.** *Ann Intern Med.* 1989; **111**(4): 273–9.

[PubMed Abstract](#) | [Publisher Full Text](#)

Neely GG, Hess A, Costigan M, *et al.*: **A genome-wide *Drosophila* screen for heat nociception identifies $\alpha 2\delta 3$ as an evolutionarily conserved pain gene.** *Cell.* 2010; **143**(4): 628–638.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Reyes-Gibby CC, Morrow PK, Buzdar A, *et al.*: **Chemotherapy-induced peripheral neuropathy as a predictor of neuropathic pain in breast cancer patients previously treated with paclitaxel.** *J Pain.* 2009; **10**(11): 1146–50.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Sahenk Z, Barohn R, New P, *et al.*: **Taxol neuropathy. Electrodiagnostic and sural nerve biopsy findings.** *Arch Neurol.* 1994; **51**(7): 726–729.

[PubMed Abstract](#) | [Publisher Full Text](#)

Seretny M, Currie GL, Sena ES, *et al.*: **Incidence, prevalence, and predictors of chemotherapy-induced peripheral neuropathy: A systematic review and meta-analysis.** *Pain.* 2014; **155**(12): 2461–70.

[PubMed Abstract](#) | [Publisher Full Text](#)

Shimozuma K, Ohashi Y, Takeuchi A, *et al.*: **Taxane-induced peripheral neuropathy and health-related quality of life in postoperative breast cancer patients undergoing adjuvant chemotherapy: N-SAS BC 02, a randomized clinical trial.** *Support Care Cancer.* 2012; **20**(12): 3355–64.

[PubMed Abstract](#) | [Publisher Full Text](#)

Tracey WD Jr, Wilson RI, Laurent G, *et al.*: **painless, a *Drosophila* gene essential for nociception.** *Cell.* 2003; **113**(2): 261–73.

[PubMed Abstract](#) | [Publisher Full Text](#)

Wani MC, Taylor HL, Wall ME, *et al.*: **Plant antitumor agents. VI. The isolation and structure of taxol, a novel antileukemic and antitumor agent from *Taxus brevifolia*.** *J Am Chem Soc.* 1971; **93**(9): 2325–7.

[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:

Version 1

Reviewer Report 22 March 2018

<https://doi.org/10.5256/f1000research.14752.r30145>

© 2018 Abdi S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Salahadin Abdi

Department of Pain Medicine, Division of Anesthesiology and Critical Care, University of Texas MD Anderson Cancer Center, Houston, TX, USA

This study examined the effects of Paclitaxel exposure on *Drosophila* larval nociception system and the authors propose that their model is suitable for high throughput screening and further mechanistic studies. The study is overall an interesting and clearly written, however, I do have the following concerns:

1. The dose response effect of thermal stimulation was only at 42 degrees. There was no discussion or explanation why this effect was not seen at 46 degrees.
2. The behavior experiment was based on thermal stimulation. I would be interested why mechanical stimulation was not chosen since mechanical sensitivity is common among patients who develop Paclitaxel induced peripheral neuropathy?
3. It is not clear to me what the timeline is between the exposure of the larvae with paclitaxel and performing microscopic studies.
4. There should be at least a short discussion about the result.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Partly

If applicable, is the statistical analysis and its interpretation appropriate?

I cannot comment. A qualified statistician is required.

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 09 Oct 2018

Greg Neely, Charles Perkins Centre and School of Life and Environmental Sciences, Camperdown, New South Wales, Australia

Thank you for your comments.

Response to comment #1: This effect is not seen at 46°C. At this temperature intensity, larvae respond rapidly (~1.5 seconds) and it is difficult to see even faster responses. To look for hyperalgesia, we instead lowered the heat stimulus intensity to 42°C, which is at the threshold for nociception in this system, and where nociceptive responses take on average ~5 seconds to elicit.

Response to comment #2: The type IV multidendritic nociceptor neurons that transduce heat nociception also transduce mechanical nociception, as these neurons are multimodal. We have tried on numerous occasions to generate reproducible data for mechanical nociception but so far in our hands this assay does not work well enough for us to feel comfortable publishing. Given the multimodal nature of type IV multidendritic nociceptor neurons, we reasoned that thermal hyperalgesia is a good readout for the overall sensitization of these sensory neurons.

Response to comment #3: The animals are born into paclitaxel containing food, and then early third instar are collected at day 6 to assess nociception or dendritic morphology. This information was provided in the methods, however we have further clarified this aspect.

Response to comment #4: We have now written a short discussion, please see discussion section.

Competing Interests: No competing interests were disclosed.

Reviewer Report 01 March 2018

<https://doi.org/10.5256/f1000research.14752.r30150>

© 2018 Claridge-Chang A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Adam Claridge-Chang

Institute of Molecular and Cell Biology (IMCB), Singapore, Singapore

The authors characterize the thermal nociception in *Drosophila* larvae that have been cultured on a range of paclitaxel concentrations. Using a heat probe to elicit the rolling defense behavior, they find that while paclitaxel has no effect on response times with a 46°C probe, it shortens probe response times when the larvae have been grown on 2.5 µm paclitaxel or above.

The authors and readers might like to consider the following comments on and questions about the 23 Jan 2018 version.

1. The Title describes a model for studying paclitaxel-induced pain, however the assay uses a heat probe to induce pain, and paclitaxel lowers the sensitivity to the probe, thus modeling the hyperalgesia component of paclitaxel CIPN. Would the Title better serve the reader if edited to focus on this side-effect specifically?
2. In **Abstract-Results**, the authors write: "We found that paclitaxel increases pain perception in a dose-dependent manner, without overt morphological changes." Changing "perception" to "sensitivity" would eliminate the baggage of the former word.
3. In **Abstract-Conclusions**: "Our simple, high throughput model can be combined with genomics approaches to identify regulators of chemotherapy-induced pain to eliminate its adverse side effects." However, they have not established that this is high-throughput by most common definitions of the term, nor do they show anywhere in the paper that it can be combined with genomics. The Conclusions would be improved if rephrased to better reflect what the data show.
4. "High-throughput" is a phrasal adjective that requires a hyphen.
5. In **Introduction** it says "This system is amenable to high throughput screening" however, this is not shown in the present manuscript nor is a reference cited in support of the statement.
6. "Kruskal-Wallis ANOVA." Correct to "Kruskal-Wallis."
7. I encourage the authors to use estimation statistics instead of significance testing. This would involve presenting and discussing the effect sizes. For example, in Figure 1c, it looks like 2.5 µm paclitaxel has the effect of reducing response time by ~1.5 s.
8. It would be nice to calculate standardized effect sizes (e.g. Cohen's d) of the paclitaxel effects; this would allow the authors and readers to estimate sample sizes needed for a screen (and thus possible throughput rates).
9. In **Results**, the authors write "Thus we establish that paclitaxel sensitizes larvae to heat pain via enhancing sensory neuron or higher order nociception, and not via inducing overt morphological changes." Is it true that enhancing sensory neuron or higher-order nociception are the only two alternatives? If not, this sentence should be rephrased.
10. The **Conclusions** section reads more like an overview of future plans for the assay. Could it be rewritten to more closely address the paper's findings?

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Partly

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

No

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 09 Oct 2018

Greg Neely, Charles Perkins Centre and School of Life and Environmental Sciences,
Camperdown, New South Wales, Australia

Thank you for your comments.

Response to comment #1: That's a reasonable point. Since first submission, we have new data that shows our model also involves peripheral neuropathy, so taken together, we have updated the title to capture this aspect and address the reviewer's comment. The new title is "**A fruit fly model for studying paclitaxel-induced peripheral neuropathy and hyperalgesia**". We hope this is acceptable.

Response to comment #2: Done.

Response to comment #3: We have revised this section and now state "Our simple system can be applied to identify regulators of chemotherapy-induced pain".

Response to comment #4: Done.

Response to comment #5: We have now included a reference for this statement.

Response to comment #6: Done.

Response to comment #7: We thank the reviewers for their comment and we have incorporated the estimation statistics into our analysis and added all the data points.

Response to comment #8: We have calculated the effect size, added it to the graphs (1C and 1D), and we have now also mentioned this in the results section. Moreover, we have changed the data representation to show all the data points.

Response to comment #9: Good point, this has been revised as suggested.

Response to comment #10: We have now added a discussion section.

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research