REVIEW

Recent advances in colonoscopy [version 1; peer review: 2 approved]

Edward Seward

Department of Gastrointestinal Services, University College London Hospitals NHS Trust, London, UK

V1 First published: 09 Jul 2019, 8(F1000 Faculty Rev):1028
<https://doi.org/10.12688/f1000research.18503.1>

Latest published: 09 Jul 2019, 8(F1000 Faculty Rev):1028
<https://doi.org/10.12688/f1000research.18503.1>

Abstract

Colonoscopy continues to evolve as equipment and techniques improve. Traditionally, colonoscopy has focused on adenoma detection, characterisation and resection as the primary aims, and there has certainly been considerable activity over the last few years in terms of addressing these important issues. This review article not only will discuss progress made in these areas but also will focus on when to colonoscope in terms of introduction of faecal immunochemical testing, how to insert with the advent of water-assisted insertion, and how to withdraw using a bundle of evidence-based techniques to improve adenoma detection. In addition, the ramifications of failing to discover polyps and of post-colonoscopy colorectal cancer are highlighted.

Keywords

colonoscopy, colorectal cancer, polyps

Open Peer Review

Approval Status

	1	2
version 1 09 Jul 2019		

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Konstantinos Triantafyllou**, National and Kapodistrian University of Athens, Athens, Greece
2. **Cesare Hassan**, Nuovo Regina Margherita Hospital, Rome, Italy

Any comments on the article can be found at the end of the article.

Corresponding author: Edward Seward (edward.seward1@nhs.net)

Author roles: Seward E: Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2019 Seward E. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Seward E. **Recent advances in colonoscopy [version 1; peer review: 2 approved]** F1000Research 2019, **8** (F1000 Faculty Rev):1028 <https://doi.org/10.12688/f1000research.18503.1>

First published: 09 Jul 2019, **8**(F1000 Faculty Rev):1028 <https://doi.org/10.12688/f1000research.18503.1>

Introduction

Colonoscopy continues to be the primary means of investigating lower gastrointestinal symptoms. Colonoscopy has been shown to reduce the risk of subsequent colorectal cancer both in screening¹ and in the symptomatic population². However, we do know that, despite its popularity, colonoscopy is not perfect, and post-colonoscopy cancer is a recognised feature. Therefore, it is worthwhile to ask the existential question, “What happens when colonoscopy goes wrong?” How to avoid this calamitous result will be discussed along with a consideration of the now-impressive data that exist for water-assisted insertion and how to optimise withdrawal as well as the prevention, management and follow-up of incomplete adenoma resection.

Whom to colonoscope: the role of faecal immunochemical testing

Faecal immunochemical testing (FIT) has been used extensively in the screening population, where it outperforms traditional guaiac faecal occult blood (gFOB) testing in terms of both sensitivity for detecting pathology but also ease of use to improve screening participation³. FIT differs from gFOB in that the former is specific to the globin portion of the haemoglobin molecule and therefore is specific to human haemoglobin, thereby reducing cross-reactivity with other mammalian forms of haemoglobin. It also provides a quantitative assessment of haemoglobin present in stool (measured in micrograms of haemoglobin per gram of stool) as opposed to the qualitative assessment in gFOB (as to whether or not a colour change has occurred). This allows a level of positivity to be chosen for a screening population (that is, a colonoscopy will be performed if the FIT level is more than 20 µg per gram of faeces); or in a symptomatic population, you may choose to make the test as sensitive as possible to decrease your chance of missing colorectal cancer (that is, any detectable faecal haemoglobin leads to a colonoscopy in the presence of colorectal symptoms). However, using FIT in the symptomatic population is challenging because although FIT has been shown to far outperform symptoms in terms of deciding who would benefit from a colonoscopy or computed tomography colonography⁴, it can also be used to rule out the need for a colonoscopy and this is more controversial. A Scottish study⁵ first highlighted the ability of FIT to be perfect (100% negative predictive value) in anticipating the absence of colorectal cancer if the FIT level was undetectable. If the FIT was undetectable in their population of both routine and urgent colorectal referrals (n = 755) who underwent both a FIT test and a colonoscopy, then no cancers were found, although about 1% had high-risk adenomas. Forty-two percent of the population had undetectable levels of faecal haemoglobin, meaning that if the decision of whether to undertake a colonoscopy had been based on the FIT result, this proportion would have been spared colonoscopy. This has obvious benefits for health systems where colonoscopy capacity acts as a constraint. It also improves convenience and safety for patients. However, subsequent publications have raised the possibility of “FIT-negative” cancer, which is colorectal cancer in the presence of undetectable levels of faecal haemoglobin^{6–9}. Despite this risk, the negative predictive value for FIT is still probably over 99%, and the studies highlighted above suggest that

FIT-negative cancers are the more clinically obvious ones (such as those accompanied by the presence of a mass, dramatic weight loss, and iron deficiency anaemia). Pathways incorporating FIT are being developed⁹, but an essential component will be safety netting to ensure that potential cancers are not missed. Defenders of the use of FIT will point to the post-colonoscopy cancer rate associated with colonoscopy, which (as will be demonstrated) is not insignificant. Two further studies of over 12,000 patients in the UK are examining the use of FIT in symptomatic patients and results are due this year, so it may be that answers will be provided shortly.

Colonoscopy insertion

Colonoscopy insertion is hampered by the discomfort that is caused by bowing and manipulation of the bowel by the colonoscope. This can be managed by using a magnetic imager that demonstrates the shape that the scope forms as it is inserted. Data on magnetic imagers have been available for over 20 years, but only recently did the largest meta-analysis to date come down in favour of their use for reducing time to caecum (even in expert hands), increasing caecal intubation rate and reducing pain scores¹⁰. A more recent unblinded study¹¹ of propofol usage in a colonoscopy population was able to demonstrate increased satisfaction scores and reduced propofol use with a magnetic imager, and as a further study has demonstrated that propofol-supported colonoscopy is associated with higher complication rates¹², it is interesting to hypothesise that a suitably powered study could demonstrate improved safety as well. The most exciting contribution to improvements in colonoscopic insertion technique has been the advent of water-assisted insertion techniques. One of these is water immersion colonoscopy, in which water is used to assist insertion along with suction of air. In this way, the sigmoid loop is shortened and straightened. Another of these techniques is water exchange colonoscopy, in which water is both channelled in through the scope and sucked out at the same time, thereby cleaning the colon on insertion and maintaining a clear view on insertion. In both techniques, water is removed and carbon dioxide is insufflated on withdrawal in order to facilitate polyp identification and removal.

Data exist to support water-assisted colonoscopy and it has been the subject of a Cochrane Review that included 16 randomised controlled trials over a total of 2933 patients and demonstrated an increase in the adenoma rate and a reduction in pain and sedation requirements¹³. A good review that describes the technique is available¹⁴, and concerns about the additional time of water-assisted colonoscopy are addressed by a review and meta-analysis¹⁵ that calculated that the procedure takes only two extra minutes for all the additional benefits for the patient.

Colonoscopy withdrawal

Colonoscopic withdrawal, where polyps are identified and removed, is actually the most important phase of colonoscopy. Although significant attention has been placed on technological advances and the role that high-definition colonoscopes play in polyp detection, a clever quality improvement trial¹⁶ demonstrated

that a bundle of four very simple measures thought, when taken together, to improve adenoma detection (taking more than 6 minutes on withdrawal, retroflexing in the rectum, using buscopan routinely to abolish peristalsis, and turning on withdrawal to optimise exposure of the colon) could be taught as a package to endoscopists. The endoscopists in the study were divided into quartiles dependent on their pre-intervention adenoma detection rate (ADR) and assessed for their adenoma detection post-intervention. The study group found that ADR significantly improves in the worse-performing endoscopist quartile (an improvement from 7.3 to 14.9%), although better-performing quartiles failed to show significant improvement. This tells us that ADR can be improved in a poorly performing group by simple educational interventions. The same group¹⁷ reported on the use of the Endocuff, a device that can be fitted on the end of a colonoscope and has multiple prongs to splay haustral folds on withdrawal, facilitating polyp detection. In that study, 1700 patients across seven hospitals took part, and adenoma detection increased significantly from 36.2 to 40.9%; the greatest benefit was in high-risk screening patients. Interestingly, the group also demonstrated a reduction in scope insertion time by 1 minute in the Endocuff group, theorising that it facilitates scope straightening. The rate of Endocuff removal, usually due to an angulated sigmoid, occurred in 4.1% because the Endocuff produced a small but appreciable effect on the colonoscope diameter.

Further recent data come from a very interesting article¹⁸ comparing Endocuff with EndoRings (another end-of-the-scope device that again is designed to increase adenoma detection), full-spectrum endoscopy (a colonoscope with side viewing capability to produce a very-wide-angle view) and standard high-definition colonoscopy. Endocuff outperformed the other three modalities in terms of ADR and adenomas per colonoscopy (APCs), particularly in the right side of the colon. This was true for both high-detector colonoscopists and those with lower ADRs, averaging an improvement in adenoma detection across all groups of 7%. These data are encouraging but require tempering from a recent meta-analysis¹⁹ of high-quality studies on the effect of Endocuff-assisted colonoscopy. Although the meta-analysis confirmed the benefit of Endocuff in adenoma detection (particularly amongst low to moderate detectors), the improvement was not as marked as previous studies suggested; the number needed to treat was 19, prompting considerations of cost. Therefore, in regard to detecting adenomas, colonoscopists need to be urged to do the basic things first²⁰ before reaching into their pockets and spending money on equipment to assist this. However the endoscopist does it, we know that there is a very strong inverse association between higher ADRs and a reduced risk of post-colonoscopy colorectal cancer (PCCRC)^{21,22} and so improving withdrawal technique and possibly using devices can only reduce the risk of this most feared complication.

A final point regarding adenoma detection on withdrawal is that it can be difficult to calculate and can be gamed²³. Focusing purely on ADR as a means of assessing the quality of a colonoscopist has been questioned as there is the anxiety that

endoscopists will be tempted to relax their search for further adenomas once one has been discovered—the so-called “one and done” phenomenon²⁴—prompting calls to have alternative key performance indicators such as APCs or significant polyps detected per 6 minutes of withdrawal time²⁵. However, only ADR has been demonstrated to predict PCCRC rate, which is the “hardest” end point for quality of colonoscopy and therefore is still regarded as the gold standard.

The reality of post-colonoscopy cancer

Every endoscopist knows the sinking feeling that occurs in a multi-disciplinary cancer meeting when the next patient up for discussion is revealed to have had an endoscopic procedure in the previous year. PCCRC is a recognised phenomenon but was not thoroughly described until the recent publication of a consensus statement from the World Endoscopy Organization²⁶. That article rigorously defined PCCRC as being between 6 and 36 months following the index colonoscopy; cases within less than 6 months were excluded to avoid counting cancers identified following a repeat colonoscopy (for inadequate bowel preparation, for instance). The article identified the PCCRC rate as an important metric of the performance of a *service* and not of an *individual* (as PCCRCs thankfully are relatively rare). A structure for evaluation of the PCCRC occurrence is suggested, and there are two basic questions: First, was there a previously resected polyp in the same segment as the cancer? In other words, could the resection have been inadequate? We already know from the CARE (Complete Adenoma Resection) study²⁷ that incomplete resection rates increase with size and sessile serrated morphology, such that almost half of large sessile serrated lesions were incompletely resected. Resection method may also be relevant, although a recent meta-analysis²⁸ found no difference between cold snare and hot snare resection in terms of polyp recurrence.

Second, was the quality of the colonoscopy adequate in regard to the extent and quality of bowel preparation? It is hoped that by defining the terms and providing a framework for prospective audit, we will gain a greater understanding of this difficult issue.

The scale of the problem is difficult to ascertain, resting as it does on the requirement for interrogation of national databases (as patients may have a colonoscopy in one hospital and have their cancer discovered in another). UK data suggest that the rate in the English National Health Service is 8.6%²⁹. The most recent data have been presented only in abstract form³⁰ but reassuringly demonstrate a fall to 7.5% from 2006 to 2012, presumably representing improvements in equipment, technique and bowel preparation. The authors made the point that variation between units demonstrated PCCRC rates of 5% in the highest-performing quintile and 11% in the lowest, which is clearly unacceptable. Falling PCCRC rates are not universal; a Canadian study³¹ failed to demonstrate a reduction in PCCRC of about 8% between 1996 and 2010. Clearly, this represents a call to arms for all colonoscopists to adopt strategies to understand why PCCRCs occur, to employ techniques to improve adenoma detection, and to ensure complete resection as well as ensuring

that adequate surveillance techniques are employed to manage the possibility of incomplete resection.

Conclusions

Since the removal of the first polyp by endoscopy, colonoscopists have been striving to develop ways of improving their techniques. As the ultimate aim is to reduce the risk of colorectal cancer or even abolish it entirely, the PCCRC rate is a very useful metric, serving to inform at a service level how effective the unit is. Just how prospective audits will inform this process and whether we will see a reduction in PCCRC remain to be seen.

New methods of water-assisted insertion are available (with encouraging data to support their use) and are easy to learn. Future studies may focus on how it improves adenoma detection, be it through colon cleansing or the magnifying effect of water. The optimum technique and whether it may just be adequate to use water insertion in the left colon only have yet to be decided. End-of-the-scope devices such as Endocuff are exciting and seem to improve every clinician's ability to spot

polyps, although perhaps the benefit does not always justify the cost in every patient. However, the clinical significance of the discovery of additional multiple tiny adenomas is not clear and this too needs further evaluation.

Colonoscopy at its inception was regarded as a “quantum leap” over what was previously available. Recent iterations of improvement have demonstrated that this progress with colonoscopy continues and innovation remains at the heart of lower gastrointestinal endoscopy.

Abbreviations

ADR, adenoma detection rate; APC, adenoma per colonoscopy; FIT, faecal immunochemical testing; gFOB, guaiac faecal occult blood; PCCRC, post-colonoscopy colorectal cancer

Grant information

The author(s) declared that no grants were involved in supporting this work.

References

- Logan RF, Patrick J, Nickerson C, *et al.*: Outcomes of the Bowel Cancer Screening Programme (BCSP) in England after the first 1 million tests. *Gut*. 2012; **61**(10): 1439–46.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Pan J, Xin L, Ma YF, *et al.*: Colonoscopy Reduces Colorectal Cancer Incidence and Mortality in Patients With Non-Malignant Findings: A Meta-Analysis. *Am J Gastroenterol*. 2016; **111**(3): 355–65.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Moss S, Mathews C, Day TJ, *et al.*: Increased uptake and improved outcomes of bowel cancer screening with a faecal immunochemical test: results from a pilot study within the national screening programme in England. *Gut*. 2017; **66**(9): 1631–44.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Cubiella J, Vega P, Salve M, *et al.*: Development and external validation of a faecal immunochemical test-based prediction model for colorectal cancer detection in symptomatic patients. *BMC Med*. 2016; **14**(1): 128.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Mowat C, Digby J, Strachan JA, *et al.*: Faecal haemoglobin and faecal calprotectin as indicators of bowel disease in patients presenting to primary care with bowel symptoms. *Gut*. 2016; **65**(9): 1463–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Godber IM, Todd LM, Fraser CG, *et al.*: Use of a faecal immunochemical test for haemoglobin can aid in the investigation of patients with lower abdominal symptoms. *Clin Chem Lab Med*. 2016; **54**(4): 595–602.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Högberg C, Karling P, Rutegård J, *et al.*: Diagnosing colorectal cancer and inflammatory bowel disease in primary care: The usefulness of tests for faecal haemoglobin, faecal calprotectin, anaemia and iron deficiency. A prospective study. *Scand J Gastroenterol*. 2016; **52**(1): 69–75.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Widiak MM, Thomas CL, Thomas MG, *et al.*: Diagnostic accuracy of faecal biomarkers in detecting colorectal cancer and adenoma in symptomatic patients. *Aliment Pharmacol Ther*. 2017; **45**(2): 354–63.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Chapman C, Bunce J, Oliver S, *et al.*: Service evaluation of faecal immunochemical testing and anaemia for risk stratification in the 2-week-wait pathway for colorectal cancer. *BJS Open*. 2019; **3**(3): 395–402.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Mark-Christensen A, Brandsborg S, Iversen LH: Magnetic endoscopic imaging as an adjuvant to elective colonoscopy: a systematic review and meta-analysis of randomized controlled trials. *Endoscopy*. 2015; **47**(3): 251–61.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Klare P, Hartrampf B, Haller B, *et al.*: Magnetic endoscope imaging for routine colonoscopy: impact on propofol dosage and patient safety - a randomized trial. *Endoscopy*. 2016; **48**(10): 916–22.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Wemli KJ, Brenner AT, Rutter CM, *et al.*: Risks Associated With Anesthesia Services During Colonoscopy. *Gastroenterology*. 2016; **150**(4): 888–94; quiz e18.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Hafner S, Zolk K, Radaelli F, *et al.*: Water infusion versus air insufflation for colonoscopy. *Cochrane Database Syst Rev*. 2015; **56**(5): CD009863.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Siau K, Beintaris I: My approach to water-assisted colonoscopy. *Frontline Gastroenterol*. 2019; **10**(2): 194–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Cadoni S, Hassan C, Frazzoni L, *et al.*: Impact of water exchange colonoscopy on endoscopy room efficiency: a systematic review and meta-analysis. *Gastrointest Endosc*. 2019; **89**(1): 159–167.e13.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Rajasekhar PT, Rees CJ, Bramble MG, *et al.*: A multicenter pragmatic study of an evidence-based intervention to improve adenoma detection: the Quality Improvement in Colonoscopy (QIC) study. *Endoscopy*. 2015; **47**(3): 217–24.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Ngu WS, Bevan R, Tsiamoulos ZP, *et al.*: Improved adenoma detection with Endocuff Vision: the ADENOMA randomised controlled trial. *Gut*. 2019; **68**(2): 280–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Rex DK, Repici A, Gross SA, *et al.*: High-definition colonoscopy versus Endocuff versus EndoRings versus full-spectrum endoscopy for adenoma detection at colonoscopy: a multicenter randomized trial. *Gastrointest Endosc*. 2018; **88**(2): 335–344.e2.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- Triantafyllou K, Gkolfakis P, Tziatzios G, *et al.*: Effect of Endocuff use on colonoscopy outcomes: A systematic review and meta-analysis. *World J Gastroenterol*. 2019; **25**(9): 1158–70.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Facciorusso A, Triantafyllou K, Murad MH, *et al.*: Compared Abilities of Endoscopic Techniques to Increase Colon Adenoma Detection Rates:

- A Network Meta-analysis.** *Clin Gastroenterol Hepatol.* 2018; pii: S1542-3565(18)31335-1.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
21. **F** Kaminski MF, Regula J, Kraszewska E, *et al.*: **Quality indicators for colonoscopy and the risk of interval cancer.** *N Engl J Med.* 2010; **362**(19): 1795–803.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
22. **F** Corley DA, Jensen CD, Marks AR, *et al.*: **Adenoma detection rate and risk of colorectal cancer and death.** *N Engl J Med.* 2014; **370**(14): 1298–306.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
23. **F** Rex DK, Ponugoti PL: **Calculating the adenoma detection rate in screening colonoscopies only: Is it necessary? Can it be gamed?** *Endoscopy.* 2017; **49**(11): 1069–74.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
24. Denis B, Sauleau EA, Gendre I, *et al.*: **The mean number of adenomas per procedure should become the gold standard to measure the neoplasia yield of colonoscopy: a population-based cohort study.** *Dig Liver Dis.* 2014; **46**(2): 176–81.
[PubMed Abstract](#) | [Publisher Full Text](#)
25. Rameshshanker R, Wilson A, Tsiamoulos Z, *et al.*: **PTU-030 Number of Significant Polyps Detected Per Six Minutes of Withdrawal Time at Colonoscopy (SP6): A New Measure of Colonoscopy Efficiency and Quality.** *Gut.* 2016; **65**(Suppl 1): A66.3-A67.
[Publisher Full Text](#)
26. **F** Rutter MD, Beintaris I, Valori R, *et al.*: **World Endoscopy Organization Consensus Statements on Post-Colonoscopy and Post-Imaging Colorectal Cancer.** *Gastroenterology.* 2018; **155**(3): 909–925.e3.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
27. **F** Pohl H, Srivastava A, Bensen SP, *et al.*: **Incomplete polyp resection during colonoscopy-results of the complete adenoma resection (CARE) study.** *Gastroenterology.* 2013; **144**(1): 74–80.e1.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
28. **F** Qu J, Jian H, Li L, *et al.*: **Effectiveness and safety of cold versus hot snare polypectomy: A meta-analysis.** *J Gastroenterol Hepatol.* 2019; **34**(1): 49–58.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
29. Morris EJ, Rutter MD, Finan PJ, *et al.*: **Post-colonoscopy colorectal cancer (PCCRC) rates vary considerably depending on the method used to calculate them: a retrospective observational population-based study of PCCRC in the English National Health Service.** *Gut.* 2015; **64**(8): 1248–56.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
30. Burr N, Rutter M, Shelton J, *et al.*: **A study of post colonoscopy colorectal cancer (PCCRC) in England.** *Gut.* 2018; **67**(Suppl 1): A184–A186.
[Reference Source](#)
31. **F** Murthy SK, Benchimol EI, Tinmouth J, *et al.*: **Temporal trends in postcolonoscopy colorectal cancer rates in 50- to 74-year-old persons: a population-based study.** *Gastrointest Endosc.* 2018; **87**(5): 1324–1334.e4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)

Open Peer Review

Current Peer Review Status:

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. Cesare Hassan

Digestive Endoscopy Unit, Nuovo Regina Margherita Hospital, Rome, Italy

Competing Interests: No competing interests were disclosed.

2. Konstantinos Triantafyllou

Hepatogastroenterology Unit, 2nd Department of Internal Medicine - Propaedeutic, Research Institute and Diabetes Center, Medical School, National and Kapodistrian University of Athens, Athens, Greece

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research